



Assignment

Derivative at a Point

Basic Level

1. If $f(x) = |x|$, then $f'(0) =$ [MNR 1982]
 (a) 0 (b) 1 (c) x (d) None of these
2. If $f(x) = \begin{cases} 1, & x < 0 \\ 1 + \sin x, & 0 \leq x < \frac{\pi}{2} \end{cases}$ then $f'(0) =$ [MP PET 1994]
 (a) 1 (b) 0 (c) ∞ (d) Does not exist
3. If $f(x) = \begin{cases} ax^2 + b; & x \leq 0 \\ x^2; & x > 0 \end{cases}$ possesses derivative at $x = 0$, then
 (a) $a = 0, b = 0$ (b) $a > 0, b = 0$ (c) $a \in \mathbb{R}, b = 0$ (d) None of these
4. The derivative of $f(x) = 3|2 + x|$ at the point $x_0 = -3$ is [Orissa JEE 2002]
 (a) 3 (b) -3 (c) 0 (d) Does not exist
5. The derivative of $y = 1 - |x|$ at $x = 0$ is [SCRA 1996]
 (a) 0 (b) 1 (c) -1 (d) Does not exist
6. The derivative of $f(x) = |x^2 - x|$ at $x = 2$ is [AMU 1999]
 (a) -3 (b) 0 (c) 3 (d) Not defined
7. The value of $\frac{d}{dx} [|x-1| + |x-5|]$ at $x = 3$ is [MP PET 2000]
 (a) -2 (b) 0 (c) 2 (d) 4
8. If $f(x)$ has a derivative at $x = a$, then $\lim_{x \rightarrow a} \frac{xf(a) - af(x)}{x - a}$ is equal to
 (a) $f(a) - af'(a)$ (b) $af(a) - f'(a)$ (c) $f(a) + f'(a)$ (d) $af(a) + f'(a)$
9. If $f(x) = x + 2$, then $f'(f(x))$ at $x = 4$ is [DCE 2001]
 (a) 8 (b) 1 (c) 4 (d) 5
10. Let $3f(x) - 2f(1/x) = x$, then $f'(2)$ is equal to [MP PET 2000]
 (a) $2/7$ (b) $1/2$ (c) 2 (d) $7/2$
11. If $f(x)$ is a differentiable function, then $\lim_{x \rightarrow a} \frac{af(x) - xf(a)}{x - a}$ is
 (a) $af'(a) - f(a)$ (b) $af(a) - f'(a)$ (c) $af'(a) + f(a)$ (d) $af(a) + f'(a)$
12. The differential coefficient of the function $|x - 1| + |x - 3|$ at the point $x = 2$ is [Rajasthan PET 2002]
 (a) -2 (b) 0 (c) 2 (d) Undefined
13. If $f(x) = |x - 3|$, then $f'(3) =$

- (a) 0 (b) 1 (c) -1 (d) Does not exist

Advance Level

14. If $y = \cot^{-1}(\cos 2x)^{1/2}$, then the value of $\frac{dy}{dx}$ at $x = \frac{\pi}{6}$ will be
 (a) $\left(\frac{2}{3}\right)^{1/2}$ (b) $\left(\frac{1}{3}\right)^{1/2}$ (c) $(3)^{1/2}$ (d) $(6)^{1/2}$
15. The values of x , at which the first derivative of the function $\left(\sqrt{x} + \frac{1}{\sqrt{x}}\right)^2$ w.r.t. x is $\frac{3}{4}$, are
 (a) ± 2 (b) $\pm \frac{1}{2}$ (c) $\pm \frac{\sqrt{3}}{2}$ (d) $\pm \frac{2}{\sqrt{3}}$
16. The number of points at which the function $f(x) = |x - 0.5| + |x - 1| + \tan x$ does not have a derivative in the interval $(0, 2)$, is
 (a) 1 (b) 2 (c) 3 (d) 4 [MNR 1995]
17. The set of all those points, where the function $f(x) = \frac{x}{1 + |x|}$ is differentiable, is
 (a) $(-\infty, \infty)$ (b) $[0, \infty)$ (c) $(-\infty, 0) \cup (0, \infty)$ (d) $(0, \infty)$
18. Let $f(x+y) = f(x)f(y)$ and $f(x) = 1 + xg(x)G(x)$ where $\lim_{x \rightarrow 0} g(x) = a$ and $\lim_{x \rightarrow 0} G(x) = b$ then $f'(x)$ is equal to
 (a) $1 + ab$ (b) ab (c) a/b (d) None of these
19. $f(x)$ is a function such that $f''(x) = -f(x)$ and $f'(x) = g(x)$ and $h(x)$ is a function such that $h(x) = [f(x)]^2 + [g(x)]^2$ and $h(5) = 11$, then the value of $h(10)$ is
 (a) 0 (b) 1 (c) 10 (d) None of these
20. Let $f(x+y) = f(x)f(y)$ for all x and y . Suppose that $f(3) = 3$ and $f'(0) = 11$, then $f'(3)$ is given by
 (a) 22 (b) 33 (c) 28 (d) None of these

Some Standard Differentiation
Basic Level

21. If $y = \frac{(1-x)^2}{x^2}$, then $\frac{dy}{dx}$ is [MP PET 1999]
 (a) $\frac{2}{x^2} + \frac{2}{x^3}$ (b) $-\frac{2}{x^2} + \frac{2}{x^3}$ (c) $-\frac{2}{x^2} - \frac{2}{x^3}$ (d) $-\frac{2}{x^3} + \frac{2}{x^2}$
22. If $2t = v^2$, then $\frac{dv}{dt}$ is equal to [MP PET 1992]
 (a) 0 (b) $1/4$ (c) $1/2$ (d) $1/v$
23. If $x = y\sqrt{1-y^2}$, then $\frac{dy}{dx} =$ [MP PET 2001]
 (a) 0 (b) x (c) $\frac{\sqrt{1-y^2}}{1-2y^2}$ (d) $\frac{\sqrt{1-y^2}}{1+2y^2}$
24. If $pv = 81$, then $\frac{dp}{dv}$ is at $v = 9$ equal to [MP PET 1999]
 (a) 1 (b) -1 (c) 2 (d) None of these

148 Differentiation

25. If $y = \sqrt{\frac{1+x}{1-x}}$, then $\frac{dy}{dx} =$ [AISSE 1981; Rajasthan PET 1995]
- (a) $\frac{2}{(1+x)^{1/2}(1-x)^{3/2}}$ (b) $\frac{1}{(1+x)^{1/2}(1-x)^{3/2}}$ (c) $\frac{1}{2(1+x)^{1/2}(1-x)^{3/2}}$ (d) $\frac{2}{(1+x)^{3/2}(1-x)^{1/2}}$
26. The derivative of $f(x) = x|x|$ is
- (a) $2x$ (b) $-2x$ (c) $2x^2$ (d) $2|x|$ [AMU 2001]
27. The derivative of $F[f\{\phi(x)\}]$ is
- (a) $F'[f\{\phi(x)\}]$ (b) $F'[f\{\phi(x)\}]f'\{\phi(x)\}$ (c) $F'[f\{\phi(x)\}]f'\{\phi(x)\}$ (d) $F'[f\{\phi(x)\}]f'\{\phi(x)\}\phi'(x)$ [Rajasthan PET 1996]
28. $\frac{d}{dx}(\sin 2x^2)$ equals
- (a) $4x \cos(2x^2)$ (b) $2 \sin x^2 \cdot \cos x^2$ (c) $4x \sin(x^2)$ (d) $4x \sin(x^2) \cdot \cos(x^2)$ [MP PET 1997]
29. If $y = \sec x^0$, then $\frac{dy}{dx} =$
- (a) $\sec x \tan x$ (b) $\sec x^0 \tan x^0$ (c) $\frac{\pi}{180} \sec x^0 \tan x^0$ (d) $\frac{180}{\pi} \sec x^0 \tan x^0$ [Kerala (Engg.) 2002]
30. If $\sin y + e^{-x \cos y} = e$, then $\frac{dy}{dx}$ at $(1, \pi)$ is
- (a) $\sin y$ (b) $-x \cos y$ (c) e (d) $\sin y - x \cos y$
31. If $y = a \sin x + b \cos x$, then $y^2 + \left(\frac{dy}{dx}\right)^2$ is a
- (a) Function of x (b) Function of y (c) Function of x and y (d) Constant [AISSE 1981; AI CBSE 1979]
32. $\frac{d}{dx}[\cos(1-x^2)^2] =$
- (a) $-2x(1-x^2)\sin(1-x^2)^2$ (b) $-4x(1-x^2)\sin(1-x^2)^2$ (c) $4x(1-x^2)\sin(1-x^2)^2$ (d) $-2(1-x^2)\sin(1-x^2)^2$
33. If $y = \cos(\sin x^2)$, then at $x = \sqrt{\frac{\pi}{2}}$, $\frac{dy}{dx} =$
- (a) -2 (b) 2 (c) $-2\sqrt{\frac{\pi}{2}}$ (d) 0
34. $\frac{d}{dx}[\sin^n x \cos nx] =$
- (a) $n \sin^{n-1} x \cos(n+1)x$ (b) $n \sin^{n-1} x \cos nx$ (c) $n \sin^{n-1} x \cos(n-1)x$ (d) $n \sin^{n-1} x \sin(n+1)x$ [DSSE 1979]
35. $\frac{d}{dx} \cos(\sin x^2) =$
- (a) $\sin(\sin x^2) \cdot \cos x^2 \cdot 2x$ (b) $-\sin(\sin x^2) \cdot \cos x^2 \cdot 2x$ (c) $-\sin(\sin x^2) \cdot \cos^2 x \cdot 2x$ (d) None of these [DSSE 1987]
36. If $y = \sin(\sqrt{\sin x + \cos x})$, then $\frac{dy}{dx} =$
- (a) $\frac{1}{2} \frac{\cos \sqrt{\sin x + \cos x}}{\sqrt{\sin x + \cos x}}$ (b) $\frac{\cos \sqrt{\sin x + \cos x}}{\sqrt{\sin x + \cos x}}$ (c) $\frac{1}{2} \frac{\cos \sqrt{\sin x + \cos x}}{\sqrt{\sin x + \cos x}} (\cos x - \sin x)$ (d) None of these [AISSE 1987]
37. If $y = \sin\left(\frac{1+x^2}{1-x^2}\right)$, then $\frac{dy}{dx} =$

- (a) $\frac{4x}{1-x^2} \cdot \cos\left(\frac{1+x^2}{1-x^2}\right)$ (b) $\frac{x}{(1-x^2)^2} \cdot \cos\left(\frac{1+x^2}{1-x^2}\right)$ (c) $\frac{x}{(1-x^2)} \cdot \cos\left(\frac{1+x^2}{1-x^2}\right)$ (d) $\frac{4x}{(1-x^2)^2} \cdot \cos\left(\frac{1+x^2}{1-x^2}\right)$
38. $\frac{d}{dx}(x^2 + \cos x)^4 =$ [DSSE 1987]
 (a) $4(x^2 + \cos x)(2x - \sin x)$ (b) $4(x^2 - \cos x)(2x - \sin x)$ (c) $4(x^2 + \cos x)^3(2x - \sin x)$ (d) $4(x^2 + \cos x)^3(2x + \sin x)$
39. $\frac{d}{dx}\left(\frac{\cot^2 x - 1}{\cot^2 x + 1}\right) =$
 (a) $-\sin 2x$ (b) $2 \sin 2x$ (c) $2 \cos 2x$ (d) $-2 \sin 2x$
40. $\frac{d}{dx} \sqrt{\frac{1 - \sin 2x}{1 + \sin 2x}} =$ [AISSE 1985; DSSE 1986]
 (a) $\sec^2 x$ (b) $-\sec^2\left(\frac{\pi}{4} - x\right)$ (c) $\sec^2\left(\frac{\pi}{4} + x\right)$ (d) $\sec^2\left(\frac{\pi}{4} - x\right)$
41. If $y = \frac{\tan x + \cot x}{\tan x - \cot x}$, then $\frac{dy}{dx} =$
 (a) $2 \tan 2x \sec 2x$ (b) $\tan 2x \sec 2x$ (c) $-\tan 2x \sec 2x$ (d) $-2 \tan 2x \sec 2x$
42. $\frac{d}{dx} \sqrt{\sec^2 x + \operatorname{cosec}^2 x} =$ [DSSE 1981]
 (a) $4 \operatorname{cosec} 2x \cdot \cot 2x$ (b) $-4 \operatorname{cosec} 2x \cdot \cot 2x$ (c) $-4 \operatorname{cosec} x \cdot \cot 2x$ (d) None of these
43. If $y = \frac{5x}{\sqrt[3]{(1-x)^2}} + \cos^2(2x+1)$, then $\frac{dy}{dx} =$ [IIT 1980]
 (a) $\frac{5(3-x)}{3(1-x)^{5/3}} - 2 \sin(4x+2)$ (b) $\frac{5(3-x)}{3(1-x)^{2/3}} - 2 \sin(4x+4)$ (c) $\frac{5(3-x)}{3(1-x)^{2/3}} - 2 \sin(2x+1)$ (d) None of these
44. If $y = \sqrt{\sin x + y}$, then $\frac{dy}{dx}$ equals to [Rajasthan PET 2001]
 (a) $\frac{\sin x}{2y-1}$ (b) $\frac{\cos x}{2y-1}$ (c) $\frac{\sin x}{2y+1}$ (d) $\frac{\cos x}{2y+1}$
45. $\frac{d}{dx} \log |x| = \dots (x \neq 0)$
 (a) $\frac{1}{x}$ (b) $-\frac{1}{x}$ (c) x (d) $-x$
46. $\frac{d}{dx} \log_{\sqrt{x}}(1/x)$ is equal to [AMU 1999]
 (a) $-\frac{1}{2\sqrt{x}}$ (b) -2 (c) $-\frac{1}{x^2\sqrt{x}}$ (d) 0
47. $\frac{d}{dx} \log(\log x) =$ [IIT 1985]
 (a) $\frac{x}{\log x}$ (b) $\frac{\log x}{x}$ (c) $(x \log x)^{-1}$ (d) None of these
48. $\frac{d}{dx}(\log \tan x) =$ [MNR 1986]
 (a) $2 \sec 2x$ (b) $2 \operatorname{cosec} 2x$ (c) $\sec 2x$ (d) $\operatorname{cosec} 2x$
49. If $y = \log x^x$, then $\frac{dy}{dx} =$ [MNR 1978]

150 Differentiation

- (a) $x^x(1 + \log x)$ (b) $\log(ex)$ (c) $\log\left(\frac{e}{x}\right)$ (d) None of these
50. Derivative of the function $f(x) = \log_5(\log_7 x)$, $x > 7$ is [Orissa JEE 2002]
 (a) $\frac{1}{x(\ln 5)(\ln 7)(\log_7 x)}$ (b) $\frac{1}{x(\ln 5)(\ln 7)}$ (c) $\frac{1}{x(\ln x)}$ (d) None of these
51. The differential coefficient of $f[\log(x)]$ when $f(x) = \log x$ is [Kurukshetra CEE 1998; DCE 2000]
 (a) $x \log x$ (b) $\frac{x}{\log x}$ (c) $\frac{1}{x \log x}$ (d) $\frac{\log x}{x}$
52. If $y = \log \frac{1 + \sqrt{x}}{1 - \sqrt{x}}$, then $\frac{dy}{dx} =$
 (a) $\frac{\sqrt{x}}{1 - x}$ (b) $\frac{1}{\sqrt{x}(1 - x)}$ (c) $\frac{\sqrt{x}}{1 + x}$ (d) $\frac{1}{\sqrt{x}(1 + x)}$
53. If $y = x^2 \log x + \frac{2}{\sqrt{x}}$, then $\frac{dy}{dx} =$
 (a) $x + 2x \log x - \frac{1}{\sqrt{x}}$ (b) $x + 2x \log x - \frac{1}{x^{3/2}}$ (c) $x + 2x \log x - \frac{2}{x^{3/2}}$ (d) None of these
54. $\frac{d}{dx} \left[\log \sqrt{\frac{1 - \cos x}{1 + \cos x}} \right] =$ [BIT Ranchi 1990]
 (a) $\sec x$ (b) $\operatorname{cosec} x$ (c) $\operatorname{cosec} \frac{x}{2}$ (d) $\sec \frac{x}{2}$
55. $\frac{d}{dx} \left\{ \log \left(\frac{e^x}{1 + e^x} \right) \right\} =$
 (a) $\frac{1}{1 - e^x}$ (b) $-\frac{1}{1 + e^x}$ (c) $-\frac{1}{1 - e^x}$ (d) None of these
56. $\frac{d}{dx} \{ \log(\sec x + \tan x) \} =$ [AISSE 1982]
 (a) $\cos x$ (b) $\sec x$ (c) $\tan x$ (d) $\cot x$
57. $\frac{d}{dx} \left[\log \left(x + \frac{1}{x} \right) \right] =$ [MP PET 1995]
 (a) $\left(x + \frac{1}{x} \right)$ (b) $\frac{\left(1 + \frac{1}{x^2} \right)}{\left(1 + \frac{1}{x} \right)}$ (c) $\frac{\left(1 - \frac{1}{x^2} \right)}{\left(x + \frac{1}{x} \right)}$ (d) $\left(1 + \frac{1}{x} \right)$
58. $\frac{d}{dx} \log(x^{10}) =$ [Rajasthan PET 1992]
 (a) x^{-10} (b) $10x$ (c) $10/x$ (d) $10x^9$
59. If $y = \log \left\{ \frac{x + \sqrt{a^2 + x^2}}{a} \right\}$, then the value of $\frac{dy}{dx}$ is
 (a) $\sqrt{a^2 - x^2}$ (b) $a\sqrt{a^2 + x^2}$ (c) $\frac{1}{\sqrt{a^2 + x^2}}$ (d) $x\sqrt{a^2 + x^2}$

60. If $y = \log(\sin^{-1} x)$, then $\frac{dy}{dx}$ equals
 (a) $\frac{1}{\sin^{-1} x \sqrt{1-x^2}}$ (b) $-\frac{1}{\sin^{-1} x \sqrt{1-x^2}}$ (c) $\frac{-2x}{\sin^{-1} x \sqrt{1-x^2}}$ (d) None of these
61. If $y = e^{(1+\log_e x)}$, then the value of $\frac{dy}{dx} =$ [MP PET 1996]
 (a) e (b) 1 (c) 0 (d) $\log_e x e^{\log_e x}$
62. If $y = e^{\sqrt{x}}$, then $\frac{dy}{dx}$ equals
 (a) $\frac{e^{\sqrt{x}}}{2\sqrt{x}}$ (b) $\frac{\sqrt{x}}{e^{\sqrt{x}}}$ (c) $\frac{x}{e^{\sqrt{x}}}$ (d) $\frac{2\sqrt{x}}{e^{\sqrt{x}}}$
63. The derivative of $y = x^{\ln x}$ is
 (a) $x^{\ln x} \ln x$ (b) $x^{\ln x-1} \ln x$ (c) $2x^{\ln x-1} \ln x$ (d) $x^{\ln x-2}$
64. Derivative of $x^6 + 6^x$ with respect to x is [Kerala (Engg.) 2002]
 (a) $12x$ (b) $x + 4$ (c) $6x^5 + 6^x \log 6$ (d) $6x^5 + x6^{x-1}$
65. $\frac{d}{dx}(e^x \log \sin 2x) =$ [AI CBSE 1985]
 (a) $e^x(\log \sin 2x + 2 \cot 2x)$ (b) $e^x(\log \cos 2x + 2 \cot 2x)$ (c) $e^x(\log \cos 2x + \cot 2x)$ (d) None of these
66. If $y = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty$, then $\frac{dy}{dx} =$ [Karnataka CET 1999]
 (a) y (b) $y - 1$ (c) $y + 1$ (d) None of these
67. $\frac{d}{dx} e^{x \sin x} =$ [DSSE 1979]
 (a) $e^{x \sin x}(x \cos x + \sin x)$ (b) $e^{x \sin x}(\cos x + x \sin x)$ (c) $e^{x \sin x}(\cos x + \sin x)$ (d) None of these
68. $\frac{d}{dx}(xe^{x^2}) =$ [DSSE 1981]
 (a) $2x^2 e^{x^2} + e^{x^2}$ (b) $x^2 e^{x^2} + e^{x^2}$ (c) $e^x \cdot 2x^2 + e^{x^2}$ (d) None of these
69. If $y = x^2 + x^{\log x}$, then $\frac{dy}{dx} =$
 (a) $\frac{x^2 + \log x \cdot x^{\log x}}{x}$ (b) $x^2 + \log x \cdot x^{\log x}$ (c) $\frac{2(x^2 + \log x \cdot x^{\log x})}{x}$ (d) None of these
70. $\frac{d}{dx}\{e^{-ax^2} \log(\sin x)\} =$ [AI CBSE 1984]
 (a) $e^{-ax^2}(\cot x + 2ax \log \sin x)$ (b) $e^{-ax^2}(\cot x + ax \log \sin x)$ (c) $e^{-ax^2}(\cot x - 2ax \log \sin x)$ (d) None of these
71. If $y = \sqrt{\frac{1+e^x}{1-e^x}}$, then $\frac{dy}{dx} =$ [AI CBSE 1987]
 (a) $\frac{e^x}{(1-e^x)\sqrt{1-e^{2x}}}$ (b) $\frac{e^x}{(1-e^x)\sqrt{1-e^x}}$ (c) $\frac{e^x}{(1-e^x)\sqrt{1+e^{2x}}}$ (d) $\frac{e^x}{(1-e^x)\sqrt{1+e^x}}$

152 Differentiation

72. $\frac{d}{dx} \{e^x \log(1+x^2)\} =$ [AI CBSE 1987]
 (a) $e^x \left[\log(1+x^2) + \frac{2x}{1+x^2} \right]$ (b) $e^x \left[\log(1+x^2) - \frac{2x}{1+x^2} \right]$ (c) $e^x \left[\log(1+x^2) + \frac{x}{1+x^2} \right]$ (d) $e^x \left[\log(1+x^2) - \frac{x}{1+x^2} \right]$
73. If $y = \frac{e^{2x} + e^{-2x}}{e^{2x} - e^{-2x}}$, then $\frac{dy}{dx} =$
 (a) $\frac{-8}{(e^{2x} - e^{-2x})^2}$ (b) $\frac{8}{(e^{2x} - e^{-2x})^2}$ (c) $\frac{-4}{(e^{2x} - e^{-2x})^2}$ (d) $\frac{4}{(e^{2x} - e^{-2x})^2}$
74. $\frac{d}{dx}(e^{x^3})$ is equal to [Rajasthan PET 1995]
 (a) $3xe^{x^3}$ (b) $3x^2e^{x^3}$ (c) $3x(e^{x^3})^2$ (d) $2x^3e^{x^3}$
75. $\frac{d}{dx}[e^{ax} \cos(bx+c)] =$ [AISSE 1989]
 (a) $e^{ax}[a \cos(bx+c) - b \sin(bx+c)]$ (b) $e^{ax}[a \sin(bx+c) - b \cos(bx+c)]$
 (c) $e^{ax}[\cos(bx+c) - \sin(bx+c)]$ (d) None of these
76. If $y = e^x \log x$, then $\frac{dy}{dx}$ is [SCRA 1996]
 (a) $\frac{e^x}{x}$ (b) $e^x \left(\frac{1}{x} + x \log x \right)$ (c) $e^x \left(\frac{1}{x} + \log x \right)$ (d) $\frac{e^x}{\log x}$
77. If $f(1) = 3$, $f'(1) = 2$, then $\frac{d}{dx} \{\log f(e^x + 2x)\}$ at $x = 0$ is [AMU 1999]
 (a) $2/3$ (b) $3/2$ (c) 2 (d) 0
78. $\frac{d}{dx}(\sin^{-1} x)$ is equal to [Rajasthan PET 1995]
 (a) $\frac{1}{\sqrt{1-x^2}}$ (b) $-\frac{1}{\sqrt{1-x^2}}$ (c) $\frac{1}{\sqrt{1+x^2}}$ (d) $\frac{1}{\sqrt{x^2-1}}$
79. If $y = \sin^{-1} \sqrt{x}$, then $\frac{dy}{dx} =$ [MP PET 1995]
 (a) $\frac{2}{\sqrt{x}\sqrt{1-x}}$ (b) $\frac{-2}{\sqrt{x}\sqrt{1-x}}$ (c) $\frac{1}{2\sqrt{x}\sqrt{1-x}}$ (d) $\frac{1}{\sqrt{1-x}}$
80. If $y = \sin^{-1} \sqrt{1-x^2}$, then $\frac{dy}{dx} =$ [AISSE 1987]
 (a) $\frac{1}{\sqrt{1-x^2}}$ (b) $\frac{1}{\sqrt{1+x^2}}$ (c) $-\frac{1}{\sqrt{1-x^2}}$ (d) $-\frac{1}{\sqrt{x^2-1}}$
81. $\frac{d}{dx} \tan^{-1} \sqrt{\frac{1+\sin x}{1-\sin x}} =$
 (a) $\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) 1 (d) -1
82. If $y = \tan^{-1} \left(\frac{\sqrt{x}-x}{1+x^{3/2}} \right)$, then $y'(1)$ is [AMU 2000]
 (a) 0 (b) $\frac{1}{2}$ (c) -1 (d) $-\frac{1}{4}$

83. Differential coefficient of $\sec^{-1} x$ is

- (a) $\frac{1}{x\sqrt{1-x^2}}$ (b) $-\frac{1}{x\sqrt{1-x^2}}$ (c) $\frac{1}{x\sqrt{x^2-1}}$ (d) $-\frac{1}{x\sqrt{x^2-1}}$

84. If $y = \cot^{-1}\left(\frac{1+x}{1-x}\right)$, then $\frac{dy}{dx} =$ [DSSE 1984]

- (a) $\frac{1}{1+x^2}$ (b) $-\frac{1}{1+x^2}$ (c) $\frac{2}{1+x^2}$ (d) $-\frac{2}{1+x^2}$

85. If $y = \tan^{-1} \sqrt{\frac{1+\cos x}{1-\cos x}}$, then $\frac{dy}{dx}$ is equal to [Roorkee 1995]

- (a) 0 (b) $-\frac{1}{2}$ (c) $\frac{1}{2}$ (d) 1

86. If $f(x) = \tan^{-1}\left(\frac{\sin x}{1+\cos x}\right)$, then $f\left(\frac{\pi}{3}\right) =$ [BIT Ranchi 1988]

- (a) $\frac{1}{2(1+\cos x)}$ (b) $\frac{1}{2}$ (c) $\frac{1}{4}$ (d) None of these

87. If $y = \sec^{-1}\left(\frac{x+1}{x-1}\right) + \sin^{-1}\left(\frac{x-1}{x+1}\right)$, then $\frac{dy}{dx} =$ [MNR 1984]

- (a) 0 (b) 1 (c) 2 (d) 3

88. If $y = \tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)$, then $\frac{dy}{dx} =$

- (a) $-\frac{1}{\sqrt{1-x^2}}$ (b) $\frac{x}{\sqrt{1-x^2}}$ (c) $\frac{1}{\sqrt{1-x^2}}$ (d) $\frac{\sqrt{1-x^2}}{x}$

89. If $y = \sin^{-1}\left(\frac{1-x^2}{1+x^2}\right)$, then $\frac{dy}{dx}$ equals [Rajasthan PET 1996; EAMCET 1991]

- (a) $\frac{2}{1-x^2}$ (b) $\frac{1}{1+x^2}$ (c) $\pm \frac{2}{1+x^2}$ (d) $-\frac{2}{1+x^2}$

90. $\frac{d}{dx} \left[\tan^{-1}\left(\frac{a-x}{1+ax}\right) \right] =$ [Karnataka CET 2001]

- (a) $-\frac{1}{1+x^2}$ (b) $\frac{1}{1+a^2} - \frac{1}{1+x^2}$ (c) $\frac{1}{1+\left(\frac{a-x}{1+ax}\right)^2}$ (d) $\frac{-1}{\sqrt{1-\left(\frac{a-x}{1+ax}\right)^2}}$

91. If $y = \tan^{-1}\left[\frac{\sin x + \cos x}{\cos x - \sin x}\right]$, then $\frac{dy}{dx}$ is [UPSEAT 2001]

- (a) $1/2$ (b) $\pi/4$ (c) 0 (d) 1

92. $\frac{d}{dx} \left(\tan^{-1} \frac{\cos x}{1+\sin x} \right) =$ [AISSE 1984, 85; MNR 1983; Rajasthan PET 1994, 96]

- (a) $-\frac{1}{2}$ (b) $\frac{1}{2}$ (c) -1 (d) 1

93. If $y = \sin^{-1} \frac{2x}{1+x^2} + \sec^{-1} \frac{1+x^2}{1-x^2}$, then $\frac{dy}{dx} =$ [Rajasthan PET 1996]

154 Differentiation

- (a) $\frac{4}{1-x^2}$ (b) $\frac{1}{1+x^2}$ (c) $\frac{4}{1+x^2}$ (d) $\frac{-4}{1+x^2}$
94. If $y = \tan^{-1} \frac{4x}{1+5x^2} + \tan^{-1} \frac{2+3x}{3-2x}$, then $\frac{dy}{dx} =$
- (a) $\frac{1}{1+25x^2} + \frac{2}{1+x^2}$ (b) $\frac{5}{1+25x^2} + \frac{2}{1+x^2}$ (c) $\frac{5}{1+25x^2}$ (d) $\frac{1}{1+25x^2}$
95. $\frac{d}{dx} \sin^{-1} \left(\frac{x^2}{\sqrt{x^4+a^4}} \right) =$
- (a) $\frac{2a^4x}{a^4+x^4}$ (b) $\frac{2a^2x}{a^4+x^4}$ (c) $\frac{2a^3x}{a^4+x^4}$ (d) $\frac{-2a^2x}{a^4+x^4}$
96. If $y = \frac{\sin^{-1} x}{\sqrt{1-x^2}}$, then
- (a) $(1-x^2) \frac{dy}{dx} - xy - 1 = 0$ (b) $(1-x^2) \frac{dy}{dx} + xy - 1 = 0$ (c) $(1-x^2) \frac{dy}{dx} + \frac{1}{2} xy - 1 = 0$ (d) None of these
97. $\frac{d}{dx} \sin^{-1}(3x-4x^3) =$ [Rajasthan PET 2003]
- (a) $\frac{3}{\sqrt{1-x^2}}$ (b) $\frac{-3}{\sqrt{1-x^2}}$ (c) $\frac{1}{\sqrt{1-x^2}}$ (d) $\frac{-1}{\sqrt{1-x^2}}$
98. $\frac{d}{dx} \sin^{-1}(2ax\sqrt{1-a^2x^2}) =$
- (a) $\frac{2a}{\sqrt{a^2-x^2}}$ (b) $\frac{a}{\sqrt{a^2-x^2}}$ (c) $\frac{2a}{\sqrt{1-a^2x^2}}$ (d) $\frac{a}{\sqrt{1-a^2x^2}}$
99. $\frac{d}{dx} \left[\sin^{-1} \left(\frac{3x}{2} - \frac{x^3}{2} \right) \right]$ equals
- (a) $\frac{3}{\sqrt{4-x^2}}$ (b) $\frac{-3}{\sqrt{4-x^2}}$ (c) $\frac{1}{\sqrt{4-x^2}}$ (d) $\frac{-1}{\sqrt{4-x^2}}$
100. If $y = \sin^{-1} \left\{ \frac{\sin x + \cos x}{\sqrt{2}} \right\}$, then $\frac{dy}{dx} =$
- (a) 1 (b) -1 (c) 0 (d) None of these
101. $\frac{d}{dx} \cos^{-1} \frac{x-x^{-1}}{x+x^{-1}} =$ [DSSE 1985; Roorkee 1963]
- (a) $\frac{1}{1+x^2}$ (b) $-\frac{1}{1+x^2}$ (c) $\frac{2}{1+x^2}$ (d) $\frac{-2}{1+x^2}$
102. $\frac{d}{dx} \left\{ \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right) \right\} =$ [AISSE 1984]
- (a) $\frac{1}{1+x^2}$ (b) $-\frac{1}{1+x^2}$ (c) $-\frac{2}{1+x^2}$ (d) $\frac{2}{1+x^2}$
103. $\frac{d}{dx} \cos^{-1} \sqrt{\frac{1+x^2}{2}} =$ [AI CBSE 1988]
- (a) $\frac{-1}{2\sqrt{1-x^4}}$ (b) $\frac{1}{2\sqrt{1-x^4}}$ (c) $\frac{-x}{\sqrt{1-x^4}}$ (d) $\frac{x}{\sqrt{1-x^4}}$

104. If $y = \tan^{-1} \left(\frac{\sqrt{a} - \sqrt{x}}{1 + \sqrt{ax}} \right)$, then $\frac{dy}{dx} =$ [AI CBSE 1988]

- (a) $\frac{1}{2(1+x)\sqrt{x}}$ (b) $\frac{1}{(1+x)\sqrt{x}}$ (c) $-\frac{1}{2(1+x)\sqrt{x}}$ (d) None of these

105. $\frac{d}{dx} \left[\tan^{-1} \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right] =$ [Roorkee 1980]

- (a) $\frac{-x}{\sqrt{1-x^4}}$ (b) $\frac{x}{\sqrt{1-x^4}}$ (c) $\frac{-1}{2\sqrt{1-x^4}}$ (d) $\frac{1}{2\sqrt{1-x^4}}$

106. If $y = (1+x^2)\tan^{-1} x - x$, then $\frac{dy}{dx} =$ [Karnataka CET 2001]

- (a) $\tan^{-1} x$ (b) $2x \tan^{-1} x$ (c) $2x \tan^{-1} x - 1$ (d) $\frac{2x}{\tan^{-1} x}$

107. If $f(x) = (x+1)\tan^{-1}(e^{-2x})$, then $f'(0)$ equals

- (a) $\frac{\pi}{6} + 5$ (b) $\frac{\pi}{2} + 1$ (c) $\frac{\pi}{4} - 1$ (d) None of these

Advance Level

108. If $y = \sec(\tan^{-1} x)$, then $\frac{dy}{dx}$ is [DCE 2002; Haryana CEE 2001]

- (a) $\frac{x}{\sqrt{1+x^2}}$ (b) $\frac{-x}{\sqrt{1+x^2}}$ (c) $\frac{x}{\sqrt{1-x^2}}$ (d) None of these

109. If $y = (1+x^{1/4})(1+x^{1/2})(1-x^{1/4})$, then $\frac{dy}{dx} =$ [MP PET 1994]

- (a) 1 (b) -1 (c) x (d) $\sqrt{x}$

110. If $f(x) = \sqrt{ax} + \frac{a^2}{\sqrt{ax}}$, then $f'(a) =$

- (a) -1 (b) 1 (c) 0 (d) a

111. $x\sqrt{1+y} + y\sqrt{1+x} = 0$, then $\frac{dy}{dx} =$

- (a) $1+x$ (b) $(1+x)^{-2}$ (c) $-(1+x)^{-1}$ (d) $-(1+x)^{-2}$

112. If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$, then $\frac{dy}{dx} =$ [MNR 1983; ISM Dhanbad 1987; Rajasthan PET 1991]

- (a) $\sqrt{\frac{1-x^2}{1-y^2}}$ (b) $\sqrt{\frac{1-y^2}{1-x^2}}$ (c) $\sqrt{\frac{x^2-1}{1-y^2}}$ (d) $\sqrt{\frac{y^2-1}{1-x^2}}$

113. Function $y = (x + \sqrt{x^2 + 1})^k$ satisfies [IIT Screening]

- (a) $(x^2 + 1)y' = k^2 y$ (b) $\sqrt{(x^2 + 1)}y' = ky$ (c) $(1 + x^2)y'' + ky' - xy = 0$ (d) $(1 + x^2)y'' + k^2 + xy' = 0$

114. The derivative of $\sqrt{\sqrt{x} + 1}$ is

- (a) $\frac{1}{\sqrt{x}(\sqrt{x} + 1)}$ (b) $\frac{-1}{\sqrt{x}\sqrt{x+1}}$ (c) $\frac{4}{\sqrt{x}(\sqrt{x} + 1)}$ (d) $\frac{1}{4\sqrt{x}(\sqrt{x} + 1)}$

156 Differentiation

- 115.** If $f(x) = \frac{1}{\sqrt{x^2+a^2} + \sqrt{x^2+b^2}}$, then $f'(x)$ is equal to [Kurukshetra CEE 1998]
- (a) $\frac{x}{(a^2-b^2)} \left[\frac{1}{\sqrt{x^2+a^2}} - \frac{1}{\sqrt{x^2+b^2}} \right]$ (b) $\frac{x}{(a^2+b^2)} \left[\frac{1}{\sqrt{x^2+a^2}} - \frac{1}{\sqrt{x^2+b^2}} \right]$
- (c) $\frac{x}{(a^2-b^2)} \left[\frac{1}{\sqrt{x^2+a^2}} + \frac{1}{\sqrt{x^2+b^2}} \right]$ (d) $(a^2-b^2) \left[\frac{1}{\sqrt{x^2+a^2}} - \frac{2}{\sqrt{x^2+b^2}} \right]$
- 116.** If $\sqrt{1-x^6} + \sqrt{1-y^6} = a^3(x^3 - y^3)$, then $\frac{dy}{dx} =$ [Roorkee 1994]
- (a) $\frac{x^2}{y^2} \sqrt{\frac{1-x^6}{1-y^6}}$ (b) $\frac{y^2}{x^2} \sqrt{\frac{1-y^6}{1-x^6}}$ (c) $\frac{x^2}{y^2} \sqrt{\frac{1-y^6}{1-x^6}}$ (d) None of these
- 117.** If $y = \sqrt{x+\sqrt{x}}$, then $y \frac{dy}{dx}$ equals
- (a) $\frac{2\sqrt{x}+1}{4\sqrt{x}}$ (b) $\frac{\sqrt{x}+1}{2\sqrt{x}}$ (c) $\frac{\sqrt{x}+1}{4x}$ (d) $\frac{x+1}{2\sqrt{x}}$
- 118.** If $y = \sqrt{\sin \sqrt{x}}$, then $\frac{dy}{dx} =$ [MP PET 1997]
- (a) $\frac{1}{2\sqrt{\cos \sqrt{x}}}$ (b) $\frac{\sqrt{\cos \sqrt{x}}}{2x}$ (c) $\frac{\cos \sqrt{x}}{4\sqrt{x} \sqrt{\sin \sqrt{x}}}$ (d) $\frac{1}{2\sqrt{\sin x}}$
- 119.** $\frac{d}{dx} \sqrt{x \sin x} =$ [AISSE 1985]
- (a) $\frac{\sin x + x \cos x}{2\sqrt{x \sin x}}$ (b) $\frac{\sin x + x \cos x}{\sqrt{x \sin x}}$ (c) $\frac{x \sin x + \cos x}{\sqrt{2 \sin x}}$ (d) $\frac{x \sin x + \cos x}{\sqrt{2x \sin x}}$
- 120.** If $y = f\left(\frac{2x-1}{x^2+1}\right)$ and $f'(x) = \sin x^2$, then $\frac{dy}{dx} =$ [IIT 1982]
- (a) $\frac{6x^2-2x+2}{(x^2+1)^2} \sin\left(\frac{2x-1}{x^2+1}\right)^2$ (b) $\frac{6x^2-2x+2}{(x^2+1)^2} \sin^2\left(\frac{2x-1}{x^2+1}\right)$
- (c) $\frac{-2x^2+2x+2}{(x^2+1)^2} \sin^2\left(\frac{2x-1}{x^2+1}\right)$ (d) $\frac{-2x^2+2x+2}{(x^2+1)^2} \sin\left(\frac{2x-1}{x^2+1}\right)^2$
- 121.** $\frac{d}{dx} \left[\frac{e^{ax}}{\sin(bx+c)} \right] =$ [AI CBSE 1983]
- (a) $\frac{e^{ax}[a \sin(bx+c) + b \cos(bx+c)]}{\sin^2(bx+c)}$ (b) $\frac{e^{ax}[a \sin(bx+c) - b \cos(bx+c)]}{\sin^2(bx+c)}$
- (c) $\frac{e^{ax}[a \sin(bx+c) - b \cos(bx+c)]}{\sin^2(bx+c)}$ (d) None of these
- 122.** If $y = b \cos \log\left(\frac{x}{n}\right)^n$, then $\frac{dy}{dx} =$
- (a) $-n b \sin \log\left(\frac{x}{n}\right)^b$ (b) $n b \sin \log\left(\frac{x}{n}\right)^n$ (c) $\frac{-nb}{x} \sin \log\left(\frac{x}{n}\right)^n$ (d) None of these
- 123.** If $y = f\left(\frac{5x+1}{10x^2-3}\right)$ and $f'(x) = \cos x$, then $\frac{dy}{dx} =$ [MP PET 1987]

(a) $\cos\left(\frac{5x+1}{10x^2-3}\right) \frac{d}{dx}\left(\frac{5x+1}{10x^2-3}\right)$

(b) $\frac{5x+1}{10x^2-3} \cos\left(\frac{5x+1}{10x^2-3}\right)$

(c) $\cos\left(\frac{5x+1}{10x^2-3}\right)$

(d) None of these

124. $\frac{d}{dx}\left(x^3 \tan^2 \frac{x}{2}\right) =$

[AISSE 1979]

(a) $x^3 \tan \frac{x}{2} \cdot \sec^2 \frac{x}{2} + 3x \tan^2 \frac{x}{2}$

(b) $x^3 \tan \frac{x}{2} \cdot \sec^2 \frac{x}{2} + 3x^2 \tan^2 \frac{x}{2}$

(c) $x^2 \tan^2 \frac{x}{2} \cdot \sec^2 \frac{x}{2} + 3x^2 \tan^2 \frac{x}{2}$

(d) None of these

125. $\frac{d}{dx}(\tan a^{1/x}) =$

(a) $\sec^2(a^{1/x}) \cdot \frac{(a^{1/x} \cdot \log a)}{x^2}$

(b) $\sec^2(a^{1/x}) \cdot (a^{1/x} \cdot \log a)$

(c) $\frac{\sec x \cdot \log a}{x^2}$

(d) $-\frac{\sec^2(a^{1/x}) \cdot (a^{1/x} \cdot \log_e a)}{x^2}$

126. If $y = \sqrt{\frac{1+\tan x}{1-\tan x}}$, then $\frac{dy}{dx} =$

(a) $\frac{1}{2} \sqrt{\frac{1-\tan x}{1+\tan x}} \cdot \sec^2\left(\frac{\pi}{4} + x\right)$

(b) $\sqrt{\frac{1-\tan x}{1+\tan x}} \cdot \sec^2\left(\frac{\pi}{4} + x\right)$

(c) $\frac{1}{2} \sqrt{\frac{1-\tan x}{1+\tan x}} \cdot \sec\left(\frac{\pi}{4} + x\right)$

(d)

127. If $A = \frac{2^x \cot x}{\sqrt{x}}$, then $\frac{dA}{dx} =$

(a) $\frac{2^{x-1} \left\{ -2x \operatorname{cosec}^2 x + \cot x \cdot \log\left(\frac{4^x}{e}\right) \right\}}{x^{3/2}}$

(b) $\frac{2^{x-1} \left\{ -2x \operatorname{cosec}^2 x + \cot x \cdot \log\left(\frac{4^x}{e}\right) \right\}}{x}$

(c) $\frac{2^x \left\{ -2x \operatorname{cosec}^2 x + \cot x \cdot \log\left(\frac{4^x}{e}\right) \right\}}{x^{3/2}}$

(d) None of these

128. Differential coefficient of $\sqrt{\sec \sqrt{x}}$ is

[MP PET 1996]

(a) $\frac{1}{4\sqrt{x}} (\sec \sqrt{x})^{3/2} \sin \sqrt{x}$

(b) $\frac{1}{4\sqrt{x}} \sec \sqrt{x} \sin \sqrt{x}$

(c) $\frac{1}{2} \sqrt{x} (\sec \sqrt{x})^{3/2} \sin \sqrt{x}$

(d) $\frac{1}{2} \sqrt{x} \sec \sqrt{x} \sin \sqrt{x}$

129. $\frac{d}{dx}\left(\frac{\sec x + \tan x}{\sec x - \tan x}\right) =$

[DSSE 1979, 81; CBSE 1981]

(a) $\frac{2 \cos x}{(1 - \sin x)^2}$

(b) $\frac{\cos x}{(1 - \sin x)^2}$

(c) $\frac{2 \cos x}{1 - \sin x}$

(d) None of these

130. If $x = f(m) \cos m - f'(m) \sin m$ and $y = f(m) \sin m + f'(m) \cos m$, then $\left(\frac{dy}{dm}\right)^2 + \left(\frac{dx}{dm}\right)^2$ equals

[AMU 1997]

(a) $[f(m) + f''(m)]^2$

(b) $[f(m) - f''(m)]^2$

(c) $\{f(m)\}^2 + \{f'(m)\}^2$

(d) $\frac{\{f(m)\}^2}{\{f'(m)\}^2}$

131. If $x = \sec \theta - \cos \theta$ and $y = \sec^n \theta - \cos^n \theta$, then

[IIT 1989]

(a) $(x^2 + 4) \left(\frac{dy}{dx}\right)^2 = n^2(y^2 + 4)$

(b)

(c) $(x^2 + 4) \left(\frac{dy}{dx}\right)^2 = x^2(y^2 + 4)$

(d) $(x^2 + 4) \left(\frac{dy}{dx}\right)^2 = (y^2 + 4)$

132. If $y = \log_{\sin x}(\tan x)$, then $\left(\frac{dy}{dx}\right)_{\pi/4} =$

158 Differentiation

- (a) $\frac{4}{\log 2}$ (b) $-4 \log 2$ (c) $\frac{-4}{\log 2}$ (d) None of these
133. If $u(x, y) = y \log x + x \log y$, then $u_x u_y - u_x \log x - u_y \log y + \log x \log y =$ [EAMCET 2003]
 (a) 0 (b) -1 (c) 1 (d) 2
134. If $y = \log x \cdot e^{(\tan x + x^2)}$, then $\frac{dy}{dx} =$
 (a) $e^{(\tan x + x^2)} \left[\frac{1}{x} + (\sec^2 x + x) \log x \right]$ (b) $e^{(\tan x + x^2)} \left[\frac{1}{x} + (\sec^2 x - x) \log x \right]$
 (c) $e^{(\tan x + x^2)} \left[\frac{1}{x} + (\sec^2 x + 2x) \log x \right]$ (d) $e^{(\tan x + x^2)} \left[\frac{1}{x} + (\sec^2 x - 2x) \log x \right]$
135. $\frac{d}{dx} \left[\log \sqrt{\sin \sqrt{e^x}} \right] =$
 (a) $\frac{1}{4} e^{x/2} \cot(e^{x/2})$ (b) $e^{x/2} \cot(e^{x/2})$ (c) $\frac{1}{4} e^x \cot(e^x)$ (d) $\frac{1}{2} e^{x/2} \cot(e^{x/2})$
136. If $y\sqrt{x^2+1} = \log\{\sqrt{x^2+1} - x\}$, then $(x^2+1)\frac{dy}{dx} + xy + 1 =$ [Roorkee 1978; Kurukshetra CEE 1998]
 (a) 0 (b) 1 (c) 2 (d) None of these
137. If $y = \log_{\cos x} \sin x$, then $\frac{dy}{dx}$ is equal to
 (a) $(\cot x \log \cos x + \tan x \log \sin x)/(\log \cos x)^2$ (b) $(\tan x \log \cos x + \cot x \log \sin x)/(\log \cos x)^2$
 (c) $(\cot x \log \cos x + \tan x \log \sin x)/(\log \sin x)^2$ (d) None of these
138. If $y = \log(x + e^{\sqrt{x}})$ then $\frac{dy}{dx} =$
 (a) $\frac{2\sqrt{x} + e^{\sqrt{x}}}{2\sqrt{x}(x + e^{\sqrt{x}})}$ (b) $\frac{2\sqrt{x} + e^{\sqrt{x}}}{2\sqrt{x}(x - e^{\sqrt{x}})}$ (c) $\frac{2\sqrt{x} - e^{\sqrt{x}}}{2\sqrt{x}(x + e^{\sqrt{x}})}$ (d) $\frac{2\sqrt{x} - e^{\sqrt{x}}}{2\sqrt{x}(x - e^{\sqrt{x}})}$
139. $\frac{d}{dx} (a^{\log_{10} \operatorname{cosec}^{-1} x}) =$ [Roorkee 1990]
 (a) $a^{\log_{10} \operatorname{cosec}^{-1} x} \cdot \frac{1}{\operatorname{cosec}^{-1} x} \cdot \frac{1}{x\sqrt{x^2-1}} \cdot \log_{10} a$ (b) $-a^{\log_{10} \operatorname{cosec}^{-1} x} \cdot \frac{1}{\operatorname{cosec}^{-1} x} \cdot \frac{1}{|x|\sqrt{x^2-1}} \cdot \log_{10} a$
 (c) $a^{\log_{10} \operatorname{cosec}^{-1} x} \cdot \frac{1}{\operatorname{cosec}^{-1} x} \cdot \frac{1}{|x|\sqrt{x^2-1}} \cdot \log_{10} a$ (d) $-a^{\log_{10} \operatorname{cosec}^{-1} x} \cdot \frac{1}{\operatorname{cosec}^{-1} x} \cdot \frac{1}{x\sqrt{x^2-1}} \cdot \log_{10} a$
140. $\frac{d}{dx} (e^{\sqrt{1-x^2}} \cdot \tan x) =$ [AI CBSE 1979]
 (a) $e^{\sqrt{1-x^2}} \left[\sec^2 x + \frac{x \tan x}{\sqrt{1-x^2}} \right]$ (b) $e^{\sqrt{1-x^2}} \left[\sec^2 x - \frac{x \tan x}{\sqrt{1-x^2}} \right]$ (c) $e^{\sqrt{1-x^2}} \left[\sec^2 x + \frac{\tan x}{\sqrt{1-x^2}} \right]$ (d) $e^{\sqrt{1-x^2}} \left[\sec^2 x - \frac{\tan x}{\sqrt{1-x^2}} \right]$
141. $10^{-x \tan x} \left[\frac{d}{dx} (10^{x \tan x}) \right]$ is equal to
 (a) $\tan x + x \sec^2 x$ (b) $\ln 10 (\tan x + x \sec^2 x)$
 (c) $\ln 10 \left(\tan x + \frac{x}{\cos^2 x} + \tan x \sec^2 x \right)$ (d) $x \tan x \ln 10$

142. If $y = \sin \left[2 \tan^{-1} \sqrt{\frac{1-x}{1+x}} \right]$, then $\frac{dy}{dx} =$

- (a) $\frac{1}{2\sqrt{1-x^2}}$ (b) $\frac{-2x}{\sqrt{1-x^2}}$ (c) $\frac{-x}{\sqrt{1-x^2}}$ (d) $\frac{x}{\sqrt{1-x^2}}$

143. $\frac{d}{dx} \left(\cos^{-1} \sqrt{\frac{1+\cos x}{2}} \right) =$

[AI CBSE 1982]

- (a) 1 (b) $\frac{1}{2}$ (c) $\frac{1}{3}$ (d) None of these

144. If $f(x) = \cos^{-1} \left[\frac{1 - (\log x)^2}{1 + (\log x)^2} \right]$, then the value of $f'(e) =$

[Karnataka CET 1999]

- (a) 1 (b) $\frac{1}{e}$ (c) $\frac{2}{e}$ (d) $\frac{2}{e^2}$

145. If $y = \frac{1}{\sqrt{a^2 - b^2}} \cos^{-1} \left[\frac{a \cos(x - \alpha) + b}{\theta} \right]$ where $\theta = a + b \cos(x - \alpha)$, then $\frac{dy}{dx} =$

[Orissa JEE 2003]

- (a) $\frac{1}{\theta}$ (b) $\frac{2}{\theta}$ (c) $\frac{1}{\theta^2}$ (d) $\frac{2}{\theta^2}$

146. $\frac{d}{dx} \tan^{-1}(\sec x + \tan x) =$

[AISSE 1985, 87; DSSE 1982, 84]

- (a) 1 (b) $1/2$ (c) $\cos x$ (d) $\sec x$

147. $\frac{d}{dx} \left[\tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}} \right] =$

[Ranchi BIT 1989; Roorkee 1989; Rajasthan PET 1996]

- (a) $-\frac{1}{2}$ (b) 0 (c) $\frac{1}{2}$ (d) 1

148. If $y = \tan^{-1} \left(\frac{x^{1/3} + a^{1/3}}{1 - x^{1/3} a^{1/3}} \right)$, then $\frac{dy}{dx} =$

[DSSE 1986]

- (a) $\frac{1}{3x^{2/3}(1+x^{2/3})}$ (b) $\frac{a}{3x^{2/3}(1+x^{2/3})}$ (c) $-\frac{1}{3x^{2/3}(1+x^{2/3})}$ (d) $-\frac{a}{3x^{2/3}(1+x^{2/3})}$

149. The differential coefficient of $\tan^{-1} \left(\frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right)$ is

[MP PET 2003]

- (a) $\sqrt{1-x^2}$ (b) $\frac{1}{\sqrt{1-x^2}}$ (c) $\frac{1}{2\sqrt{1-x^2}}$ (d) x

150. $\frac{d}{dx} \left[\sin^2 \cot^{-1} \frac{1}{\sqrt{\frac{1+x}{1-x}}} \right]$ equals

[EAMCET 1996]

- (a) 0 (b) $\frac{1}{2}$ (c) $-\frac{1}{2}$ (d) 1

151. If $f'(x) = \sin(\log x)$ and $y = f\left(\frac{2x+3}{3-2x}\right)$ then $\frac{dy}{dx}$ equals

160 Differentiation

- (a) $\sin(\log x) \cdot \frac{1}{x \log x}$ (b) $\frac{12}{(3-2x)^2} \sin\left(\log\left(\frac{2x+3}{3-2x}\right)\right)$ (c) $\sin\left(\log\left(\frac{2x+3}{3-2x}\right)\right)$ (d) None of these
152. If $y = \tan^{-1}\left\{\frac{3a^2x - x^3}{a(a^2 - 3x^2)}\right\}$ then $\frac{dy}{dx}$ equals [IIT 1969; Rajasthan PET 1998]
- (a) $\frac{3}{a^2 + x^2}$ (b) $\frac{a}{a^2 + x^2}$ (c) $\frac{3a}{a^2 + x^2}$ (d) $\frac{3x}{a^2 + x^2}$
153. If $y = \sinh^{-1}(\tan x)$, then the value of $\frac{dy}{dx}$ is
- (a) $\sin x$ (b) $\cos x$ (c) $\sec x$ (d) $\operatorname{cosec} x$
154. $\frac{d}{dx}[\sinh^{-1}x]^x$ equals
- (a) $\frac{x}{\sqrt{1+x^2} \cdot \sinh^{-1}x} + \log(\sinh^{-1}x)$ (b) $(\sinh^{-1}x)^{x-1} \frac{1}{\sqrt{1+x^2}}$
- (c) $(\sinh^{-1}x)^x \left[\frac{x}{\sqrt{1+x^2} \cdot \sinh^{-1}x} + \log(\sinh^{-1}x) \right]$ (d) None of these
155. If $y = \sin^{-1}(x\sqrt{1-x} + \sqrt{x}\sqrt{1-x^2})$, then $\frac{dy}{dx} =$ [Roorkee 1981]
- (a) $\frac{-2x}{\sqrt{1-x^2}} + \frac{1}{2\sqrt{x-x^2}}$ (b) $\frac{-1}{\sqrt{1-x^2}} - \frac{1}{2\sqrt{x-x^2}}$ (c) $\frac{1}{\sqrt{1-x^2}} + \frac{1}{2\sqrt{x-x^2}}$ (d) None of these
156. If $y = \tan^{-1} \frac{x}{1+\sqrt{1-x^2}} + \sin\left\{2 \tan^{-1} \sqrt{\frac{1-x}{1+x}}\right\}$, then $\frac{dy}{dx} =$
- (a) $\frac{x}{\sqrt{1-x^2}}$ (b) $\frac{1-2x}{\sqrt{1-x^2}}$ (c) $\frac{1-2x}{2\sqrt{1-x^2}}$ (d) $\frac{1}{1+x^2}$
157. If $y = \cot^{-1}\left[\frac{\sqrt{1+x^2}+1}{x}\right]$ then $\frac{dy}{dx} =$
- (a) $\frac{1}{2} \cdot \frac{1}{1+x^2}$ (b) $\frac{1}{2} \cdot \frac{1}{1-x^2}$ (c) $\frac{2}{1+x^2}$ (d) $\frac{2}{1-x^2}$
158. If $y = \tan^{-1}\left(\frac{2^{x+1}}{1-4^x}\right)$, then $\frac{dy}{dx} =$
- (a) $\frac{2^{x+1} \log_e 2}{4^x}$ (b) $\frac{2^{x+1} \log_e 2}{1+4^x}$ (c) $\frac{2^{x+1} \log_e 2}{1-4^x}$ (d) $\frac{2^{x+1} \log_2 e}{1-4^x}$
159. If $y = \tan^{-1}\left(\frac{2 \cdot a^x}{1-a^{2x}}\right)$, then $\frac{dy}{dx} =$
- (a) $\frac{2 \cdot a^x \log a}{1-a^{2x}}$ (b) $\frac{2 \cdot a^x \log a}{1+a^{2x}}$ (c) $2 \cdot a^x \log a$ (d) $\frac{2 \cdot a^x \log a}{a^{2x}-1}$
160. If $y = x \cdot e^{\cos^{-1}x} + \sec(2x-1)$, then $\frac{dy}{dx}$ equals [Rajasthan PET 1986]

(a) $e^{\cos^{-1}x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) + \sec(2x-1) \cdot \tan(2x-1)$

(b) $e^{\cos^{-1}x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) - \sec(2x-1) \cdot \tan(2x-1)$

(c) $e^{\cos^{-1}x} \left(1 - \frac{x}{\sqrt{1-x^2}} \right) + 2 \sec(2x-1) \cdot \tan(2x-1)$

(d) None of these

161. If $y = \tan^{-1} \left(\frac{a+b \tan x}{b-a \tan x} \right)$, then $\frac{dy}{dx} =$

(a) 1

(b) -1

(c) x

(d) $\frac{1}{1+x^2}$

Methods of differentiation

Basic Level

162. If $x^3 + 8xy + y^3 = 64$, then $\frac{dy}{dx} =$

(a) $-\frac{3x^2+8y}{8x+3y^2}$

(b) $\frac{3x^2+8y}{8x+3y^2}$

(c) $\frac{3x+8y^2}{8x^2+3y}$

(d) None of these

163. If $\sin^2 x + 2 \cos y + xy = 0$, then $\frac{dy}{dx} =$

[AI CBSE 1980]

(a) $\frac{y+2 \sin x}{2 \sin y+x}$

(b) $\frac{y+\sin 2x}{2 \sin y-x}$

(c) $\frac{y+2 \sin x}{\sin y+x}$

(d) None of these

164. If $y \sec x + \tan x + x^2 y = 0$, then $\frac{dy}{dx} =$

[DSSE 1981; CBSE 1981]

(a) $\frac{2xy + \sec^2 x + y \sec x \tan x}{x^2 + \sec x}$

(b) $-\frac{2xy + \sec^2 x + \sec x \tan x}{x^2 + \sec x}$

(c) $-\frac{2xy + \sec^2 x + y \sec x \tan x}{x^2 + \sec x}$

(d) None of these

165. If $\sin(xy) + \frac{x}{y} = x^2 - y$, then $\frac{dy}{dx} =$

(a) $\frac{y[2xy - y^2 \cos(xy) - 1]}{xy^2 \cos(xy) + y^2 - x}$

(b) $\frac{[2xy - y^2 \cos(xy) - 1]}{xy^2 \cos(xy) + y^2 - x}$

(c) $-\frac{y[2xy - y^2 \cos(xy) - 1]}{xy^2 \cos(xy) + y^2 - x}$

(d) None of these

166. If $3 \sin(xy) + 4 \cos(xy) = 5$, then $\frac{dy}{dx} =$

[EAMCET 1994]

(a) $-\frac{y}{x}$

(b) $\frac{3 \sin(xy) + 4 \cos(xy)}{3 \cos(xy) - 4 \sin(xy)}$

(c) $\frac{3 \cos(xy) + 4 \sin(xy)}{4 \cos(xy) - 3 \sin(xy)}$

(d) None of these

167. If $x^2 e^y + 2xy e^x + 13 = 0$, then $\frac{dy}{dx} =$

[Rajasthan PET 1987]

(a) $\frac{2xe^{y-x} + 2y(x+1)}{x(xe^{y-x} + 2)}$

(b) $\frac{2xe^{x-y} + 2y(x+1)}{x(xe^{y-x} + 2)}$

(c) $-\frac{2xe^{y-x} + 2y(x+1)}{x(xe^{y-x} + 2)}$

(d) None of these

162 Differentiation

- 168.** If $\sin y = x \sin(a+y)$, then $\frac{dy}{dx} =$ [Karnataka CET 2000; UPSEAT 2001]
- (a) $\frac{\sin^2(a+y)}{\sin(a+2y)}$ (b) $\frac{\sin^2(a+y)}{\cos(a+2y)}$ (c) $\frac{\sin^2(a+y)}{\sin a}$ (d) $\frac{\sin^2(a+y)}{\cos a}$
- 169.** If $y = x^x$, then $\frac{dy}{dx} =$ [AISSE 1984; DSSE 1982; MNR 1979; SCRA 1996; Rajasthan PET 1996; Kerala (Engg.) 2002]
- (a) $x^x \log ex$ (b) $x^x \left(1 + \frac{1}{x}\right)$ (c) $(1 + \log x)$ (d) $x^x \log x$
- 170.** If $y^x + x^y = a^b$, then $\frac{dy}{dx} =$
- (a) $-\frac{y x^{y-1} + y^x \log y}{x y^{x-1} + x^y \log x}$ (b) $\frac{y x^{y-1} + y^x \log y}{x y^{x-1} + x^y \log x}$ (c) $-\frac{y x^{y-1} + y^x}{x y^{x-1} + x^y}$ (d) $\frac{y x^{y-1} + y^x}{x y^{x-1} + x^y}$
- 171.** If $y = \sqrt{\frac{(x-a)(x-b)}{(x-c)(x-d)}}$, then $\frac{dy}{dx} =$
- (a) $\frac{y}{2} \left[\frac{1}{x-a} + \frac{1}{x-b} - \frac{1}{x-c} - \frac{1}{x-d} \right]$ (b) $y \left[\frac{1}{x-a} + \frac{1}{x-b} - \frac{1}{x-c} - \frac{1}{x-d} \right]$
- (c) $\frac{1}{2} \left[\frac{1}{x-a} + \frac{1}{x-b} - \frac{1}{x-c} - \frac{1}{x-d} \right]$ (d) None of these
- 172.** $\frac{d}{dx}(x^{\log_e x}) =$ [MP PET 1993]
- (a) $2x^{(\log_e x - 1)} \cdot \log_e x$ (b) $x^{(\log_e x - 1)}$ (c) $\frac{2}{x} \log_e x$ (d) $x^{(\log_e x - 1)} \cdot \log_e x$
- 173.** If $x^y = y^x$, then $\frac{dy}{dx} =$ [DSSE 1986; MP PET 1997]
- (a) $\frac{y(x \log_e y + y)}{x(y \log_e x + x)}$ (b) $\frac{y(x \log_e y - y)}{x(y \log_e x - x)}$ (c) $\frac{x(x \log_e y - y)}{y(y \log_e x - x)}$ (d) $\frac{x(x \log_e y + y)}{y(y \log_e x + x)}$
- 174.** If $y = x^{\sin x}$, then $\frac{dy}{dx} =$ [DSSE 1983, 84]
- (a) $\frac{x \cos x \cdot \log x + \sin x}{x} \cdot x^{\sin x}$ (b) $\frac{y[x \cos x \cdot \log x + \cos x]}{x}$
- (c) $y[x \sin x \cdot \log x + \cos x]$ (d) None of these
- 175.** $\frac{d}{dx} \{(\sin x)^x\} =$ [DSSE 1985, 87; AISSE 1983]
- (a) $\left[\frac{x \cos x + \sin x \log \sin x}{\sin x} \right]$ (b) $(\sin x)^x \left[\frac{x \cos x + \sin x \log \sin x}{\sin x} \right]$
- (c) $(\sin x)^x \left[\frac{x \sin x + \sin x \log \sin x}{\sin x} \right]$ (d) None of these
- 176.** If $2^x + 2^y = 2^{x+y}$, then the value of $\frac{dy}{dx}$ at $x = y = 1$ is [Karnataka CET 2000]
- (a) 0 (b) -1 (c) 1 (d) 2
- 177.** If $x^y = e^{x-y}$, then $\frac{dy}{dx} =$ [MP PET 1987; MNR 1984; Roorkee 1954; Ranchi BIT 1991; Rajasthan PET 2000]

- (a) $\log x \cdot [\log(ex)]^{-2}$ (b) $\log x [\log(ex)]^2$ (c) $\log x \cdot (\log x)^2$ (d) None of these
178. $\frac{d}{dx} \{(\sin x)^{\log x}\} =$ [DSSE 1984]
 (a) $(\sin x)^{\log x} \left[\frac{1}{x} \log \sin x + \cot x \right]$ (b) $(\sin x)^{\log x} \left[\frac{1}{x} \log \sin x + \cot x \log x \right]$
 (c) $(\sin x)^{\log x} \left[\frac{1}{x} \log \sin x + \cot x \right]$ (d) None of these
179. $y = (\tan x)^{(\tan x)^{\tan x}}$, then at $x = \frac{\pi}{4}$, the value of $\frac{dy}{dx} =$ [West Bengal JEE 1990]
 (a) 0 (b) 1 (c) 2 (d) None of these
180. If $x^p y^q = (x+y)^{p+q}$, then $\frac{dy}{dx} =$ [Rajasthan PET 1999; UPSEAT 2001]
 (a) $\frac{y}{x}$ (b) $-\frac{y}{x}$ (c) $\frac{x}{y}$ (d) $-\frac{x}{y}$
181. If $y = (\tan x)^{\cot x}$, then $\frac{dy}{dx} =$
 (a) $y \operatorname{cosec}^2 x (1 - \log \tan x)$ (b) $y \operatorname{cosec}^2 x (1 + \log \tan x)$ (c) $y \operatorname{cosec}^2 x (\log \tan x)$ (d) None of these
182. If $y = \frac{e^x \log x}{x^2}$, then $\frac{dy}{dx} =$ [AI CBSE 1982]
 (a) $\frac{e^x [1 + (x+2) \log x]}{x^3}$ (b) $\frac{e^x [1 - (x-2) \log x]}{x^4}$ (c) $\frac{e^x [1 - (x-2) \log x]}{x^3}$ (d) $\frac{e^x [1 + (x-2) \log x]}{x^3}$
183. If $y = \frac{e^{2x} \cos x}{x \sin x}$, then $\frac{dy}{dx} =$
 (a) $\frac{e^{2x} [(2x-1) \cot x - x \operatorname{cosec}^2 x]}{x^2}$ (b) $\frac{e^{2x} [(2x+1) \cot x - x \operatorname{cosec}^2 x]}{x^2}$
 (c) $\frac{e^{2x} [(2x-1) \cot x + x \operatorname{cosec}^2 x]}{x^2}$ (d) None of these
184. If $y = \frac{\sqrt{x} (2x+3)^2}{\sqrt{x+1}}$, then $\frac{dy}{dx} =$ [AISSE 1986]
 (a) $y \left[\frac{1}{2x} + \frac{4}{2x+3} - \frac{1}{2(x+1)} \right]$ (b) $y \left[\frac{1}{3x} + \frac{4}{2x+3} + \frac{1}{2(x+1)} \right]$ (c) $y \left[\frac{1}{3x} + \frac{4}{2x+3} + \frac{1}{(x+1)} \right]$ (d)
185. If $y = \frac{2(x - \sin x)^{3/2}}{\sqrt{x}}$, then $\frac{dy}{dx} =$
 (a) $\frac{2(x - \sin x)^{3/2}}{\sqrt{x}} \left[\frac{3}{2} \cdot \frac{1 - \cos x}{1 - \sin x} - \frac{1}{2x} \right]$ (b) $\frac{2(x - \sin x)^{3/2}}{\sqrt{x}} \left[\frac{3}{2} \cdot \frac{1 - \cos x}{x - \sin x} - \frac{1}{2x} \right]$
 (c) $\frac{2(x - \sin x)^{1/2}}{\sqrt{x}} \left[\frac{3}{2} \cdot \frac{1 - \cos x}{x - \sin x} - \frac{1}{2x} \right]$ (d) None of these
186. $\frac{d}{dx} [(x-2)^x] =$ [Rajasthan PET 1992]
 (a) $(x-2)^x [x + \log(x-2)]$ (b) $(x-2)^{x-1} [(x-2) \log(x-2) + x]$ (c) $(x-2)^{x-1} [x + \log(x-2)]$ (d) None of these
187. The derivative of x^{a^x} is

164 Differentiation

- (a) $x^{a^x} \left[\frac{a^x}{x} + a^x \log a \log x \right]$ (b) $x^{a^x} [a^x + xa^x \log x]$ (c) $x^{a^x} [xa^x + a^x \log x]$ (d) None of these
- 188.** If $x = a \sin 2\theta(1 + \cos 2\theta)$, $y = b \cos 2\theta(1 - \cos 2\theta)$, then $\frac{dy}{dx} =$ **[Kurukshetra CEE 1998]**
- (a) $\frac{b \tan \theta}{a}$ (b) $\frac{a \tan \theta}{b}$ (c) $\frac{a}{b \tan \theta}$ (d) $\frac{b}{a \tan \theta}$
- 189.** If $x = a \left(\cos t + \log \tan \frac{t}{2} \right)$, $y = a \sin t$, then $\frac{dy}{dx} =$ **[Rajasthan PET 1997; MP PET 2001]**
- (a) $\tan t$ (b) $-\tan t$ (c) $\cot t$ (d) $-\cot t$
- 190.** If $x = \sin^{-1}(3t - 4t^3)$ and $y = \cos^{-1}(\sqrt{1 - t^2})$, then $\frac{dy}{dx}$ is equal to
- (a) $1/2$ (b) $2/5$ (c) $3/2$ (d) $1/3$
- 191.** If $x = \frac{2t}{1+t^2}$, $y = \frac{1-t^2}{1+t^2}$, then $\frac{dy}{dx}$ equals **[Rajasthan PET 1999]**
- (a) $\frac{2t}{t^2+1}$ (b) $\frac{2t}{t^2-1}$ (c) $\frac{2t}{1-t^2}$ (d) None of these
- 192.** If $x = \frac{3at}{1+t^3}$, $y = \frac{3at^2}{1+t^3}$, then $\frac{dy}{dx} =$
- (a) $\frac{t(2+t^3)}{1-2t^3}$ (b) $\frac{t(2-t^3)}{1-2t^3}$ (c) $\frac{t(2+t^3)}{1+2t^3}$ (d) $\frac{t(2-t^3)}{1+2t^3}$
- 193.** If $x = a(t + \sin t)$ and $y = a(1 - \cos t)$, then $\frac{dy}{dx}$ equals **[Rajasthan PET 1996; MP PET 2002]**
- (a) $\tan(t/2)$ (b) $\cot(t/2)$ (c) $\tan 2t$ (d) $\tan t$
- 194.** If $x = a \cos^4 \theta$, $y = a \sin^4 \theta$, then $\frac{dy}{dx}$ at $\theta = \frac{3\pi}{4}$ **[Kerala (Engg.) 2002]**
- (a) -1 (b) 1 (c) $-a^2$ (d) a^2
- 195.** If $x = 2 \cos t - \cos 2t$, $y = 2 \sin t - \sin 2t$, then at $t = \frac{\pi}{4}$, $\frac{dy}{dx} =$
- (a) $\sqrt{2} + 1$ (b) $\sqrt{2} - 1$ (c) $\frac{\sqrt{2}+1}{2}$ (d) None of these
- 196.** If $\tan y = \frac{2t}{1-t^2}$ and $\sin x = \frac{2t}{1+t^2}$, then $\frac{dy}{dx} =$ **[Rajasthan PET 1994]**
- (a) $\frac{2}{1+t^2}$ (b) $\frac{1}{1+t^2}$ (c) 1 (d) 2
- 197.** If $x = at^2$, $y = 2at$ then $\frac{dy}{dx}$ at $t = 2$ **[Rajasthan PET 1992]**
- (a) 2 (b) 4 (c) $1/2$ (d) $1/4$
- 198.** If $x = t^2 + t + 1$ and $y = \sin \frac{\pi}{2}t + \cos \frac{\pi}{2}t$ then at $t = 1$ $\frac{dy}{dx}$ equals
- (a) $-\pi/6$ (b) $\pi/2$ (c) $-\pi/4$ (d) $\pi/3$
- 199.** If $y = e^{x+e^{x+e^{x+\dots}}}$, then $\frac{dy}{dx} =$

(a) $\frac{y}{1-y}$ (b) $\frac{1}{1-y}$ (c) $\frac{y}{1+y}$ (d) $\frac{y}{y-1}$

200. If $y = (\sin x)^{(\sin x)^{(\sin x)^{\dots \infty}}}$, then $\frac{dy}{dx} =$

(a) $\frac{y^2 \cot x}{1-y \log \sin x}$ (b) $\frac{y^2 \cot x}{1+y \log \sin x}$ (c) $\frac{y \cot x}{1-y \log \sin x}$ (d) $\frac{y \cot x}{1+y \log \sin x}$

201. The differential equation satisfied by the function $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$, is [MP PET 1998]

(a) $(2y-1)\frac{dy}{dx} - \sin x = 0$ (b) $(2y-1)\cos x + \frac{dy}{dx} = 0$
(c) $(2y-1)\cos x - \frac{dy}{dx} = 0$ (d) $(2y-1)\frac{dy}{dx} - \cos x = 0$

202. If $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots \infty}}}$, then $\frac{dy}{dx} =$

(a) $\frac{x}{2y-1}$ (b) $\frac{x}{2y+1}$ (c) $\frac{1}{x(2y-1)}$ (d) $\frac{1}{x(1-2y)}$

203. If $y = \sqrt{f(x) + \sqrt{f(x) + \sqrt{f(x) + \dots \infty}}}$, then the value of $(2y-1)\frac{dy}{dx}$ is

(a) $f(x)$ (b) $f'(x)$ (c) $2f'(x)$ (d) None of these

Advance Level

204. If $x^2 + y^2 = t - \frac{1}{t}$, $x^4 + y^4 = t^2 + \frac{1}{t^2}$, then $\frac{dy}{dx}$ equals

[Rajasthan PET 1999]

(a) $1/xy^3$ (b) $1/x^3y$ (c) $-1/x^3y$ (d) $-1/xy^3$

205. If $f'(x) = \sin(\log x)$ and $y = f\left(\frac{2x+3}{3-2x}\right)$, then $\frac{dy}{dx} =$

[BIT Ranchi 1986]

(a) $\frac{9 \cos(\log x)}{x(3-2x)^2}$ (b) $\frac{9 \cos\left(\log \frac{2x+3}{3-2x}\right)}{x(3-2x)^2}$ (c) $\frac{9 \sin\left(\log \frac{2x+3}{3-2x^2}\right)}{(3-2x)^2}$ (d) None of these

206. $\frac{dy}{dx}$ of $\log(xy) = x^2 + y^2$ is

(a) $\frac{y(2x^2-1)}{x(1-2y^2)}$ (b) $\frac{y(2x^2+1)}{x(1+2y^2)}$ (c) $\frac{x(2x^2-1)}{y(2y^2-1)}$ (d) $\frac{y(2x^2-1)}{x(2y^2-1)}$

207. $(x-y)e^{x/(x-y)} = k$, then

(a) $(y-2x)\frac{dy}{dx} + 3x - 2y = 0$ (b) $y\frac{dy}{dx} + x - 2y = 0$ (c) $a\left(y\frac{dy}{dx} + x - 2y\right) = 1$ (d) None of these

208. If $y = (x^x)^x$, then $\frac{dy}{dx} =$

(a) $(x^x)^x(1+2\log x)$ (b) $(x^x)^x(1+\log x)$ (c) $x(x^x)^x(1+2\log x)$ (d) $x(x^x)^x(1+\log x)$

209. If $y = (x \log x)^{\log \log x}$, then $\frac{dy}{dx} =$

[Roorkee 1981]

(a) $(x \log x)^{\log \log x} \left\{ \frac{1}{x \log x} (\log x + \log \log x) + (\log \log x) \left(\frac{1}{x} + \frac{1}{x \log x} \right) \right\}$

166 Differentiation

(b) $(x \log x)^{x \log x} \log \log x \left[\frac{2}{\log x} + \frac{1}{x} \right]$

(c) $(x \log x)^{x \log x} \frac{\log \log x}{x} \left[\frac{1}{\log x} + 1 \right]$

(d) None of these

210. If $y = \left(1 + \frac{1}{x}\right)^x$, then $\frac{dy}{dx} =$

[BIT Ranchi 1992]

(a) $\left(1 + \frac{1}{x}\right)^x \left[\log \left(1 + \frac{1}{x}\right) - \frac{1}{1+x} \right]$

(b) $\left(1 + \frac{1}{x}\right)^x \left[\log \left(1 + \frac{1}{x}\right) \right]$

(c) $\left(x + \frac{1}{x}\right)^x \left[\log(x-1) - \frac{x}{1+x} \right]$

(d) $\left(x + \frac{1}{x}\right)^x \left[\log \left(1 + \frac{1}{x}\right) + \frac{1}{1+x} \right]$

211. If $y = x^{(x^x)}$, then $\frac{dy}{dx} =$

[AISSE 1989]

(a) $y[x^x(\log ex) \cdot \log x + x^x]$

(b) $y[x^x(\log ex) \cdot \log x + x]$

(c) $y[x^x(\log ex) \cdot \log x + x^{x-1}]$

(d) $y[x^x(\log_e x) \cdot \log x + x^{x-1}]$

212. If $y = \frac{a^{\cos^{-1} x}}{1 + a^{\cos^{-1} x}}$ and $z = a^{\cos^{-1} x}$, then $\frac{dy}{dz} =$

[MP PET 1994]

(a) $\frac{1}{1 + a^{\cos^{-1} x}}$

(b) $-\frac{1}{1 + a^{\cos^{-1} x}}$

(c) $\frac{1}{(1 + a^{\cos^{-1} x})^2}$

(d) None of these

213. Let the function $y = f(x)$ be given by $x = t^5 - 5t^3 - 20t + 7$ and $y = 4t^3 - 3t^2 - 18t + 3$, where $t \in (-2, 2)$. Then $f'(x)$ at $t = 1$ is

(a) $\frac{5}{2}$

(b) $\frac{2}{5}$

(c) $\frac{7}{5}$

(d) None of these

214. If $y = \sqrt{x}^{\sqrt{x}^{\sqrt{x}^{\dots \infty}}}$, then $\frac{dy}{dx} =$

(a) $\frac{y^2}{2x - 2y \log x}$

(b) $\frac{y^2}{2x + \log x}$

(c) $\frac{y^2}{2x + 2y \log x}$

(d) None of these

215. If $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots}}}$, then $\frac{dy}{dx}$ equals

(a) $\frac{y}{2y - x}$

(b) $\frac{y}{2y + x}$

(c) $\frac{y}{y - 2x}$

(d) $\frac{y}{y + 2x}$

216. If $y = \frac{x}{a + \frac{x}{b + \frac{x}{a + \frac{x}{b + \dots}}}}$, then $\frac{dy}{dx}$ equals

(a) $\frac{b}{a(b + 2y)}$

(b) $\frac{b}{b + 2y}$

(c) $\frac{a}{b(b + 2y)}$

(d) None of these

217. If $y = \frac{\sin x}{1 + \frac{\cos x}{1 + \frac{\sin x}{1 + \cos x \dots \infty}}}$, then

$\frac{dy}{dx} =$

(a) $\frac{(1+y)\cos x + \sin x}{1 + 2y + \cos x - \sin x}$

(b) $\frac{(1+y)\cos x - \sin x}{1 + 2y + \cos x + \sin x}$

(c) $\frac{(1+y)\cos x + \sin x}{1 + 2y + \cos x + \sin x}$

(d) None of these

218. If $f(x) = \frac{1}{1-x}$, then the derivative of the composite function $f[f(x)]$ is equal to [Orissa JEE 2003]

- (a) 0 (b) $1/2$ (c) 1 (d) 2

219. If $u = f(x^3)$, $v = g(x^2)$, $f'(x) = \cos x$ and $g'(x) = \sin x$ then $\frac{du}{dv}$ is

- (a) $\frac{3}{2}x \cdot \cos x^3 \cdot \operatorname{cosec} x^2$ (b) $\frac{2}{3} \sin x^3 \cdot \sec x^2$ (c) $\tan x$ (d) None of these

220. Let $f(x) = e^x$, $g(x) = \sin^{-1} x$ and $h(x) = f(g(x))$, then $h'(x)/h(x) =$

- (a) $e^{\sin^{-1} x}$ (b) $1/\sqrt{1-x^2}$ (c) $\sin^{-1} x$ (d) $1/(1-x^2)$

Differentiation of a Function with Respect to Another Function

Basic Level

221. The derivative of $\sin^2 x$ with respect to $\cos^2 x$ is [DCE 2002]

- (a) $\tan^2 x$ (b) $\tan x$ (c) $-\tan x$ (d) None of these

222. The differential of e^{x^3} with respect to $\log x$ is [KCET 2002]

- (a) e^{x^3} (b) $3x^2 e^{x^3}$ (c) $3x^3 e^{x^3}$ (d) $3x^2 e^{x^3} + 3x^2$

223. The differential coefficient of x^6 with respect to x^3 is [EAMCET 1988; UPSEAT 2000]

- (a) $5x^2$ (b) $3x^3$ (c) $5x^5$ (d) $2x^3$

224. The rate of change of $\sqrt{x^2 + 16}$ with respect to $\frac{x}{x-1}$ at $x = 3$, will be [MP PET 1987]

- (a) $-\frac{24}{5}$ (b) $\frac{24}{5}$ (c) $\frac{12}{5}$ (d) $-\frac{12}{5}$

225. Differential coefficient of $\sin^{-1} \frac{1-x}{1+x}$ w.r.t. $\sqrt{x}$ is [Roorkee 1984]

- (a) $\frac{1}{2\sqrt{x}}$ (b) $\frac{\sqrt{x}}{\sqrt{1-x}}$ (c) 1 (d) None of these

226. Differential coefficient of $\sec^{-1} \frac{1}{2x^2-1}$ w.r.t. $\sqrt{1-x^2}$ at $x = \frac{1}{2}$ is

- (a) 2 (b) 4 (c) 6 (d) 1

227. Differential coefficient of $\sin^{-1} x$ w.r.t. $\cos^{-1} \sqrt{1-x^2}$ is [MNR 1983; AMU 2002]

- (a) 1 (b) $\frac{1}{1+x^2}$ (c) 2 (d) None of these

228. The differential coefficient of $\tan^{-1} \sqrt{x}$ with respect to $\sqrt{x}$ is

- (a) $\frac{1}{\sqrt{1+x}}$ (b) $\frac{1}{2x\sqrt{1+x}}$ (c) $\frac{1}{2\sqrt{x(1+x)}}$ (d) $\frac{1}{1+x}$

229. Derivative of $\sec^{-1} \left\{ \frac{1}{2x^2-1} \right\}$ w.r.t. $\sqrt{1+3x}$ at $x = -\frac{1}{3}$ is [EAMCET 1991]

- (a) 0 (b) $1/2$ (c) $1/3$ (d) None of these

230. Differential coefficient of $\cos^{-1}(\sqrt{x})$ with respect to $\sqrt{1-x}$ is

168 Differentiation

- (a) $\sqrt{x}$ (b) $-\sqrt{x}$ (c) $\frac{1}{\sqrt{x}}$ (d) $-\frac{1}{\sqrt{x}}$
231. Differential coefficient of $\tan^{-1} \sqrt{\frac{1-x^2}{1+x^2}}$ w.r.t. $\cos^{-1}(x^2)$ is
 (a) $\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) 1 (d) 0
232. If $u = \tan^{-1} \left\{ \frac{\sqrt{1+x^2}-1}{x} \right\}$ and $v = 2 \tan^{-1} x$, then $\frac{du}{dv}$ is equal to
 (a) 4 (b) 1 (c) $\frac{1}{4}$ (d) $-\frac{1}{4}$
233. The derivative of $\sin^{-1} \left(\frac{2x}{1+x^2} \right)$ w.r.t. $\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$ is [Karnataka CET 2000]
 (a) -1 (b) 1 (c) 2 (d) 4
234. Differential coefficient of $\tan^{-1} \left(\frac{x}{1+\sqrt{1-x^2}} \right)$ w.r.t. $\sin^{-1} x$, is
 (a) $\frac{1}{2}$ (b) 1 (c) 2 (d) $\frac{3}{2}$
235. The derivative of $\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$ w.r.t. $\cot^{-1} \left(\frac{1-3x^2}{3x-x^2} \right)$ is [Karnataka CET 2003]
 (a) 1 (b) $\frac{3}{2}$ (c) $\frac{2}{3}$ (d) $\frac{1}{2}$
236. The differential coefficient of $e^{\sin^{-1} x}$ with respect to $\sin^{-1} x$ is
 (a) $\cos^{-1} x$ (b) $e^{\cos^{-1} x}$ (c) $e^{\sin^{-1} x}$ (d) $\sin^{-1} x$

Advance Level

237. Differential coefficient of $\frac{\tan^{-1} x}{1+\tan^{-1} x}$ w.r.t. $\tan^{-1} x$ is
 (a) $\frac{1}{1+\tan^{-1} x}$ (b) $\frac{-1}{1+\tan^{-1} x}$ (c) $\frac{1}{(1+\tan^{-1} x)^2}$ (d) $\frac{-1}{2(1+\tan^{-1} x)^2}$
238. The derivative of $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ with respect to $\tan^{-1} \left(\frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$ at $x = 0$, is
 (a) $\frac{1}{8}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) 1
239. Differentiation of $\tan^{-1} \left(\frac{\sqrt{1-x^2}}{x} \right)$ with respect to $\cos^{-1}(2x\sqrt{1-x^2})$ is
 (a) $\frac{1}{2}$ (b) $-\frac{1}{2}$ (c) 1 (d) -1
240. Differentiation of $\sin^{-1}(2ax\sqrt{1-a^2x^2})$ with respect to $\sqrt{1-a^2x^2}$ is
 (a) 2 (b) ax (c) $\frac{2}{ax}$ (d) $-\frac{2}{ax}$

241. Differentiation of $\tan^{-1}\left(\frac{1+ax}{1-ax}\right)$ with respect to $\sqrt{1+a^2x^2}$ is

- (a) $\frac{1}{ax\sqrt{1+ax}}$ (b) $\frac{1}{\sqrt{1+ax}}$ (c) $\frac{1}{ax\sqrt{1+a^2x^2}}$ (d) $\frac{1}{ax\sqrt{1-a^2x^2}}$

242. The value of derivative of $\tan^{-1}\left(\frac{2x\sqrt{1-x^2}}{1-2x^2}\right)$ w.r.t. to $\sec^{-1}\left(\frac{1}{2x^2-1}\right)$ at $x = \frac{1}{2}$ equals

- (a) 1 (b) -1 (c) 0 (d) None of these

Successive Differentiation or Higher Order Derivatives

Basic Level

243. If $y = (x^2 - 1)^m$, then the $(2m)^{\text{th}}$ differential coefficient of y is

- (a) m (b) $(2m)!$ (c) $2m$ (d) $m!$

244. The n^{th} derivative of xe^x vanishes when

[AMU 1999]

- (a) $x = 0$ (b) $x = -1$ (c) $x = -n$ (d) $x = n$

245. If $x^p y^q = (x+y)^{p+q}$, then $\frac{d^2y}{dx^2} =$

[West Bengal JEE 1992]

- (a) 0 (b) 1 (c) 2 (d) None of these

246. If $y = A \cos nx + B \sin nx$, then $\frac{d^2y}{dx^2} =$

[Karnataka CET 1996]

- (a) n^2y (b) $-y$ (c) $-n^2y$ (d) None of these

247. If $x = a \sin \theta$ and $y = b \cos \theta$, then $\frac{d^2y}{dx^2}$ is

[UPSEAT 2002]

- (a) $\frac{a}{b^2} \sec^2 \theta$ (b) $\frac{-b}{a} \sec^2 \theta$ (c) $\frac{-b}{a^2} \sec^3 \theta$ (d) $\frac{-b}{a^2 \sec^3 \theta}$

248. If $y = a \cos(\log x) + \sin(\log x)$, then

- (a) $x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} + y = 0$ (b) $x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} - y = 0$ (c) $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = 0$ (d) $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = 0$

249. If $e^y + xy = e$, then the value of $\frac{d^2y}{dx^2}$ for $x = 0$, is

[Kurukshetra CEE 2002]

- (a) $\frac{1}{e}$ (b) $\frac{1}{e^2}$ (c) $\frac{1}{e^3}$ (d) None of these

250. If $y = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \dots$, then $\frac{d^2y}{dx^2} =$

[Karnataka CET 2003]

- (a) x (b) $-x$ (c) $-y$ (d) y

251. If $y = ax^{n+1} + bx^{-n}$, then $x^2 \frac{d^2y}{dx^2} =$

- (a) $n(n-1)y$ (b) $n(n+1)y$ (c) ny (d) n^2y

170 Differentiation

252. If $y = a + bx^2$; a, b arbitrary constants, then

[EAMCET 1994]

- (a) $\frac{d^2y}{dx^2} = 2xy$ (b) $x \frac{d^2y}{dx^2} = \frac{dy}{dx}$ (c) $x \frac{d^2y}{dx^2} - \frac{dy}{dx} + y = 0$ (d) $x \frac{d^2y}{dx^2} = 2xy$

253. If $y = x \log \left(\frac{x}{a+bx} \right)$, then $x^3 \frac{d^2y}{dx^2} =$

[West Bengal JEE 1991; Roorkee 1976]

- (a) $x \frac{dy}{dx} - y$ (b) $\left(x \frac{dy}{dx} - y \right)^2$ (c) $y \frac{dy}{dx} - x$ (d) $\left(y \frac{dy}{dx} - x \right)^2$

254. $\frac{d^2}{dx^2} (2 \cos x \cos 3x) =$

[Rajasthan PET 2003]

- (a) $2^2(\cos 2x + 2^2 \cos 4x)$ (b) $2^2(\cos 2x - 2^2 \cos 4x)$ (c) $2^2(-\cos 2x + 2^2 \cos 4x)$ (d) $-2^2(\cos 2x + 2^2 \cos 4x)$

255. If $x = t^2$, $y = t^3$, then $\frac{d^2y}{dx^2} =$

- (a) $3/2$ (b) $3/(4t)$ (c) $3/(2t)$ (d) $3t/2$

256. If $y = ae^{mx} + be^{-mx}$, then $\frac{d^2y}{dx^2} - m^2y =$

[MP PET 1987]

- (a) $m^2(ae^{mx} - be^{-mx})$ (b) 1 (c) 0 (d) None of these

257. If $y = x^2 e^{mx}$, where m is a constant, then $\frac{d^3y}{dx^3} =$

[MP PET 1987]

- (a) $me^{mx}(m^2x^2 + 6mx + 6)$ (b) $2m^3xe^{mx}$ (c) $me^{mx}(m^2x^2 + 2mx + 2)$ (d) None of these

258. If f be a polynomial, then the second derivative of $f(e^x)$ is

- (a) $f'(e^x)$ (b) $f''(e^x)e^x + f'(e^x)$ (c) $f''(e^x)e^{2x} + f'(e^x)$ (d) $f''(e^x)e^{2x} + f'(e^x)e^x$

259. If $y = ae^x + be^{-x} + c$ where a, b, c are parameters then $y''' =$

- (a) y (b) y' (c) 0 (d) y''

260. If $y = a \cos(\log x) + b \sin(\log x)$ where a, b are parameters then $x^2y'' + xy' =$

[EAMCET 2002]

- (a) y (b) $-y$ (c) $2y$ (d) $-2y$

261. If $y = x^3 \log \log_e(1+x)$ then $y''(0)$ equals

[AMU 1999]

- (a) 0 (b) -1 (c) $6 \log_e 2$ (d) 6

262. $\frac{d^2x}{dy^2}$ is equal to

[AMU 2001]

- (a) $\frac{1}{(dy/dx)^2}$ (b) $\frac{(d^2y/dx^2)}{(dy/dx)^2}$ (c) $\frac{d^2y}{dx^2}$ (d) $\frac{(-d^2y/dx^2)}{(dy/dx)^2}$

263. If $x = e^t \sin t$, $y = e^t \cos t$, t is a parameter, then $\frac{d^2y}{dx^2}$ at $(1, 1)$ is equal to

[AMU 2001]

- (a) $-1/2$ (b) $-1/4$ (c) 0 (d) $1/2$

264. $\frac{d^n}{dx^n}(\sin 2x) =$

- (a) $\sin\left(\frac{n\pi}{2} + x\right)$ (b) $2^n \sin\left(\frac{n\pi}{2} + 2x\right)$ (c) $2^n \sin\left(\frac{\pi}{2} + 2x\right)$ (d) None of these

265. $\frac{d^n}{dx^n}(\log x) =$

[Rajasthan PET 2002]

- (a) $\frac{(n-1)!}{x^n}$ (b) $\frac{n!}{x^n}$ (c) $\frac{(n-2)!}{x^n}$ (d) $(-1)^{n-1} \frac{(n-1)!}{x^n}$

266. $\frac{d^n}{dx^n}(e^{2x} + e^{-2x}) =$

- (a) $e^{2x} + (-1)^n e^{-2x}$ (b) $2^n(e^{2x} - e^{-2x})$ (c) $2^n[e^{2x} + (-1)^n e^{-2x}]$ (d) None of these

267. If $y = \sin x \sin 3x$, then $y_n =$

- (a) $\frac{1}{2} \left[\cos\left(2x + n\frac{\pi}{2}\right) - \cos\left(4x + n\frac{\pi}{2}\right) \right]$ (b) $\frac{1}{2} \left[2^n \cos\left(2x + n\frac{\pi}{2}\right) - 4^n \cos\left(4x + n\frac{\pi}{2}\right) \right]$
 (c) $\frac{1}{2} \left[4^n \cos\left(4x + n\frac{\pi}{2}\right) - 2^n \cos\left(2x + n\frac{\pi}{2}\right) \right]$ (d) None of these

268. The n^{th} derivative of $\frac{x}{1-x}$ is

- (a) $\frac{(-1)^n n!}{(1-x)^{n+1}}$ (b) $\frac{n!}{(1-x)^{n+1}}$ (c) $\frac{(-1)^n}{(1-x)^{n+1}}$ (d) $\frac{1}{(1-x)^{n+1}}$

269. If $y = \sin^2 x$, then value of y_n is

- (a) $2^n \cos\left(2x + \frac{n\pi}{2}\right)$ (b) $-2^n \cos\left(2x + \frac{n\pi}{2}\right)$ (c) $-2^{n-1} \cos\left(2x + \frac{n\pi}{2}\right)$ (d) None of these

270. If $y = \sin 2x \cos 2x$, then value of y_n is

- (a) $2^{2n-1} \sin\left(4x + \frac{n\pi}{2}\right)$ (b) $2^{2n} \sin\left(4x + \frac{n\pi}{2}\right)$ (c) $2^{2n-1} \cos\left(4x + \frac{n\pi}{2}\right)$ (d) None of these

271. If $y = e^{6-5x}$, then the value of y_n is

- (a) $5^n e^{6-5x}$ (b) $(-5)^n e^{6-5x}$ (c) $5^{n-1} e^{6-5x}$ (d) $(-5)^{n-1} e^{6-5x}$

272. If $y = 8^x$, then the value of y_n is

- (a) $\frac{8^x}{\log_e 8}$ (b) $\frac{8^x}{(\log_e 8)^n}$ (c) $8^x \log_e 8$ (d) $8^x (\log_e 8)^n$

273. $D^n[f(ax+b)]$ is equal to

- (a) $n! f_n(ax+b)$ (b) $a^n f_n(ax+b)$ (c) $(n-1)! a^n f_n(ax+b)$ (d) 0

274. If $y = x^{n-1} \log x$, then which of the following statement is true

- (a) $xy_n = n!$ (b) $xy_n = (n-1)!$ (c) $xy_n = (n-2)!$ (d) $x^2 y_n = n!$

172 Differentiation

- 275.** If $x = f_1(t)$ and $y = f_2(t)$, then $\frac{d^2y}{dx^2} =$
- (a) $\frac{f_1' f_2'' - f_2' f_1''}{(f_1')^2}$ (b) $\frac{f_1' f_2'' - f_2' f_1''}{(f_1')^3}$ (c) $\frac{f_1''(t)}{f_2''(t)}$ (d) $\frac{-f_1''(t)}{f_2''(t)}$
- 276.** If $y^2 = p(x)$ is a polynomial of degree three, then $2 \frac{d}{dx} \left\{ y^3 \cdot \frac{d^2y}{dx^2} \right\} =$ [IIT 1988; Rajasthan PET 2000]
- (a) $p'''(x) + p'(x)$ (b) $p''(x) \cdot p'''(x)$ (c) $p(x) \cdot p'''(x)$ (d) Constant
- 277.** If $x = a \cos \theta$, $y = b \sin \theta$, then $\frac{d^3y}{dx^3}$ is equal to
- (a) $-\frac{3b}{a^3} \operatorname{cosec}^4 \theta \cot^4 \theta$ (b) $\frac{3b}{a^3} \operatorname{cosec}^4 \theta \cot \theta$ (c) $-\frac{3b}{a^3} \operatorname{cosec}^4 \theta \cot \theta$ (d) None of these
- 278.** $\frac{d^{20}y}{dx^{20}} (2 \cos x \cos 3x) =$ [EAMCET 1994]
- (a) $2^{20} (\cos 2x - 2^{20} \cos 4x)$ (b) $2^{20} (\cos 2x + 2^{20} \cos 4x)$ (c) $2^{20} (\sin 2x + 2^{20} \sin 4x)$ (d) $2^{20} (\sin 2x - 2^{20} \sin 4x)$
- 279.** If $u = x^2 + y^2$ and $x = s + 3t$, $y = 2s - t$, then $\frac{d^2u}{ds^2} =$ [Orissa JEE 2002]
- (a) 12 (b) 32 (c) 36 (d) 10
- 280.** If $y = \sin x + e^x$, then $\frac{d^2x}{dy^2} =$ [KCET 1999; UPSEAT 2001; Haryana CEE 2002]
- (a) $(-\sin x + e^x)^{-1}$ (b) $\frac{\sin x - e^x}{(\cos x + e^x)^2}$ (c) $\frac{\sin x - e^x}{(\cos x + e^x)^3}$ (d) $\frac{\sin x + e^x}{(\cos x + e^x)^3}$
- 281.** If $x = at^2$, $y = 2at$, then $\frac{d^2y}{dx^2} =$ [Karnataka CET 1993]
- (a) $-\frac{1}{t^2}$ (b) $\frac{1}{2at^3}$ (c) $-\frac{1}{t^3}$ (d) $-\frac{1}{2at^3}$
- 282.** If $I_n = \frac{d^n}{dx^n} (x^n \log x)$, then $I_n - nI_{n-1} =$ [EAMCET 2003]
- (a) n (b) $n - 1$ (c) $n !$ (d) $(n - 1)!$
- 283.** If $y = (\sin^{-1} x)^2 + k \sin^{-1} x$ then which is true
- (a) $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2$ (b) $(1 + x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2$ (c) $(1 - x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 2$ (d) $(1 + x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 2$
- 284.** If $y = e^{\tan^{-1} x}$ then which is true
- (a) $(1 + x^2)y_2 + (2x - 1)y_1 = 0$ (b) $(1 + x^2)y_2 + (2x + 1)y_1 = 0$ (c) $(1 + x^2)y_2 - (2x - 1)y_1 = 0$ (d) $(1 + x^2)y_2 - (2x + 1)y_1 = 0$
- 285.** The function $u = e^x \sin x$, $v = e^x \cos x$ satisfy the equation
- (a) $v \frac{du}{dv} = u \frac{dv}{dx} + u^2 + v^2$ (b) $\frac{d^2u}{dx^2} = 2v$ (c) $\frac{d^2v}{dx^2} = -2u$ (d) None of these

286. If $x^2 + y^2 = a^2$ and $k = \frac{1}{a}$, then k is equal to

- (a) $\frac{y''}{\sqrt{1+y'}}$ (b) $\frac{|y''|}{\sqrt{(1+y'^2)^3}}$ (c) $\frac{2y''}{\sqrt{1+y'^2}}$ (d) $\frac{y''}{2\sqrt{(1+y'^2)^3}}$

287. If $(a+bx)e^{y/x} = x$, then the value of $x^3 \frac{d^2y}{dx^2}$ is

- (a) $\left(y \frac{dy}{dx} - x\right)^2$ (b) $\left(x \frac{dy}{dx} - y\right)^2$ (c) $x \frac{dy}{dx} - y$ (d) None of these

288. If $y = [\log(x + \sqrt{x^2 + 1})]^2$ then which is correct

- (a) $(1+x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 2$ (b) $(1+x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2$ (c) $(1+x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 0$ (d) $(1+x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 0$

289. If $y = \frac{1}{x^2 - a^2}$, then $\frac{d^2y}{dx^2}$ equals

- (a) $\frac{3x^2 + a^2}{(x^2 - a^2)^3}$ (b) $\frac{3x^2 + a^2}{(x^2 - a^2)^4}$ (c) $\frac{2(3x^2 + a^2)}{(x^2 - a^2)^3}$ (d) $\frac{2(3x^2 + a^2)}{(x^2 - a^2)^4}$

290. If $y^{1/m} + y^{-1/m} = 2x$, then $(x^2 - 1)y_2 + xy_1$ is equal to

- (a) m^2y (b) $-m^2y$ (c) $\pm m^2y$ (d) $\pm my$

291. If $y = (\sin^{-1} x)^2 + (\cos^{-1} x)^2$, then $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx}$ is equal to

- (a) 4 (b) 3 (c) 1 (d) 0

292. If $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$, then $xy_2 + \frac{1}{2}y_1 - \frac{1}{4}y$ is equal to

- (a) 0 (b) 1 (c) -1 (d) 2

293. If $y = \sin 2x$ then $\frac{d^6y}{dx^6}$ at $x = \frac{\pi}{2}$ is equal to

- (a) -64 (b) 0 (c) 64 (d) None of these

294. $\frac{d^n}{dx^n} \cos^2 x =$

- (a) $2^{n-1} \cos\left(2x + \frac{\pi}{2}\right)$ (b) $2^{n-1} \cos\left(2x - \frac{\pi}{2}\right)$ (c) $2^{n-1} \cos\left(2x + \frac{n\pi}{2}\right)$ (d) $2^{n-1} \cos\left(2x - \frac{n\pi}{2}\right)$

295. If $y = \cos^4 x$, then y_n is equal to

- (a) $2^{2n-3} \cos\left(4x + \frac{n\pi}{2}\right) + 2^{n-1} \cos\left(2x + \frac{n\pi}{2}\right)$ (b) $2^{2n-3} \cos\left(2x + \frac{n\pi}{2}\right) + 2^{n-1} \cos\left(4x + \frac{n\pi}{2}\right)$
(c) $\cos\left(4x + \frac{n\pi}{2}\right) + \cos\left(2x + \frac{n\pi}{2}\right)$ (d) None of these

296. If $y = \sin^2 x \sin 2x$ then y_n is equal to

- (a) $2^{n-1} \sin\left(2x + \frac{n\pi}{2}\right) + 4^{n-1} \sin\left(4x + \frac{n\pi}{2}\right)$ (b) $2^{n-1} \sin\left(2x + \frac{n\pi}{2}\right) + 4^{n-1} \sin\left(4x + \frac{n\pi}{2}\right)$

174 Differentiation

(c) $2 \sin\left(2x + \frac{n\pi}{2}\right) + \sin\left(4x + \frac{n\pi}{2}\right)$

(d) None of these

n^{th} Derivative Using Partial Fractions

Basic Level

297. n^{th} derivative of $\frac{1}{3x^2 - 5x + 2}$ is

(a) $(-1)^n n! \left\{ \frac{1}{(x-1)^{n+1}} + \frac{3^{n+1}}{(3x-2)^{n+1}} \right\}$

(b) $n! \left\{ \frac{1}{(x-1)^{n+1}} - \frac{3^{n+1}}{(3x-2)^{n+1}} \right\}$

(c) $(-1)^n n! \left\{ \frac{1}{(x-1)^{n+1}} - \frac{3^{n+1}}{(3x-2)^{n+1}} \right\}$

(d) None of these

298. n^{th} derivative of $\frac{1}{x^2 + 5x + 6}$ is

(a) $(-1)^n n! \left[\frac{1}{(x+2)^{n+1}} + \frac{1}{(x+3)^{n+1}} \right]$ (b) $(-1)^n n! \left[\frac{1}{(x+3)^{n+1}} - \frac{1}{(x+2)^{n+1}} \right]$ (c) $(-1)^n n! \left[\frac{1}{(x+2)^{n+1}} - \frac{1}{(x+3)^{n+1}} \right]$ (d) None of these

Advance Level

299. n^{th} derivative of $\frac{2x+3}{5x+7}$

(a) $\frac{(-1)^n n! 5^{n-1}}{(5x+7)^{n+1}}$

(b) $\frac{(-1)^n n! 5^{n-1}}{(5x+7)^{n-1}}$

(c) $\frac{(-1)^n n! 5^{n+1}}{(5x+7)^{n+1}}$

(d) $\frac{(-1)^n n! 5^{n+1}}{(5x+7)^{n-1}}$

300. n^{th} derivative of $\frac{1}{x^2 - a^2}$ is

(a) $\frac{(-1)^n n!}{2a} [(x-a)^{n+1} - (x+a)^{n+1}]$

(b) $\frac{(-1)^n n!}{2a} [(x+a)^{n+1} - (x-a)^{n+1}]$

(c) $\frac{(-1)^n n!}{2a} \left[\frac{1}{(x-a)^{n+1}} - \frac{1}{(x+a)^{n+1}} \right]$

(d) $\frac{(-1)^n n!}{2a} [(x-a)^{n+1} + (x+a)^{n+1}]$

Differentiation of Determinants

Basic Level

301. If $f(x) = \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix}$, then $f'(x)$ is

(a) x^2

(b) $6x$

(c) $6x^2$

(d) 1

302. If $f(\theta) = \begin{vmatrix} \sec \theta & \tan^2 \theta & 1 \\ \theta \sec x & \tan x & x \\ 1 & \tan x - \tan \theta & 0 \end{vmatrix}$, then $f'(\theta)$ is

- (a) 0 (b) -1 (c) Independent of θ (d) None of these

303. Let f, g, h and k be differentiable in (a, b) and F is defined as $F(x) = \begin{vmatrix} f(x) & g(x) \\ h(x) & k(x) \end{vmatrix}$ for all $x \in (a, b)$ then F' is given by

- (a) $\begin{vmatrix} f & g \\ h & k \end{vmatrix} + \begin{vmatrix} f & g \\ h' & k \end{vmatrix}$ (b) $\begin{vmatrix} f' & g' \\ h & k \end{vmatrix} + \begin{vmatrix} f & g \\ h' & k' \end{vmatrix}$ (c) $\begin{vmatrix} f & g' \\ h & k' \end{vmatrix} + \begin{vmatrix} f' & g \\ h & k \end{vmatrix}$ (d) $\begin{vmatrix} f & g \\ h' & k' \end{vmatrix} + \begin{vmatrix} f' & g \\ h & k \end{vmatrix}$

Advance Level

304. $f(x) = \begin{vmatrix} x^3 & x^2 & 3x^2 \\ 1 & -6 & 4 \\ p & p^2 & p^3 \end{vmatrix}$, here p is a constant, then $\frac{d^3 f(x)}{dx^3}$ is

- (a) Proportional to x^2 (b) Proportional to x (c) Proportional to x^3 (d) A constant

305. If $y = \sin px$ and y_n is the n^{th} derivative of y , then $\begin{vmatrix} y & y_1 & y_2 \\ y_3 & y_4 & y_5 \\ y_6 & y_7 & y_8 \end{vmatrix}$ is equal to [AMU 2002]

- (a) 1 (b) 0 (c) -1 (d) None of these

Differentiation of Integral Functions

Basic Level

306. Let $f(t) = \log(t)$, then $\frac{d}{dx} \left(\int_{x^2}^{x^3} f(t) dt \right)$

- (a) Has a value 0 when $x = 0$ (b) Has a value 0 when $x = 1$ and $x = \frac{4}{9}$
 (c) Has a value $9e^2 - 4$ when $x = e$ (d) Has a differential coefficient $27e - 8$ for $x = e$

307. If $f(x) = \int_0^x t \sin t dt$, then $f'(x) =$

- (a) $x \sin x$ (b) $x \cos x$ (c) $\sin x + \cos x$ (d) $x^2/2$

Advance Level

308. If $F(x) = \frac{1}{x^2} \int_4^x (4t^2 - 2F'(t)) dt$, then $F'(4)$ equals

- (a) $32/9$ (b) $64/3$ (c) $64/9$ (d) None of these

Leibnitz's theorem

Basic Level

176 Differentiation

309. If $y = x \sin x$, then at $x = 0$ the value of y_{15} equal to

- (a) 0 (b) -15 (c) 15! (d) -(15)!

Advance Level

310. If $y = xe^x$ then the value of y_n is

- (a) $(n+1)e^x$ (b) $(x+1)e^x$ (c) $(x+n)e^x$ (d) $(x-n)e^x$

Miscellaneous Problems

Basic Level

311. Given that $d/dx f(x) = f'(x)$. The relationship $f(a+b) = f'(a) + f'(b)$ is valid if $f(x)$ is equal to

- (a) x (b) x^2 (c) x^3 (d) x^4

312. $f(x)$ and $g(x)$ are two differentiable function on $[0, 2]$ such that $f''(x) - g''(x) = 0$, $f'(1) = 2$, $g'(1) = 4$, $f(2) = 3$, $g(2) = 9$, then $f(x) - g(x)$ at $x = 3/2$ is

- (a) 0 (b) 2 (c) 10 (d) -5

313. If $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$, then $\frac{y'}{y} =$

[IIT 1998]

- (a) $\left(\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x}\right)$ (b) $\left(\frac{a}{a+x} + \frac{b}{b+x} + \frac{c}{c+x}\right)$ (c) $\frac{1}{x} \left(\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x}\right)$ (d) $\frac{1}{y} \left(\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x}\right)$

314. If $y = \frac{1}{1+x^{a-b}+x^{a-c}} + \frac{1}{1+x^{b-a}+x^{b-c}} + \frac{1}{1+x^{c-a}+x^{c-b}}$, then $\frac{dy}{dx}$ equals

- (a) $ax^{-1} + bx^{-1} + cx^{-1}$ (b) 0 (c) 1 (d) $a+b+c$

315. Let $f(x)$ be a polynomial function of the second degree. If $f(1) = f(-1)$ and a_1, a_2, a_3 are in A.P. then $f'(a_1), f'(a_2), f'(a_3)$ are in

- (a) A.P. (b) G.P. (c) H.P. (d) None of these

Answer Sheet

Assignment (Basic & Advance Level)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
d	d	c	b	d	c	b	a	b	b	a	b	d	a	a	c	a	d	c	b
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
d	d	c	b	b	d	d	a	c	c	d	c	d	a	b	c	d	c	d	d
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
d	b	a	b	a	d	c	b	b	a	c	b	b	b	d	b	c	c	c	a
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
a	a	c	c	a	a	a	a	c	c	a	a	a	b	a	c	c	a	c	c
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
a	d	c	b	b	b	a	c	c	a	d	a	c	c	b	a	a	c	a	a
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
d	d	c	c	a	b	c	a	b	c	d	b	b	d	a	c	b	c	a	d
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
c	c	a	b	d	a	a	a	a	a	a	c	c	c	a	a	b	a	b	b
141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
b	c	b	b	a	b	c	a	c	b	c	c	c	c	c	c	c	d	a	c
161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
a	a	b	c	b	a	c	c	a	a	a	a	b	a	b	b	a	b	c	a
181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
a	d	a	a	b	b	a	a	a	d	b	b	a	a	a	c	c	a	a	a
201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220
d	c	b	b	d	a	b	c	a	a	c	c	b	d	b	a	b	c	a	b
221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
d	c	d	d	d	b	a	d	a	c	a	c	b	a	c	a	c	b	b	d
241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260
c	b	b	c	a	c	c	d	b	d	b	b	b	d	b	c	a	d	b	b
261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280
a	d	a	b	d	c	b	b	a	a	b	d	b	b	b	c	c	b	d	c
281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300
b	d	a	a	b	b	c	a	c	a	a	a	a	c	c	b	c	c	a	c
301	302	303	304	305	306	307	308	309	310	311	312	313	314	315					
c	c	b	d	b	b	a	a	d	c	b	d	c	b	a					