

Class XI Session 2024-25
Subject - Mathematics
Sample Question Paper - 8

Time Allowed: 3 hours

Maximum Marks: 80

General Instructions:

1. This Question paper contains - five sections A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

Section A

1. $\sin \frac{\pi}{12} = ?$ [1]
a) $\frac{(\sqrt{3}+1)}{2\sqrt{2}}$ b) $\frac{(\sqrt{3}-1)}{2\sqrt{2}}$
c) $\frac{(2\sqrt{3}+1)}{3\sqrt{2}}$ d) $\frac{-(\sqrt{3}-1)}{2\sqrt{2}}$
2. If $n(A) = 10$, $n(B) = 6$ and $n(C) = 5$ for three disjoint sets A, B and C, then $n(A \cup B \cup C) =$ [1]
a) 11 b) 21
c) 1 d) 9
3. The mean of the series x_1, x_2, \dots, x_n is \bar{X} . If x_2 is replaced by λ , then what is the new mean? [1]
a) $\frac{\bar{X}-x_2-\lambda}{n}$ b) $\frac{n\bar{X}-x_2-\lambda}{n}$
c) $\frac{\bar{X}-x_2+\lambda}{n}$ d) $\bar{X} - x_2 + \lambda$
4. If $f(x) = x \sin x$, then $f' \left(\frac{\pi}{2} \right)$ is equal to [1]
a) 1 b) $\frac{1}{2}$
c) -1 d) 0
5. The coordinates of the foot of perpendicular from (0, 0) upon the line $x + y = 2$ are [1]
a) (1, 1) b) (1, -2)
c) (-1, 2) d) (1, 2)
6. The length of the foot of perpendicular drawn from the point P (3, 4, 5) on y-axis is [1]
a) $\sqrt{34}$ b) 10

- c) $\sqrt{113}$ d) $5\sqrt{2}$
7. Mark the correct answer for $3i^{34} + 5i^{27} - 2i^{38} + 5i^{41} = ?$ [1]
 a) 1 b) -1
 c) -4i d) 10i
8. A fair dice is rolled n times. The number of all the possible outcomes is [1]
 a) 6n b) n^6
 c) 6^n d) $6+n$
9. If $y = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$, then $\frac{dy}{dx} =$ [1]
 a) y^2 b) $y + 1$
 c) y d) $y - 1$
10. If $\frac{3\pi}{4} < \alpha < \pi$, then $\sqrt{2 \cot \alpha + \frac{1}{\sin^2 \alpha}}$ is equal to [1]
 a) $-1 + \cot \alpha$ b) $-1 - \cot \alpha$
 c) $1 - \cot \alpha$ d) $1 + \cot \alpha$
11. Each set X_r contains 5 elements and each set Y_r contains 2 elements and $\bigcup_{r=1}^{20} x_r = S = \bigcup_{r=1}^n Y_r$. If each element of S belong to exactly 10 of the X_r 's and to exactly 4 of the Y_r 's, then n is [1]
 a) 10 b) 20
 c) 50 d) 100
12. In the expansion of $(x + a)^n$, if the sum of odd terms be P and the sum of even terms be Q, then $4PQ = ?$ [1]
 a) $(x + a)^n - (x - a)^n$ b) $(x + a)^{2n} - (x - a)^{2n}$
 c) $(x + a)^n + (x - a)^n$ d) $(x + a)^{2n} + (x - a)^{2n}$
13. If $(1 - x + x^2)^n = a_0 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$, then $a_0 + a_2 + a_4 + \dots + a_{2n}$ equals. [1]
 a) $3^n + \frac{1}{2}$ b) $\frac{3^n+1}{2}$
 c) $\frac{3^n-1}{2}$ d) $\frac{1-3^n}{2}$
14. If x is a real number and $|x| < 3$, then [1]
 a) $-3 < x < 3$ b) $x \geq -3$
 c) $x \geq 3$ d) $-3 \leq x \leq 3$
15. Which of the following is a set? [1]
 A. A collection of vowels in English alphabets is a set.
 B. The collection of most talented writers of India is a set.
 C. The collection of most difficult topics in Mathematics is a set.
 D. The collection of good cricket players of India is a set.
- a) B b) D
 c) A d) C
16. If $3 \sin x + 4 \cos x = 5$, then $4 \sin x - 3 \cos x =$ [1]

OR

Using g binomial theorem, expand $\{(x+y)^5 + (x-y)^5\}$ and hence find the value of $\{(\sqrt{2}+1)^5 + (\sqrt{2}-1)^5\}$

29. Evaluate the following limits: $\lim_{x \rightarrow 0} \frac{2x}{\sqrt{a+x} - \sqrt{a-x}}$. [3]

OR

Differentiate e^{ax+b} from first principle.

30. The sum of three numbers a, b, c in A.P. is 18. If a and b are each increased by 4 and c is increased by 36, the new numbers form a G.P. Find a, b, c. [3]

OR

If A.M. and G.M. of roots of a quadratic equation are 8 and 5 respectively then obtain the quadratic equation.

31. Are the $E = \{x : x \in \mathbb{Z}, x^2 \leq 4\}$ and $F = \{x : x \in \mathbb{Z}, x^2 = 4\}$ pairs of equal set? [3]

Section D

32. The mean and standard deviation of 100 observations were calculated as 40 and 5.1, respectively by a student who took by mistake 50 instead of 40 for one observation. What are the correct mean and standard deviation? [5]
33. Find the lengths major and minor axes, coordinates of the vertices, coordinates of the foci, eccentricity, and length of the latus rectum of the ellipse $25x^2 + 4y^2 = 100$. [5]

OR

A visitor with sign board 'DO NOT LITTER' is moving on a circular path in an exhibition. During the movement he stops at points represented by (3, -2) and (-2, 0). Also, centre of the circular path is on the line $2x - y = 3$. What is the equation of the path? What message he wants to give to the public?

34. Solve the following system of linear inequalities [5]

$$\frac{4x}{3} - \frac{9}{4} < x + \frac{3}{4} \text{ and } \frac{7x-1}{3} - \frac{7x+2}{6} > x.$$

35. Prove that: $\cot x + \cot\left(\frac{\pi}{3} + x\right) + \cot\left(\frac{2\pi}{3} + x\right) = 3 \cot 3x$. [5]

OR

If $A + B + C = \pi$, prove that $\frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C} = 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$

Section E

36. Read the following text carefully and answer the questions that follow: [4]

Method to Find the Sets When Cartesian Product is Given

For finding these two sets, we write first element of each ordered pair in first set say A and corresponding second element in second set B (say).

Number of Elements in Cartesian Product of Two Sets

If there are p elements in set A and q elements in set B, then there will be pq elements in $A \times B$ i.e. if $n(A) = p$ and $n(B) = q$, then $n(A \times B) = pq$.

- The Cartesian product $A \times A$ has 9 elements among which are found (-1, 0) and (0, 1). Find the set A and the remaining elements of $A \times A$. (1)
- A and B are two sets given in such a way that $A \times B$ contains 6 elements. If three elements of $A \times B$ are (1, 3), (2, 5) and (3, 3), then find the remaining elements of $A \times B$. (1)
- If the set A has 3 elements and set B has 4 elements, then find the number of elements in $A \times B$. (2)

OR

If $A \times B = \{(a, 1), (b, 3), (a, 3), (b, 1), (a, 2), (b, 2)\}$. Find A and B. (2)

37. Read the following text carefully and answer the questions that follow: [4]

On her vacation, Priyanka visits four cities. Delhi, Lucknow, Agra, Meerut in a random order.



Meerut



New Delhi



Agra



Lucknow

- i. What is the probability that she visits Delhi before Lucknow? (1)
- ii. What is the probability she visit Delhi before Lucknow and Lucknow before Agra? (1)
- iii. What is the probability she visits Delhi first and Lucknow last? (2)

OR

What is the probability she visits Delhi either first or second? (2)

38. Two complex numbers $Z_1 = a + ib$ and $Z_2 = c + id$ are said to be equal, if $a = c$ and $b = d$. [4]

- i. If $(x + iy)(2 - 3i) = 4 + i$ then find the value of (x, y) . (1)
- ii. If $\frac{(1+i)^2}{2-i} = x + iy$, then find the value of $x + y$. (1)
- iii. If $\left(\frac{1-i}{1+i}\right)^{100} = a + ib$, then find the values of a and b . (2)

OR

If $(a - 2, 2b + 1) = (b - 1, a + 2)$, then find the real values of a and b . (2)

Solution

Section A

1.

(b) $\frac{(\sqrt{3}-1)}{2\sqrt{2}}$

Explanation: $\sin \frac{\pi}{12} = \sin \left(\frac{\pi}{4} - \frac{\pi}{6} \right) = \sin \frac{\pi}{4} \cos \frac{\pi}{6} - \cos \frac{\pi}{4} \sin \frac{\pi}{6}$
 $= \left(\frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2} \right) - \left(\frac{1}{\sqrt{2}} \times \frac{1}{2} \right) = \frac{(\sqrt{3}-1)}{2\sqrt{2}}$
2.

(b) 21

Explanation: Since A, B, C are disjoint
 $\therefore n(A \cup B \cup C) = n(A) + n(B) + n(C)$
 $= 10 + 6 + 5 = 21$
3.

(b) $\frac{n\bar{X}-x_2-\lambda}{n}$

Explanation: We know, $\bar{X} = \frac{x_1 + x_2 + \dots + x_n}{n} \Rightarrow x_1 + x_2 + \dots + x_n = n\bar{X}$
 $\Rightarrow x_1 + x_2 + \dots + x_n = n\bar{X} - x_2$
 $\Rightarrow x_1 + x_3 + \dots + x_n + \lambda = n\bar{X} - x_2 + \lambda$
 $\Rightarrow \text{Mean} = \frac{\text{Sum of all values}}{\text{Total number of values}} = \frac{x_1 + x_3 + \dots + x_n + \lambda}{n}$
 $= \frac{n\bar{X}-x_2-\lambda}{n}$
4.

(a) 1

Explanation: $f'(x) = x \cos x + \sin x$
 So, $f'\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \cos \frac{\pi}{2} + \sin \frac{\pi}{2} = 1$
5.

(a) (1, 1)

Explanation: The equation of the line perpendicular to the given line is $x - y + k = 0$
 Since it passes through the origin,
 $0 - 0 + k = 0$
 Therefore, $k = 0$
 Hence the equation of the line is $x - y = 0$
 On solving these two equations we get $x = 1$ and $y = 1$
 The point of intersection of these two lines is (1, 1)
 Hence the coordinates of the foot of the perpendicular is (1, 1)
6.

(a) $\sqrt{34}$

Explanation: Let l be the foot of the perpendicular from point P on the y-axis. Therefore, its x and z-coordinates are zero, i.e., (0, 4, 0). Therefore, the distance between the points (0, 4, 0) and (3, 4, 5) is $\sqrt{9 + 25} = \sqrt{34}$.
7.

(b) -1

Explanation: $3i^{34} + 5i^{27} - 2i^{38} + 5i^{41} = 3 \times (i^4)^8 \times i^2 + 5 \times (i^4)^6 \times i^3 - 2 \times (i^4)^9 \times i^2 + 5 \times (i^4)^{10} \times i$
 $= 3 \times 1 \times (-1) + 5 \times 1 \times (-i) - 2 \times 1 \times (-1) + 5 \times 1 \times i$
 $= -3 - 5i + 2 + 5i = -1$
8.

(c) 6^n

Explanation: Each time there are 6 possibilities, therefore for n times there are 6^n possibilities.
9.

(c) y

Explanation: $y = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

$$\begin{aligned} \text{Differentiating both sides with respect to } x, \text{ we get } \frac{dy}{dx} &= \frac{d}{dx} \left(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) \\ &= \frac{d}{dx} (1) + \frac{d}{dx} \left(\frac{x}{1!} \right) + \frac{d}{dx} \left(\frac{x^2}{2!} \right) + \frac{d}{dx} \left(\frac{x^3}{3!} \right) + \frac{d}{dx} \left(\frac{x^4}{4!} \right) + \dots \\ &= \frac{d}{dx} (1) + \frac{1}{1!} \frac{d}{dx} (x) + \frac{1}{2!} \frac{d}{dx} (x^2) + \frac{1}{3!} \frac{d}{dx} (x^3) + \frac{1}{4!} \frac{d}{dx} (x^4) + \dots \\ &= 0 + \frac{1}{1!} \times 1 + \frac{1}{2!} \times 2x + \frac{1}{3!} \times 3x^2 + \frac{1}{4!} \times 4x^3 + \dots \quad (y = x^2 \Rightarrow \frac{dy}{dx} = nx^{n-1}) \\ &= 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \left[\frac{x}{n!} = \frac{1}{(n-1)!} \right] \\ &= y \\ \therefore \frac{dy}{dx} &= y \end{aligned}$$

10.

(b) $-1 - \cot \alpha$

Explanation: We have:

$$\begin{aligned} &\sqrt{2 \cot \alpha + \frac{1}{\sin^2 \alpha}} \\ &= \sqrt{\frac{2 \cos \alpha}{\sin \alpha} + \frac{1}{\sin^2 \alpha}} \\ &= \sqrt{\frac{2 \sin \alpha \cos \alpha + 1}{\sin^2 \alpha}} \\ &= \sqrt{\frac{2 \sin \alpha \cos \alpha + \sin^2 \alpha + \cos^2 \alpha}{\sin^2 \alpha}} \\ &= \sqrt{\frac{(\sin \alpha + \cos \alpha)^2}{\sin^2 \alpha}} \\ &= \sqrt{(1 + \cot \alpha)^2} \\ &= |1 + \cot \alpha| \\ &= -(1 + \cot \alpha) \quad \left[\text{When } \frac{3\pi}{4} < \alpha < \pi, \cot \alpha < -1 \Rightarrow \cot \alpha + 1 < 0 \right] \\ &= -1 - \cot \alpha \end{aligned}$$

11.

(b) 20

Explanation: The correct answer is (B)

Since, $n(X_r) = 5, \bigcup_{r=1}^{20} X_r = S$, we obtain $n(S) = 100$

But each element of S belong to exactly 10 of the X_r 's

Thus, $\frac{100}{10} = 10$ are the number of distinct elements in S.

Also each element of S belong to exactly 4 of the Y_r 's and each Y_r 's contain 2 elements. If S has n number of Y_r in it.

Then $\frac{2n}{4} = 10$

which gives $n = 20$

12.

(b) $(x+a)^{2n} - (x-a)^{2n}$

Explanation: $P + Q = (x+a)^n$ and $P - Q = (x-a)^n$

$$\Rightarrow 4PQ = (P+Q)^2 - (P-Q)^2 = (x+a)^{2n} - (x-a)^{2n}$$

13.

(b) $\frac{3^{n+1}}{2}$

Explanation: $(1 - x + x^2)^n = a_0 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n} \dots (1)$

Put $x=1$ in (1), we get

$$1 = a_0 + a_1 + a_2 + a_3 + \dots + a_{2n} \dots (2)$$

Put $x=-1$ in (1), we get

$$3^n = a_0 - a_1 + a_2 - a_3 + \dots + a_{2n} \dots (3)$$

Adding (1) and (2), we get

$$3^n + 1 = 2(a_0 + a_2 + a_4 + \dots + a_{2n})$$

$$\text{Thus, } a_0 + a_2 + a_4 + \dots + a_{2n} = \frac{3^n + 1}{2}$$

14. (a) $-3 < x < 3$

Explanation: We have $|x| < a \Leftrightarrow -a < x < a$

15.

(c) A

Explanation: The set is {a, e, i, o, u}

16.

(d) 0

Explanation: $3 \sin x + 4 \cos x = 5$

$$\frac{3}{5} \sin x + \frac{4}{5} \cos x = 1$$

$$\text{Let } \cos \alpha = \frac{3}{5} \text{ and } \sin \alpha = \frac{4}{5}$$

$$\therefore \cos \alpha \sin x + \sin \alpha \cos x = 1$$

$$\Rightarrow \sin(\alpha + x) = \sin \frac{\pi}{2}$$

$$\Rightarrow \alpha + x = \frac{\pi}{2}$$

$$\Rightarrow x = \frac{\pi}{2} - \alpha \dots (i)$$

We have to find the value of $4 \sin x - 3 \cos x$

$$4 \sin\left(\frac{\pi}{2} - \alpha\right) - 3 \cos\left(\frac{\pi}{2} - \alpha\right) \dots \text{From eq. (i)}$$

$$= 4 \cos \alpha - 3 \sin \alpha$$

$$= 4 \times \frac{3}{5} - 3 \times \frac{4}{5} \left(\because \cos \alpha = \frac{3}{5} \text{ and } \sin \alpha = \frac{4}{5} \right)$$

$$0$$

17.

(b) -1

Explanation: Given, $\lim_{x \rightarrow \pi} \frac{\sin x}{x - \pi} = \lim_{x \rightarrow \pi} \frac{\sin(\pi - x)}{-(\pi - x)}$

$$= -1 \left[\because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \text{ and } \pi - x \rightarrow 0 \Rightarrow x \rightarrow \pi \right]$$

18. (a) $2^8 - 2$

Explanation: $({}^7C_0 + {}^7C_1) + ({}^7C_1 + {}^7C_2) + ({}^7C_2 + {}^7C_3) + ({}^7C_3 + {}^7C_4) + ({}^7C_4 + {}^7C_5) + ({}^7C_5 + {}^7C_6) + ({}^7C_6 + {}^7C_7)$

$$= 1 + 2 \times {}^7C_1 + 2 \times {}^7C_2 + 2 \times {}^7C_3 + 2 \times {}^7C_4 + 2 \times {}^7C_5 + 2 \times {}^7C_6 + 1$$

$$= 1 + 2 \times {}^7C_1 + 2 \times {}^7C_2 + 2 \times {}^7C_3 + 2 \times {}^7C_4 + 2 \times {}^7C_5 + 2 \times {}^7C_6 + 1$$

$$= 2 + 2^2 ({}^7C_1 + {}^7C_2 + {}^7C_3)$$

$$= 2 + 2^2 \left(7 + \frac{7}{2} \times 6 + \frac{7}{3} \times \frac{6}{2} \times 5 \right)$$

$$= 2 + 252$$

$$= 254$$

$$= 2^8 - 2$$

19. (a) Both A and R are true and R is the correct explanation of A.

Explanation: Assertion The collection of all natural numbers less than 100, is a well-defined collection. So, it is a set.

20.

(c) A is true but R is false.

Explanation: Assertion Let a be the first term and $r (|r| < 1)$ be the common ratio of the GP.

\therefore The GP is a, ar, ar^2, \dots

According to the question,

$$T_1 + T_2 = 5 \Rightarrow a + ar = 5 \Rightarrow a(1 + r) = 5$$

$$\text{and } T_n = 3(T_{n+1} + T_{n+2} + T_{n+3} + \dots)$$

$$\Rightarrow ar^{n-1} = 3(ar^n + ar^{n+1} + ar^{n+2} + \dots)$$

$$\Rightarrow ar^{n-1} = 3ar^n(1 + r + r^2 + \dots)$$

$$\Rightarrow 1 = 3r\left(\frac{1}{1-r}\right)$$

$$\Rightarrow 1 - r = 3r$$

$$\Rightarrow r = \frac{1}{4}$$

Reason: Given, 3, 6, 9, 12 ...

Here, a = 3, d = 6 - 3 = 3

$$\therefore T_{10} = a + (10 - 1)d$$

$$= 3 + 9 \times 3$$

$$= 3 + 27 = 30$$

Section B

21. According to the question, we can state,

We know the square of a real number is never negative.

Clearly, $f(x)$ takes real values only when $x + 1 \geq 0$

$$= x > -1$$

$$\therefore x \in [-1, \infty)$$

Thus, domain of $f = (-1, \infty)$

Similarly, $g(x)$ takes real values only when $9 - x^2 \geq 0$

$$= 9 > x^2$$

$$= x^2 < 9$$

$$= x^2 - 9 < 0$$

$$= x^2 - 32 < 0$$

$$= (x + 3)(x - 3) < 0$$

$$= x \geq -3 \text{ and } x < 3$$

$$x \in [-3, 3]$$

Thus, domain of $g = [-3, 3]$

i.f + g

We know $(f + g)(x) = f(x) + g(x)$

$$\therefore (f + g)(x) = \sqrt{x + 1} + \sqrt{9 - x^2}$$

Domain of $f + g = \text{Domain of } f \cap \text{Domain of } g$

$$= \text{Domain of } f + g = [-1, \infty) \cap [-3, 3]$$

Domain of $f + g = [-1, 3]$

Thus, $f + g : [-1, 3] \rightarrow \mathbb{R}$ is given by $(f + g)(x) = \sqrt{x + 1} + \sqrt{9 - x^2}$

OR

$$\text{Here we have, } f(x) = \frac{|x-4|}{x-4}$$

We need to find where the function is defined.

To find the domain of the function $f(x)$ we need to equate the denominator of the function to 0

Therefore,

$$x - 4 = 0 \text{ or } x = 4$$

It means that the denominator is zero when $x = 4$

So, the domain of the function is the set of all the real numbers except 4

The domain of the function, $D_{\{f(x)\}} = (-\infty, 4) \cup (4, \infty)$

The numerator is an absolute function of the denominator.

So, for any value of x from the domain set, we always get either +1 or -1 as the output.

So, the range of the function is a set containing -1 and +1

Therefore, the range of the function, $R_{f(x)} = \{-1, 1\}$

22. Given, $\lim_{x \rightarrow 1} \frac{x^3 - 1}{x - 1}$

If we put $x = 1$, then expression $\frac{x^3 - 1}{x - 1}$ becomes the indeterminate form $\frac{0}{0}$. Therefore, $(x - 1)$ is a common factor of $(x^3 - 1)$ and $(x - 1)$.

Factorising the numerator and denominator, we have

$$\lim_{x \rightarrow 1} \frac{x^3 - 1}{x - 1} \left[\frac{0}{0} \text{ form} \right]$$

$$= \lim_{x \rightarrow 1} \frac{(x - 1)(x^2 + x + 1)}{(x - 1)}$$

$$= \lim_{x \rightarrow 1} (x^2 + x + 1) = 1^2 + 1 + 1 = 3$$

23. i. It is given that

$$: P(A) = 0.25, P(A \text{ or } B) = 0.5 \text{ and } P(B) = 0.4$$

To find : $P(A \text{ and } B)$

Formula used : $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$

Substituting the value in the above formula we get,

$$0.5 = 0.25 + 0.4 - P(A \text{ and } B)$$

$$0.5 = 0.65 - P(A \text{ and } B)$$

$$P(A \text{ and } B) = 0.65 - 0.5$$

$$P(A \text{ and } B) = 0.15$$

ii. Given : $P(A) = 0.25$, $P(A \text{ and } B) = 0.15$ (from part (i))

To find : $P(A \text{ and } \bar{B})$

$$\text{Formula used : } P(A \text{ and } \bar{B}) = P(A) - P(A \text{ and } B)$$

Substituting the value in the above formula we get,

$$P(A \text{ and } \bar{B}) = 0.25 - 0.15$$

$$P(A \text{ and } \bar{B}) = 0.10$$

$$P(A \text{ and } \bar{B}) = 0.10$$

OR

Given that $P(A) = 0.5$, $P(B) = 0.4$ and $P(A \cup B) = 0.8$

Applying the general addition rule,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\therefore 0.8 = 0.5 + 0.4 - P(A \cap B)$$

$$\Rightarrow P(A \cap B) = 0.9 - 0.8 = 0.1$$

$$\therefore P(A \cap B) < P(A) \text{ and } P(A \cap B) < P(B)$$

Thus the given probabilities are consistently defined.

24. Let H be the set of students who know Hindi and E be the set of students who know English.

Here $n(H) = 100$, $n(E) = 50$ and $n(H \cap E) = 25$

We know that $n(H \cup E) = n(H) + n(E) - n(H \cap E)$


$$= 100 + 50 - 25 = 125.$$

25. Here, it is given: Point (0,0) and line $7x + 24y = 50$

We have to find: The length of the perpendicular from the origin to the line $7x + 24y = 50$

We know that the length of the perpendicular from P (m,n) to the line $ax + by + c = 0$ is given by,

$$D = \frac{|am+bn+c|}{\sqrt{a^2+b^2}}$$



The given equation of the line is $7x + 24y - 50 = 0$

Here $m = 0$ and $n = 0$, $a = 7$, $b = 24$, $c = -50$

$$D = \frac{|7(0)+24(0)-50|}{\sqrt{7^2+24^2}}$$

$$D = \frac{|0+0-50|}{\sqrt{49+576}} = \frac{|-50|}{\sqrt{625}} = \frac{|-50|}{25} = \frac{50}{25} = 2$$

$$D = 2$$

Therefore, the length of perpendicular from the origin to the line $7x + 24y = 50$ is 2 units.

Section C

26. Given: Six persons are to be arranged in a row.

Assume six seats, now in the first seat, any one of six members can be seated, so the total number of possibilities is 6C_1

Similarly, in the second seat, any one of five members can be seated, so the total number of possibilities is 5C_1

In the third seat, any one of four members can be seated, so the total number of possibilities is 4C_1

In the fourth seat, any one of three members can be seated, so the total number of possibilities is 3C_1

In the fifth seat, any one of two members can be seated, so the total number of possibilities is 2C_1

In the sixth seat, only one remaining person can be seated, so the total number of possibilities is 1C_1

Hence the total number of possible outcomes = ${}^6C_1 \times {}^5C_1 \times {}^4C_1 \times {}^3C_1 \times {}^2C_1 \times {}^1C_1 = 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 720$

27. The general point on yz plane is D(0, y, z).

Consider this point is equidistant to the points A(3, 2, -1), B(1, -1, 0) and C(2, 1, 2).

$\therefore AD = BD$

$$\sqrt{(0-3)^2 + (y-2)^2 + (z+1)^2} = \sqrt{(0-1)^2 + (y+1)^2 + (z-0)^2}$$

Squaring both sides,

$$(0-3)^2 + (y-2)^2 + (z+1)^2 = (0-1)^2 + (y+1)^2 + (z-0)^2$$

$$9 + y^2 - 4y + 4 + z^2 + 2z + 1 = 1 + y^2 + 2y + 1 + z^2$$

$$-6y + 2z + 12 = 0 \dots(1)$$

Also, $AD = CD$

$$\sqrt{(0-3)^2 + (y-2)^2 + (z+1)^2} = \sqrt{(0-2)^2 + (y-1)^2 + (z-2)^2}$$

Squaring both sides,

$$(0-3)^2 + (y-2)^2 + (z+1)^2 = (0-2)^2 + (y-1)^2 + (z-2)^2$$

$$9 + y^2 - 4y + 4 + z^2 + 2z + 1 = 4 + y^2 - 2y + 1 + z^2 - 4z + 4$$

$$-2y + 6z + 5 = 0 \dots(2)$$

By solving equation (1) and (2) we get

$$y = \frac{31}{16} \quad z = \frac{-3}{16}$$

The point which is equidistant to the points A(3, 2, -1), B(1, -1, 0) and C(2, 1, 2) is $(\frac{31}{16}, \frac{-3}{16})$.

$$\begin{aligned} 28. (x+1)^6 + (x-1)^6 &= [{}^6C_0 x^6 + {}^6C_1 x^5 + {}^6C_2 x^4 + {}^6C_3 x^3 + {}^6C_4 x^2 + {}^6C_5 x + {}^6C_6] \\ &+ [{}^6C_0 x^6 + {}^6C_1 x^5(-1) + {}^6C_2 x^4(-1)^2 + {}^6C_3 x^3(-1)^3 + {}^6C_4 x^2(-1)^4 + {}^6C_5 x(-1)^5 + {}^6C_6(-1)^6] \\ &= [x^6 + 6x^5 + 15x^4 + 20x^3 + 15x^2 + 6x + 1] + [x^6 - 6x^5 + 15x^4 - 20x^3 + 15x^2 - 6x + 1] \\ &= 2x^6 + 30x^4 + 30x^2 + 2 \\ &= 2(x^6 + 15x^4 + 15x^2 + 1) \end{aligned}$$

We have

$$\begin{aligned} (x+y)^5 + (x-y)^5 &= 2 [{}^5C_0 x^5 + {}^5C_2 x^3 y^2 + {}^5C_4 x^1 y^4] \\ &= 2 (x^5 + 10x^3 y^2 + 5x y^4) \end{aligned}$$

Putting $x = \sqrt{2}$ and $y = 1$, we get

$$\begin{aligned} (\sqrt{2}+1)^5 + (\sqrt{2}-1)^5 &= 2 \left[(\sqrt{2})^5 + 10(\sqrt{2})^3 + 5\sqrt{2} \right] \\ &= 2 [4\sqrt{2} + 20\sqrt{2} + 5\sqrt{2}] \\ &= 58\sqrt{2} \end{aligned}$$

$$29. \text{ Given: } \lim_{x \rightarrow 0} \frac{2x}{\sqrt{a+x} - \sqrt{a-x}}$$

Rationalizing the given equation,

$$= \lim_{x \rightarrow 0} \frac{2x}{(\sqrt{a+x} - \sqrt{a-x})} \cdot \frac{(\sqrt{a+x} + \sqrt{a-x})}{(\sqrt{a+x} + \sqrt{a-x})}$$

Formula: $(a+b)(a-b) = a^2 - b^2$

$$= \lim_{x \rightarrow 0} \frac{2x(\sqrt{a+x} + \sqrt{a-x})}{a+x-a+x}$$

$$= \lim_{x \rightarrow 0} \frac{2x(\sqrt{a+x} + \sqrt{a-x})}{2x}$$

$$= \lim_{x \rightarrow 0} \frac{(\sqrt{a+x} + \sqrt{a-x})}{1}$$

Now we can see that the indeterminate form is removed, so substituting x as 0

$$\text{Therefore, } \lim_{x \rightarrow 0} \frac{2x}{\sqrt{a+x} - \sqrt{a-x}} = \sqrt{a} + \sqrt{a} = 2\sqrt{a}$$

OR

We need to find derivative of $f(x) = e^{ax+b}$

Derivative of a function $f(x)$ is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \text{ \{where h is a very small positive number\}}$$

\therefore derivative of $f(x) = e^{ax+b}$ is given as

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{e^{a(x+h)+b} - e^{ax+b}}{h}$$

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{e^{ax+b} e^{ah} - e^{ax+b}}{h}$$

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{e^{ax+b} (e^{ah} - 1)}{h}$$

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} e^{ax+b} \times \lim_{h \rightarrow 0} \frac{e^{ah} - 1}{h}$$

As one of the limits $\times \lim_{h \rightarrow 0} \frac{e^{ah} - 1}{h}$ can't be evaluated by directly putting the value of h as it will take $\frac{0}{0}$ form.

So we need to take steps to find its value.

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} e^{ax+b} \times \lim_{h \rightarrow 0} \frac{e^{ah} - 1}{ah} \times a$$

$$\text{Use the formula: } \lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \log_e e = 1$$

$$\Rightarrow f'(x) = e^{ax+b} \times (a)$$

$$\Rightarrow f'(x) = ae^{ax+b}$$

Hence,

$$\text{Derivative of } f(x) = e^{ax+b} = ae^{ax+b}$$

30. Let the first term of the A.P. be a and the common difference be d.

$$\therefore a = a, b = a + d \text{ and } c = a + 2d$$

$$a + b + c = 18$$

$$\Rightarrow 3a + 3d = 18$$

$$\Rightarrow a + d = 6 \dots (i)$$

Now, according to the question, $a + 4$, $a + d + 4$ and $a + 2d + 36$ are in G.P.

$$\therefore (a + d + 4)^2 = (a + 4)(a + 2d + 36)$$

$$\Rightarrow (6 - d + 4)^2 = (6 - d + 4)(6 - d + 2d + 36) \text{ [using (i)]}$$

$$\Rightarrow (10)^2 = (10 - d)(42 + d)$$

$$\Rightarrow 100 = 420 + 10d - 42d - d^2$$

$$\Rightarrow d^2 + 32d - 320 = 0$$

$$\Rightarrow (d + 40)(d - 8) = 0$$

$$\Rightarrow d = 8, -40$$

Now, substituting $d = 8, -40$ in equation (i), we obtain, $a = -2, 46$, respectively.

For $a = -2$ and $d = 8$, we obtain

$$a = -2, b = 6, c = 14$$

And for $a = 46$ and $d = -40$, we obtain

$$a = 46, b = 6, c = -34$$

OR

Let a and b be the roots of required quadratic equation.

$$\text{Then A.M.} = \frac{a+b}{2} = 8$$

$$\Rightarrow$$

$$a + b = 16$$

$$\text{And G.M.} = \sqrt{ab} = 5$$

$$\Rightarrow ab = 25$$

Now, Quadratic equation $x^2 - (\text{Sum of roots})x + (\text{Product of roots}) = 0$

$$\Rightarrow x^2 - (a + b)x + ab = 0$$

$$\Rightarrow x^2 - 16x + 25 = 0$$

Therefore, required equation is $x^2 - 16x + 25 = 0$

31. We know two sets A and B are said to be equal if they have exactly the same elements & we write $A = B$

We have, $E = \{x : x \in Z, x^2 \leq 4\}$

Here, $x \in Z$ and $x^2 \leq 4$

If $x = -2$, then $x^2 = (-2)^2 = 4 = 4$

If $x = -1$, then $x^2 = (-1)^2 = 1 < 4$

If $x = 0$, then $x^2 = (0)^2 = 0 < 4$

If $x = 1$, then $x^2 = (1)^2 = 1 < 4$

If $x = 2$, then $x^2 = (2)^2 = 4 = 4$

Therefore, $E = \{-2, -1, 0, 1, 2\}$

and $F = \{x : x \in Z, x^2 = 4\}$

Here, $x \in Z$ and $x^2 = 4$

If $x = -2$, then $x^2 = (-2)^2 = 4 = 4$

If $x = 2$, then $x^2 = (2)^2 = 4 = 4$

Therefore, $F = \{-2, 2\}$

$\therefore E \neq F$ because the elements in the both the sets are not equal.

Section D

32. We have, $n = 100$, $\bar{x} = 40$ and $\sigma = 5.1$

$$\therefore \bar{x} = \frac{1}{n} \sum x_i$$

$$\Rightarrow \sum x_i = n\bar{x} = 100 \times 40 = 4000$$

\therefore Incorrect $\sum x_i = 4000$

and,

$$\sigma = 5.1$$

$$\Rightarrow \sigma^2 = 26.01$$

$$\Rightarrow \frac{1}{n} \sum x_i^2 - (\text{mean})^2 = 26.01$$

$$\Rightarrow \frac{1}{100} \sum x_i^2 - 1600 = 26.01$$

$$\Rightarrow \sum x_i^2 = 1626.01 \times 100$$

\therefore Incorrect $\sum x_i^2 = 162601$

To correct the $\sum x_i$, we need to subtract the incorrect observation 50 and add correct observation is 40.

We have, incorrect $\sum x_i = 4000$

\therefore Correct $\sum x_i = 4000 - 50 + 40 = 3990$

and,

Similarly, to obtain correct $\sum x_i^2$ we need to subtract 50^2 and add 40^2 to incorrect one.

Incorrect $\sum x_i^2 = 162601$

\therefore Correct $\sum x_i^2 = 162601 - 50^2 + 40^2 = 161701$

Now, Correct mean = $\frac{3990}{100} = 39.90$

Correct variance = $\frac{1}{100} (\text{Correct } \sum x_i^2) - (\text{Correct mean})^2$

$$\Rightarrow \text{Correct variance} = \frac{161701}{100} - \left(\frac{3990}{100}\right)^2$$

$$\Rightarrow \text{Correct variance} = \frac{161701 \times 100 - (3990)^2}{(100)^2}$$

$$\Rightarrow \text{Correct variance} = \frac{16170100 - 15920100}{10000} = 25$$

\therefore Correct standard deviation = $\sqrt{25} = 5$

33. Given that:

$$25x^2 + 4y^2 = 100$$

after divide by 100 to both the sides, we get

$$\frac{25}{100}x^2 + \frac{4}{100}y^2 = 1 \Rightarrow \frac{x^2}{4} + \frac{y^2}{25} = 1 \dots (i)$$

Now, above equation is of the form,

$$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1 \dots (ii)$$

Comparing eq. (i) and (ii), we get

$$a^2 = 25 \text{ and } b^2 = 4 \Rightarrow a = \sqrt{25} \text{ and } b = \sqrt{4} \Rightarrow a = 5 \text{ and } b = 2$$

i. Length of major axes

$$\therefore \text{Length of major axes} = 2a = 2 \times 5 = 10 \text{ units}$$

ii. Length of minor axes

$$\text{Length of minor axes} = 2b = 2 \times 2 = 4 \text{ units}$$

iii. Coordinates of the vertices

$$\therefore \text{Coordinates of the vertices} = (0, a) \text{ and } (0, -a) = (0, 5) \text{ and } (0, -5)$$

iv. Coordinates of the foci

As we know that,

$$\text{Coordinates of foci} = (0, \pm c) \text{ where } c^2 = a^2 - b^2$$

Now

$$c^2 = 25 - 4 \Rightarrow c^2 = 21 \Rightarrow c = \sqrt{21} \dots (iii)$$

$$\therefore \text{Coordinates of foci} = (0, \pm\sqrt{21})$$

v. Eccentricity

$$\text{As we know that, Eccentricity} = \frac{c}{a} \Rightarrow e = \frac{\sqrt{21}}{5}$$

vi. Length of the Latus Rectum

$$\text{As we know that Length of Latus Rectum} = \frac{2b^2}{a} = \frac{2 \times (2)^2}{5} = \frac{8}{5}$$

OR

Let the equation of circle whose centre (- g, - f) be

$$x^2 + y^2 + 2gx + 2fy + c = 0 \dots (i)$$

Since, it passes through points (3, - 2) and (- 2, 0)

$$\therefore (3)^2 + (- 2)^2 + 2g(3) + 2f(- 2) + c = 0$$

$$\text{and } (- 2)^2 + (0)^2 + 2g(- 2) + 2f(0) + c = 0$$

$$\Rightarrow 9 + 4 + 6g - 4f + c = 0$$

$$\text{and } 4 + 0 - 4g + 0 + c = 0$$

$$\Rightarrow 6g - 4f + c = - 13$$

$$\text{and } c = 4g - 4 \dots (ii)$$

$$\therefore 6g - 4f + (4g - 4) = - 13$$

$$\Rightarrow 10g - 4f = - 9 \dots (iii)$$

Also, centre (- g, - f) lies on the line $2x - y = 3$

$$\therefore - 2g + f = 3 \dots (iv)$$

On solving Eqs. (iii) and (iv), we get

$$g = \frac{3}{2} \text{ and } f = 6$$

On putting the values of g and f in Eq. (ii), we get

$$c = 4\left(\frac{3}{2}\right) - 4 = 6 - 4 = 2$$

On putting the values of g, f and c in Eq. (i), we get

$$x^2 + y^2 + 2\left(\frac{3}{2}\right)x + 2(6)y + 2 = 0$$

$$\Rightarrow x^2 + y^2 + 3x + 12y + 2 = 0$$

which is the required equation of the path

The message which he wants to give to the public is 'Keep your place clean'.

$$34. \text{ We have, } \frac{4x}{3} - \frac{9}{4} < x + \frac{3}{4} \dots (i)$$

$$\text{and } \frac{7x-1}{3} - \frac{7x+2}{6} > x \dots (ii)$$

From inequality (i), we get

$$\frac{4x}{3} - \frac{9}{4} < x + \frac{3}{4} \Rightarrow \frac{16x-27}{12} < \frac{4x+3}{4}$$

$$\Rightarrow 16x - 27 < 12x + 9 \text{ [multiplying both sides by 12]}$$


$$\Rightarrow 16x - 27 + 27 < 12x + 9 + 27 \text{ [adding 27 on both sides]}$$

$$\Rightarrow 16x < 12x + 36$$

$$\Rightarrow 16x - 12x < 12x + 36 - 12x \text{ [subtracting 12x from bot sides]}$$

$$\Rightarrow 4x < 36 \Rightarrow x < 9 \text{ [dividing both sides by 4]}$$

Thus, any value of x less than 9 satisfies the inequality. So, the solution of inequality (i) is given by $x \in (-\infty, 9)$



From inequality (ii) we get,

$$\frac{7x-1}{3} - \frac{7x+2}{6} > x \Rightarrow \frac{14x-2-7x-2}{6} > x$$

$$\Rightarrow 7x - 4 > 6x \text{ [multiplying by 6 on both sides]}$$

$$\Rightarrow 7x - 4 + 4 > 6x + 4 \text{ [adding 4 on both sides]}$$


$$\Rightarrow 7x > 6x + 4$$

$$\Rightarrow 7x - 6x > 6x + 4 - 6x \text{ [subtracting 6x from both sides]}$$


$$\therefore x > 4$$

Thus, any value of x greater than 4 satisfies the inequality.

So, the solution set is $x \in (4, \infty)$



The solution set of inequalities (i) and (ii) are represented graphically on number line as given below:



Clearly, the common value of x lie between 4 and 9.

Hence, the solution of the given system is, $4 < x < 9$ i.e., $x \in (4, 9)$

35. We have to prove $\cot x + \cot\left(\frac{\pi}{3} + x\right) + \cot\left(\frac{2\pi}{3} + x\right) = 3 \cot 3x$.

$$\text{LHS} = \cot x + \cot\left(\frac{\pi}{3} + x\right) + \cot\left(\frac{2\pi}{3} + x\right)$$

We know,

$$\cot\left(\frac{2\pi}{3} + x\right) = \cot\left(\pi - \left(\frac{\pi}{3} - x\right)\right) = -\cot\left(\frac{\pi}{3} - x\right) \dots (\text{as } -\cot\theta = \cot(180^\circ - \theta))$$

Hence the above LHS becomes

$$= \cot x + \cot\left(\frac{\pi}{3} + x\right) - \cot\left(\frac{\pi}{3} - x\right)$$

$$= \frac{1}{\tan x} + \frac{1}{\tan\left(\frac{\pi}{3} + x\right)} - \frac{1}{\tan\left(\frac{\pi}{3} - x\right)}$$

$$= \frac{1}{\tan x} + \left(\frac{1 - \tan x \tan \frac{\pi}{3}}{\tan \frac{\pi}{3} + \tan x}\right) - \left(\frac{1 + \tan x \tan \frac{\pi}{3}}{\tan \frac{\pi}{3} - \tan x}\right) \dots \left[\because \tan(A + B) = \left(\frac{\tan A + \tan B}{1 - \tan A \tan B}\right) \text{ and } \tan(A - B) = \left(\frac{\tan A - \tan B}{1 + \tan A \tan B}\right)\right]$$

$$= \frac{1}{\tan x} + \left(\frac{1 - \sqrt{3} \tan x}{\sqrt{3} + \tan x}\right) - \left(\frac{1 + \sqrt{3} \tan x}{\sqrt{3} - \tan x}\right)$$

$$= \frac{1}{\tan x} + \left(\frac{(1 - \sqrt{3} \tan x)(\sqrt{3} - \tan x) - (1 + \sqrt{3} \tan x)(\sqrt{3} + \tan x)}{(\sqrt{3} + \tan x)(\sqrt{3} - \tan x)}\right)$$

$$= \frac{1}{\tan x} + \left(\frac{(\sqrt{3} - \tan x - 3 \tan x + \sqrt{3} \tan^2 x) - (\sqrt{3} + 3 \tan x + \tan x + \sqrt{3} \tan^2 x)}{(3 - \tan^2 x)}\right)$$

$$= \frac{1}{\tan x} + \left(\frac{(0 - 4 \tan x - 4 \tan x + 0)}{(3 - \tan^2 x)}\right)$$

$$= \frac{1}{\tan x} - \left(\frac{8 \tan x}{(3 - \tan^2 x)}\right)$$

$$= \left(\frac{(3 - \tan^2 x) - 8 \tan^2 x}{\tan x (3 - \tan^2 x)}\right) = \left(\frac{(3 - \tan^2 x) - 8 \tan^2 x}{\tan x (3 - \tan^2 x)}\right)$$

$$= 3 \left(\frac{1 - 3 \tan^2 x}{(3 \tan x - \tan^3 x)}\right)$$

$$= 3 \times \frac{1}{\tan 3x} \dots (\text{as } \tan 3x = \frac{3 \tan x - \tan^3 x}{1 - 3 \tan^2 x})$$

$$= \cot 3x$$

$$\text{LHS} = \text{RHS}$$

Hence proved.

OR

Here it is given that, $A + B + C = \pi$

We need to prove that, $\frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C} = 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$

Proof: Taking LHS, we have,

$$L. H. S = \frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C}$$

Where, $\sin 2A + \sin 2B + \sin 2C = 2\sin A \cos A + 2\sin(B + C)\cos(B - C)$

[By using, $\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$

and $\sin 2A = 2\sin A \cos A$]

Since $A + B + C = \pi$

$$\Rightarrow B + C = 180 - A$$

$$\therefore \sin 2A + \sin 2B + \sin 2C = 2\sin A \cos A + 2\sin(\pi - A)\cos(B - C)$$

$$= 2\sin A \cos A + 2\sin A \cos(B - C)$$

$$= 2\sin A \{\cos A + \cos(B - C)\}$$

$$(\text{but } \cos A = \cos \{ 180 - (B + C) \} = -\cos(B + C))$$

And now using

$$\cos A - \cos B = 2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{-A+B}{2}\right)$$

$$\text{So, } \sin 2A + \sin 2B + \sin 2C = 2\sin A \{2\sin B \sin C\}$$

$$= 4\sin A \sin B \sin C$$

$$= 32 \sin \frac{A}{2} \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{B}{2} \sin \frac{C}{2} \cos \frac{C}{2}$$

Now, take denominator we have

$$\sin A + \sin B + \sin C = \sin A + \left\{ 2 \sin\left(\frac{B+C}{2}\right) \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= \sin A + \left\{ 2 \sin\left(\frac{\pi-A}{2}\right) \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= \sin A + \left\{ 2 \cos\left(\frac{A}{2}\right) \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= 2 \sin \frac{A}{2} \cos \frac{A}{2} + \left\{ 2 \cos\left(\frac{A}{2}\right) \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= 2 \cos \frac{A}{2} \left\{ \sin \frac{A}{2} + \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= 2 \cos \frac{A}{2} \left\{ \cos\left(\frac{B+C}{2}\right) + \cos\left(\frac{B-C}{2}\right) \right\}$$

$$= 2 \cos \frac{A}{2} \left\{ 2 \cos\left(\frac{B}{2}\right) \cos\left(\frac{C}{2}\right) \right\}$$

$$= 4 \cos \frac{A}{2} \cos\left(\frac{B}{2}\right) \cos\left(\frac{C}{2}\right)$$

Therefore,

$$L. H. S = \frac{\sin 2A + \sin 2B + \sin 2C}{\sin A + \sin B + \sin C} = \frac{32 \sin \frac{A}{2} \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{B}{2} \sin \frac{C}{2} \cos \frac{C}{2}}{4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}}$$

$$= 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$$

= R.H.S

Section E

36. i. $n(A \times A) = 9$

$$\Rightarrow n(A) \subset n(A) = 9 \Rightarrow n(A) = 3$$

$$(-1, 0) \in A \times A \Rightarrow -1 \in A, 0 \in A$$

$$(0, 1) \in A \times A \Rightarrow 0 \in A, 1 \in A$$

$$\Rightarrow -1, 0, 1 \in A$$

$$\text{Also, } n(A) = 3 \Rightarrow A = \{-1, 0, 1\}$$

$$\text{Hence, } A = \{-1, 0, 1\}$$

$$\text{Also, } A \times A = \{-1, 0, 1\} \times \{-1, 0, 1\}$$

$$= \{(-1, -1), (-1, 0), (-1, 1), (0, -1), (0, 0), (0, 1), (1, -1), (1, 0), (1, 1)\}$$

Hence, the remaining elements of $A \times A$ are

$$(-1, -1), (-1, 1), (0, -1), (0, 0), (1, -1), (1, 0) \text{ and } (1, 1).$$

ii. Given, $(A \times B) = 6$ and $(A \times B) = \{(1, 3), (2, 5), (3, 3)\}$

We know that Cartesian product of set $A = \{a, b\}$ & $B = \{c, d\}$ is $A \times B = \{(a, c), (a, d), (b, c), (b, d)\}$

Therefore, $A = \{1, 2, 3\}$ & $B = \{3, 5\}$

$$\Rightarrow A \times B = \{(1, 3), (1, 5), (2, 3), (2, 5), (3, 3), (3, 5)\}$$

Thus, remaining elements are $A \times B = \{(1, 5), (2, 3), (3, 5)\}$

iii. If the set A has 3 elements and set B has 4 elements, then the number of elements in $A \times B = 12$

OR

Clearly, A is the set of all first entries in ordered pairs in $A \times B$ and B is the set of all second entries in ordered pairs in $A \times B$

$\therefore A = \{a, b\}$ and $B = \{1, 2, 3\}$

37. i. Let the Priyanka visits four cities Delhi, Lucknow, Agra, Meerut are respectively A, B, C and D. Number of way's in which Priyanka can visit four cities A, B, C and D is 4! i.e. 24

$\therefore n(S) = 24$

Clearly, sample space for this experiment is

$$S = \left\{ \begin{array}{l} ABCD, ABDC, ACBD, ACDB, ADBC, ADCB \\ BACD, BADC, BCAD, BCDA, BDAC, BDC A \\ CABD, CADB, CBAD, CBDA, CDAB, CDBA, \\ DABC, DACB, DCAB, DCBA, DBAC, DBC A \end{array} \right\}$$

Let E_1 be the event that Priyanka visits A before B.

Then,

$E_1 = \{ABCD, ABDC, ACBD, ACDB, ADBC, ADCB, CABD, CADB, CDAB, DABC, DACB, DCAB\}$

$\Rightarrow n(E_1) = 12$

$\therefore P(\text{she visits A before B}) = P(E_1) = \frac{n(E_1)}{n(S)} = \frac{12}{24} = \frac{1}{2}$

- ii. Let the Priyanka visits four cities Delhi, Lucknow, Agra, Meerut are respectively A, B, C and D. Number of way's in which Priyanka can visit four cities A, B, C and D is 4! i.e. 24

$\therefore n(S) = 24$

Clearly, sample space for this experiment is

$$S = \left\{ \begin{array}{l} ABCD, ABDC, ACBD, ACDB, ADBC, ADCB \\ BACD, BADC, BCAD, BCDA, BDAC, BDC A \\ CABD, CADB, CBAD, CBDA, CDAB, CDBA, \\ DABC, DACB, DCAB, DCBA, DBAC, DBC A \end{array} \right\}$$

$E_1 = \{ABCD, ABDC, ACBD, ACDB, ADBC, ADCB, CABD, CADB, CDAB, DABC, DACB, DCAB\}$

$\Rightarrow n(E_1) = 12$

$\therefore P(\text{she visits A before B}) = P(E_1) = \frac{n(E_1)}{n(S)} = \frac{12}{24} = \frac{1}{2}$

- iii. Let the Priyanka visits four cities Delhi, Lucknow, Agra, Meerut are respectively A, B, C and D. Number of way's in which Priyanka can visit four cities A, B, C and D is 4! i.e. 24

$\therefore n(S) = 24$

Clearly, sample space for this experiment is

$$S = \left\{ \begin{array}{l} ABCD, ABDC, ACBD, ACDB, ADBC, ADCB \\ BACD, BADC, BCAD, BCDA, BDAC, BDC A \\ CABD, CADB, CBAD, CBDA, CDAB, CDBA, \\ DABC, DACB, DCAB, DCBA, DBAC, DBC A \end{array} \right\}$$

Let E_3 be the event that she visits A first and B last.

Then,

$E_3 = \{ACDB, ADCB\}$

$n(E_3) = 2$

$\therefore P(\text{she visits A first and B last}) = P(E_3)$

$= \frac{n(E_3)}{n(S)} = \frac{2}{24} = \frac{1}{12}$

OR

Let the Priyanka visits four cities Delhi, Lucknow, Agra, Meerut are respectively A, B, C and D. Number of way's in which Priyanka can visit four cities A, B, C and D is 4! i.e. 24

$\therefore n(S) = 24$

Clearly, sample space for this experiment is

$$S = \left\{ \begin{array}{l} ABCD, ABDC, ACBD, ACDB, ADBC, ADCB \\ BACD, BADC, BCAD, BCDA, BDAC, BDC A \\ CABD, CADB, CBAD, CBDA, CDAB, CDBA, \\ DABC, DACB, DCAB, DCBA, DBAC, DBC A \end{array} \right\}$$

Let E_4 be the event that she visits A either first or second. Then,

$$E_4 = \{ABCD, ABDC, ACBD, ACDB, ADBC, ADCB, BACD, BADC, CABD, CADB, DABC, DACB\}$$

$$\Rightarrow n(E_4) = 12$$

Hence, P(she visits A either first or second)

$$= P(E_4) = \frac{n(E_4)}{n(S)} = \frac{12}{24} = \frac{1}{2}$$

38. i. $(x + iy)(2 - 3i) = 4 + i$

$$2x - (3x)i + (2y)i - 3yi^2 = 4 + i$$

$$2x + 3y + (2y - 3x)i = 4 + i$$

Comparing the real & imaginary parts,

$$2x + 3y = 4 \dots(i)$$

$$2y - 3x = 1 \dots(ii)$$

Solving eq (i) & eq (ii), $4x + 6y = 8$

$$-9x + 6y = 3$$

$$13x = 5 \Rightarrow x = \frac{5}{13}$$

$$y = \frac{14}{13}$$

$$\therefore (x, y) = \left(\frac{5}{13}, \frac{14}{13}\right)$$

ii. $x + iy = \frac{(1+i)^2}{2-i}$

$$x + iy = \frac{(1+i)^2}{2-i} = \frac{1+2i+i^2}{2-i} = \frac{2i}{2-i} = \frac{2i(2+i)}{(2-i)(2+i)} = \frac{4i+2i^2}{4-i^2}$$

$$= \frac{4i-2}{4+1} = \frac{-2}{5} + \frac{4i}{5}$$

$$\Rightarrow x = \frac{-2}{5}, y = \frac{4}{5} \Rightarrow x + y = \frac{-2}{5} + \frac{4}{5} = \frac{2}{5}$$

iii. We have $\left(\frac{1-i}{1+i}\right)^{100} = a + bi$

$$\Rightarrow \left(\frac{1-i}{1+i} \times \frac{1-i}{1-i}\right)^{100} = a + bi$$

$$\Rightarrow \left(\frac{1+i^2-2i}{1-i^2}\right)^{100} = a + bi$$

$$\Rightarrow \left(\frac{1-1-2i}{1+1}\right)^{100} = a + bi$$

$$\Rightarrow \left(\frac{-2i}{2}\right)^{100} = a + bi$$

$$\Rightarrow (-i)^{100} = a + bi$$

$$\Rightarrow i^{100} = a + bi$$

$$\Rightarrow (i^4)^{25} = a + bi$$

$$\Rightarrow (1)^{25} = a + bi$$

$$\Rightarrow 1 = a + bi$$

$$\Rightarrow 1 + 0i = a + bi$$

Comparing the real and imaginary parts,

We have $a = 1, b = 0$

Hence $(a, b) = (1, 0)$

OR

Given, $(a - 2, 2b + 1) = (b - 1, a + 2)$

Comparing x coordinates of both the sides, we get,

$$a - 2 = b - 1$$

$$\therefore a - b = 1 \dots(1)$$

Comparing y coordinates of both the sides, we get,

$$2b + 1 = a + 2$$

$$\therefore a - 2b = -1 \dots(2)$$

Subtracting equation (2) from (1), we get,

$$(a - a) + (-b - (-2b)) = 1 - (-1)$$

$$\therefore (-b + 2b) = 1 + 1$$

$$\therefore b = 2$$

Put this value in equation (1), we get,

$$a - 2 = 1$$

$$\therefore a = 3$$