## Chemistry

# CISCE

# Academic Year: 2023-2024 (English Medium) Date & Time: 11th March 2024, 11:00 am

## Duration: 2h

## Marks: 100

- 1. Answers to this Paper must be written on the paper provided separately.
- 2. You will not be allowed to write during the first 15 minutes.
- 3. This time is to be spent in reading the question paper.
- 4. The time given at the head of this Paper is the time allowed for writing the answers.
- 5. Section A is compulsory. Attempt any four questions from Section B.
- 6. The intended marks for questions or parts of questions are given in brackects[].

SECTION-A (40 Marks) (Attempt all questions from this Section.)

Q1. Choose the correct answers to the questions from the given options. (Do not copy the questions, write the correct answers only.)

- **1.1.** The unsaturated hydrocarbons undergo \_\_\_\_\_\_.
  - 1. A substitution reaction
  - 2. An oxidation reaction
  - 3. An addition reaction
  - 4. None of the above
  - 5. Redox reaction

## Solution

The unsaturated hydrocarbons undergo <u>an addition reaction</u>.

## **Explanation:**

Because double and triple bonds in unsaturated hydrocarbons are broken down into single bonds, they are subject to addition reactions.

**1.2.** In the 2<sup>nd</sup> period Neon has maximum Ionization Potential because \_\_\_\_\_.

- 1. It has unstable electronic configuration.
- 2. It easily accepts electrons.
- 3. It easily loses electrons.
- 4. The outer most shell is completely filled.

## Solution

In the 2<sup>nd</sup> period Neon has maximum Ionization Potential because <u>the outer most</u> <u>shell is completely filled</u>.

## **Explanation:**

Neon's valence shell is completely filled, making it extremely stable and requiring more energy to remove an electron, giving it the highest ionisation potential in the second period.

**1.3.** Copper, zinc, and Tin are the metals alloyed to form \_\_\_\_\_.

- 1. Duralumin
- 2. Brass
- 3. Bronze
- 4. Solder

## Solution

Copper, zinc, and Tin are the metals alloyed to form **bronze**.

## Explanation:

Duralumin consists of 90% Al and 4% Cu, while bronze is an alloy with 80% Cu, 4% Zn, and 16% Sn. Cu and Zn combine to form brass. Pb and Sn are alloyed using solder.

**1.4.** The metal hydroxide which reacts with both acids and alkalis to form salt and water is \_\_\_\_\_.

1. Calcium hydroxide

- 2. Magnesium hydroxide
- 3. Aluminium hydroxide
- 4. Ferric hydroxide

## Solution

The metal hydroxide which reacts with both acids and alkalis to form salt and water is **aluminium hydroxide**.

## **Explanation:**

Because Al (OH)<sub>3</sub> is amphoteric in nature, it can generate salt and water as well as behave as a base with a strong acid.

**1.5.** Reaction of an alcohol with a carboxylic acid in the presence of concentrated  $H_2SO_4$  is termed as \_\_\_\_\_.

- 1. Halogenation
- 2. Esterification
- 3. Hydrogenation
- 4. Dehydrohalogenation

## Solution

Reaction of an alcohol with a carboxylic acid in the presence of concentrated H<sub>2</sub>SO<sub>4</sub> is termed as <u>esterification</u>.

## **Explanation:**

The process that produces an ester when an alcohol reacts with carboxylic acid in the presence of  $H_2SO_4$  is known as an esterification reaction.

$$\begin{array}{c} CH_{3}CH_{2}OH + \underset{Acetic \ acid}{HOOC} - CH_{3} \longrightarrow \underset{Ethyl \ Ethanoate}{CH_{3}CH_{2}} - O - C - CH_{3} + H_{2}O \\ & (ester) \end{array}$$

**1.6.** Conversion of ethanol to ethene by the action of concentrated sulphuric acid is an example of \_\_\_\_\_\_.

1. Dehydration

- 2. Dehydrogenation
- 3. Dehydrohalogenation
- 4. Hydrolysis

### Solution

Conversion of ethanol to ethene by the action of concentrated sulphuric acid is an example of <u>dehydration</u>.

### **Explanation:**

Conc. H<sub>2</sub>SO<sub>4</sub> is a useful dehydrator since it turns alcohol into an alkene by removing the water molecule.

$$\begin{array}{c|c} \mathbf{H} & \mathbf{OH} \\ | & | \\ \mathbf{H} - \mathbf{C} - \mathbf{C} - \mathbf{H} \xrightarrow{\mathbf{Conc} \cdot}_{\mathbf{H}_2 \mathbf{SO}_4} \mathbf{H} - \mathbf{C} = \mathbf{C} - \mathbf{H} + \mathbf{H}_2 \mathbf{O} \\ \\ | & | & | \\ \mathbf{H} & \mathbf{H} \\ \mathbf{H} & \mathbf{H} \\ \mathbf{E} \text{thanol} & \mathbf{H} \\ \end{array}$$

**1.7.** The oxidizing agent in the equation  $S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O$  is \_\_\_\_\_.

- 1. Sulphur
- 2. Sulphuric acid
- 3. Sulphur dioxide
- 4. Water

## Solution

The oxidizing agent in the equation  $S + 2H_2SO_4 \rightarrow 3SO_2 + 2H_2O$  is <u>sulphuric acid</u>.

## **Explanation:**

The reaction is as follows:

 $S + 2H_2SO_4 \text{ (conc.)} \rightarrow 3SO_2 + 2H_2O$ 

The oxidizing agent in this process is concentrated  $H_2SO_4$ . While being converted to  $SO_2$ , it oxidizes sulphur (S) to produce sulphur dioxide  $SO_2$ .

**1.8.** Electron Affinity is maximum in \_\_\_\_\_.

- 1. Mg
- 2. Ar
- 3. Li
- 4. Br

#### Solution

Electron Affinity is maximum in **Br**.

## **Explanation:**

From left to right in the periodic table, electron affinity generally increases because atoms become more prone or unstable to gaining an electron in order to achieve a stable electronic configuration, such as that of noble gases. Because they need one electron to complete their octet, halogens such as bromine (Br) have an excessively high electron affinity. Metals like magnesium (Mg) and lithium (Li) have lesser propensities to lose electrons than noble gases like argon (Ar), which have entire outer shells.

**1.9.** The compound that is not a constituent of the electrolytic mixture used in Hall-Heroult's process is \_\_\_\_\_.

- 1. Al<sub>2</sub>O<sub>3</sub>
- 2. NaAlO<sub>2</sub>
- 3. Na<sub>3</sub>AlF<sub>6</sub>
- 4. CaF<sub>2</sub>

## Solution

The compound that is not a constituent of the electrolytic mixture used in Hall-Heroult's process is <u>NaAlO<sub>2</sub></u>.

## **Explanation:**

Aluminum from aluminum oxide is extracted using the Hall-Heroult technique (alumina, Al<sub>2</sub>O<sub>3</sub>). In order to lower the melting point of alumina and increase the conductivity of the solution, an electrolytic mixture made primarily of molten

alumina, a mixture of cryolite (Na<sub>3</sub>AlF<sub>6</sub>), and additional fluxes such as calcium fluoride (CaF<sub>2</sub>) is employed in this process.

**1.10.** On passing ammonia gas over heated copper oxide for some time, a reddishbrown residue is left behind. What property of ammonia is demonstrated here?

- 1. Basic property
- 2. Oxidising property
- 3. Reducing property
- 4. Acidic property

## Solution

Reducing property

## **Explanation:**

 ${
m CuO}$  functions as a reducing agent because it is reduced to  ${
m Cu}$  (reddish brown ppt.) in the presence of ammonia.

$$3\,\mathrm{CuO} + 2\,\mathrm{NH}_3 \longrightarrow 3\,\mathrm{Cu} + \mathrm{N}_2 + 3\,\mathrm{H}_2\mathrm{O}$$

**1.11.** Rotten egg smell is due to the liberation of \_\_\_\_\_.

- 1. HCI gas
- $2. H_2S gas$
- 3.  $Cl_2$  gas
- 4. SO<sub>2</sub> gas

## Solution

Rotten egg smell is due to the liberation of  $H_2S$  gas.

## Explanation:

Hydrogen sulfide (H<sub>2</sub>S) release is frequently the cause of the stench of rotting eggs. Hydrogen sulphide is a colorless gas with a foul odour reminiscent of rotten eggs. As they break down organic waste, microorganisms naturally create it. 1.12. Ammonia gas is collected by downward displacement of air since ammonia is

- 1. Very slightly soluble in water
- 2. Heavier than air
- 3. Lighter than air
- 4. Insoluble in water

### Solution

Ammonia gas is collected by downward displacement of air since ammonia is **lighter than air**.

#### **Explanation:**

Ammonia gas is collected in an inverted gas jar by displacing air downward due to its lighter density. Ammonia is very soluble and cannot be collected over water.

1.13. Which of the following would occupy 22.4 litres at S.T.P.?

- 1. 32 g of oxygen gas
- 2. 2 moles of hydrogen gas
- 3.  $6.022 \times 10^{23}$  molecules of ammonia

[Atomic weights: O = 16, H = 1, N = 14]

- 1. 1 and 2
- 2. 1 and 3
- 3. 2 and 3
- 4. 1, 2 and 3

## Solution

1 and 3

## **Explanation:**

Gram molecular = 1 mole

=  $6.022 \times 10^{23}$  molecules

= 22.4 L

 $\therefore$  Molecular Mass of O<sub>2</sub> gas = 32 g

```
= 6.022 \times 10^{23} molecules
```

= 22.4 L

Similary,  $6.022 \times 10^{23}$  molecule of NH<sub>3</sub> = 22.4 L

Since 1 mole of any gas occupies 22.4 litres at STP, 2 moles of hydrogen gas would occupy  $2 \times 22.4 = 44.8$  litres, which is more than 22.4 litres.

**1.14.** In the molecule of water, oxygen atom has \_\_\_\_\_.

- 1. One shared pair of electrons
- 2. Three shared pairs of electrons
- 3. Two lone pairs of electrons
- 4. One lone pair of electrons

#### Solution

In the molecule of water, oxygen atom has two lone pairs of electrons.

#### **Explanation:**

A water molecule has 2H atoms and 10 atoms. O has an electrical configuration of 2, 6 and requires 2 electrons to complete its octet.

 $\therefore$  it shares  $2e^{-}$  with  $2H^{-}$  atoms



**1.15.** A mineral from which the metal can be extracted economically and conveniently is known as \_\_\_\_\_.

- 1. Matrix
- 2. Ore
- 3. Flux
- 4. Alloy

### Solution

A mineral from which the metal can be extracted economically and conveniently is known as <u>ore</u>.

### **Explanation:**

An ore is a mineral that allows for profitable metal extraction.

## Q2.

**2.1.** The following sketch represents the electroplating of an Iron cup with Nickel metal.

Study the diagram and answer the following questions:



- a. During electroplating, the iron cup is placed at the cathode. Why?
- b. Name the ion that must be present in the electrolyte.
- c. State one condition that is necessary to ensure that the deposit is smooth, firm and even.
- d. Write the reaction taking place at the cathode.
- e. What change would you observe at the anode?

## Solution

- a. In electroplating, the iron cup is put at the cathode because the cathode is a negative terminal that attracts metals that are positively charged. This leads to the reduction and formation of metal ions.
- b. The electrolyte used is a water-based solution of NiSO<sub>4</sub>, so the ions formed are Ni<sup>2+</sup>, H<sup>+</sup>, SO<sub>4</sub><sup>2-</sup>, OH<sup>-</sup>.
- c. To ensure smooth deposition, current should be passed slowly and over a longer period of time.

- d. Cathode:  $Ni^{2+} + 2e^- \rightarrow Ni$  (Reduction)
- e. The anode, a Ni plate, is etched after the Ni ions finish in the electrolyte.

**2.2.** Match the Column A with Column B.

Column A		Column B	
(a)	Water	1.	Lithium
(b)	Alkali metal	2.	lodine
(c)	Halogen	3.	Covalent compound
(d)	Calcium oxide	4.	Acetic acid
(e)	Weak acid	5.	lonic compound
		6.	Sulphuric acid

### Solution

Column A		Column B	
(a)	Water	3.	Covalent compound
(b)	Alkali metal	1.	Lithium
(c)	Halogen	2.	Iodine
(d)	Calcium oxide	5.	Ionic compound
(e)	Weak acid	4.	Acetic acid

**2.3.** Complete the following sentences by choosing the correct answer from the brackets:

**2.3.** (a) The salt prepared by the method of direct combination is \_\_\_\_\_.

- 1. Iron (II) chloride (FeCl<sub>2</sub>)
- 2. Iron (III) chloride (FeCl<sub>3</sub>)

#### Solution

The salt prepared by the method of direct combination is Iron (III) chloride (FeCl<sub>3</sub>).

2.3. (b) The metallic oxide which can be reduced by using common reducing agents is

1. Fe<sub>2</sub>O<sub>3</sub>

2.  $AI_2O_3$ 

#### Solution

The metallic oxide which can be reduced by using common reducing agents is  $Fe_2O_3$ .

2.3. (c) The metal nitrate which on thermal decomposition forms a black residue is

- 1. zinc nitrate
- 2. copper nitrate

### Solution

The metal nitrate which on thermal decomposition forms a black residue is <u>copper</u> <u>nitrate</u>.

**2.3.** (d) During the electrolysis of copper sulphate solution, if \_\_\_\_\_\_ is used as electrodes, the colour of the electrolyte does not fade.

- 1. copper
- 2. platinum

## Solution

During the electrolysis of copper sulphate solution, if <u>copper</u> is used as electrodes, the colour of the electrolyte does not fade.

**2.3.** (e) The process of heating the concentrated ore in a limited supply or absence of air is \_\_\_\_\_.

- 1. Roasting
- 2. Calcination

## Solution

The process of heating the concentrated ore in a limited supply or absence of air is **calcination**.

**2.4.** (a) State the term for the following:

The group obtained by removing one hydrogen atom from the parent alkane.

## Solution

Alkyl group

**2.4.** (b) State the term for the following:

Two metal plates or wires through which the current enters and leaves the electrolytic cell.

## Solution

## Electrodes

2.4. (c) State the term for the following:

The amount of substance which contains the same number of units as the number of atoms in carbon-12.

## Solution

## Mole

2.4. (d) State the term for the following:

The tendency of an atom to pull a shared pair of electrons towards itself in a compound.

## Solution

Electronegativity

2.4. (e) State the term for the following:

The formula which represents the simplest ratio between the atoms of elements present in a compound.

## Solution

Empirical formula

**2.5.** (a)

**1.** Give the IUPAC name of the organic compound represented by the structural formula given below:

 $\begin{array}{cccccccccc} H & Cl & H & H & H \\ & | & | & | & | & | \\ H - C - C - C - C - C - C - H \\ & | & | & | & | \\ H & H & Cl & H & H \end{array}$ 

## Solution

The IUPAC name of the given organic compound is 2, 3-Dichloropentane.

**2.** Give the IUPAC name of the organic compound represented by the structural formula given below:

$$\begin{array}{cccccc} H & H & O \\ & | & | & || \\ H - C - C - C - C - OH \\ & | & | \\ H & H \end{array}$$

#### Solution

The IUPAC name of the given organic compound is propan-1-oic acid.

**2.5.** (b)

**1.** Draw the structural diagram for the following organic compound:

3-methyl pentane

#### Solution

**2.** Write the structural formula of propyne.

Draw the structural diagram for the following organic compound:

## Solution

The structural formula of propyne is as follows:

$$\begin{array}{c} \mathrm{H} \\ | \\ \mathrm{H} - \mathrm{C} - \mathrm{C} \equiv \mathrm{C} - \mathrm{H} \\ | \\ \mathrm{H} \end{array}$$

**3.** Give the structural formula of the following organic compound:

Methanal

## Solution

$$H - C = O$$
  
|  
H

SECTION-B (40 Marks) (Attempt any four questions from this Section.)

Q3.

**3.1.** (a) Rewrite the following statement by adding the correct word, as shown in the example:

## Example:

Given Statement: Ammonia changes moist red litmus to blue.

Correct Statement: <u>Aqueous</u> ammonia changes moist red litmus to blue.

Sulphuric acid acts as a dehydrating agent.

## Solution

Concentrated sulphuric acid acts as a dehydrating agent.

### **Explanation:**

Conc.  $H_2SO_4$  has a strong affinity for water and so reduces it from the molecule, whereas dilute acid dissolves in water.

**3.1.** (b) Rewrite the following statement by adding the correct word, as shown in the example:

#### Example:

Given Statement: Ammonia changes moist red litmus to blue.

Correct Statement: <u>Aqueous</u> ammonia changes moist red litmus to blue.

Ammonia reacts with chlorine to give ammonium chloride and nitrogen.

### Solution

Excess ammonia reacts with chlorine to give ammonium chloride and nitrogen.

## **Explanation:**

Ammonia reacts in two ways:

1. If ammonia is in excess,

 $\underset{(\mathrm{Excess})}{\mathrm{NH}_3} + \mathrm{Cl}_2 \longrightarrow \underset{\substack{\mathrm{Ammonium}\\\mathrm{chloride}}}{\mathrm{NH}_4\mathrm{Cl}} + \underset{\mathrm{Nitrogen}}{\mathrm{N}_2}$ 

2. If ammonia is in limited supply,

 $\underset{(\mathrm{Limited})}{\mathrm{NH}_3} + \underset{(\mathrm{Excess})}{\mathrm{Cl}_2} \longrightarrow \underset{(\mathrm{Nitrogen\ trichloride})}{\mathrm{NCl}_3} + \underset{(\mathrm{Hydrogen\ chloride})}{\mathrm{HCl}(g)}$ 

**3.2.** (a) Identify only the anion present in the following compound:

The compound, on heating, produces a colourless, odourless gas which turns lime water milky and has no effect on acidified potassium dichromate solution.

## Solution

The anion present is carbonate  $CO_3^{2-}$ . Heating the appropriate component causes  $CO_2$  to be released, turning lime water milky and not affecting acidified potassium dichromate solutions.

**3.2.** (b) Identify only the anion present in the following compound:

The solution of the compound which on treating with concentrated sulphuric acid and freshly prepared ferrous sulphate solution produces a brown ring.

### Solution

The anion present is nitrate  $NO_3^-$ . The nitrate-containing chemical undergoes the following reactions to generate a brown ring.

**3.3.** (a) Mohan has three solutions P, Q and R having a pH of 13, 5 and 2 respectively. Which of the above solutions P, Q or R will react with magnesium to liberate hydrogen gas?

## Solution

Highly acidic with pH-2, 'R' interacts with metal to create salt and release H<sub>2</sub> gas.

 $Mg + R \longrightarrow Salt of acid + H_2$ 

**3.3.** (b) Mohan has three solutions P, Q and R having a pH of 13, 5 and 2 respectively. Which of the above solutions P, Q or R will liberate ammonia gas when it reacts with ammonium chloride?

## Solution

'P' reacts with ammonium chloride to produce ammonia due to the fact that it is highly basic at pH-13.

 $NH_4CI + NaOH \rightarrow NH_3 + NaCI + H_2O$ 

**3.3.** (c) Mohan has three solutions P, Q and R having a pH of 13, 5 and 2 respectively. Which of the above solutions P, Q or R will contain molecules as well as ions?

## Solution

With a pH of 5, 'Q' is a weak acid. Because it is partially dissociated, it contains both molecules and ions, making it unionised.

**3.4.** The following table is related to an industrial process of an acid.

Name of the process	Reactant	Catalyst	Final product
(a)	SO <sub>2</sub> + O <sub>2</sub>	(b)	(C)

Identify (a), (b) and (c).

#### Solution

Name of the process	Reactant	Catalyst	Final product
(a) Contact process	SO <sub>2</sub> + O <sub>2</sub>	(b) <b>Vanadium oxide</b>	(c) <b>Sulphuric acid</b>

The contact process is an industrial method used to produce sulphuric acid through a sequence of reactions.

- 1. Combustion of Sulphur:  $\mathrm{S}+\mathrm{O}_{2}\left(\mathrm{g}\right)\longrightarrow\mathrm{SO}_{2}\left(\mathrm{g}\right)$
- 2. Formation of SO<sub>3</sub>: SO<sub>2</sub> + O<sub>2</sub>  $\xrightarrow[2-3]{V_2O_5}{450^{\circ}C}$  SO<sub>3</sub> (g)
- 3. Formation of Oleum:  $SO_2 + H_2SO_4 \longrightarrow H_2S_2O_7$

## Formation of Sulphuric acid

$$\mathrm{H_2S_2O_7} + \mathrm{H_2O} \longrightarrow 2\,\mathrm{H_2SO_4}$$

## Q4.

**4.1.** (a) Define the term.

Molar volume

#### Solution

One mole of any gaseous molecules occupies 22.4 dm<sup>3</sup> (litre) or 22400 cm<sup>3</sup> (ml) at standard temperature and pressure (STP). This volume is known as the molar volume.

"The molar volume of a gas can be defined as the volume occupied by one mole of a gas at standard temperature and pressure."

**4.1.** (b) Define normal salt.

## Solution

Normal salts are the salts formed by the complete replacement of the ionizable hydrogen atoms of an acid by a metallic or ammonium ion. For example: Sodium chloride (NaCl).

**4.2.** (a) Draw the electron dot structure of Methane molecule.

[Atomic number: N = 7, C = 6, H = 1]

#### Solution

Methane:  $CH_4$ , C = 2, 4 H = 1



**4.2.** (b) Draw the electron dot structure of Nitrogen molecule.

[Atomic number: N = 7, C = 6, H = 1]

#### Solution

Nitrogen:  $N_2$ , N = 2, 5

$$(N = N)$$
 :  $N = N$ :

**4.3.** (a) Complete and balance the following equation:

 $AI_2O_3 + NaOH \longrightarrow$ 

Solution

 $\begin{array}{c} Al_2O_3 + 2\,NaOH \longrightarrow 2\,NaAlO_2 + H_2O \\ & \text{Sodium aluminate} \end{array}$ 

**4.3.** (b) Complete and balance the following equation:

$$C_2H_5COONa + NaOH \xrightarrow{\Delta}_{CaO}$$

Solution

$$\mathrm{C_{2}H_{5}COONa} + \mathrm{NaOH} \xrightarrow[\mathrm{CaO}]{\Delta} \underset{\mathrm{Ethane}}{\overset{\Delta}{\longrightarrow}} \mathrm{C_{2}H_{6}} + \mathrm{Na_{2}CO_{3}}$$

**4.3.** (c) Complete and balance the following equation:

$$C_2H_4Br_2 + alcoholic \text{ KOH} \stackrel{\Delta}{\longrightarrow}$$

Solution

$$\mathrm{C_2H_4Br_2} + 2\,\mathrm{alc}\cdot\mathrm{KOH} \stackrel{\Delta}{ o}\mathrm{CH} = \mathrm{CH} + 2\,\mathrm{KBr} + 2\,\mathrm{H_2O}$$

**4.4.** Choose the organic compound from the list given below to answer the following questions:

**4.4.** (a) The compound which does not have a double bond in its structure.

- 1. Ethene
- 2. Ethanoic acid
- 3. Ethanol
- 4. Methanal

#### Solution

Ethanol

#### **Explanation:**

The structural formula clearly represents all single bonds.

$$\begin{array}{cccc} H & H \\ & | & | \\ H - C - C - O - H \\ & | & | \\ H & H \end{array}$$

**4.4.** (b) The compound in its pure form turns into an ice like solid on cooling.

- 1. Ethene
- 2. Ethanoic acid
- 3. Ethanol
- 4. Methanal

#### Solution

Ethanoic acid

### **Explanation:**

Melting point of ethanoic acid is 289.5 K, or 16°C. It so freezes below this temperature and is also known as glacial acetic acid.

**4.4.** (c) The compound which is used for artificial ripening of fruits.

- 1. Ethene
- 2. Ethanoic acid
- 3. Ethanol
- 4. Methanal

#### Solution

Ethene

## **Explanation:**

Fruits start their ripening process with gaseous ethylene emissions. So, a chemical called ethephon (2-chloroethyl phosphonic acid) is also used to make fruits ripen faster. This chemical gets inside the fruit and breaks down into ethylene calcium carbide.

Q5.

**5.1.** (a) Name the main constituent metal in the following alloy:

Duralumin

Name the main metal used in making the alloy given below:

Duralumin

#### Solution

Aluminium

### **Explantion:**

Duralurnin is actually a composition with 95% Al, 4% Copper, 0.5% Mg and 0.5% Mn.

**5.1.** (b) Name the main constituent metal in the following alloy: Stainless steel

Name the main metal used in making of the alloy given below:

Stainless steel

## Solution

Iron

### **Explanation:**

Iron and carbon make up most of stainless steel, in minor amounts. Therefore, steel is just stainless steel with a 10% Ni and a 15% Cr addition.

Stainless steel

- Fe (73%)
- Cr (18%)
- Ni (8%)
- C (1%)

**5.2.** (a) Differentiate between the following pairs based on the odourless gas which turns lime water milky and the criteria given:

Sulphuric acid and Nitric acid (using barium chloride solution)

## Solution

Sulphuric Acid	Nitric Acid
When sulphuric acid reacts with barium	Barium chloride does not react with
chloride, a white ppt. of barium sulphate is formed.	nitric acid.
$BaCl_2 + H_2SO_4 \longrightarrow BaSO_4 + 2HCl$	

**5.2.** (b) Differentiate between the following pairs based on the criteria given:

Unsaturated and Saturated hydrocarbons (type of bond present)

#### Solution

Unsaturated Hydrocarbon	Saturated Hydrocarbon
Unsaturated hydrocarbons are those where carbon atoms self-catenate through double and triple bonds ( $\sigma$ and $\pi$ bonds).	Saturated hydrocarbons are those where carbon atoms self catenate through a single bond (σ bond).
E.g. H - C = C - H $ $	E.g. H H 
$ \begin{array}{c} \mathbf{H} & \mathbf{H} \\ \mathbf{C} \equiv \mathbf{C} \\   &   \\ \mathbf{H} & \mathbf{H} \end{array} $	$\begin{array}{c c} \mathbf{H} - \mathbf{C} - \mathbf{C} - \mathbf{H} \\ &   &   \\ \mathbf{H} & \mathbf{H} \end{array}$

**5.3.** (a) Calcium carbonate react with dilute hydrochloric acid as given below:

 $CaCO_3 + 2HCI \longrightarrow CaCI_2 + H_2O + CO_2$ 

What is the mass of 5 moles of calcium carbonate? (Relative molecular mass of calcium carbonate is 100)

#### Solution

1 mole of  $CaCO_3 = 100 g$ 

5 moles of CaCO<sub>3</sub>  $\rightarrow$  5 × 100 = 500 g

Hence, the mass of 5 moles of CaCO<sub>3</sub> will be 500 g

**5.3.** (b) Calcium carbonate react with dilute hydrochloric acid as given below:

 $CaCO_3 + 2HCI \longrightarrow CaCI_2 + H_2O + CO_2$ 

How many moles of HCl will react with 5 moles of calcium carbonate?

## Solution

2 moles of HCl are used for 1 mole of  $CaCO_3$ 

i.e., CaCO<sub>3</sub> HCl

 $1 mole \rightarrow 2 mole$ 

5mole  $\rightarrow$  5 × 2 = 10 moles

Hence 10 moles of HCl will react with 5 moles of calcium carbonate.

**5.3.** (c) Calcium carbonate react with dilute hydrochloric acid as given below:

 $CaCO_3 + 2HCI \rightarrow CaCl_2 + H_2O + CO_2$ 

What is the volume of carbon dioxide liberated at S.T.P. at the same time?

#### Solution

100 g of CaCO<sub>3</sub>  $\longrightarrow$  22.41 of CO<sub>2</sub> 500 g of CaCO<sub>3</sub> =  $\frac{22.4 \times 500}{100}$ = 112.0

 $\therefore$  112.0 of  $\mathrm{CO}_2$ 

Hence, 112 l of  $CO_2$  is liberated from 5 moles of calcium carbonate.

**5.4.** (a) Identify the gas evolved in the following reaction:

Methane undergoes complete combustion.

#### Solution

Carbon dioxide

#### **Explanation:**

Methane burns completely to produce carbon dioxide and water.

 $\mathsf{CH}_4 + 2\mathsf{O}_2 \longrightarrow \mathsf{CO}_2 + 2\mathsf{H}_2\mathsf{O}$ 

5.4. (b) Identify the gas evolved in the following reaction:

Copper carbonate is heated.

#### Solution

Carbon dioxide

**Explanation:** 

On heating, green copper carbonate breaks down to produce black copper oxide and releases CO<sub>2</sub>.

$${
m CuCO_3} \stackrel{\Delta}{\longrightarrow} {
m CuO} + {
m CO_2}$$

**5.4.** (c) Identify the gas evolved in the following reaction:

MnO<sub>2</sub> reacts with concentrated HCl.

### Solution

Chlorine

## **Explanation:**

Manganese oxide reacts with cone. HCl, releasing greenish colored chlorine and forming manganese chloride.

 $MnO_2 + HCI_2 \longrightarrow MnCI_2 + CI_2 + 2 H_2O$ 

Q6.

**6.1.** (a)

X -  $HCl \rightleftharpoons H^{1+} + Cl^-$  (in solution state)

 ${\bf Y}$  -  $PbBr_2 \rightleftharpoons Pb^{2+} + 2\,Br^{1-}$  (in molten state)

From the above reaction X or Y, identify the reaction which exhibit:

electrolytic dissociation

## Solution

'Y' Electrolytic dissociation of PbBr<sub>2</sub> occurs when energy in the form of heat induces molecules to generate ions, breaking the electrostatic interaction between ions.

**6.1.** (b)

X -  $HCl \rightleftharpoons H^{1+} + Cl^-$  (in solution state)

 $extsf{Y}$  -  $PbBr_2 \rightleftharpoons Pb^{2+} + 2 \, Br^{1-}$  (in molten state)

From the above reaction X or Y, identify the reaction which exhibit:

lonization

#### Solution

'X' HCl is a powerful acid; therefore, in solution, it completely dissociates and generates independent ions that are not held together by any force.

**6.2.** (a) Give a reason for Inert gases do not form ions.

### Solution

Inert gases do not produce ions because their outermost shell is completely filled, resulting in a stable electronic state. As a result, atoms cannot absorb or lose electrons in order to create ions.

**6.2.** (b) Give reason for the following:

Covalent compounds have a low melting and boiling point.

#### Solution

Covalent compounds are held together by modest intramolecular forces. As a result, breaking the bonds between two or more molecules requires only a minimal amount of energy.

Therefore, these compounds have low melting and boiling points.

**6.3.** (a) Arrange the following as per the instruction given in the bracket:

Carbon, Fluorine, Beryllium (decreasing order of atomic size).

#### Solution

Beryllium > Carbon > Fluorine

#### **Explanation:**

As we proceed along a period, the size of the atom decreases due to increased nuclear pull.

**6.3.** (b) Arrange the following as per the instruction given in the bracket:

Sulphuric acid, Phosphoric acid, Acetic acid (increasing order of number of replaceable H atoms per molecule).

#### Solution

Acetic acid < sulphuric acid < phosphoric acid.

#### **Explanation:**

Acetic acid is CH<sub>3</sub>COOH when it ionizes, it has only one H<sup>+</sup> ion (CH<sub>3</sub>COOH – CH<sub>3</sub>COO<sup>-</sup> + H<sup>+</sup>) While sulphuric has 2 replaceable Hydrogen atoms (H<sub>2</sub>SO<sub>4</sub>  $\rightarrow$  2H<sup>+</sup> + SO<sub>4</sub><sup>2-</sup>). Phosporic acid is H<sub>3</sub>PO<sub>5</sub> which ionises to give 3-H atoms (H<sub>3</sub>PO<sub>4</sub> = 3H<sup>+</sup> + PO<sub>4</sub><sup>3-</sup>).

**6.3.** (c) Arrange the following as per the instruction given in the bracket:

Potassium, Lithium, Sodium (increasing order of ionization potential).

## Solution

Lithium < Sodium < Potassium.

## **Explanation:**

Li's electrical configuration 2, 1 will result in least ionization potential. Atomic number 11 for sodium corresponds to 2, 8, 1. With an atomic number of 19, potasium exhibits electrical configuration as 2, 8, 8, 1. This indicates that the atomic radii change from Li to Na to Kand from shell count. As size grows, nuclear attraction reduces and the distance from nucleus rises; so, electron can readily exit with great energy. Greater and smaller the atom's size will be the nuclear pull.

## 6.4. (a) Identify the following:

An element in Period 1 which can be placed in both Group 1 and Group 17 of the Periodic Table.

## Solution

Hydrogen

## **Explanation:**

Hydrogen has an electronic configuration of 1, which allows it to take  $e^-$  and complete its duplet as halogen (Group 17), as well as donate 1  $e^-$  as alkali metals (Group 1).

6.4. (b) Identify the following:

The element having electronic configuration 2, 8, 6.

## Solution

Sulphur

#### **Explanation:**

The atomic number of S is 16. The electronic arrangements are 2, 8, 6.

**6.4.** (c) Identify the following:

The most electronegative element of Period 3.

## Solution

Chlorine

## **Explanation:**

Electronegativity grows from left to right during a certain duration.

## Q7.

**7.1.** Rita was given an unknown salt for identification. She prepared a solution of the salt and divided it into two parts.

- To the first part of the salt solution, she added a few drops of ammonium hydroxide and obtained a reddish-brown precipitate.
- To the second part of the salt solution, she added a few drops of silver nitrate solution and obtained a white precipitate.

Name:

- a. The cation present and
- b. The anion present in the salt given for identification.

## Solution

a. Cation 
$$Fe^{3+}$$
 (ferric ion) When  $Fe^{3+}$  reacts with  $NH_4OH$  it forms ferric hydroxide.

 ${\rm Fe}^{3+} + 3\,{\rm NH_4OH} \longrightarrow {\rm Fe}({\rm OH})_3 + 3\,{\rm NH_4}^+ \\ {\rm (Reddish\ brown\ ppt \,\cdot\,)}$ 

b. Anion:  $\mathrm{Cl}^-$  (Chloride ion)

When  $\mathrm{Cl}^-$  reacts with silver nitrate, it forms a white ppt. of silver chloride.

 $\begin{array}{c} Cl^- + AgNO_3 - AgCl + NO_3^- \\ & \text{Silver chloride} \end{array}$ 

**7.2.** Fill in the blanks by choosing the correct answer from the bracket:

**7.2.** (a) Carbon tetrachloride is a \_\_\_\_\_ covalent molecule.

- 1. Polar
- 2. Non-polar

## Solution

Carbon tetrachloride is a **non-poplar** covalent molecule.

7.2. (b) During electrolysis of acidulated water, the gas liberated at the anode is

- 1. Oxygen
- 2. Hydrogen

## Solution

During electrolysis of acidulated water, the gas liberated at the anode is oxygen.

7.3. Ammonia burns in oxygen, as shown below.

 $4NH_3+3O_2 \longrightarrow 2N_2+6~H_2O$ 

If 240 cc of ammonia is burnt in 300 cc of oxygen, find out the composition of the resultant gaseous mixture at room temperature.

## Solution

```
4NH_3+3O_2 \longrightarrow 2N_2+6~H_2O (vapour)
```

By Gay Lussac's Law

For N<sub>2</sub>:

 $4 \mbox{ vol. of } NH_3 \mbox{ results } 2 \mbox{ vol. of } N_2$ 

240 cc of NH<sub>3</sub> will result = 
$$\frac{2 \times 240}{4}$$

= 120 cc of  $N_2$ 

For H<sub>2</sub>O (vapour phase):

4 Vol. of  $NH_3$  results 6 vol. of  $H_2O$ 

240 cc of NH<sub>3</sub> will result =  $\frac{240 \times 6}{4}$ 

= 360 cc of H<sub>2</sub>O

For O<sub>2</sub>:

4 vol. of  $NH_3$  reacts with 3 vol. of  $H_2O$ 

240 cc of NH<sub>3</sub> will react with = 
$$\frac{3 \times 240}{4}$$

= 180 cc of O<sub>2</sub>

**7.4.** The following table shows the electronic configuration of the atoms A, B, C and D.

Element	A	В	C	D
Electronic configuration	2, 8, 8, 2	2, 6	2, 8, 7	2, 4

- a. Write the formula of the compound formed between:
  - 1. A and B
  - 2. D and C
- b. Which of the above elements will exhibit catenation?

## Solution

(a) (1)

	Electronic Configuration	Valency
Α:	2, 8, 8, 2	+ 2
B:	2, 6	- 2

Formula: AB

(2)

	Electronic Configuration	Valency
D:	2, 4	± 4
C :	2, 8, 7	- 1

Formula: DC<sub>4</sub>

(b) 'D' will exhibit catenation.

Q8.

**8.1.** Choose the correct answer from the list given below:

**8.1.** (a) The ore which can be concentrated by magnetic separation.

- 1. Zinc blende
- 2.  $C_2H_2$
- 3. Calamine
- 4. CH
- 5. Haematite

## Solution

Haematite

## **Explanation:**

Haematite, an iron ore, is magnetically attractive.

8.1. (b) Empirical formula of Ethyne.

- 1. Zinc blende
- 2.  $C_2H_2$
- 3. Calamine
- 4. CH
- 5. Haematite

## Solution

СН

## **Explanation:**

 $C_2H_2$  is a molecular formula that represents the simple ratio in which atoms are joined; consequently, the empirical formula is CH.

**8.2.** (a) Give a balanced equation for the following reaction:

Copper reacts with concentrated nitric acid.

## Solution

$$\begin{array}{c} \mathrm{Cu} + 4 \,\mathrm{HNO}_3 \longrightarrow \mathrm{Cu}(\mathrm{NO}_3)_2 \, + 2 \,\mathrm{NO}_2 + 2 \,\mathrm{H}_2\mathrm{O} \\ _{\mathrm{Copper \ nitrate}} \end{array}$$

**8.2.** (b) Write the equation for the reaction:

Aluminum, Nitride and Water.

Give balanced equation for the following reaction:

Aluminium nitride is treated with warm water

#### Solution

$$AlN + 3\,H_2O \longrightarrow \underset{Aluminium \ hydroxide}{Al(OH)_3} + \underset{Ammonia}{NH_3}$$

**8.3.** Match the salts underlined in Column A with the most suitable method of preparation given in Column B.

Column A	Column B
(a) from ZnCl <sub>2</sub> from Zn	1. Precipitation
(b) from KNO₃ from KOH.	2. Direct combination
(c) from $CaCO_3$ from $CaCl_2$ .	3. Displacement reaction
	4. Neutralization

#### Solution

Column A	Column B
(a) from ZnCl <sub>2</sub> from Zn	3. Displacement reaction
(b) from KNO <sub>3</sub> from KOH.	4. Neutralization
(c) from $CaCO_3$ from $CaCl_2$ .	1. Precipitation

#### **Explanation:**

a. Displacement Reaction

 $Zn + HCI \longrightarrow ZnCI_2 + H_2$ 

Displace H from HCl because zinc is more reactive than hydrogen.

- b. Neutralization Reaction  $KOH + HNO_3 \rightarrow KNO_3 + H_2O$ Base + Acid  $\rightarrow$  Salt + water; Neutralization reaction
- c. Precipitation Reaction
   CaCl<sub>2</sub> + Na<sub>2</sub>CO<sub>3</sub> → 2 NaCl + CaCO<sub>3</sub>↓
   Sodium chloride and a calcium carbonate precipitate follow from the double displacement reaction.

**8.4.** (a) Hydrogen chloride gas is prepared in the laboratory by the action of concentrated sulphuric acid on sodium chloride.

Give a balanced chemical equation for the above reaction.

## Solution

 $2 \text{ NaCl} + \text{H}_2\text{SO}_4 \longrightarrow 2 \text{ HCl} + \text{Na}_2\text{SO}_4$ 

Sodium's greater reactivity than hydrogen replaces 'H' from acid to produce matching salts and strong volatile acid (HCl).

**8.4.** (b) Hydrogen chloride gas is prepared in the laboratory by the action of concentrated sulphuric acid on sodium chloride.

State the method of collection of the gas formed above.

## Solution

Upward displacement of water.

**8.4.** (c) Hydrogen chloride gas is prepared in the laboratory by the action of concentrated sulphuric acid on sodium chloride.

What is the property of sulphuric acid that makes it a suitable reagent for the reaction?

## Solution

Sulphuric acid's low volatility and high boiling point qualify it as a suitable reagent for this reaction.