

BINARY OPERATIONS (XII, R.S. AGARWAL)

EXERCISE 3-A [Pg. No. 74]

1. Let $*$ be a binary operation on the set I of all integers, defined by $a * b = 3a + 4b - 2$.
Find the value of $4 * 5$

Sol. $a * b = 3a + 4b - 2$

$$\Rightarrow 4 * 5 = 3 \times 4 + 4 \times 5 - 2 = 30$$

2. The binary operation $*$ on R is defined by $a * b = 2a + b$. Find $(2 * 3) * 4$

Sol. Given, $a * b = 2a + b \forall a \text{ and } b \in R$.

Now, $(2 * 3) * 4$

$$= (2 \times 2 + 3) * 4 = 12 * 4 = 2 \times 12 + 4 = 28$$

3. Let $*$ be a binary operation on the set of all nonzero real numbers, defined by $a * b = \frac{ab}{5}$. Find the value of x given that $2 * (x * 5) = 10$

Sol. Given:- $a * b = \frac{ab}{5} \quad \therefore 2 * (x * 5) = 10$

$$\Rightarrow 2 * \left(\frac{x \times 5}{5} \right) = 10 \Rightarrow 2 * x = 10 \Rightarrow \frac{2 \times x}{5} = 10 \Rightarrow x = 25$$

4. Let $x : R \times R \rightarrow R$ be a binary operation given by $a * b = a + 4b^2$. Then, compute $(-5) * (2 * 0)$

Sol. Given:- $a * b = a + 4b^2$

Now, $(-5) * (2 * 0)$

$$= (-5) * (2 + 4 \times 0^2) = -5 * 2 = -5 + 4 \times (-2)^2 = 11$$

5. Let $*$ be a binary operation on the set Q of all rational numbers given as $a * b = (2a - b)^2$ for all $a, b \in Q$. Find $3 * 5$ and $5 * 3$. Is $3 * 5 = 5 * 3$?

Sol. Given : $\rightarrow a * b = (2a - b)^2$

$$\text{Now, } 3 * 5 = (2 \times 3 - 5)^2 = 1$$

$$\text{And } 5 * 3 = (2 \times 5 - 3)^2 = 49$$

Here, $3 * 5 \neq 5 * 3$

6. Let $*$ be a binary operation on N given by $a * b = \text{L.C.M. of } a \text{ and } b$. Find the value of $20 * 16$

Is $*$ (i) commutative (ii) associative?

Sol. Given : $\rightarrow a * b = \text{L.C.M. of } a \text{ and } b$

$$\therefore 20 * 16 = \text{L.C.M. of } 20 \text{ and } 16 = 80.$$

Commutatively: \rightarrow

Let, a and $b \in N$

$\therefore \text{L.C.M. at } a \text{ and } b = \text{L.C.M. of } b \text{ and } a$

$$\Rightarrow (a * b) = (b * a) \forall a \text{ \& } b \in N$$

Hence, $*$ is commutative.

Associatively : \rightarrow

$$(a * b) * c$$

$$\begin{aligned}
 &= \{\text{L.C.M of } a \text{ and } b\} * c \\
 &= \text{L.C.M of } [\{\text{L.C.M of } a \text{ and } b\} \text{ and } C] = \text{L.C.M of } [\text{L.C.M of } a, b \text{ and } c] \\
 &= \text{L.C.M of } [a \text{ and } \{\text{L.C.M of } b \text{ and } c\}] = a * \{\text{L.C.M of } b \text{ and } c\} \\
 &= a * (b * c)
 \end{aligned}$$

Here, $a * (b * c) = (a * b) * c \forall a, b, c \in N$ Hence, $*$ is associative.

7. If $*$ be the binary operation on the set Z of all integers defined by $a * b = (a + 3b^2)$, find $2 * 4$

Sol. Given: $\rightarrow a * b = a + 3b^2$

$$\therefore 2 * 4 = 2 + 3 \times (4)^2 = 50$$

8. show that $*$ on Z^+ defined by $a * b = |a - b|$ is not binary operation

Sol. On Z^+ , $*$ is defined by $a * b = |a - b|$, it is seen that for $a, a \in Z^+$.

$$a * a = |a - a| = 0 \notin Z^+, \text{ hence } * \text{ is not a binary operation}$$

9. Let $*$ on Z^+ defined by $a * b = a^b$ is neither commutative nor associative

Sol. **Commutativity:** Let $a, b \in N$, then $a * b = a^b$ and $b * a = b^a$.

a^b and b^a are not equal for every $a, b \in N$.

$\Rightarrow a * b \neq b * a \Rightarrow *$ is not commutative only.

Associativity: Let $a, b, c \in N$, then

$$(a * b) * c = a^b * c = (a^b)^c = a^{bc} \quad \dots(1)$$

$$\text{and } a * (b * c) = a * b^c = (a)^{b^c} \quad \dots(2)$$

From (1) & (2), $(a * b) * c \neq a * (b * c) \Rightarrow *$ is associative on N .

10. Let $a * b = \text{L.C.M}(a, b)$ for all values of $a, b \in N$

(i) Find $(12 * 16)$

(ii) show that $*$ is commutative on N

(iii) Find the identity element in N

(iv) Find all invertible elements in N

Sol. (i) $12 * 16 = \text{L.C.M}(12, 16) = 48$

(ii) Let, $a \& b \in N$

$$a * b = \text{L.C.M}(a, b)$$

$$= \text{L.C.M}(b, a) = b * a \forall a, b \in N$$

Hence, $*$ is commutative.

(iii) Let, C be the identity element

$$\because C \text{ is the identity element } \therefore C * a = a \forall a \in N$$

$$\Rightarrow \text{L.C.M}(8, 9) = a \Rightarrow C = 1$$

(iv) Let, a be an orbitary element on N

$$\text{Now, } a * b = 1 \text{ for } a^{-1} = b$$

$$\Rightarrow \text{L.C.M}(a, b) = 1 \Rightarrow a = b = 1$$

$\Rightarrow 1$ is the only element in N , Which is invertible.

11. Let Q^+ be the set of all positive rational numbers

(i) show that the operation $*$ on Q^+ defined by $a * b = \frac{1}{2}(a + b)$ is a binary operation

(ii) show that $*$ is commutative

(iii) show that $*$ is not associative

Sol. (i) Let, $a \& b \in Q^+$

$$\Rightarrow (a+b) \in Q^+ \Rightarrow \frac{1}{2}(a+b) \in Q^+ \Rightarrow a * b \in Q^+$$

Hence, $*$ on Q^+ is a binary operation.

(ii) Let, a & $b \in Q^+$

$$a * b = \frac{1}{2}(a+b) = \frac{1}{2}(b+a) = b * a \quad \forall a \text{ & } b \in Q^+$$

Hence, $*$ is commutative on Q^+

$$(iii) 10 * (2 * 6) = 10 * \frac{1}{2}(2+6) = 10 * 4 = \frac{10+4}{2} = 7$$

$$(10 * 2) * 6 = \frac{1}{2}(10+2) * 6 = 6 * 6 = \frac{6+6}{2} = 6$$

$\therefore 10 * (2 * 6) \neq (10 * 2) * 6 \therefore *$ is not associative.

12. Show that the set $A = (-1, 0, 1)$ is not closed for addition.

Sol. We have, $1 \in A$, $1 \in A$ and $1+1=2 \notin A$. Hence, A is not closed for addition.

13. $*$ on $R - \{-1\}$, defined by $(a * b) = \frac{a}{(b+1)}$ is neither commutative nor associative

Sol. **Commutativity** : Let $a, b \in R - \{-1\}$, then $a * b = \frac{a}{b+1}$ and $b * a = \frac{b}{a+1}$

$$\Rightarrow a * b \neq b * a \Rightarrow * \text{ is not commutative on } R - \{-1\}.$$

Associativity : Let $a, b, c \in R - \{-1\}$, then

$$(a * b) * c = \left(\frac{a}{b+1} \right) * c = \frac{\frac{a}{b+1}}{c+1} = \frac{a}{(b+1)(c+1)} \quad \dots(1)$$

$$\text{and } a * (b * c) = a * \left(\frac{b}{c+1} \right) = \frac{a}{\frac{b}{c+1} + 1} = \frac{a(c+1)}{b+c+1} \quad \dots(2)$$

From (1) and (2), $(a * b) * c \neq a * (b * c) \Rightarrow *$ is not associative on $R - \{-1\}$

14. For all $a, b \in R$, we defined $a * b = |a - b|$

Show that $*$ is commutative but not associative

Sol. Let, a & $b \in R$.

$$a * b = |a - b|$$

$$= |b - a| = b * a$$

$$\therefore * \text{ is commutative. } (2 * 3) * 4 = |2 - 3| * 4$$

$$= 1 * 4 = |1 - 4| = 3$$

$$\text{and, } 2 * (3 * 4) = 2 * (3 - 4)$$

$$= 2 * 1 = |2 - 1| = 1$$

$$\therefore (2 * 3) * 4 \neq 2 * (3 * 4) \therefore * \text{ is not associative}$$

15. For all $a, b \in N$, we defined $a * b = a^3 + b^3$

Show that $*$ is commutative but not associative

Sol. Let, a and $b \in N$

$$a * b = a^3 + b^3$$

$$= b^3 + a^3 \quad \forall a, b \in N$$

$$\therefore * \text{ is commutative. } (1 * 2) * 3 = (1^3 + 2^3) * 3$$

$$= 9 * 3 = 9^3 + 3^3 = 729 + 27 = 756$$

$$1 * (2 * 3) = 1 * (2^2 + 3^2)$$

$$= 1 * 35 = 1^3 + 35^3 = 1 + 42875 = 42876$$

$$\therefore (1 * 2) * 3 \neq 1 * (2 * 3) \therefore f \text{ is not associative}$$

16. Let X be a non-empty set and $*$ be a binary operation on $P(X)$, the power set of X , defined by $A * B = A \cap B$ for all $A, B \in P(X)$.

(i) Find the identity element in $(P(X), *)$ (ii) Show that X is the only invertible element in $P(X)$.

Sol. (i) Since $A \cap X = A$ for all A in $P(X)$. $\therefore X$ is the identity element.

(ii) Let A be invertible in $P(X)$ and let B be its inverse.

Then, $A \cap B = X$. This is possible only when $A = B = X$.

$\therefore X$ is the only invertible element in $P(X)$ and its inverse is X .

17. A binary operation $*$ on the set $\{0, 1, 2, 3, 4, 5\}$ is defined as $a * b = \begin{cases} a + b; & \text{if } a + b < 6 \\ a + b - 6; & \text{if } a + b \geq 6 \end{cases}$

Show that 0 is the identity for this operation and each element a has an inverse $(6 - a)$

Sol. Here, $a * 0 = a + 0 \begin{cases} \because a < 6 \\ \therefore a + 0 < 6 \end{cases}$

$$\Rightarrow a * 0 = a$$

Hence, 0 is the identity element

Inverse element of $a : \rightarrow$

$$\text{Let, } b = a^{-1}$$

$$\therefore a * b = 0$$

$$\Rightarrow a + b = 0 \text{ if } a + b < 6$$

$$\text{or, } a + b - 6 = 0 \text{ if } a + b \geq 6$$

$$\text{If, } a + b = 0 \Rightarrow a = -b$$

$$\text{clearly, If } a = 0 \text{ then, } a^{-1} = 0$$

$$\text{If, } a + b - 6 = 0 \text{ then, } b = 6 - a$$

$$\therefore a^{-1} = 6 - a, \forall a \in \{1, 2, 3, 4, 5\}$$

EXERCISE 3-B [Pg.No. 78]

1. Define $*$ on N by $m * n = \text{lcm}(m, n)$. Show that $*$ is a binary operation which is commutative as well as associative.

Sol. Let m and $n \in N$, then $m * n = \text{lcm}(m, n) = \text{lcm}(n, m) = n * m$

Hence, $*$ is commutative binary operation.

$$\text{and } (m * n) * p = (\text{lcm}(m, n)) * p = \text{lcm of } (\text{lcm of } (m, n) \text{ and } p) = \text{lcm of } (m, n, p)$$

$$\text{and } m * (n * p) = m * (\text{lcm of } (n, p)) = \text{lcm of } (m \text{ and } \text{lcm of } (n, p)) = \text{lcm of } (m, n, p)$$

$$\therefore (m * n) * p = m * (n * p). \text{ Hence, the operation is associative.}$$

2. Define $*$ on Z by $a * b = a - b + ab$. Show that $*$ is a binary operation on Z which is neither commutative nor associative.

Sol. Commutativity : Let us take two elements 1 and 2 of Z .

Then, $1*2 = 1 - 2 + 1 \times 2 = 1$ and $2*1 = 2 - 1 + 2 \times 1 = 3 \quad \therefore 1*2 \neq 2*1$

Hence, the binary operation is not commutative.

Associativity : $2, 3, 4 \in \mathbb{Z}$

$$(2*3)*4 = (2 - 3 + 2 \times 3)*4 = 5*4 = 5 - 4 + 5 \times 4 = 21$$

$$\text{and } 2*(3*4) = 2*(3 - 4 + 12) = 2*11 = 2 - 11 + 2 \times 11 = 13$$

$\Rightarrow (2*3)*4 \neq 2*(3*4)$. Hence, binary operation is not associative.

3. Define $*$ on \mathbb{Z} by $a*b = a + b - ab$. Show that $*$ is a binary operation on \mathbb{Z} which is commutative as well as associative.

Sol. Let $a, b \in \mathbb{Z}$, then by definition $a*b = a + b - ab$ and $b*a = b + a - ba$

Hence, $a*b = b*a$.

For Associativity : Let $a, b, c \in \mathbb{Z}$.

$$\begin{aligned} \text{Now, } (a*b)*c &= (a + b - ab)*c = a + b - ab + c - (a + b - ab)c \\ &= a + b + c - ab - bc - ca + abc \quad \dots(1) \end{aligned}$$

$$\begin{aligned} \text{and } a*(b*c) &= a*(b + c - bc) = a + b + c - bc - a(b + c - bc) \\ &= a + b + c - ab - bc - ca + abc \quad \dots(2) \end{aligned}$$

Hence, by (1) and (2), $(a*b)*c = a*(b*c)$

Hence, the binary operation on \mathbb{Z} is associative.

4. Consider a binary operation on $\mathbb{Q} - \{1\}$, defined by $a*b = a + b - ab$

(i) Find the identity element in $\mathbb{Q} - \{1\}$ (ii) Show that each $a \in \mathbb{Q} - \{1\}$ has its inverse

Sol. (i) Let e be the identity element.

$$\text{Then, } a*e = a \quad \forall a \in \mathbb{Q} - \{1\}$$

$$\Rightarrow a + e - ae = a \quad \Rightarrow e(1 - a) = 0 \quad \Rightarrow e = 0 \in \mathbb{Q} - \{1\}$$

$$\text{Now, } a*0 = a + 0 = a$$

$$\text{and, } 0*a = 0 + a = a$$

thus, 0 is the identity element in $\mathbb{Q} - \{1\}$

(ii) Let $a \in \mathbb{Q} - \{1\}$ and Let, $a^{-1} = b$,

$$\text{Now, } a*b = 0$$

$$\Rightarrow a + b - ab = 0 \Rightarrow a = ab - b \Rightarrow a = (a - 1) \cdot b \Rightarrow b = \frac{a}{a - 1} \in \mathbb{Q} - \{1\} \Rightarrow a^{-1} = \frac{a}{a - 1} \in \mathbb{Q} - \{1\}$$

Hence, each $a \in \mathbb{Q} - \{1\}$ has its inverse.

5. Let \mathbb{Q}_0 be the set of all non-zero rational numbers. Let $*$ be a binary operation on \mathbb{Q}_0 , defined by

$$a*b = \frac{ab}{4} \text{ for all } a, b \in \mathbb{Q}_0$$

(i) Show that $*$ is commutative and associative (ii) Find the identity element in \mathbb{Q}_0

(iii) Find the inverse of an element a in \mathbb{Q}_0

Sol. Let $a*b = \frac{ab}{4}$

$$(i) \text{ For all } a, b, c \in \mathbb{Q}_0, \text{ we have } a*b = \frac{ab}{4} = \frac{ba}{4} = b*a \text{ and } (a*b)*c = \frac{ab}{4} * c = \frac{\frac{ab}{4} * c}{4} = \frac{(ab)c}{16}$$

Also, $a*(b*c) = a*\frac{bc}{4} = \frac{a\left(\frac{bc}{4}\right)}{4} = \frac{a(bc)}{16}$. But $(ab)c = a(bc)$. Hence, $(a*b)*c = a*(b*c)$

(ii) Let e be the identity element and let $a \in Q_0$. Then $a*e = a \Rightarrow \frac{ae}{4} = a \Rightarrow e = 4$

$\therefore 4$ is the identity element in Q .

(iii) Let $a \in Q_0$ and let its inverse be b , then, $a*b = e \Rightarrow \frac{ab}{4} = 4 \Rightarrow b = \frac{16}{a} \in Q_0$

Thus, each $a \in Q_0$ has $\frac{16}{a}$ as its inverse.

6. On the set Q^+ of all positive rational numbers, define an operation $*$ on Q^+ by $a*b = \frac{ab}{2}$ for all

$a, b \in Q^+$

Show that

(i) $*$ is a binary operation on Q^+ (ii) $*$ is commutative (iii) $*$ is associative

Find the identity element in Q^+ for $*$

What is the inverse of $a \in Q^+$?

Sol. Let, a and $b \in Q^+ \therefore a$ and $b \in Q^+$

$$\Rightarrow a \cdot b \in Q^+ \Rightarrow \frac{a \cdot b}{2} \in Q^+ \Rightarrow a * b \in Q^+$$

Hence, $*$ is binary operation on Q^+

Commutativity: \rightarrow

Let, a and $b \in Q^+$

$$\text{Now, } a * b = \frac{ab}{2} = \frac{ba}{2} = b * a \quad \forall a \& b \in Q^+$$

Hence, $*$ is commutative on Q^+

Associativity: \rightarrow

Let, a, b and $c \in Q^+$

$$\text{Now, } a * (b * c) = a * \left(\frac{bc}{2}\right)$$

$$= \frac{a\left(\frac{ab}{2}\right)}{2} = \frac{abc}{4} \quad \text{and, } (a * b) * c = \frac{ab}{2} * c$$

$$= \frac{\frac{ab}{2} \cdot c}{2} = \frac{abc}{4}$$

$$\therefore (a * b) * c = a * (b * c), \quad \forall a, b \& c \in Q^+$$

Hence, $*$ is associative on Q^+

Identity element:-

Let, $e \in Q^+$ be the identity element $\therefore a * e = a$ and $e * a = a$

$$\Rightarrow \frac{a \cdot e}{2} = a \text{ and } \frac{e \cdot a}{2} = a \Rightarrow e = 2 \in Q^+$$

Hence, $e = 2$ is the identity element on Q^+

Inverse of a :-

Let, $a^{-1} = b$

$$\text{Now, } a * b = 2 \Rightarrow \frac{a \cdot b}{2} = 2 \Rightarrow b = \frac{4}{a} \Rightarrow a^{-1} = \frac{4}{a}$$

7. Let Q^+ be the set of all positive rational numbers.

(i) Show that the operation $*$ on Q^+ defined by $a * b = \frac{1}{2}(a + b)$ is a binary operation

(ii) Show that $*$ is commutative

(iii) Show that $*$ is not associative

Sol. (i) On Q^+ , $*$ defined by $a * b = \frac{a+b}{2}$

It is seen that for each $a, b \in Q^+$, there is a unique element $\frac{a+b}{2}$ in Q^+ .

This means that $*$ carries each pair (a, b) to a unique element $a * b = \frac{a+b}{2}$ in Q^+ .

Therefore, $*$ is a binary operation.

(ii) **Commutative** : $a * b = \frac{a+b}{2} = \frac{b+a}{2} = b * a$, $a * b = b * a$, which shows $*$ is commutative.

(iii) **Associative** : $(a * b) * c = \left(\frac{a+b}{2}\right) * c = \frac{\left(\frac{a+b}{2}\right) + c}{2} = \frac{a+b+2c}{4}$

$$a * (b * c) = a * \left(\frac{b+c}{2}\right) = \frac{a + \left(\frac{b+c}{2}\right)}{2} = \frac{2a+b+c}{4}$$

Now, $\frac{a+b+2c}{4} \neq \frac{2a+b+c}{4} \Rightarrow (a * b) * c \neq a * (b * c)$, hence, $*$ is not associative.

8. Let Q be the set of all rational numbers. Define an operation $*$ on $Q - \{-1\}$ by $a * b = a + b + ab$.

Show that :

(i) $*$ is a binary operation on $Q - \{-1\}$ (ii) $*$ is commutative

(iii) $*$ is associative

(iv) zero is the identity element in $Q - \{-1\}$ for $*$

(v) $a^{-1} = \left(\frac{-a}{1+a}\right)$, where $a \in Q - \{-1\}$

Sol. (i) On Q^+ , $*$ is defined by $a * b = a + b + ab$, it is seen that for each $a, b \in Q^+$ there is unique element $a + b + ab$ in Q^+ . This means that $*$ carries each pair (a, b) to a unique element $a * b = a + b + ab$ in Q^+ . Therefore, $*$ is a binary operation.

Note that $a + b + ab = -1$ is not possible.

\therefore If $a + b + ab = -1 \Rightarrow (1+a)(1+b) = 0 \Rightarrow a = -1$ or $b = -1$, which is not possible.

\therefore Both cannot be -1 .

(ii) **Commutativity** : For any $a, b \in R - \{-1\}$,

We have $a * b = a + b + ab$ and $b * a = b + a + ba$ and as $a + b + ab = b + a + ba$

$\therefore a * b = b * a$. So, $*$ is commutative on $R - \{-1\}$.

(iii) **Associativity** : For any $a, b, c \in R - \{-1\}$ we have $(a * b) * c = (a + b + ab) * c$

$$(a * b) * c = (a + b + ab) + c + (a + b + ab)c$$

$$\Rightarrow (a * b) * c = a + b + c + ab + bc + ac + abc \quad \dots(1)$$

$$\text{and } a * (b * c) = a * (b + c + bc)$$

$$a * (b * c) = a + (b + c + bc) + a(b + c + bc) = a + b + c + ab + bc + ca + abc \quad \dots(2)$$

From (1) and (2), we have, $(a * b) * c = a * (b * c)$ for all $a, b, c \in R - \{-1\}$

So, $*$ is associative on $R - \{-1\}$.

(iv) Existence of identity : Let e be the identity element. Then, $a * e = a = e * a$ for all $a \in R - \{-1\}$

$$\Rightarrow a + e + ae = a \text{ and } e + a + ea = a \text{ for all } a \in R - \{-1\}$$

$$\Rightarrow e(1+a) = 0 \text{ for all } a \in R - \{-1\} \Rightarrow e = 0$$

Also, $0 \in R - \{-1\}$. So, 0 is the identity element for $*$ defined on $R - \{-1\}$.

Existence of inverse : Let $a \in R - \{-1\}$ and let b be the inverse of a .

$$\text{Then, } a * b = e = b * a \Rightarrow a * b = e$$

$$\Rightarrow a + b + ab = 0 \quad [\because \text{Identity element is } 0]$$

$$\Rightarrow b = \frac{-a}{a+1}, \text{ Since, } a \in R - \{-1\} \quad [\because a \neq -1 \Rightarrow a+1 \neq 0, \text{ hence, } \frac{-a}{a+1} \text{ is defined}]$$

Hence, every element of $R - \{-1\}$ is invertible and the inverse of an element a is $\frac{-a}{a+1}$.

9. Let $A = N \times N$. Define $*$ on A by $(a, b) * (c, d) = (a + c, b + d)$

Show that :

(i) A is closed for $*$

(ii) $*$ is commutative

(iii) $*$ is associative

(iv) identity element does not exist in A

Sol. (i) Let $(a, b) \in A$ and $(c, d) \in A$, then $a, b, c, d \in N$

$$(a, b) * (c, d) = (a + c, b + d) \in A \quad [\because a + c \in N, b + d \in N]$$

$\therefore A$ is closed for $*$.

(ii) Commutativity : Let $(a, b), (c, d) \in A$,

$$\text{then } (a, b) * (c, d) = (a + c, b + d) \text{ and } (c, d) * (a, b) = (c + a, d + b)$$

$$\because a + c = c + a \text{ and } b + d = d + b \text{ for all } a, b, c, d \in N$$

$$\therefore (a + c, b + d) = (c + a, d + b) \text{ for all } a, b, c, d \in N$$

$$\Rightarrow (a, b) * (c, d) = (c, d) * (a, b) \text{ for all } (a, b), (c, d) \in N \times N = A$$

(iii) Associativity : For any $(a, b), (c, d), (e, f) \in A$, we have, $**$

$$\{(a, b) * (c, d)\} * (e, f) = (a + c, b + d) * (e, f) = ((a + c) + e, (b + d) + f)$$

$$= (a + (c + e), b + (d + f)) \quad [\because \text{Addition is associative on } N]$$

$$= (a, b) * (c + e, d + f) = (a, b) * \{(c, d) * (e, f)\}$$

So, $**$ is associative on A .

(iv) Let (x, y) be the identity element in A . Then $(a, b) * (x, y) = (a, b)$ for all $(a, b) \in A$

$$\Rightarrow (a + x, b + y) = (a, b) \text{ for all } (a, b) \in A \Rightarrow a + x = a, b + y = b \text{ for all } a, b \in N$$

$\Rightarrow x=0, y=0$, clearly $(0, 0) \in A$. Hence, identity element does not exist in A .

10. Let $A = \{1, -1, i, -i\}$ be the set of four 4th roots of unity. Prepare the composition table for multiplication on A and show that

- (i) A is closed for multiplication (ii) multiplication is associative on A
 (iii) multiplication is commutative on A (iv) 1 is the multiplicative identity
 (v) every element in A has its multiplicative inverse

Sol.

\bullet	1	-1	i	-i
1	1	-1	i	-i
-1	-1	1	-i	i
i	i	-i	-1	1
-i	-i	i	1	-1

- (i) Clearly every element of table belongs to the set A . Hence A is closed for multiplication.
 (ii) Clearly $a(bc) = (ab)c$ is satisfied for all $a, b, c \in A$. Hence multiplication is Associative.
 (iii) Clearly the table is symmetrical about the diagonal line.
 Hence multiplication is commutative for A .
 (iv) As $1.1=1$, $1.(-1)=-1$, $1.i=i$ and $1.(-i)=-i$ and $1.1=1$, $-1.1=-1$, $i.1=i$ and $-i.1=-i$.
 Hence 1 is multiplicative identity.
 (v) As 1 is present in every row and columns of product.
 Hence every element in A has its multiplicative inverse.