

Chapter

7

Three Dimensional Co-ordinate Geometry

CONTENTS

System of Co-ordinates

7.1	Co-ordinates of a point in space
7.2	Distance formula
7.3	Section formulas
7.4	Triangle
7.5	Volume of tetrahedron
7.6	Direction cosines and Direction ratio
7.7	Projection
7.8	Angle between two lines

The Straight Line

7.9	Straight line in space
7.10	Equation of line passing through two given points
7.11	Changing unsymmetrical form to symmetrical form
7.12	Angle between two lines
7.13	Reduction of cartesian form of the equation of a line to vector form and vice-versa
7.14	Intersection of two lines
7.15	Foot of perpendicular from a point
7.16	Shortest distance between two lines

The Plane

7.17	Definition of plane and its equations
7.18	Equation of plane passing through the given point
7.19	Foot of perpendicular from a point to a given plane
7.20	Angle between two planes
7.21	Equation of planes bisecting angle between two given planes
7.22	Image of a point in a plane
7.23	Coplaner lines

Line and Plane

7.24	Equation of plane through a given line
7.25	Transformation from unsymmetric form of the equation of line to the symmetric form
7.26	Intersection point of a line and plane
7.27	Angle between line and plane
7.28	Projection of a line on a plane

Sphere

7.29	General equation of sphere
7.30	Equation of sphere of various types
7.31	Section of a sphere by a plane
7.32	Condition of tangency of a plane to a sphere
7.33	Intersection of straight line and a sphere
7.34	Angle of intersection of two spheres

Assignment (Basic and Advance Level)

Answer Sheet of Assignment



Pierre Fermat

Rene' Descartes (1596-1650 A.D.), the father of analytical geometry, essentially dealt with plane geometry only in 1637. The same is true of his coinventor Pierre Fermat (1601-1665 A.D.)

Descartes had the idea of co-ordinates in three dimensions but did not develop it.

J.Bernoulli (1667-1748 A.D.) in a letter of 1715 A.D. to Leibnitz introduced the three co-ordinate planes which we use today. It was Antoine Parent (1666-1716 A.D.), who gave a systematic development of analytical solid geometry for the first time in a paper presented to the French Academy in 1700 A.D.

L.Euler (1707-1783 A.D.) took up systematically the three dimensional co-ordinate geometry.


It was not until the middle of the nineteenth century that geometry was extended to more than three dimensions, the well-known application of which is in the Space-Time Continuum of Einstein's Theory of Relativity.

Three Dimensional Co-ordinate Geometry

System of Co-ordinates

7.1 Co-ordinates of a Point in Space

(1) **Cartesian Co-ordinates** : Let O be a fixed point, known as origin and let OX , OY and OZ be three mutually perpendicular lines, taken as x -axis, y -axis and z -axis respectively, in such a way that they form a right-handed system.




The planes XOY , YOZ and ZOX are known as xy -plane, yz -plane and zx -plane respectively.

Let P be a point in space and distances of P from yz , zx and xy -planes be x , y , z respectively (with proper signs), then we say that co-ordinates of P are (x, y, z) .

Also $OA = x$, $OB = y$, $OC = z$.

The three co-ordinate planes (XOY , YOZ and ZOX) divide space into eight parts and these parts are called octants.



Signs of co-ordinates of a point : The signs of the co-ordinates of a point in three dimension follow the convention that all distances measured along or parallel to OX , OY , OZ will be positive and distances moved along or parallel to OX' , OY' , OZ' will be negative.

The following table shows the signs of co-ordinates of points in various octants :


Octant co-ordinate	$OXYZ$	$OX'YZ$	$OXY'Z$	$OX'Y'Z$	$OXYZ'$	$OX'YZ'$	$OXY'Z'$	$OX'Y'Z'$
x	+	−	+	−	+	−	+	−
y	+	+	−	−	+	+	−	−
z	+	+	+	+	−	−	−	−

(2) Other methods of defining the position of any point P in space :

(i) **Cylindrical co-ordinates** : If the rectangular cartesian co-ordinates of P are (x, y, z) , then those of N are (u, ϕ, z) and we can easily have the following relations : $x = u \cos \phi$, $y = u \sin \phi$ and $z = z$.

Hence, $u^2 = x^2 + y^2$ and $\phi = \tan^{-1}(y/x)$.

Cylindrical co-ordinates of $P \equiv (u, \phi, z)$



(ii) **Spherical polar co-ordinates** : The measures of quantities r , θ , ϕ are known as spherical or three dimensional polar co-ordinates of the point P . If the rectangular cartesian co-ordinates of P are (x, y, z) then

$$z = r \cos \theta, u = r \sin \theta \therefore x = u \cos \phi = r \sin \theta \cos \phi, y = u \sin \phi = r \sin \theta \sin \phi \text{ and } z = r \cos \theta$$

$$\text{Also } r^2 = x^2 + y^2 + z^2 \text{ and } \tan \theta = \frac{u}{z} = \frac{\sqrt{x^2 + y^2}}{z}; \tan \phi = \frac{y}{x}$$

Note : \square The co-ordinates of a point on xy -plane is $(x, y, 0)$, on yz -plane is $(0, y, z)$ and on zx -plane is $(x, 0, z)$

\square The co-ordinates of a point on x -axis is $(x, 0, 0)$, on y -axis is $(0, y, 0)$ and on z -axis is $(0, 0, z)$

\square **Position vector of a point** : Let $\mathbf{i}, \mathbf{j}, \mathbf{k}$ be unit vectors along OX, OY and OZ respectively. Then position vector of a point $P(x, y, z)$ is $\overrightarrow{OP} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$.

7.2 Distance Formula

(1) **Distance formula** : The distance between two points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ is given by

$$AB = \sqrt{[(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2]}$$


(2) **Distance from origin** : Let O be the origin and $P(x, y, z)$ be any point, then $OP = \sqrt{(x^2 + y^2 + z^2)}$.

(3) **Distance of a point from co-ordinate axes** : Let $P(x, y, z)$ be any point in the space. Let PA, PB and PC be the perpendiculars drawn from P to the axes OX, OY and OZ respectively.

$$\text{Then, } PA = \sqrt{(y^2 + z^2)}$$

$$PB = \sqrt{(z^2 + x^2)}$$

$$PC = \sqrt{(x^2 + y^2)}$$



[MP PET 2003]

Example: 1 The distance of the point $(4, 3, 5)$ from the y -axis is

(a) $\sqrt{34}$

(b) 5

(c) $\sqrt{41}$

(d) $\sqrt{15}$

Solution: (c) Distance $= \sqrt{x^2 + z^2} = \sqrt{16 + 25} = \sqrt{41}$

Example: 2 The points $(5, -4, 2)$, $(4, -3, 1)$, $(7, -6, 4)$ and $(8, -7, 5)$ are the vertices of

[Rajasthan PET 2002]

(a) A rectangle

(b) A square

(c) A parallelogram

(d) None of these

Solution: (c) Let the points be $A(5, -4, 2)$, $B(4, -3, 1)$, $C(7, -6, 4)$ and $D(8, -7, 5)$.

$$AB = \sqrt{1+1+1} = \sqrt{3}, CD = \sqrt{1+1+1} = \sqrt{3}, BC = \sqrt{9+9+9} = 3\sqrt{3}, AD = \sqrt{9+9+9} = 3\sqrt{3}$$

$$\text{Length of diagonals } AC = \sqrt{4+4+4} = 2\sqrt{3}, BD = \sqrt{16+16+16} = 4\sqrt{3}$$


$$\text{i.e., } AC \neq BD$$

Hence, A, B, C, D are vertices of a parallelogram

7.3 Section Formulas

(1) **Section formula for internal division** : Let $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ be two points. Let R be a point on the line segment joining P and Q such that it divides the join of P and Q internally in the ratio $m_1 : m_2$. Then the co-ordinates of R are

$$\left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2}, \frac{m_1 z_2 + m_2 z_1}{m_1 + m_2} \right).$$



(2) **Section formula for external division** : Let $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ be two points, and let R be a point on PQ produced, dividing it externally in the ratio $m_1 : m_2$ ($m_1 \neq m_2$). Then the co-ordinates of R are

$$\left(\frac{m_1 x_2 - m_2 x_1}{m_1 - m_2}, \frac{m_1 y_2 - m_2 y_1}{m_1 - m_2}, \frac{m_1 z_2 - m_2 z_1}{m_1 - m_2} \right).$$

Note : \square **Co-ordinates of the midpoint** : When division point is the mid-point of PQ then ratio will be

$$1 : 1, \text{ hence co-ordinates of the mid point of } PQ \text{ are } \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right).$$

\square **Co-ordinates of the general point** : The co-ordinates of any point lying on the line joining points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ may be taken as $\left(\frac{kx_2 + x_1}{k+1}, \frac{ky_2 + y_1}{k+1}, \frac{kz_2 + z_1}{k+1} \right)$, which divides PQ in the ratio $k : 1$. This is called general point on the line PQ .

Example: 3 If the x -co-ordinate of a point P on the join of $Q(2, 2, 1)$ and $R(5, 1, -2)$ is 4, then its z -co-ordinate is

[Rajasthan PET 2003]

- (a) 2 (b) 1 (c) -1 (d) -2

Solution: (c) Let the point P be $\left(\frac{5k+2}{k+1}, \frac{k+2}{k+1}, \frac{-2k+1}{k+1} \right)$. \because Given that $\frac{5k+2}{k+1} = 4 \Rightarrow k = 2 \therefore z\text{-co-ordinate of } P = \frac{-2(2)+1}{2+1} = -1$

7.4 Triangle

(1) Co-ordinates of the centroid

(i) If $(x_1, y_1, z_1), (x_2, y_2, z_2)$ and (x_3, y_3, z_3) are the vertices of a triangle, then co-ordinates of its centroid are $\left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$.


(ii) If $(x_r, y_r, z_r); r = 1, 2, 3, 4$, are vertices of a tetrahedron, then co-ordinates of its centroid are $\left(\frac{x_1 + x_2 + x_3 + x_4}{4}, \frac{y_1 + y_2 + y_3 + y_4}{4}, \frac{z_1 + z_2 + z_3 + z_4}{4} \right)$.

(iii) If $G(\alpha, \beta, \gamma)$ is the centroid of $\triangle ABC$, where A is (x_1, y_1, z_1) , B is (x_2, y_2, z_2) , then C is $(3\alpha - x_1 - x_2, 3\beta - y_1 - y_2, 3\gamma - z_1 - z_2)$.

(2) **Area of triangle** : Let $A(x_1, y_1, z_1)$, $B(x_2, y_2, z_2)$ and $C(x_3, y_3, z_3)$ be the vertices of a triangle, then

$$\Delta_x = \frac{1}{2} \begin{vmatrix} y_1 & z_1 & 1 \\ y_2 & z_2 & 1 \\ y_3 & z_3 & 1 \end{vmatrix}, \Delta_y = \frac{1}{2} \begin{vmatrix} x_1 & z_1 & 1 \\ x_2 & z_2 & 1 \\ x_3 & z_3 & 1 \end{vmatrix}, \Delta_z = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

Now, area of $\triangle ABC$ is given by the relation $\Delta = \sqrt{\Delta_x^2 + \Delta_y^2 + \Delta_z^2}$.



$$\text{Also, } \Delta = \frac{1}{2} |\vec{AB} \times \vec{AC}| = \frac{1}{2} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix}$$

(3) **Condition of collinearity** : Points $A(x_1, y_1, z_1)$, $B(x_2, y_2, z_2)$ and $C(x_3, y_3, z_3)$ are collinear

$$\text{If } \frac{x_1 - x_2}{x_2 - x_3} = \frac{y_1 - y_2}{y_2 - y_3} = \frac{z_1 - z_2}{z_2 - z_3}$$

7.5 Volume of Tetrahedron

$$\text{Volume of tetrahedron with vertices } (x_r, y_r, z_r); r = 1, 2, 3, 4, \text{ is } V = \frac{1}{6} \begin{vmatrix} x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \\ x_4 & y_4 & z_4 & 1 \end{vmatrix}$$

Example: 4 If centroid of tetrahedron $OABC$, where A, B, C are given by $(a, 2, 3)$, $(1, b, 2)$ and $(2, 1, c)$ respectively be $(1, 2, -1)$, then distance of $P(a, b, c)$ from origin is equal to

- (a) $\sqrt{107}$ (b) $\sqrt{14}$ (c) $\sqrt{107/14}$ (d) None of these

Solution: (a) $(1, 2, -1)$ is the centroid of the tetrahedron

$$\therefore 1 = \frac{0+a+1+2}{4} \Rightarrow a=1, 2 = \frac{0+2+b+1}{4} \Rightarrow b=5, -1 = \frac{0+3+2+c}{4} \Rightarrow c=-9.$$

$$\therefore (a, b, c) = (1, 5, -9). \text{ Its distance from origin} = \sqrt{1+25+81} = \sqrt{107}$$

Example: 5 If vertices of triangle are $A(1, -1, 2)$, $B(2, 0, -1)$ and $C(0, 2, 1)$, then the area of triangle is

[Rajasthan PET 2000]

- (a) $\sqrt{6}$ (b) $2\sqrt{6}$ (c) $3\sqrt{6}$ (d) $4\sqrt{6}$

Solution: (b) $\Delta = \frac{1}{2} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2-1 & 0+1 & -1-2 \\ 0-2 & 2-0 & 1+1 \end{vmatrix} = \frac{1}{2} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -3 \\ -2 & 2 & 2 \end{vmatrix} = \frac{1}{2} |\mathbf{i}(8) - \mathbf{j}(-4) + \mathbf{k}(4)|$

$$= |4\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}| = \sqrt{16+4+4} = \sqrt{24} = 2\sqrt{6}$$

Example: 6 The points $(5, 2, 4)$, $(6, -1, 2)$ and $(8, -7, k)$ are collinear, if k is equal to

[Kurukshetra CEE 2000]

- (a) -2 (b) 2 (c) 3 (d) -1

Solution: (a) If given points are collinear, then

$$\frac{x_1 - x_2}{x_2 - x_3} = \frac{y_1 - y_2}{y_2 - y_3} = \frac{z_1 - z_2}{z_2 - z_3} \Rightarrow \frac{5-6}{6-8} = \frac{2+1}{-1+7} = \frac{4-2}{2-k} \Rightarrow \frac{-1}{-2} = \frac{3}{6} = \frac{2}{2-k} \Rightarrow \frac{1}{2} = \frac{2}{2-k} \Rightarrow k = -2$$

7.6 Direction cosines and Direction ratio


(1) Direction cosines

(i) The cosines of the angle made by a line in anticlockwise direction with positive direction of co-ordinate axes are called the direction cosines of that line.

If α, β, γ be the angles which a given directed line makes with the positive direction of the x, y, z co-ordinate axes respectively, then $\cos \alpha, \cos \beta, \cos \gamma$ are called the direction cosines of the given line and are generally denoted by l, m, n respectively.

Thus, $l = \cos \alpha$, $m = \cos \beta$ and $n = \cos \gamma$.

By definition, it follows that the direction cosine of the axis of x are respectively $\cos 0^\circ, \cos 90^\circ, \cos 90^\circ$ i.e. $(1, 0, 0)$. Similarly direction cosines of the axes of y and z are respectively $(0, 1, 0)$ and $(0, 0, 1)$.



Relation between the direction cosines : Let OP be any line through the origin O which has direction cosines l, m, n .

n. Let $P = (x, y, z)$ and $OP = r$. Then $OP^2 = x^2 + y^2 + z^2 = r^2$ (i)


From P draw PA, PB, PC perpendicular on the co-ordinate axes, so that $OA = x, OB = y, OC = z$. Also, $\angle POA = \alpha, \angle POB = \beta$ and $\angle POC = \gamma$.

From triangle AOP , $l = \cos \alpha = \frac{x}{r} \Rightarrow x = lr$

Similarly $y = mr$ and $z = nr$.

Hence from (i), $r^2(l^2 + m^2 + n^2) = x^2 + y^2 + z^2 = r^2 \Rightarrow l^2 + m^2 + n^2 = 1$

or, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$, or, $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 2$



Note : \square If $OP = r$ and the co-ordinates of point P be (x, y, z) , then d.c.'s of line OP are $x/r, y/r, z/r$.

- \square Direction cosines of $\mathbf{r} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ are $\frac{a}{|\mathbf{r}|}, \frac{b}{|\mathbf{r}|}, \frac{c}{|\mathbf{r}|}$.
- \square Since $-1 \leq \cos x \leq 1, \forall x \in \mathbb{R}$, hence values of l, m, n are such real numbers which are not less than -1 and not greater than 1 . Hence d.c.'s $\in [-1, 1]$.
- \square The direction cosines of a line parallel to any co-ordinate axis are equal to the direction cosines of the co-ordinate axis.
- \square The number of lines which are equally inclined to the co-ordinate axes is 4.
- \square If l, m, n are the d.c.'s of a line, then the maximum value of $lmn = \frac{1}{3\sqrt{3}}$.

Important Tips

- \hookrightarrow The angles α, β, γ are called the direction angles of line AB .
- \hookrightarrow The d.c.'s of line BA are $\cos(\pi - \alpha), \cos(\pi - \beta)$ and $\cos(\pi - \gamma)$ i.e., $-\cos \alpha, -\cos \beta, -\cos \gamma$.
- \hookrightarrow Angles α, β, γ are not coplanar.
- \hookrightarrow $\alpha + \beta + \gamma$ is not equal to 360° as these angles do not lie in same plane.
- \hookrightarrow If $P(x, y, z)$ be a point in space such that $\mathbf{r} = \overrightarrow{OP}$ has d.c.'s l, m, n then $x = l|\mathbf{r}|, y = m|\mathbf{r}|, z = n|\mathbf{r}|$.
- \hookrightarrow Projection of a vector \mathbf{r} on the co-ordinate axes are $l|\mathbf{r}|, m|\mathbf{r}|, n|\mathbf{r}|$.
- \hookrightarrow $\mathbf{r} = |\mathbf{r}|(\hat{\mathbf{i}} + m\hat{\mathbf{j}} + n\hat{\mathbf{k}})$ and $\hat{\mathbf{r}} = \hat{\mathbf{i}} + m\hat{\mathbf{j}} + n\hat{\mathbf{k}}$

(2) Direction ratio

(i) Three numbers which are proportional to the direction cosines of a line are called the direction ratio of that line. If a, b, c are three numbers proportional to direction cosines l, m, n of a line, then a, b, c are called its direction ratios. They are also called direction numbers or direction components.

Hence by definition, we have $\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = k$ (say) $\Rightarrow l = ak, m = bk, n = ck$

$$\Rightarrow l^2 + m^2 + n^2 = (a^2 + b^2 + c^2) = k^2 \Rightarrow k = \pm \frac{1}{\sqrt{a^2 + b^2 + c^2}}$$

$$l = \pm \frac{a}{\sqrt{a^2 + b^2 + c^2}}, m = \pm \frac{b}{\sqrt{a^2 + b^2 + c^2}}, n = \pm \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

where the sign should be taken all positive or all negative.

Note : \square Direction ratios are not unique, whereas d.c.'s are unique. i.e., $a^2 + b^2 + c^2 \neq 1$

(ii) Let $\mathbf{r} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ be a vector. Then its d.r.'s are a, b, c


If a vector \mathbf{r} has d.r.'s a, b, c then $\mathbf{r} = \frac{|\mathbf{r}|}{\sqrt{a^2 + b^2 + c^2}}(a\mathbf{i} + b\mathbf{j} + c\mathbf{k})$

(iii) **D.c.'s and d.r.'s of a line joining two points :** The direction ratios of line PQ joining $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ are $x_2 - x_1 = a$, $y_2 - y_1 = b$ and $z_2 - z_1 = c$ (say).

Then direction cosines are,

$$l = \frac{(x_2 - x_1)}{\sqrt{\sum(x_2 - x_1)^2}}, m = \frac{(y_2 - y_1)}{\sqrt{\sum(x_2 - x_1)^2}}, n = \frac{(z_2 - z_1)}{\sqrt{\sum(x_2 - x_1)^2}}$$

$$\text{i.e., } l = \frac{x_2 - x_1}{PQ}, m = \frac{y_2 - y_1}{PQ}, n = \frac{z_2 - z_1}{PQ}.$$



Example: 7

A line makes the same angle θ with each of the x and z -axis. If the angle β , which it makes with y -axis, is such that $\sin^2 \beta = 3 \sin^2 \theta$, then $\cos^2 \theta$ equals [AIEEE 2004]

- (a) $\frac{2}{5}$ (b) $\frac{3}{5}$ (c) $\frac{1}{5}$ (d) $\frac{2}{3}$

Solution: (b)

We know that, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$. Since line makes angle θ with x and z -axis and angle β with y -axis.

$$\Rightarrow \cos^2 \theta + \cos^2 \beta + \cos^2 \theta = 1 \Rightarrow -(2 \cos^2 \theta - 1) = \cos^2 \beta \dots\dots(i)$$

$$\text{Given that } \sin^2 \beta = 3 \sin^2 \theta \dots\dots(ii)$$

$$\text{From (i) and (ii), } 1 = 3 \sin^2 \theta - 2 \cos^2 \theta + 1 \Rightarrow 0 = 3(1 - \cos^2 \theta) - 2 \cos^2 \theta \Rightarrow 5 \cos^2 \theta = 3 \Rightarrow \cos^2 \theta = 3/5$$

Example: 8

Direction cosines of the line that makes equal angles with the three axes in a space are

[Kurukshetra CEE 1995]

- (a) $\pm \frac{1}{3}, \pm \frac{1}{3}, \pm \frac{1}{3}$ (b) $\pm \frac{6}{7}, \pm \frac{2}{3}, \pm \frac{3}{7}$ (c) $\pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}$ (d) $\pm \sqrt{\frac{1}{7}}, \pm \sqrt{\frac{3}{14}}, \pm \sqrt{\frac{1}{14}}$

Solution: (c)

$$\because l^2 + m^2 + n^2 = 1 \Rightarrow \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\text{Now, } \alpha = \beta = \gamma$$

$$\Rightarrow 3 \cos^2 \alpha = 1 \Rightarrow \cos \alpha = \pm 1/\sqrt{3} \text{ i.e., } l = m = n = \pm 1/\sqrt{3}.$$

$$\text{Hence required d.c.'s are } \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}.$$

Example: 9

A line which makes angle 60° with y -axis and z -axis, then the angle which it makes with x -axis is

[Rajasthan PET 2002; DCE 1996]

- (a) 45° (b) 60° (c) 75° (d) 30°

Solution: (a)

$$\text{Given that } \beta = \gamma = 60^\circ \text{ i.e., } m = \cos \beta = \cos 60^\circ = 1/2, n = \cos \gamma = \cos 60^\circ = 1/2$$

$$\because l^2 + m^2 + n^2 = 1 \Rightarrow l^2 = 1 - \frac{1}{4} - \frac{1}{4} = \frac{1}{2} \Rightarrow l = \frac{1}{\sqrt{2}} \Rightarrow \cos \alpha = \frac{1}{\sqrt{2}} \Rightarrow \alpha = 45^\circ$$

Example: 10

A line passes through the points $(6, -7, -1)$ and $(2, -3, 1)$. The direction cosines of line, so directed that the angle made by it with the positive direction of x -axis is acute, are

- (a) $\frac{2}{3}, \frac{-2}{3}, \frac{-1}{3}$ (b) $\frac{-2}{3}, \frac{2}{3}, \frac{1}{3}$ (c) $\frac{2}{3}, \frac{-2}{3}, \frac{1}{3}$ (d) $\frac{2}{3}, \frac{2}{3}, \frac{1}{3}$

Solution: (a)

Let l, m, n be the d.c.'s of a given line.

Then, as it makes an acute angle with x -axis, therefore $l > 0$.

$$\text{Direction ratios} = 4, -4, -2 \text{ or } 2, -2, -1 \text{ and Direction cosines} = \frac{2}{3}, \frac{-2}{3}, \frac{-1}{3}.$$

Example: 11

If the direction cosines of a line are $\left(\frac{1}{c}, \frac{1}{c}, \frac{1}{c}\right)$, then

[DCE 2000; Pb. CET 1996, 98]

(a) $c > 0$

(b) $c = \pm\sqrt{3}$

(c) $0 < c < 1$

(d) $c > 2$

Solution: (b) We know that $l^2 + m^2 + n^2 = 1 \Rightarrow \frac{1}{c^2} + \frac{1}{c^2} + \frac{1}{c^2} = 1 \Rightarrow \frac{3}{c^2} = 1 \Rightarrow c = \pm\sqrt{3}$.

Example: 12 If \mathbf{r} is a vector of magnitude 21 and has d.r.'s 2, -3, 6. Then \mathbf{r} is equal to

(a) $6\mathbf{i} - 9\mathbf{j} + 18\mathbf{k}$

(b) $6\mathbf{i} + 9\mathbf{j} + 18\mathbf{k}$

(c) $6\mathbf{i} - 9\mathbf{j} - 18\mathbf{k}$

(d) $6\mathbf{i} + 9\mathbf{j} - 18\mathbf{k}$


Solution: (a) D.r.'s of \mathbf{r} are 2, -3, 6. Therefore, its d.c.'s are $l = \frac{2}{7}, m = \frac{-3}{7}, n = \frac{6}{7}$

$$\therefore \mathbf{r} = |\mathbf{r}| (\hat{\mathbf{i}} + m\hat{\mathbf{j}} + n\hat{\mathbf{k}}) = 21 \left[\frac{2}{7}\hat{\mathbf{i}} - \frac{3}{7}\hat{\mathbf{j}} + \frac{6}{7}\hat{\mathbf{k}} \right] = 6\mathbf{i} - 9\mathbf{j} + 18\mathbf{k}.$$

7.7 Projection

(1) **Projection of a point on a line :** The projection of a point P on a line AB is the foot N of the perpendicular PN from P on the line AB .


N is also the same point where the line AB meets the plane through P and perpendicular to AB .





(2) **Projection of a segment of a line on another line and its length :** The projection of the segment AB of a given line on another line CD is the segment $A'B'$ of CD where A' and B' are the projections of the points A and B on the line CD .

The length of the projection $A'B'$.

$$A'B' = AN = AB \cos \theta$$



(3) **Projection of a line joining the points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ on another line whose direction cosines are l, m and n :** Let PQ be a line segment where $P \equiv (x_1, y_1, z_1)$ and $Q = (x_2, y_2, z_2)$ and AB be a given line with d.c.'s as l, m, n . If the line segment PQ makes angle θ with the line AB , then



$$\begin{aligned} \text{Projection of } PQ \text{ is } P'Q' &= PQ \cos \theta = (x_2 - x_1) \cos \alpha + (y_2 - y_1) \cos \beta + (z_2 - z_1) \cos \gamma \\ &= (x_2 - x_1)l + (y_2 - y_1)m + (z_2 - z_1)n \end{aligned}$$

Important Tips

☞ For x -axis, $l = 1, m = 0, n = 0$.

Hence, projection of PQ on x -axis $= x_2 - x_1$, Projection of PQ on y -axis $= y_2 - y_1$ and Projection of PQ on z -axis $= z_2 - z_1$

☞ If P is a point (x_1, y_1, z_1) , then projection of OP on a line whose direction cosines are l, m, n , is $l_1x_1 + m_1y_1 + n_1z_1$, where O is the origin.

☞ If l_1, m_1, n_1 and l_2, m_2, n_2 are the d.c.'s of two concurrent lines, then the d.c.'s of the lines bisecting the angles between them are proportional to $l_1 \pm l_2, m_1 \pm m_2, n_1 \pm n_2$.

Example: 13 If A, B, C, D are the points $(3, 4, 5)$ $(4, 6, 3)$, $(-1, 2, 4)$ and $(1, 0, 5)$, then the projection of CD on AB is

[Orissa JEE 2002; Rajasthan PET 2002]

- (a) $\frac{3}{4}$ (b) $\frac{-4}{3}$ (c) $\frac{3}{5}$ (d) None of these

Solution: (b) Let l, m, n be the direction cosines of AB

$$\text{Then } l = \frac{4-3}{\sqrt{(4-3)^2 + (6-4)^2 + (3-5)^2}} = \frac{1}{3}, m = \frac{6-4}{3} = \frac{2}{3}. \text{ Similarly } n = \frac{-2}{3}$$

$$\therefore \text{ The projection of } CD \text{ on } AB = \left[1 - (-1) \left(\frac{1}{3} \right) \right] + [0 - 2] \left(\frac{2}{3} \right) + [5 - 4] \left(-\frac{2}{3} \right) = \frac{2}{3} - \frac{4}{3} + \left(-\frac{2}{3} \right) = -\frac{4}{3}$$

Example: 14 The projection of a line on co-ordinate axes are 2, 3, 6. Then the length of the line is

[Orissa JEE 2002]

- (a) 7 (b) 5 (c) 1 (d) 11

Solution: (b) Let AB be the line and its direction cosines be $\cos \alpha, \cos \beta, \cos \gamma$. Then the projection of line AB on the co-ordinate axes are $AB \cos \alpha, AB \cos \beta, AB \cos \gamma$. $\therefore AB \cos \alpha = 2, AB \cos \beta = 3, AB \cos \gamma = 6$


$$\Rightarrow AB^2(\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma) = 2^2 + 3^2 + 6^2 = 49 \Rightarrow AB^2(1) = 49 \Rightarrow AB = 7$$

7.8 Angle between Two lines

(1) Cartesian form : Let θ be the angle between two straight lines AB and AC whose direction cosines are l_1, m_1, n_1 and l_2, m_2, n_2 respectively, is given by $\cos \theta = l_1 l_2 + m_1 m_2 + n_1 n_2$.

If direction ratios of two lines a_1, b_1, c_1 and a_2, b_2, c_2 are given, then angle

between two lines is given by $\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \cdot \sqrt{a_2^2 + b_2^2 + c_2^2}}$.



Particular results : We have, $\sin^2 \theta = 1 - \cos^2 \theta = (l_1^2 + m_1^2 + n_1^2)(l_2^2 + m_2^2 + n_2^2) - (l_1 l_2 + m_1 m_2 + n_1 n_2)^2$
 $= (l_1 m_2 - l_2 m_1)^2 + (m_1 n_2 - m_2 n_1)^2 + (n_1 l_2 - n_2 l_1)^2$

$$\Rightarrow \sin \theta = \pm \sqrt{\sum (l_1 m_2 - l_2 m_1)^2}, \text{ which is known as Lagrange's identity.}$$

The value of $\sin \theta$ can easily be obtained by the following form. $\sin \theta = \sqrt{\begin{vmatrix} l_1 & m_1 \\ l_2 & m_2 \end{vmatrix}^2 + \begin{vmatrix} m_1 & n_1 \\ m_2 & n_2 \end{vmatrix}^2 + \begin{vmatrix} n_1 & l_1 \\ n_2 & l_2 \end{vmatrix}^2}$

When d.r.'s of the lines are given if a_1, b_1, c_1 and a_2, b_2, c_2 are d.r.'s of given two lines, then angle θ between them

is given by $\sin \theta = \frac{\sqrt{\sum (a_1 b_2 - a_2 b_1)^2}}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$

Condition of perpendicularity : If the given lines are perpendicular, then $\theta = 90^\circ$ i.e. $\cos \theta = 0$

$$\Rightarrow l_1 l_2 + m_1 m_2 + n_1 n_2 = 0 \text{ or } a_1 a_2 + b_1 b_2 + c_1 c_2 = 0$$

Condition of parallelism : If the given lines are parallel, then $\theta = 0^\circ$ i.e. $\sin \theta = 0$

$$\Rightarrow (l_1 m_2 - l_2 m_1)^2 + (m_1 n_2 - m_2 n_1)^2 + (n_1 l_2 - n_2 l_1)^2 = 0, \text{ which is true, only when}$$

$$l_1 m_2 - l_2 m_1 = 0, m_1 n_2 - m_2 n_1 = 0 \text{ and } n_1 l_2 - n_2 l_1 = 0$$

$$\Rightarrow \frac{l_1}{l_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2}.$$

$$\text{Similarly, } \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}.$$

Note : \square The angle between any two diagonals of a cube is $\cos^{-1}\left(\frac{1}{3}\right)$.

\square The angle between a diagonal of a cube and the diagonal of a faces of the cube is $\cos^{-1}\left(\sqrt{\frac{2}{3}}\right)$.

\square If a straight line makes angles $\alpha, \beta, \gamma, \delta$ with the diagonals of a cube, then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{4}{3}$$

\square If the edges of a rectangular parallelopiped be a, b, c , then the angles between the two diagonals are

$$\cos^{-1}\left[\frac{\pm a^2 \pm b^2 \pm c^2}{a^2 + b^2 + c^2}\right]$$

(2) **Vector form** : Let the vector equations of two lines be $\mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b}_1$ and $\mathbf{r} = \mathbf{a}_2 + \lambda \mathbf{b}_2$

As the lines are parallel to the vectors \mathbf{b}_1 and \mathbf{b}_2 respectively, therefore angle between the lines is same as the angle between the vectors \mathbf{b}_1 and \mathbf{b}_2 . Thus if θ is the angle between the given lines, then $\cos \theta = \frac{\mathbf{b}_1 \cdot \mathbf{b}_2}{|\mathbf{b}_1| |\mathbf{b}_2|}$.

Note : \square If the lines are perpendicular, then $\mathbf{b}_1 \cdot \mathbf{b}_2 = 0$.

\square If the lines are parallel, then \mathbf{b}_1 and \mathbf{b}_2 are parallel, therefore $\mathbf{b}_1 = \lambda \mathbf{b}_2$ for some scalar λ .

Example: 15 If d.c.'s of two lines are proportional to $(2, 3, -6)$ and $(3, -4, 5)$, then the acute angle between them is [MP PET 2003]

- (a) $\cos^{-1}\left(\frac{49}{36}\right)$ (b) $\cos^{-1}\left(\frac{18\sqrt{2}}{35}\right)$ (c) 90° (d) $\cos^{-1}\left(\frac{18}{35}\right)$

Solution: (b) D.c.'s of two lines are proportional to $(2, 3, -6)$ and $(3, -4, 5)$
i.e. d.r.'s are $(2, 3, -6)$ and $(3, -4, 5)$

$$\therefore \cos \theta = \frac{2(3) + 3(-4) + (-6)5}{\sqrt{2^2 + 3^2 + (-6)^2} \sqrt{3^2 + (-4)^2 + 5^2}} = \frac{6 - 12 - 30}{\sqrt{49} \cdot \sqrt{50}} = \frac{-36}{7.5\sqrt{2}} \Rightarrow \cos \theta = \frac{-18\sqrt{2}}{35}$$

$$\text{Taking acute angle, } \theta = \cos^{-1}\left(\frac{18\sqrt{2}}{35}\right)$$

Example: 16 If the direction ratio of two lines are given by $3lm - 4ln + mn = 0$ and $l + 2m + 3n = 0$, then the angle between the lines is [EAMCET 2003]

- (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$

Solution: (a) We have, $l + 2m + 3n = 0$ (i)
 $3lm - 4ln + mn = 0$ (ii)

From equation (i), $l = -(2m + 3n)$

Putting the value of l in equation (ii)

$$\Rightarrow 3(-2m - 3n)m + mn - 4(-2m - 3n)n = 0 \Rightarrow -6m^2 - 9mn + mn + 8mn + 12n^2 = 0 \Rightarrow 6m^2 - 12n^2 = 0$$

$$\Rightarrow m^2 - 2n^2 = 0 \Rightarrow m + \sqrt{2}n = 0 \text{ or } m - \sqrt{2}n = 0$$

$$l + 2m + 3n = 0 \quad \text{.....(i)} \quad 0.l + m + \sqrt{2}n = 0 \quad \text{.....(iii)} \quad 0.l + m - \sqrt{2}n = 0 \quad \text{.....(iv)}$$

$$\text{From equation (i) and equation (iii), } \frac{l}{2\sqrt{2} - 3} = \frac{m}{-\sqrt{2}} = \frac{n}{1}$$

$$\text{From equation (i) and equation (iv), } \frac{l}{-2\sqrt{2} - 3} = \frac{m}{\sqrt{2}} = \frac{n}{1}$$

Thus, the direction ratios of two lines are $2\sqrt{2} - 3, -\sqrt{2}, 1$ and $-2\sqrt{2} - 3, \sqrt{2}, 1$

$(l_1, m_1, n_1) = (2\sqrt{2} - 3, -\sqrt{2}, 1)$, $(l_2, m_2, n_2) = (-2\sqrt{2} - 3, \sqrt{2}, 1)$, $l_1 l_2 + m_1 m_2 + n_1 n_2 = 0$. Hence, the angle between them $\pi/2$.

Example: 17

If a line makes angles $\alpha, \beta, \gamma, \delta$ with four diagonals of a cube, then the value of $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + \sin^2 \delta$ is

- (a) $\frac{4}{3}$ (b) 1 (c) $\frac{8}{3}$ (d) $\frac{7}{3}$

Solution: (c)

Let side of the cube = a

Then OG, BE and AD, CF will be four diagonals.

d.r.'s of $OG = a, a, a = 1, 1, 1$

d.r.'s of $BE = -a, -a, a = 1, 1, -1$

d.r.'s of $AD = -a, a, a = -1, 1, 1$


d.r.'s of $CF = a, -a, a = 1, -1, 1$

Let d.r.'s of line be l, m, n . Therefore angle between line and diagonal

$$\cos \alpha = \frac{l+m+n}{\sqrt{3}}, \cos \beta = \frac{l+m-n}{\sqrt{3}}, \cos \gamma = \frac{-l+m+n}{\sqrt{3}}, \cos \delta = \frac{l-m+n}{\sqrt{3}}$$

$$\Rightarrow \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{1}{3} [(l+m+n)^2 + (l+m-n)^2 + (-l+m+n)^2 + (l-m+n)^2] = \frac{4}{3}$$

$$\Rightarrow \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + \sin^2 \delta = \frac{8}{3}$$

**Example: 18**

If l_1, m_1, n_1 and l_2, m_2, n_2 are d.c.'s of two lines inclined to each other at an angle θ , then the d.c.'s of the internal bisectors of angle between these lines are

- (a) $\frac{l_1+l_2}{2 \sin \theta / 2}, \frac{m_1+m_2}{2 \sin \theta / 2}, \frac{n_1+n_2}{2 \sin \theta / 2}$ (b) $\frac{l_1+l_2}{2 \cos \theta / 2}, \frac{m_1+m_2}{2 \cos \theta / 2}, \frac{n_1+n_2}{2 \cos \theta / 2}$
 (c) $\frac{l_1-l_2}{2 \sin \theta / 2}, \frac{m_1-m_2}{2 \sin \theta / 2}, \frac{n_1-n_2}{2 \sin \theta / 2}$ (d) $\frac{l_1-l_2}{2 \cos \theta / 2}, \frac{m_1-m_2}{2 \cos \theta / 2}, \frac{n_1-n_2}{2 \cos \theta / 2}$

Solution: (b)


Let OA and OB be two lines.

D.c.'s of OA is (l_1, m_1, n_1) and OB is (l_2, m_2, n_2) .

Let $OA = OB = 1$.

Then the co-ordinates of A and B are (l_1, m_1, n_1) and (l_2, m_2, n_2) .

Let OC be the bisector of $\angle AOB$.



Then C is the mid-point of AB and so its co-ordinates are $\left(\frac{l_1+l_2}{2}, \frac{m_1+m_2}{2}, \frac{n_1+n_2}{2} \right)$.

\therefore D.r.'s of line OC are $\left(\frac{l_1+l_2}{2}, \frac{m_1+m_2}{2}, \frac{n_1+n_2}{2} \right)$

$$\begin{aligned} \text{We have, } OC &= \sqrt{\left(\frac{l_1+l_2}{2} \right)^2 + \left(\frac{m_1+m_2}{2} \right)^2 + \left(\frac{n_1+n_2}{2} \right)^2} = \frac{1}{2} \sqrt{l_1^2 + m_1^2 + n_1^2 + l_2^2 + m_2^2 + n_2^2 + 2(l_1 l_2 + m_1 m_2 + n_1 n_2)} \\ &= \frac{1}{2} \sqrt{1+1+2 \cos \theta} = \frac{1}{2} \sqrt{2(2 \cos^2 \theta / 2)} = \cos \theta / 2. \end{aligned}$$

D.r.'s of line OC are $\frac{l_1+l_2}{2 \cos \theta / 2}, \frac{m_1+m_2}{2 \cos \theta / 2}, \frac{n_1+n_2}{2 \cos \theta / 2}$.

Example: 19

The angle between the lines $\mathbf{r} = (4\mathbf{i} - \mathbf{j}) + s(2\mathbf{i} + \mathbf{j} - 3\mathbf{k})$ and $\mathbf{r} = (\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + t(\mathbf{i} - 3\mathbf{j} + 2\mathbf{k})$ is

[Tamilnadu (Engg.) 2002]

- (a) $\frac{3\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{2\pi}{3}$ (d) $\frac{\pi}{6}$

Solution: (b)

We have, $\mathbf{r} = (4\mathbf{i} - \mathbf{j}) + s(2\mathbf{i} + \mathbf{j} - 3\mathbf{k})$ and $\mathbf{r} = (\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + t(\mathbf{i} - 3\mathbf{j} + 2\mathbf{k})$

We know that, $\cos \theta = \frac{\mathbf{b}_1 \cdot \mathbf{b}_2}{|\mathbf{b}_1| |\mathbf{b}_2|}$, $\cos \theta = \frac{(2\mathbf{i} + \mathbf{j} - 3\mathbf{k}) \cdot (\mathbf{i} - 3\mathbf{j} + 2\mathbf{k})}{\sqrt{4+1+9} \sqrt{1+9+4}} = \frac{2-3-6}{\sqrt{14} \cdot \sqrt{14}} = \frac{-7}{14}$, $\cos \theta = -\frac{1}{2}$

Hence, acute angle $\theta = \cos^{-1} \left(\frac{1}{2} \right)$ i.e. $\theta = \frac{\pi}{3}$

The Straight Line


7.9 Straight line in Space

Every equation of the first degree represents a plane. Two equations of the first degree are satisfied by the co-ordinates of every point on the line of intersection of the planes represented by them. Therefore, the two equations together represent that line. Therefore $ax + by + cz + d = 0$ and $a'x + b'y + c'z + d' = 0$ together represent a straight line.

(1) Equation of a line passing through a given point

(i) **Cartesian form or symmetrical form** : Cartesian equation of a straight line passing through a fixed point (x_1, y_1, z_1) and having direction ratios a, b, c is $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$.

(ii) **Vector form** : Vector equation of a straight line passing through a fixed point with position vector \mathbf{a} and parallel to a given vector \mathbf{b} is $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$.



Important Tips

- ☞ The parametric equations of the line $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$ are $x = x_1 + a\lambda, y = y_1 + b\lambda, z = z_1 + c\lambda$, where λ is the parameter.
- ☞ The co-ordinates of any point on the line $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$ are $(x_1 + a\lambda, y_1 + b\lambda, z_1 + c\lambda)$, where $\lambda \in \mathbb{R}$.
- ☞ Since the direction cosines of a line are also direction ratios, therefore equation of a line passing through (x_1, y_1, z_1) and having direction cosines l, m, n is $\frac{x - x_1}{l} = \frac{y - y_1}{m} = \frac{z - z_1}{n}$.
- ☞ Since x, y and z -axes pass through the origin and have direction cosines $1, 0, 0; 0, 1, 0$ and $0, 0, 1$ respectively. Therefore, the equations are x -axis : $\frac{x - 0}{1} = \frac{y - 0}{0} = \frac{z - 0}{0}$ or $y = 0$ and $z = 0$.
 y -axis : $\frac{x - 0}{0} = \frac{y - 0}{1} = \frac{z - 0}{0}$ or $x = 0$ and $z = 0$; z -axis : $\frac{x - 0}{0} = \frac{y - 0}{0} = \frac{z - 0}{1}$ or $x = 0$ and $y = 0$.
- ☞ In the symmetrical form of equation of a line, the coefficients of x, y, z are unity.

7.10 Equation of Line passing through Two given points

(i) **Cartesian form** : If $A(x_1, y_1, z_1), B(x_2, y_2, z_2)$ be two given points, the equations to the line AB are

$$\frac{x - x_1}{x_2 - x_1} = \frac{y - y_1}{y_2 - y_1} = \frac{z - z_1}{z_2 - z_1}$$


The co-ordinates of a variable point on AB can be expressed in terms of a parameter λ in the form

$$x = \frac{\lambda x_2 + x_1}{\lambda + 1}, y = \frac{\lambda y_2 + y_1}{\lambda + 1}, z = \frac{\lambda z_2 + z_1}{\lambda + 1}$$

λ being any real number different from -1 . In fact, (x, y, z) are the co-ordinates of the point which divides the join of A and B in the ratio $\lambda : 1$.

(ii) **Vector form** : The vector equation of a line passing through two points with position vectors \mathbf{a} and \mathbf{b} is

$$\mathbf{r} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$$



7.11 Changing Unsymmetrical form to Symmetrical form

The unsymmetrical form of a line $ax + by + cz + d = 0, a'x + b'y + c'z + d' = 0$

Can be changed to symmetrical form as follows :
$$\frac{x - \frac{bd' - b'd}{ab' - a'b}}{bc' - b'c} = \frac{y - \frac{da' - d'a}{ab' - a'b}}{ca' - c'a} = \frac{z}{ab' - a'b}$$

Example: 20

The equation to the straight line passing through the points (4, -5, -2) and (-1, 5, 3) is

[MP PET 2003]

(a) $\frac{x-4}{1} = \frac{y+5}{-2} = \frac{z+2}{-1}$ (b) $\frac{x+1}{1} = \frac{y-5}{2} = \frac{z-3}{-1}$ (c) $\frac{x}{-1} = \frac{y}{5} = \frac{z}{3}$ (d) $\frac{x}{4} = \frac{y}{-5} = \frac{z}{-2}$

Solution: (a)

We know that equation of a straight line is of the form $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$

D.r.'s of the line = $(-1-4, 5+5, 3+2)$ i.e., $(-5, 10, 5)$ or $(-1, 2, 1)$.

Hence the equation is $\frac{x-4}{-1} = \frac{y+5}{2} = \frac{z+2}{1}$ i.e., $\frac{x-4}{1} = \frac{y+5}{-2} = \frac{z+2}{-1}$

Example: 21

The d.c.'s of the line $6x - 2 = 3y + 1 = 2z - 2$ are

(a) $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$ (b) $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$ (c) 1, 2, 3 (d) None of these

Solution: (b)

We have $6x - 2 = 3y + 1 = 2z - 2 \Rightarrow \frac{6x - (2/6)}{1} = \frac{3y + (1/3)}{1} = \frac{2(z-1)}{1}$

$\Rightarrow \frac{x - (1/3)}{1/6} = \frac{y + (1/3)}{1/3} = \frac{z-1}{1/2} \Rightarrow \frac{x - (1/3)}{1} = \frac{y + (1/3)}{2} = \frac{z-1}{3}$

d.r.'s of line are (1, 2, 3). Hence d.c.'s of line are $(1/\sqrt{14}, 2/\sqrt{14}, 3/\sqrt{14})$

Example: 22

The vector equation of line through the point A(3, 4, -7) and B(1, -1, 6) is

[Pb. CET 1999]

(a) $\mathbf{r} = (3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k}) + \lambda(\mathbf{i} - \mathbf{j} + 6\mathbf{k})$ (b) $\mathbf{r} = (\mathbf{i} - \mathbf{j} + 6\mathbf{k}) + \lambda(3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k})$
 (c) $\mathbf{r} = (3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k}) + \lambda(-2\mathbf{i} - 5\mathbf{j} + 13\mathbf{k})$ (d) $\mathbf{r} = (\mathbf{i} - \mathbf{j} + 6\mathbf{k}) + \lambda(4\mathbf{i} + 3\mathbf{j} - \mathbf{k})$

Solution: (c)

Position vector of A is $\mathbf{a} = 3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k}$ and that of B is $\mathbf{b} = \mathbf{i} - \mathbf{j} + 6\mathbf{k}$

We know that equation of line in vector form, $\mathbf{r} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$, $\mathbf{r} = (3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k}) + \lambda(-2\mathbf{i} - 5\mathbf{j} + 13\mathbf{k})$.

7.12 Angle between Two lines

Let the cartesian equations of the two lines be

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \quad \dots(i) \quad \text{and} \quad \frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2} \quad \dots(ii)$$

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

Condition of perpendicularity : If the lines are perpendicular, then $a_1 a_2 + b_1 b_2 + c_1 c_2 = 0$

Condition of parallelism : If the lines are parallel, then $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$.

332 Three Dimensional Co-ordinate Geometry

Example: 23 If the lines $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$ and $\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5}$ are at right angles, then $k =$

[MP PET 1997, 2001; DCE 1997, 99]

- (a) -10 (b) $10/7$ (c) $-10/7$ (d) $-7/10$

Solution: (a) We have $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$ and $\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5}$

Since lines are \perp to each other. So, $a_1a_2 + b_1b_2 + c_1c_2 = 0$

$$(-3)(3k) + (2k)(1) + (2)(-5) = 0 \Rightarrow -9k + 2k - 10 = 0 \Rightarrow -7k = 10 \Rightarrow k = -10/7.$$

Example: 24 The lines $x = ay + b$, $z = cy + d$ and $x = a'y + b'$, $z = c'y + d'$ are perpendicular to each other, if [IIT 1984; AIEEE 2003]

- (a) $ad' + cc' = 1$ (b) $ad' + cc' = -1$ (c) $ac + d'c' = 1$ (d) $ac + d'c' = -1$

Solution: (b) We have, $x = ay + b$, $z = cy + d$

$$\frac{x-b}{a} = y, \frac{z-d}{c} = y \Rightarrow \frac{x-b}{a} = \frac{y-0}{1} = \frac{z-d}{c} \quad \dots\dots(i)$$

and $x = a'y + b'$, $z = c'y + d'$

$$\frac{x-b'}{a'} = y, \frac{z-d'}{c'} = y \Rightarrow \frac{x-b'}{a'} = \frac{y-0}{1} = \frac{z-d'}{c'} \quad \dots\dots(ii)$$

\therefore Given, lines (i) and (ii) are perpendicular

$$\therefore a(a') + 1(1) + c(c') = 0, \quad ad' + cc' = -1$$

Example: 25 The direction ratio of the line which is perpendicular to the lines $\frac{x-7}{2} = \frac{y+17}{-3} = \frac{z-6}{1}$ and $\frac{x+5}{1} = \frac{y+3}{2} = \frac{z-4}{-2}$ are

[Pb. CET 1999]

- (a) $\langle 4, 5, 7 \rangle$ (b) $\langle 4, -5, 7 \rangle$ (c) $\langle 4, -5, -7 \rangle$ (d) $\langle -4, 5, 7 \rangle$

Solution: (a) Let d.r.'s of line be l, m, n .

\therefore line is perpendicular to given line

$$\therefore 2l - 3m + n = 0 \quad \dots\dots(i)$$

$$l + 2m - 2n = 0 \quad \dots\dots(ii)$$

From equation (i) and (ii)

$$\frac{l}{6-2} = \frac{m}{1+4} = \frac{n}{4+3} \text{ or } \frac{l}{4} = \frac{m}{5} = \frac{n}{7}. \text{ Hence, d.r.'s of line } (\langle 4, 5, 7 \rangle)$$

7.13 Reduction of Cartesian form of the Equation of a line to Vector form and Vice versa

Cartesian to vector : Let the Cartesian equation of a line be $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} \quad \dots\dots(i)$

This is the equation of a line passing through the point $A(x_1, y_1, z_1)$ and having direction ratios a, b, c . In vector form this means that the line passes through point having position vector $\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}$ and is parallel to the vector $\mathbf{m} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$. Thus, the vector form of (i) is $\mathbf{r} = \mathbf{a} + \lambda\mathbf{m}$ or $\mathbf{r} = (x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}) + \lambda(a\mathbf{i} + b\mathbf{j} + c\mathbf{k})$, where λ is a parameter.

Vector to cartesian : Let the vector equation of a line be $\mathbf{r} = \mathbf{a} + \lambda\mathbf{m} \quad \dots\dots(ii)$

Where $\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}$, $\mathbf{m} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ and λ is a parameter.

To reduce (ii) to Cartesian form we put $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and equate the coefficients of \mathbf{i}, \mathbf{j} and \mathbf{k} as discussed below.

Putting $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, $\mathbf{a} = x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}$ and $\mathbf{m} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ in (ii), we obtain

$$x\mathbf{i} + y\mathbf{j} + z\mathbf{k} = (x_1\mathbf{i} + y_1\mathbf{j} + z_1\mathbf{k}) + \lambda(a\mathbf{i} + b\mathbf{j} + c\mathbf{k})$$

Equating coefficients of \mathbf{i}, \mathbf{j} and \mathbf{k} , we get $x = x_1 + a\lambda, y = y_1 + b\lambda, z = z_1 + c\lambda$ or $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} = \lambda$

Example: 26 The cartesian equations of a line are $6x - 2 = 3y + 1 = 2z - 2$. The vector equation of the line is

- (a) $\mathbf{r} = \left(\frac{1}{3}\mathbf{i} - \frac{1}{3}\mathbf{j} + \mathbf{k}\right) + \lambda(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ (b) $\mathbf{r} = (3\mathbf{i} - 3\mathbf{j} + \mathbf{k}) + \lambda(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$
 (c) $\mathbf{r} = (\mathbf{i} + \mathbf{j} + \mathbf{k}) + \lambda(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ (d) None of these

Solution: (a) The given line is $6x - 2 = 3y + 1 = 2z - 2 \Rightarrow \frac{x - 1/3}{1} = \frac{y + 1/3}{2} = \frac{z - 1}{3}$

This show that the given line passes through $(1/3, -1/3)$ and has direction ratio 1, 2, 3.

Position vector $\mathbf{a} = \frac{1}{3}\mathbf{i} - \frac{1}{3}\mathbf{j} + \mathbf{k}$ and is parallel to vector $\mathbf{b} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. Hence, $\mathbf{r} = \left(\frac{1}{3}\mathbf{i} - \frac{1}{3}\mathbf{j} + \mathbf{k}\right) + \lambda(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$.

7.14 Intersection of Two lines

Determine whether two lines intersect or not. In case they intersect, the following algorithm is used to find their point of intersection.

Algorithm for cartesian form : Let the two lines be $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1}$ (i)

And $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2}$ (ii)

Step I : Write the co-ordinates of general points on (i) and (ii). The co-ordinates of general points on (i) and (ii) are given by $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1} = \lambda$ and $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2} = \mu$ respectively.

i.e., $(a_1\lambda + x_1, b_1\lambda + y_1, c_1\lambda + z_1)$ and $(a_2\mu + x_2, b_2\mu + y_2, c_2\mu + z_2)$

Step II : If the lines (i) and (ii) intersect, then they have a common point.

$a_1\lambda + x_1 = a_2\mu + x_2, b_1\lambda + y_1 = b_2\mu + y_2$ and $c_1\lambda + z_1 = c_2\mu + z_2$.

Step III : Solve any two of the equations in λ and μ obtained in step II. If the values of λ and μ satisfy the third equation, then the lines (i) and (ii) intersect, otherwise they do not intersect.

Step IV : To obtain the co-ordinates of the point of intersection, substitute the value of λ (or μ) in the co-ordinates of general point (s) obtained in step I.

Example: 27 If the line $\frac{x - 1}{2} = \frac{y + 1}{3} = \frac{z - 1}{4}$ and $\frac{x - 3}{1} = \frac{y - k}{2} = \frac{z}{1}$ intersect, then $k =$ [IIT Screening 2004]

- (a) 2/9 (b) 9/2 (c) 0 (d) -1

Solution: (b) We have, $\frac{x - 1}{2} = \frac{y + 1}{3} = \frac{z - 1}{4} = r_1$ (Let)

$x = 2r_1 + 1, y = 3r_1 - 1, z = 4r_1 + 1$ i.e. point is $(2r_1 + 1, 3r_1 - 1, 4r_1 + 1)$ and $\frac{x - 3}{1} = \frac{y - k}{2} = \frac{z}{1} = r_2$ (Let)

i.e. point is $(r_2 + 3, 2r_2 + k, r_2)$.

If the lines are intersecting, then they have a common point.

$\Rightarrow 2r_1 + 1 = r_2 + 3, 3r_1 - 1 = 2r_2 + k, 4r_1 + 1 = r_2$

On solving, $r_1 = -3/2, r_2 = -5$

Hence, $k = 9/2$.

Example: 28 A line with direction cosines proportional to 2, 1, 2 meets each of the lines $x = y + a = z$ and $x + a = 2y = 2z$. The co-ordinates of each of the points of intersection are given by [AIEEE 2004]

- (a) $(2a, 3a, 3a)$ (b) $(2a, a, a)$ (c) $(3a, 2a, 3a)$ (d) $(3a, 3a, 3a)$

334 Three Dimensional Co-ordinate Geometry

Solution: (b) Given lines are $\frac{x}{1} = \frac{y+a}{1} = \frac{z}{1} = \lambda$ (say) \therefore Point is $P(\lambda, \lambda - a, \lambda)$

and $\frac{x+a}{1} = \frac{y}{1/2} = \frac{z}{1/2}$ i.e. $\frac{x+a}{2} = \frac{y}{1} = \frac{z}{1} = \mu$ (say)

\therefore Point $Q(2\mu - a, \mu, \mu)$


Since d.r.'s of given lines are 2, 1, 2 and d.r.'s of $PQ = (2\mu - a - \lambda, \mu - \lambda + a, \mu - \lambda)$

According to question, $\frac{2\mu - a - \lambda}{2} = \frac{\mu - \lambda + a}{1} = \frac{\mu - \lambda}{2}$

Then $\lambda = 3a$, $\mu = a$. Therefore, points of intersection are $P(3a, 2a, 3a)$ and $Q(a, a, a)$.

Alternative method : Check by option $x = y + a = z$ i.e. $3a = 2a + a = 3a$

$\Rightarrow a = a = a$ and $x + a = 2y = 2z$ i.e. $a + a = 2a = 2a \Rightarrow a = a = a$. Hence (b) is correct.




7.15 Foot of perpendicular from a point $A(\alpha, \beta, \gamma)$ to the line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$

(1) Cartesian form

Foot of perpendicular from a point $A(\alpha, \beta, \gamma)$ to the line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$: If P be the foot of perpendicular, then P is $(lr + x_1, mr + y_1, nr + z_1)$. Find the direction ratios of AP and apply the condition of perpendicularity of AP and the given line. This will give the value of r and hence the point P which is foot of perpendicular.


Length and equation of perpendicular : The length of the perpendicular is the distance AP and its equation is the line joining two known points A and P .



Note : \square The length of the perpendicular is the perpendicular

distance of given point from that line.


Reflection or image of a point in a straight line : If the perpendicular PL from point P on the given line be produced to Q such that $PL = QL$, then Q is known as the image or reflection of P in the given line. Also, L is the foot of the perpendicular or the projection of P on the line.



(2) Vector form

Perpendicular distance of a point from a line : Let L is the foot of perpendicular drawn from $P(\vec{\alpha})$ on the line $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$. Since \mathbf{r} denotes the position vector of L , So, let the position vector of L be $\mathbf{a} + \lambda \mathbf{b}$.


$$\text{Then } \vec{PL} = \mathbf{a} - \vec{\alpha} + \lambda \mathbf{b} = (\mathbf{a} - \vec{\alpha}) - \left(\frac{(\mathbf{a} - \vec{\alpha}) \cdot \mathbf{b}}{|\mathbf{b}|^2} \right) \mathbf{b}$$



The length PL , is the magnitude of \vec{PL} , and required length of perpendicular.

Image of a point in a straight line : Let $Q(\vec{\beta})$ is the image of P in $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$

Then, $\vec{\beta} = 2\mathbf{a} - \left(\frac{2(\mathbf{a} - \vec{\alpha}) \cdot \mathbf{b}}{|\mathbf{b}|^2} \right) \mathbf{b}$

**Example: 29**

The co-ordinates of the foot of the perpendicular drawn from the point $A(1, 0, 3)$ to the join of the points $B(4, 7, 1)$ and $C(3, 5, 3)$ are [Rajasthan PET 2001]

- (a) $(5/3, 7/3, 17/3)$ (b) $(5, 7, 17)$ (c) $(5/3, -7/3, 17/3)$ (d) $(-5/3, 7/3, -17/3)$

Solution: (a)

Equation of BC , $\frac{x-4}{-1} = \frac{y-7}{-2} = \frac{z-1}{2}$

i.e. $\frac{x-4}{1} = \frac{y-7}{2} = \frac{z-1}{-2} = r$ (say)

Any point on the given line is $D(r+4, 2r+7, -2r+1)$


Then, d.r.'s of $AD = (r+4-1, 2r+7-0, -2r+1-3)$

i.e. d.r.'s of $AD = (r+3, 2r+7, -2r-2)$ and d.r.'s of $BC = (-1, -2, 2)$

Since AD is \perp to given line,

$$\therefore (-1)(r+3) + (2r+7)(-2) + (2)(-2r-2) = 0 \Rightarrow -r-3-4r-14-4r-4 = 0 \Rightarrow -9r-21 = 0 \Rightarrow r = -7/3$$

$\therefore D$ is $\{4 - (7/3), 7 - (14/3), (14/3) + 1\}$ i.e. D is $(5/3, 7/3, 17/3)$.

**Example: 30**

The image of the point $(1, 6, 3)$ in the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$ is

- (a) $(1, 0, 7)$ (b) $(-1, 0, 7)$ (c) $(1, 0, -7)$ (d) None of these

Solution: (a)

Let $P(1, 6, 3)$ be the given point, and let L be the foot of the perpendicular from P to the given line. The co-ordinates of a general

point on the given line are given by $\frac{x-0}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda$

i.e. $x = \lambda, y = 2\lambda + 1, z = 3\lambda + 2$.

Let the co-ordinates of L be $(\lambda, 2\lambda + 1, 3\lambda + 2)$ (i)

So, direction ratios of PL are $\lambda - 1, 2\lambda + 1 - 6, 3\lambda + 2 - 3$ i.e. $\lambda - 1, 2\lambda - 5, 3\lambda - 1$.

Direction ratios of the given line are $1, 2, 3$ which is perpendicular to PL .


$$\therefore (\lambda - 1) \cdot 1 + (2\lambda - 5) \cdot 2 + (3\lambda - 1) \cdot 3 = 0 \Rightarrow 14\lambda - 14 = 0 \Rightarrow \lambda = 1$$

So, co-ordinates of L are $(1, 3, 5)$. Let $Q(x_1, y_1, z_1)$ be the image of $P(1, 6, 3)$ in the given line.

Then L is the mid-point of PQ .

$$\therefore \frac{x_1 + 1}{2} = 1, \frac{y_1 + 6}{2} = 3 \text{ and } \frac{z_1 + 3}{2} = 5 \Rightarrow x_1 = 1, y_1 = 0 \text{ and } z_1 = 7.$$

Hence the image of $P(1, 6, 3)$ in the given line is $(1, 0, 7)$.




Example: 31 The length of the perpendicular from the origin to line $\mathbf{r} = (4\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}) + \lambda(3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k})$ is

[AMU 1992]

- (a) $2\sqrt{5}$ (b) 2 (c) $5\sqrt{2}$ (d) 6

Solution: (d) $\vec{a} = 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}$

$$\vec{PL} = (\mathbf{a} - \vec{a}) - \left(\frac{(\mathbf{a} - \vec{a}) \cdot \mathbf{b}}{|\mathbf{b}|^2} \right) \mathbf{b}$$



$$\vec{PL} = (4\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}) - \left[\frac{(4\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}) \cdot (3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k})}{9 + 16 + 25} \right] (3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k}) = 4\mathbf{i} + 2\mathbf{j} + 4\mathbf{k} - \left(\frac{12 + 8 - 20}{50} \right) (3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k})$$

$$\vec{PL} = 4\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}$$

The length of PL is magnitude of \vec{PL} i.e., Length of perpendicular $= |\vec{PL}| = \sqrt{16 + 4 + 16} = 6$.

Example: 32 The image of point $(1, 2, 3)$ in the line $\mathbf{r} = (6\mathbf{i} + 7\mathbf{j} + 7\mathbf{k}) + \lambda(3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$ is

- (a) $(5, -8, 15)$ (b) $(5, 8, -15)$ (c) $(-5, -8, -15)$


Solution: (d) Given that, $\mathbf{a} = 6\mathbf{i} + 7\mathbf{j} + 7\mathbf{k}$, $\mathbf{b} = 3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$ and $\vec{a} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$

$$\text{Then, } \vec{\beta} = 2\mathbf{a} - \left(\frac{2(\mathbf{a} - \vec{a}) \cdot \mathbf{b}}{|\mathbf{b}|^2} \right) \mathbf{b} - \vec{a}$$

$$= 2(6\mathbf{i} + 7\mathbf{j} + 7\mathbf{k}) - \left(\frac{2(5\mathbf{i} + 5\mathbf{j} + 4\mathbf{k}) \cdot (3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})}{9 + 4 + 4} \right) (3\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}) - (\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$$

On solving, $\vec{\beta} = 5\mathbf{i} + 8\mathbf{j} + 15\mathbf{k}$. Thus $\vec{\beta}$ is the position vector of Q , which is the image of P in given line.


Hence image of point $(1, 2, 3)$ in the given line is $(5, 8, 15)$.



7.16 Shortest distance between two straight lines

(1) **Skew lines** : Two straight lines in space which are neither parallel nor intersecting are called skew lines.

Thus, the skew lines are those lines which do not lie in the same plane.



(2) **Line of shortest distance** : If l_1 and l_2 are two skew lines, then the straight line which is perpendicular to each of these two non-intersecting lines is called the “line of shortest distance.”

Note : \square There is one and only one line perpendicular to each of lines l_1 and l_2 .

(3) **Shortest distance between two skew lines**

(i) **Cartesian form** : Let two skew lines be $\frac{x-x_1}{l_1} = \frac{y-y_1}{m_1} = \frac{z-z_1}{n_1}$ and $\frac{x-x_2}{l_2} = \frac{y-y_2}{m_2} = \frac{z-z_2}{n_2}$

Therefore, the shortest distance between the lines is given by

$$d = \frac{\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix}}{\sqrt{(m_1 n_2 - m_2 n_1)^2 + (n_1 l_2 - l_1 n_2)^2 + (l_1 m_2 - l_2 m_1)^2}}$$

(ii) **Vector form** : Let l_1 and l_2 be two lines whose equations are $l_1 : \mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b}_1$ and $l_2 : \mathbf{r} = \mathbf{a}_2 + \mu \mathbf{b}_2$ respectively. Then, Shortest distance $PQ = \left| \frac{(\mathbf{b}_1 \times \mathbf{b}_2) \cdot (\mathbf{a}_2 - \mathbf{a}_1)}{|\mathbf{b}_1 \times \mathbf{b}_2|} \right| = \frac{|\mathbf{b}_1 \mathbf{b}_2 (\mathbf{a}_2 - \mathbf{a}_1)|}{|\mathbf{b}_1 \times \mathbf{b}_2|}$

(4) **Shortest distance between two parallel lines** : The shortest distance between the parallel lines $\mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b}$ and $\mathbf{r} = \mathbf{a}_2 + \mu \mathbf{b}$ is given by $d = \frac{|(\mathbf{a}_2 - \mathbf{a}_1) \times \mathbf{b}|}{|\mathbf{b}|}$.

(5) **Condition for two lines to be intersecting i.e. coplanar**

(i) **Cartesian form** : If the lines $\frac{x-x_1}{l_1} = \frac{y-y_1}{m_1} = \frac{z-z_1}{n_1}$ and $\frac{x-x_2}{l_2} = \frac{y-y_2}{m_2} = \frac{z-z_2}{n_2}$ intersect, then

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0.$$

(ii) **Vector form** : If the lines $\mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b}_1$ and $\mathbf{r} = \mathbf{a}_2 + \lambda \mathbf{b}_2$ intersect, then the shortest distance between them is zero. Therefore, $[\mathbf{b}_1 \mathbf{b}_2 (\mathbf{a}_2 - \mathbf{a}_1)] = 0 \Rightarrow [(\mathbf{a}_2 - \mathbf{a}_1) \mathbf{b}_1 \mathbf{b}_2] = 0 \Rightarrow (\mathbf{a}_2 - \mathbf{a}_1) \cdot (\mathbf{b}_1 \times \mathbf{b}_2) = 0$

Important Tips

- ☞ Skew lines are non-coplanar lines.
- ☞ Parallel lines are not skew lines.
- ☞ If two lines intersect, the shortest distance (SD) between them is zero.
- ☞ Length of shortest distance between two lines is always taken to be positive.
- ☞ Shortest distance between two skew lines is perpendicular to both the lines.

(6) **To determine the equation of line of shortest distance** : To find the equation of line of shortest distance, we use the following procedure :

(i) From the given equations of the straight lines,

$$\text{i.e. } \frac{x-a_1}{l_1} = \frac{y-b_1}{m_1} = \frac{z-c_1}{n_1} = \lambda \text{ (say)} \quad \dots\dots(i)$$

$$\text{and } \frac{x-a_2}{l_2} = \frac{y-b_2}{m_2} = \frac{z-c_2}{n_2} = \mu \text{ (say)} \quad \dots\dots(ii)$$

Find the co-ordinates of general points on straight lines (i) and (ii) as

$$(a_1 + \lambda l_1, b_1 + \lambda m_1, c_1 + \lambda n_1) \text{ and } (a_2 + \mu l_2, b_2 + \mu m_2, c_2 + \mu n_2).$$

(ii) Let these be the co-ordinates of P and Q , the two extremities of the length of shortest distance. Hence, find the direction ratios of PQ as $(a_2 + \mu l_2) - (a_1 + \lambda l_1)$, $(b_2 + \mu m_2) - (b_1 + \lambda m_1)$, $(c_2 + \mu n_2) - (c_1 + \lambda n_1)$.

(iii) Apply the condition of PQ being perpendicular to straight lines (i) and (ii) in succession and get two equations connecting λ and μ . Solve these equations to get the values of λ and μ .

(iv) Put these values of λ and μ in the co-ordinates of P and Q to determine points P and Q .

(v) Find out the equation of the line passing through P and Q , which will be the line of shortest distance.

Note : □ The same algorithm may be observed to find out the position vector of P and Q , the two extremities of the shortest distance, in case of vector equations of straight lines. Hence, the line of shortest distance, which passes through P and Q , can be obtained.

338 Three Dimensional Co-ordinate Geometry

Example: 33 The shortest distance between the lines $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and $\frac{x-2}{3} = \frac{y-4}{4} = \frac{z-5}{5}$ is [Kerala (Engg.)2001; DCE 1993]

- (a) $\frac{1}{6}$ (b) $\frac{1}{\sqrt{6}}$ (c) $\frac{1}{\sqrt{3}}$ (d) $\frac{1}{3}$

Solution: (b)
$$\text{S.D.} = \frac{\begin{vmatrix} 2-1 & 4-2 & 5-3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}}{\sqrt{(15-16)^2 + (12-10)^2 + (8-9)^2}} = \frac{\begin{vmatrix} 1 & 2 & 2 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}}{\sqrt{1+1+4}} = \frac{1}{\sqrt{6}}.$$

Example: 34 The shortest distance between the lines $\mathbf{r} = (\mathbf{i} + \mathbf{j} - \mathbf{k}) + \lambda(3\mathbf{i} - \mathbf{j})$ and $\mathbf{r} = (4\mathbf{i} - \mathbf{k}) + \mu(2\mathbf{i} + 3\mathbf{k})$ is [Pb. CET 1995]

- (a) 6 (b) 0 (c) 2 (d) 4

Solution: (b)
$$\text{S.D.} = \frac{|(\mathbf{b}_1 \times \mathbf{b}_2) \cdot (\mathbf{a}_2 - \mathbf{a}_1)|}{|\mathbf{b}_1 \times \mathbf{b}_2|} = \frac{|[(3\mathbf{i} - \mathbf{j}) \times (2\mathbf{i} + 3\mathbf{k})] \cdot (3\mathbf{i} - \mathbf{j})|}{|(3\mathbf{i} - \mathbf{j}) \times (2\mathbf{i} + 3\mathbf{k})|} = \frac{|(-3\mathbf{i} - 9\mathbf{j} + 2\mathbf{k}) \cdot (3\mathbf{i} - \mathbf{j})|}{\sqrt{9+81+4}} = \frac{-9+9+0}{\sqrt{94}}.$$

Hence, S.D. = 0

Example: 35 The line $\frac{x-2}{1} = \frac{y-3}{1} = \frac{z-4}{-k}$ and $\frac{x-1}{k} = \frac{y-4}{2} = \frac{z-5}{1}$ are coplanar, if [AIEEE 2003]

- (a) $k = 0$ or -1 (b) $k = 0$ or 1 (c) $k = 0$ or -3 (d) $k = 3$ or -3

Solution: (c) Lines are coplanar, if

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} 1-2 & 4-3 & 5-4 \\ 1 & 1 & -k \\ k & 2 & 1 \end{vmatrix} = 0 \Rightarrow k^2 + 3k = 0 \Rightarrow k(k+3) = 0 \Rightarrow k = 0, k = -3$$

Example: 36 The lines $\mathbf{r} = \mathbf{a} + \lambda(\mathbf{b} \times \mathbf{c})$ and $\mathbf{r} = \mathbf{b} + \mu(\mathbf{c} \times \mathbf{a})$ will intersect if

- (a) $\mathbf{a} \times \mathbf{c} = \mathbf{b} \times \mathbf{c}$ (b) $\mathbf{a} \cdot \mathbf{c} = \mathbf{b} \cdot \mathbf{c}$ (c) $\mathbf{b} \times \mathbf{a} = \mathbf{c} \times \mathbf{a}$ (d) None of these

Solution: (b) If lines are intersecting, then

$$\begin{aligned} (\mathbf{a}_2 - \mathbf{a}_1) \cdot (\mathbf{b}_1 \times \mathbf{b}_2) &= 0 \Rightarrow \mathbf{b} \cdot (\mathbf{a} - \mathbf{b}) \cdot [(\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a})] = 0 \\ \Rightarrow (\mathbf{a} - \mathbf{b}) \cdot [(\mathbf{b} \times \mathbf{c}) \cdot \mathbf{a} - (\mathbf{b} \times \mathbf{c} \cdot \mathbf{c}) \mathbf{a}] &= 0 \Rightarrow (\mathbf{a} - \mathbf{b}) \cdot [(\mathbf{b} \times \mathbf{c} \cdot \mathbf{a}) \mathbf{c}] = 0 \\ \Rightarrow [(\mathbf{a} - \mathbf{b}) \cdot \mathbf{c}] \mathbf{a} \cdot \mathbf{b} \cdot \mathbf{c} &= 0 \Rightarrow (\mathbf{a} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{c})(\mathbf{a} \cdot \mathbf{b} \cdot \mathbf{c}) = 0 \Rightarrow \mathbf{a} \cdot \mathbf{c} - \mathbf{b} \cdot \mathbf{c} = 0 \Rightarrow \mathbf{a} \cdot \mathbf{c} = \mathbf{b} \cdot \mathbf{c} \end{aligned}$$

Example: 37 If the straight lines $x = 1 + s, y = 3 - \lambda s, z = 1 + \lambda s$ and $x = \frac{t}{2}, y = 1 + t, z = 2 - t$, with parameters s and t respectively, are coplanar, then λ equals [AIEEE 2004]

- (a) 0 (b) -1 (c) $-\frac{1}{2}$ (d) -2

Solution: (d) We have $\frac{x-1}{1} = \frac{y+3}{-\lambda} = \frac{z-1}{\lambda} = s$ and $\frac{2x}{1} = \frac{y-1}{1} = \frac{z-2}{-1} = t$

i.e. $\frac{x-0}{1} = \frac{y-1}{2} = \frac{z-2}{-2} = \frac{t}{2}$

Since, lines are co-planar,

Then,
$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} -1 & 4 & 1 \\ 1 & -\lambda & \lambda \\ 1 & 2 & -2 \end{vmatrix} = 0$$

On solving, $\lambda = -2$.

The Plane

7.17 Definition of plane and its equations

If point $P(x, y, z)$ moves according to certain rule, then it may lie in a 3-D region on a surface or on a line or it may simply be a point. Whatever we get, as the region of P after applying the rule, is called locus of P . Let us discuss about the plane or curved surface. If Q be any other point on its locus and all points of the straight line PQ lie on it, it is a plane. In other words if the straight line PQ , however small and in whatever direction it may be, lies completely on the locus, it is a plane, otherwise any curved surface.

(1) **General equation of plane** : Every equation of first degree of the form $Ax + By + Cz + D = 0$ represents the equation of a plane. The coefficients of x, y and z i.e. A, B, C are the direction ratios of the normal to the plane.

(2) **Equation of co-ordinate planes**

XOY -plane : $z = 0$



YOZ -plane : $x = 0$

ZOX -plane : $y = 0$

(3) **Vector equation of plane**


(i) Vector equation of a plane through the point $A(\mathbf{a})$ and perpendicular to the vector \mathbf{n} is $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ or $\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$

Note : □ The above equation can also be written as $\mathbf{r} \cdot \mathbf{n} = d$, where $d = \mathbf{a} \cdot \mathbf{n}$. This is known as the scalar product form of a plane.




(4) **Normal form** : Vector equation of a plane normal to unit vector $\hat{\mathbf{n}}$ and at a distance d from the origin is $\mathbf{r} \cdot \hat{\mathbf{n}} = d$.

Note : □ If \mathbf{n} is not a unit vector, then to reduce the equation $\mathbf{r} \cdot \mathbf{n} = d$ to normal form we divide both sides by $|\mathbf{n}|$ to obtain $\mathbf{r} \cdot \frac{\mathbf{n}}{|\mathbf{n}|} = \frac{d}{|\mathbf{n}|}$ or $\mathbf{r} \cdot \hat{\mathbf{n}} = \frac{d}{|\mathbf{n}|}$.



(5) **Equation of a plane passing through a given point and parallel to two given vectors** : The equation of the plane passing through a point having position vector \mathbf{a} and parallel to \mathbf{b} and \mathbf{c} is $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$, where λ and μ are scalars.



(6) **Equation of plane in various forms**

(i) **Intercept form** : If the plane cuts the intercepts of length a, b, c on co-ordinate axes, then its equation is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1.$$

(ii) **Normal form** : Normal form of the equation of plane is $lx + my + nz = p$,

where l, m, n are the d.c.'s of the normal to the plane and p is the length of perpendicular from the origin.

(7) Equation of plane in particular cases

(i) Equation of plane through the origin is given by $Ax + By + Cz = 0$.

i.e. if $D = 0$, then the plane passes through the origin.

(8) Equation of plane parallel to co-ordinate planes or perpendicular to co-ordinate axes

(i) Equation of plane parallel to YOZ -plane (or perpendicular to x -axis) and at a distance ' a ' from it is $x = a$.

(ii) Equation of plane parallel to ZOX -plane (or perpendicular to y -axis) and at a distance ' b ' from it is $y = b$.

(iii) Equation of plane parallel to XOY -plane (or perpendicular to z -axis) and at a distance ' c ' from it is $z = c$.

Important Tips

☞ Any plane perpendicular to co-ordinate axis is evidently parallel to co-ordinate plane and vice versa.

☞ A unit vector perpendicular to the plane containing three points A, B, C is $\frac{\overrightarrow{AB} \times \overrightarrow{AC}}{|\overrightarrow{AB} \times \overrightarrow{AC}|}$.

(9) Equation of plane perpendicular to co-ordinate planes or parallel to co-ordinate axes

(i) Equation of plane perpendicular to YOZ -plane or parallel to x -axis is $By + Cz + D = 0$.

(ii) Equation of plane perpendicular to ZOX -plane or parallel to y axis is $Ax + Cz + D = 0$.

(iii) Equation of plane perpendicular to XOY -plane or parallel to z -axis is $Ax + By + D = 0$.

(10) Equation of plane passing through the intersection of two planes

(i) **Cartesian form** : Equation of plane through the intersection of two planes

$P = a_1x + b_1y + c_1z + d_1 = 0$ and $Q = a_2x + b_2y + c_2z + d_2 = 0$ is $P + \lambda Q = 0$, where λ is the parameter.

(ii) **Vector form** : The equation of any plane through the intersection of planes $\mathbf{r} \cdot \mathbf{n}_1 = d_1$ and $\mathbf{r} \cdot \mathbf{n}_2 = d_2$ is $\mathbf{r} \cdot (\mathbf{n}_1 + \lambda \mathbf{n}_2) = d_1 + \lambda d_2$, where λ is an arbitrary constant.

(11) Equation of plane parallel to a given plane

(i) **Cartesian form** : Plane parallel to a given plane $ax + by + cz + d = 0$ is $ax + by + cz + d' = 0$, i.e. only constant term is changed.

(ii) **Vector form** : Since parallel planes have the common normal, therefore equation of plane parallel to plane $\mathbf{r} \cdot \mathbf{n} = d_1$ is $\mathbf{r} \cdot \mathbf{n} = d_2$, where d_2 is a constant determined by the given condition.

7.18 Equation of plane passing through the given point

(1) **Equation of plane passing through a given point** : Equation of plane passing through the point (x_1, y_1, z_1) is $A(x - x_1) + B(y - y_1) + C(z - z_1) = 0$, where A, B and C are d.r.'s of normal to the plane.

(2) **Equation of plane through three points** : The equation of plane passing through three non-collinear

points (x_1, y_1, z_1) , (x_2, y_2, z_2) and (x_3, y_3, z_3) is
$$\begin{vmatrix} x & y & z & 1 \\ x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \end{vmatrix} = 0 \quad \text{or} \quad \begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0.$$

7.19 Foot of perpendicular from a point $A(\alpha, \beta, \gamma)$ to a given plane $ax + by + cz + d = 0$

If AP be the perpendicular from A to the given plane, then it is parallel to the normal, so that its equation is

$$\frac{x - \alpha}{a} = \frac{y - \beta}{b} = \frac{z - \gamma}{c} = r \quad (\text{say})$$

Any point P on it is $(ar + \alpha, br + \beta, cr + \gamma)$. It lies on the given plane and we find the value of r and hence the point P .

(1) Perpendicular distance

(i) **Cartesian form** : The length of the perpendicular from the point $P(x_1, y_1, z_1)$ to the plane $ax + by + cz + d = 0$

is $\left| \frac{ax_1 + by_1 + cz_1 + d}{\sqrt{a^2 + b^2 + c^2}} \right|$.

Note : \square The distance between two parallel planes is the algebraic difference of perpendicular distances on the planes from origin.

\square Distance between two parallel planes $Ax + By + Cz + D_1 = 0$ and $Ax + By + Cz + D_2 = 0$ is $\frac{D_2 - D_1}{\sqrt{A^2 + B^2 + C^2}}$.

(ii) **Vector form** : The perpendicular distance of a point having position vector \mathbf{a} from the plane $\mathbf{r} \cdot \mathbf{n} = d$ is given

by $p = \frac{|\mathbf{a} \cdot \mathbf{n} - d|}{|\mathbf{n}|}$

(2) **Position of two points w.r.t. a plane** : Two points $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ lie on the same or opposite sides of a plane $ax + by + cz + d = 0$ according to $ax_1 + by_1 + cz_1 + d$ and $ax_2 + by_2 + cz_2 + d$ are of same or opposite signs. The plane divides the line joining the points P and Q externally or internally according to P and Q are lying on same or opposite sides of the plane.

7.20 Angle between two planes

(1) **Cartesian form** : Angle between the planes is defined as angle between normals to the planes drawn from any point. Angle between the planes $a_1x + b_1y + c_1z + d_1 = 0$ and $a_2x + b_2y + c_2z + d_2 = 0$ is

$$\cos^{-1} \left(\frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{(a_1^2 + b_1^2 + c_1^2)(a_2^2 + b_2^2 + c_2^2)}} \right)$$

Note : \square If $a_1a_2 + b_1b_2 + c_1c_2 = 0$, then the planes are perpendicular to each other.

\square If $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, then the planes are parallel to each other.

(2) **Vector form** : An angle θ between the planes $\mathbf{r}_1 \cdot \mathbf{n}_1 = d_1$ and $\mathbf{r}_2 \cdot \mathbf{n}_2 = d_2$ is given by $\cos \theta = \pm \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|}$.

7.21 Equation of planes bisecting angle between two given planes

(1) **Cartesian form** : Equations of planes bisecting angles between the planes $a_1x + b_1y + c_1z + d_1 = 0$ and

$a_2x + b_2y + c_2z + d = 0$ are $\frac{a_1x + b_1y + c_1z + d_1}{\sqrt{(a_1^2 + b_1^2 + c_1^2)}} = \pm \frac{a_2x + b_2y + c_2z + d_2}{\sqrt{(a_2^2 + b_2^2 + c_2^2)}}$.

Note : \square If angle between bisector plane and one of the plane is less than 45° , then it is acute angle bisector, otherwise it is obtuse angle bisector.

- If $a_1a_2 + b_1b_2 + c_1c_2$ is negative, then origin lies in the acute angle between the given planes provided d_1 and d_2 are of same sign and if $a_1a_2 + b_1b_2 + c_1c_2$ is positive, then origin lies in the obtuse angle between the given planes.

(2) **Vector form :** The equation of the planes bisecting the angles between the planes $\mathbf{r} \cdot \mathbf{n}_1 = d_1$ and $\mathbf{r} \cdot \mathbf{n}_2 = d_2$ are $\frac{|\mathbf{r} \cdot \mathbf{n}_1 - d_1|}{|\mathbf{n}_1|} = \frac{|\mathbf{r} \cdot \mathbf{n}_2 - d_2|}{|\mathbf{n}_2|}$ or $\frac{\mathbf{r} \cdot \mathbf{n}_1 - d_1}{|\mathbf{n}_1|} = \pm \frac{\mathbf{r} \cdot \mathbf{n}_2 - d_2}{|\mathbf{n}_2|}$ or $\mathbf{r} \cdot (\hat{\mathbf{n}}_1 \pm \hat{\mathbf{n}}_2) = \frac{d_1}{|\mathbf{n}_1|} \pm \frac{d_2}{|\mathbf{n}_2|}$.

7.22 Image of a point in a plane

Let P and Q be two points and let π be a plane such that

- Line PQ is perpendicular to the plane π , and
- Mid-point of PQ lies on the plane π .

Then either of the point is the image of the other in the plane π .

To find the image of a point in a given plane, we proceed as follows

- Write the equations of the line passing through P and normal to the given plane as


$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}.$$

- Write the co-ordinates of image Q as $(x_1 + ar, y_1 + br, z_1 + cr)$.

- Find the co-ordinates of the mid-point R of PQ .

- Obtain the value of r by putting the co-ordinates of R in the equation of the plane.

- Put the value of r in the co-ordinates of Q .



7.23 Coplanar lines

Lines are said to be coplanar if they lie in the same plane or a plane can be made to pass through them.

- Condition for the lines to be coplanar**

- Cartesian form :** If the lines $\frac{x - x_1}{l_1} = \frac{y - y_1}{m_1} = \frac{z - z_1}{n_1}$ and $\frac{x - x_2}{l_2} = \frac{y - y_2}{m_2} = \frac{z - z_2}{n_2}$ are coplanar

Then
$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0.$$

The equation of the plane containing them is
$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0 \text{ or } \begin{vmatrix} x - x_2 & y - y_2 & z - z_2 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0.$$

- Vector form :** If the lines $\mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b}_1$ and $\mathbf{r} = \mathbf{a}_2 + \lambda \mathbf{b}_2$ are coplanar, then $[\mathbf{a}_1 \mathbf{b}_1 \mathbf{b}_2] = [\mathbf{a}_2 \mathbf{b}_1 \mathbf{b}_2]$ and the equation of the plane containing them is $[\mathbf{r} \mathbf{b}_1 \mathbf{b}_2] = [\mathbf{a}_1 \mathbf{b}_1 \mathbf{b}_2]$ or $[\mathbf{r} \mathbf{b}_1 \mathbf{b}_2] = [\mathbf{a}_2 \mathbf{b}_1 \mathbf{b}_2]$.

Note: □ Every pair of parallel lines is coplanar.

- ❑ Two coplanar lines are either parallel or intersecting.
 ❑ The three sides of a triangle are coplanar.

Important Tips

☞ **Division by plane :** The ratio in which the line segment PQ , joining $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$, is divided by plane $ax + by + cz + d = 0$ is

$$= -\left(\frac{ax_1 + by_1 + cz_1 + d}{ax_2 + by_2 + cz_2 + d}\right).$$

☞ **Division by co-ordinate planes :** The ratio in which the line segment PQ , joining $P(x_1, y_1, z_1)$ and $Q(x_2, y_2, z_2)$ is divided by co-ordinate planes are as follows :

(i) By yz -plane : $-x_1/x_2$ (ii) By zx -plane : $-y_1/y_2$ (iii) By xy -plane : $-z_1/z_2$

Example: 38 The xy -plane divides the line joining the points $(-1, 3, 4)$ and $(2, -5, 6)$ [Rajasthan PET 2000]

- (a) Internally in the ratio 2 : 3 (b) Internally in the ratio 3 : 2
 (c) Externally in the ratio 2 : 3 (d) Externally in the ratio 3 : 2

Solution: (c) Required ratio $= -\frac{z_1}{z_2} = -\left(\frac{4}{6}\right) = -\frac{2}{3}$

$\therefore xy$ -plane divide externally in the ratio 2 : 3.

Example: 39 The ratio in which the plane $x - 2y + 3z = 17$ divides the line joining the point $(-2, 4, 7)$ and $(3, -5, 8)$ is [AISSSE 1988]

- (a) 10 : 3 (b) 3 : 1 (c) 3 : 10 (d) 10 : 1

Solution: (c) Required ratio $= -\left(\frac{ax_1 + by_1 + cz_1 + d}{ax_2 + by_2 + cz_2 + d}\right) = -\left(\frac{-2 - 8 + 21 - 17}{3 + 10 + 24 - 17}\right) = \frac{6}{20} = \frac{3}{10}$.

Example: 40 The equation of the plane, which makes with co-ordinate axes a triangle with its centroid (α, β, γ) , is [MP PET 2004]

- (a) $\alpha x + \beta y + \gamma z = 3$ (b) $\frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 1$ (c) $\alpha x + \beta y + \gamma z = 1$ (d) $\frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 3$

Solution: (d) We know that $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ (i)

Centroid $\left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3}\right)$ i.e. $\alpha = a/3, \beta = b/3, \gamma = c/3 \Rightarrow a = 3\alpha, b = 3\beta, c = 3\gamma$

From equation (i), $\frac{x}{3\alpha} + \frac{y}{3\beta} + \frac{z}{3\gamma} = 1$

$\therefore \frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 3$.

Example: 41 The equation of plane passing through the points $(2, 2, 1)$ and $(9, 3, 6)$ and perpendicular to the plane $2x + 6y + 6z = 1$ is [AISSSE 1984; Tamilnadu (Engg.) 2002]

- (a) $3x + 4y + 5z = 9$ (b) $3x + 4y + 5z = 0$ (c) $3x + 4y - 5z = 9$ (d) None of these

Solution: (c) We know that, equation of plane is $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$

It passes through $(2, 2, 1)$

$\therefore a(x - 2) + b(y - 2) + c(z - 1) = 0$ (i)

Plane (i) also passes through $(9, 3, 6)$ and is perpendicular to the plane $2x + 6y + 6z = 1$

$\therefore 7a + b + 5c = 0$ (ii)

and $2a + 6b + 6c = 0$ (iii)

$$\frac{a}{6 - 30} = \frac{b}{10 - 42} = \frac{c}{42 - 2} \text{ or } \frac{a}{-24} = \frac{b}{-32} = \frac{c}{40}$$

$$\text{or } \frac{a}{3} = \frac{b}{4} = \frac{c}{-5} = k \text{ (say)}$$

$$\text{From equation (i), } 3k(x-2) + 4k(y-2) + (-5)k(z-1) = 0$$

$$\text{Hence, } 3x + 4y - 5z = 9.$$

Example: 42 The equation of the plane containing the line $\mathbf{r} = \mathbf{a} + k\mathbf{b}$ and perpendicular to the plane $\mathbf{r} \cdot \mathbf{n} = q$ is

$$(a) \quad (\mathbf{r} - \mathbf{b}) \cdot (\mathbf{n} \times \mathbf{a}) = 0 \quad (b) \quad (\mathbf{r} - \mathbf{a}) \cdot (\mathbf{n} \times (\mathbf{a} \times \mathbf{b})) = 0 \quad (c) \quad (\mathbf{r} - \mathbf{a}) \cdot (\mathbf{n} \times \mathbf{b}) = 0 \quad (d) \quad (\mathbf{r} - \mathbf{b}) \cdot (\mathbf{n} \times (\mathbf{a} \times \mathbf{b})) = 0$$

Solution: (c) Since the required plane contains the line $\mathbf{r} = \mathbf{a} + k\mathbf{b}$ and is perpendicular to the plane $\mathbf{r} \cdot \mathbf{n} = q$.

\therefore It passes through the point \mathbf{a} and parallel to vectors \mathbf{b} and \mathbf{n} . Hence, it is perpendicular to the vector $\mathbf{N} = \mathbf{n} \times \mathbf{b}$.

$$\therefore \text{Equation of the required plane is } (\mathbf{r} - \mathbf{a}) \cdot \mathbf{N} = 0 \Rightarrow (\mathbf{r} - \mathbf{a}) \cdot (\mathbf{n} \times \mathbf{b}) = 0.$$

Example: 43 The equation of the plane through the intersection of the planes $x + 2y + 3z - 4 = 0$, $4x + 3y + 2z + 1 = 0$ and passing through the origin will be [MP PET 1997; Kerala (Engg.) 2001; AISSCE 1983]

$$(a) \quad x + y + z = 0 \quad (b) \quad 17x + 14y + 11z = 0 \quad (c) \quad 7x + 4y + z = 0 \quad (d) \quad 17x + 14y + z = 0$$

Solution: (b) Any plane through the given planes is $(x + 2y + 3z - 4) + k(4x + 3y + 2z + 1) = 0$

It passes through $(0, 0, 0)$

$$\therefore -4 + k = 0 \Rightarrow k = 4$$

$$\therefore \text{Required plane is } (x + 2y + 3z - 4) + 4(4x + 3y + 2z + 1) = 0 \Rightarrow 17x + 14y + 11z = 0.$$

Example: 44 The vector equation of the plane passing through the origin and the line of intersection of plane $\mathbf{r} \cdot \mathbf{a} = \lambda$ and $\mathbf{r} \cdot \mathbf{b} = \mu$ is

$$(a) \quad \mathbf{r} \cdot (\lambda \mathbf{a} - \mu \mathbf{b}) = 0 \quad (b) \quad \mathbf{r} \cdot (\lambda \mathbf{b} - \mu \mathbf{a}) = 0 \quad (c) \quad \mathbf{r} \cdot (\lambda \mathbf{a} + \mu \mathbf{b}) = 0 \quad (d) \quad \mathbf{r} \cdot (\lambda \mathbf{b} + \mu \mathbf{a}) = 0$$

Solution: (b) The equation of a plane through the line of intersection of plane $\mathbf{r} \cdot \mathbf{a} = \lambda$ and $\mathbf{r} \cdot \mathbf{b} = \mu$ can be written as $\mathbf{r} \cdot (\mathbf{a} + k\mathbf{b}) = \lambda + k\mu$

.....(i)

This passes through the origin, therefore putting the value of k in (i),

$$\mathbf{r} \cdot (\mu \mathbf{a} - \lambda \mathbf{b}) = 0 \Rightarrow \mathbf{r} \cdot (\lambda \mathbf{b} - \mu \mathbf{a}) = 0.$$

Example: 45 Angle between two planes $x + 2y + 2z = 3$ and $-5x + 3y + 4z = 9$ is

[IIT Screening 2004]

$$(a) \quad \cos^{-1} \frac{3\sqrt{2}}{10} \quad (b) \quad \cos^{-1} \frac{19\sqrt{2}}{30} \quad (c) \quad \cos^{-1} \frac{9\sqrt{2}}{20} \quad (d) \quad \cos^{-1} \frac{3\sqrt{2}}{5}$$

Solution: (a) We know that, $\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = \frac{1(-5) + 2(3) + 2(4)}{\sqrt{1+4+4} \sqrt{25+9+16}} = \frac{9}{3.5\sqrt{2}} = \frac{3\sqrt{2}}{10}$

$$\text{i.e. } \theta = \cos^{-1} \left(\frac{3\sqrt{2}}{10} \right).$$

Example: 46 Distance between two parallel planes $2x + y + 2z = 8$ and $4x + 2y + 4z + 5 = 0$ is

[AIEEE 2004]

$$(a) \quad \frac{9}{2} \quad (b) \quad \frac{5}{2} \quad (c) \quad \frac{7}{2} \quad (d) \quad \frac{3}{2}$$

Solution: (c) We have $2x + y + 2z - 8 = 0$

.....(i)

$$\text{and } 4x + 2y + 4z + 5 = 0 \text{ or } 2x + y + 2z + 5/2 = 0$$

.....(ii)

$$\text{Distance between the planes} = \frac{(5/2) + 8}{\sqrt{4+1+4}} = \frac{21}{2.3} = \frac{7}{2}.$$

Example: 47 A tetrahedron has vertices at $O(0, 0, 0)$, $A(1, 2, 1)$, $B(2, 1, 3)$ and $C(-1, 1, 2)$. Then the angle between the faces OAB and ABC will be [MNR 1994; UPSEAT 2000; AIEEE 2003]

$$(a) \quad \cos^{-1} \left(\frac{19}{35} \right) \quad (b) \quad \cos^{-1} \left(\frac{17}{31} \right) \quad (c) \quad 30^\circ \quad (d) \quad 90^\circ$$

Solution: (a) Angle between two plane faces is equal to the angle between the normals n_1 and n_2 to the planes. \mathbf{n}_1 , the normal to the face

$$OAB \text{ is given by } \overrightarrow{OA} \times \overrightarrow{OB} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 1 \\ 2 & 1 & 3 \end{vmatrix} = 5\mathbf{i} - \mathbf{j} - 3\mathbf{k} \quad \text{.....(i)}$$

\mathbf{n}_2 , the normal to the face ABC , is given by $\overrightarrow{AB} \times \overrightarrow{AC}$.

$$\mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 2 \\ -2 & -1 & 1 \end{vmatrix} = \mathbf{i} - 5\mathbf{j} - 3\mathbf{k} \quad \text{.....(ii)}$$

If θ be the angle between \mathbf{n}_1 and \mathbf{n}_2 , Then $\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1| |\mathbf{n}_2|} = \frac{5 \cdot 1 + 5 + 9}{\sqrt{35} \sqrt{35}}$

$$\cos \theta = \frac{19}{35} \Rightarrow \theta = \cos^{-1} \left(\frac{19}{35} \right).$$

Example: 48

The distance of the point $(2, 1, -1)$ from the plane $x - 2y + 4z = 9$ is

[Kerala (Engg.) 2001]

- (a) $\frac{\sqrt{13}}{21}$ (b) $\frac{13}{21}$ (c) $\frac{13}{\sqrt{21}}$ (d) $\sqrt{\frac{13}{21}}$

Solution: (c)

$$\text{Distance of the plane from } (2, 1, -1) = \left| \frac{2 - 2(1) + 4(-1) - 9}{\sqrt{1 + 4 + 16}} \right| = \frac{13}{\sqrt{21}}.$$

Example: 49

A unit vector perpendicular to plane determined by the points $P(1, -1, 2)$, $Q(2, 0, -1)$ and $R(0, 2, 1)$ is

[IIT 1994]

- (a) $\frac{2\mathbf{i} - \mathbf{j} + \mathbf{k}}{\sqrt{6}}$ (b) $\frac{2\mathbf{i} + \mathbf{j} + \mathbf{k}}{\sqrt{6}}$ (c) $\frac{-2\mathbf{i} + \mathbf{j} + \mathbf{k}}{\sqrt{6}}$ (d) $\frac{2\mathbf{i} + \mathbf{j} - \mathbf{k}}{\sqrt{6}}$

Solution: (b)

We know that, $\frac{\overrightarrow{PQ} \times \overrightarrow{PR}}{|\overrightarrow{PQ} \times \overrightarrow{PR}|}$

$$\overrightarrow{PQ} = \mathbf{i} + \mathbf{j} - 3\mathbf{k}, \quad \overrightarrow{PR} = -\mathbf{i} + 3\mathbf{j} - \mathbf{k}$$

$$\overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -3 \\ -1 & 3 & -1 \end{vmatrix} = 8\mathbf{i} + 4\mathbf{j} + 4\mathbf{k} \quad \text{and} \quad |\overrightarrow{PQ} \times \overrightarrow{PR}| = 4\sqrt{6}$$

Hence, the unit vector is $\frac{4(2\mathbf{i} + \mathbf{j} + \mathbf{k})}{4\sqrt{6}}$ i.e. $\frac{2\mathbf{i} + \mathbf{j} + \mathbf{k}}{\sqrt{6}}$.

Example: 50

The perpendicular distance from origin to the plane through the point $(2, 3, -1)$ and perpendicular to vector $3\mathbf{i} - 4\mathbf{j} + 7\mathbf{k}$ is

- (a) $\frac{13}{\sqrt{74}}$ (b) $-\frac{13}{\sqrt{74}}$ (c) 13 (d) None of these

Solution: (a)

We know, the equation of the plane is $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$

$$\text{or } (\mathbf{r} - (2\mathbf{i} + 3\mathbf{j} - \mathbf{k})) \cdot (3\mathbf{i} - 4\mathbf{j} + 7\mathbf{k}) = 0 \Rightarrow (x\mathbf{i} + y\mathbf{j} + z\mathbf{k} - 2\mathbf{i} - 3\mathbf{j} + \mathbf{k}) \cdot (3\mathbf{i} - 4\mathbf{j} + 7\mathbf{k}) = 0 \Rightarrow 3x - 4y + 7z + 13 = 0$$

$$\text{Hence, perpendicular distance of the plane from origin} = \frac{13}{\sqrt{3^2 + (-4)^2 + 7^2}} = \frac{13}{\sqrt{74}}.$$

Example: 51

If $P = (0, 1, 0)$, $Q = (0, 0, 1)$, then projection of PQ on the plane $x + y + z = 3$ is

[EAMCET 2002]

- (a) $\sqrt{3}$ (b) 3 (c) $\sqrt{2}$ (d) 2

Solution: (c)

Given plane is $x + y + z - 3 = 0$. From point P and Q draw PM and QN perpendicular on the given plane and $QR \perp MP$.


$$|MP| = \left| \frac{0 + 1 + 0 - 3}{\sqrt{1^2 + 1^2 + 1^2}} \right| = \frac{2}{\sqrt{3}}$$

$$|NQ| = \frac{2}{\sqrt{3}}$$

$$|PQ| = \sqrt{(0-0)^2 + (0-1)^2 + (1-0)^2} = \sqrt{2}$$

$$|RP| = |MP| - |MR| = |MP| - |NQ| = 0 \quad (\text{i.e. } R \text{ and } P \text{ are the same point})$$

$$\therefore |NM| = |QR| = \sqrt{PQ^2 - RP^2} = \sqrt{(\sqrt{2})^2 - 0} = \sqrt{2}$$



Example: 52

The reflection of the point $(2, -1, 3)$ in the plane $3x - 2y - z = 9$ is

[AMU 1995]

- (a) $\left(\frac{26}{7}, \frac{15}{7}, \frac{17}{7} \right)$ (b) $\left(\frac{26}{7}, \frac{-15}{7}, \frac{17}{7} \right)$ (c) $\left(\frac{15}{7}, \frac{26}{7}, \frac{-17}{7} \right)$ (d) $\left(\frac{26}{7}, \frac{17}{7}, \frac{-15}{7} \right)$

Solution: (b)

Let P be the point $(2, -1, 3)$ and Q be its reflection in the given plane.

Then, PQ is perpendicular to the given plane

Hence, d.r.'s of PQ are 3, -2, 1 and consequently, equations of PQ are $\frac{x-2}{3} = \frac{y+1}{-2} = \frac{z-3}{-1}$

Any point on this line is $(3r+2, -2r-1, -r+3)$

Let this point be Q . Then midpoint of $PQ = \left(\frac{3r+2+2}{2}, \frac{-2r-1-1}{2}, \frac{-r+3+3}{2} \right) = \left(\frac{3r+4}{2}, -r-1, \frac{-r+6}{2} \right)$

This point lies in given plane i.e. $3\left(\frac{3r+4}{2}\right) - 2(-r-1) - \left(\frac{-r+6}{2}\right) = 9 \Rightarrow 9r+12+4r+4+r-6=9 \Rightarrow 14r=8 \Rightarrow r=\frac{4}{7}$

Hence, the required point Q is $\left(3\left(\frac{4}{7}\right)+2, -2\left(\frac{4}{7}\right)-1, \frac{-4}{7}+3 \right) = \left(\frac{26}{7}, \frac{-15}{7}, \frac{17}{7} \right)$.

Example: 53

A non-zero vector \mathbf{a} is parallel to the line of intersection of the plane determined by the vectors $\mathbf{i}, \mathbf{i} + \mathbf{j}$ and the plane determined by the vectors $\mathbf{i} - \mathbf{j}, \mathbf{i} + \mathbf{k}$. The angle between \mathbf{a} and the vector $\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$ is [IIT 1996]

- (a) $\frac{\pi}{4}$ or $\frac{3\pi}{4}$ (b) $\frac{2\pi}{4}$ or $\frac{3\pi}{4}$ (c) $\frac{\pi}{2}$ or $\frac{3\pi}{2}$ (d) None of these

Solution: (a)

Equation of plane containing \mathbf{i} and $\mathbf{i} + \mathbf{j}$ is

$$[\mathbf{r}-\mathbf{i}, \mathbf{i}, \mathbf{i}+\mathbf{j}] = 0 \Rightarrow (\mathbf{r}-\mathbf{i}) \cdot [\mathbf{i} \times (\mathbf{i}+\mathbf{j})] = 0 \Rightarrow [(x-1)\mathbf{i} + y\mathbf{j} + z\mathbf{k}] \cdot \mathbf{k} = 0 \Rightarrow z = 0 \quad \dots\dots(i)$$

Equation of plane containing $\mathbf{i} - \mathbf{j}$ and $\mathbf{i} + \mathbf{k}$ is

$$\Rightarrow [\mathbf{r}-(\mathbf{i}-\mathbf{j}), \mathbf{i}-\mathbf{j}, \mathbf{i}+\mathbf{k}] = 0 \Rightarrow (\mathbf{r}-\mathbf{i}+\mathbf{j}) \cdot [(\mathbf{i}-\mathbf{j}) \times (\mathbf{i}+\mathbf{k})] = 0 \Rightarrow x+y-z=0 \quad \dots\dots(ii)$$

Let $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$. Since \mathbf{a} is parallel to (i) and (ii)

$$a_3 = 0, \quad a_1 + a_2 - a_3 = 0 \Rightarrow a_1 = -a_2, \quad a_3 = 0$$

Thus a vector in the direction of \mathbf{a} is $\mathbf{u} = \mathbf{i} - \mathbf{j}$. If θ is the angle between \mathbf{a} and $\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$.

$$\text{Then } \cos \theta = \pm \frac{1(1) + (-1)(-2)}{\sqrt{1+1}\sqrt{1+4+4}} = \pm \frac{3}{\sqrt{2} \cdot 3} \Rightarrow \cos \theta = \pm \frac{1}{\sqrt{2}} \Rightarrow \theta = \pi/4 \text{ or } 3\pi/4$$

Example: 54

The d.r.'s of normal to the plane through $(1, 0, 0)$ and $(0, 1, 0)$ which makes an angle $\pi/4$ with plane $x+y=3$, are [AIEEE 2002]

- (a) $1, \sqrt{2}, 1$ (b) $1, 1, \sqrt{2}$ (c) $1, 1, 2$ (d) $\sqrt{2}, 1, 1$

Solution: (b)

Let d.r.'s of normal to plane (a, b, c)

$$a(x-1) + b(y-0) + c(z-0) = 0 \quad \dots\dots(i)$$

It passes through $(0, 1, 0)$. $\therefore a+b=0 \Rightarrow b=-a$. D.r.'s of normal is (a, a, c) and d.r.'s of given plane is $(1, 1, 0)$

$$\therefore \cos \pi/4 = \frac{a+a+0}{\sqrt{a^2+a^2+c^2}\sqrt{2}} \Rightarrow 4a^2 = 2a^2 + c^2 \Rightarrow \sqrt{2}a = c$$

Then, d.r.'s of normal $(a, a, \sqrt{2}a)$ or $(1, 1, \sqrt{2})$.

Line and plane

7.24 Equation of plane through a given line

(1) If equation of the line is given in symmetrical form as $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$, then equation of plane is

$$a(x-x_1) + b(y-y_1) + c(z-z_1) = 0 \quad \dots\dots(i)$$

where a, b, c are given by $al + bm + cn = 0 \quad \dots\dots(ii)$

(2) If equation of line is given in general form as $a_1x + b_1y + c_1z + d_1 = 0 = a_2x + b_2y + c_2z + d_2$, then the equation of plane passing through this line is $(a_1x + b_1y + c_1z + d_1) + \lambda(a_2x + b_2y + c_2z + d_2) = 0$.

(3) **Equation of plane through a given line parallel to another line :** Let the d.c.'s of the other line be l_2, m_2, n_2 . Then, since the plane is parallel to the given line, normal is perpendicular.

$$\therefore al_2 + bm_2 + cn_2 = 0 \quad \dots\dots(iii)$$

Hence, the plane from (i), (ii) and (iii) is
$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0.$$

7.25 Transformation from unsymmetric form of the equation of line to the symmetric form

If $P \equiv a_1x + b_1y + c_1z + d_1 = 0$ and $Q \equiv a_2x + b_2y + c_2z + d_2 = 0$ are equations of two non-parallel planes, then these two equations taken together represent a line. Thus the equation of straight line can be written as $P = 0 = Q$. This form is called unsymmetrical form of a line.

To transform the equations to symmetrical form, we have to find the d.r.'s of line and co-ordinates of a point on the line.

7.26 Intersection point of a line and plane

To find the point of intersection of the line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$ and the plane $ax + by + cz + d = 0$.

The co-ordinates of any point on the line

$\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$ are given by

$$\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n} = r \text{ (say) or } (x_1 + lr, y_1 + mr, z_1 + nr) \quad \dots(i)$$

If it lies on the plane $ax + by + cz + d = 0$, then

$$a(x_1 + lr) + b(y_1 + mr) + c(z_1 + nr) + d = 0 \Rightarrow (ax_1 + by_1 + cz_1 + d) + r(al + bm + cn) = 0$$

$$\therefore r = -\frac{(ax_1 + by_1 + cz_1 + d)}{al + bm + cn}.$$

Substituting the value of r in (i), we obtain the co-ordinates of the required point of intersection.

Algorithm for finding the point of intersection of a line and a plane

Step I : Write the co-ordinates of any point on the line in terms of some parameters r (say).

Step II : Substitute these co-ordinates in the equation of the plane to obtain the value of r .

Step III : Put the value of r in the co-ordinates of the point in step I.

7.27 Angle between line and plane

(1) **Cartesian form :** The angle θ between the line $\frac{x-\alpha}{l} = \frac{y-\beta}{m} = \frac{z-\gamma}{n}$, and the plane

$$ax + by + cz + d = 0, \text{ is given by } \sin \theta = \frac{al + bm + cn}{\sqrt{(a^2 + b^2 + c^2)}\sqrt{(l^2 + m^2 + n^2)}}.$$

(i) The line is perpendicular to the plane if and only if $\frac{a}{l} = \frac{b}{m} = \frac{c}{n}$.



(ii) The line is parallel to the plane if and only if $al + bm + cn = 0$.

(iii) The line lies in the plane if and only if $al + bm + cn = 0$ and $a\alpha + b\beta + c\gamma + d = 0$.

(2) **Vector form :** If θ is the angle between a line $\mathbf{r} = (\mathbf{a} + \lambda\mathbf{b})$ and the plane $\mathbf{r} \cdot \mathbf{n} = d$, then $\sin \theta = \frac{|\mathbf{b} \cdot \mathbf{n}|}{|\mathbf{b}| |\mathbf{n}|}$.

(i) **Condition of perpendicularity :** If the line is perpendicular to the plane, then it is parallel to the normal to the plane. Therefore \mathbf{b} and \mathbf{n} are parallel.

So, $\mathbf{b} \times \mathbf{n} = 0$ or $\mathbf{b} = \lambda\mathbf{n}$ for some scalar λ .



(ii) **Condition of parallelism** : If the line is parallel to the plane, then it is perpendicular to the normal to the plane. Therefore \mathbf{b} and \mathbf{n} are perpendicular. So, $\mathbf{b} \cdot \mathbf{n} = 0$.

(iii) If the line $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$ lies in the plane $\mathbf{r} \cdot \mathbf{n} = d$, then (i) $\mathbf{b} \cdot \mathbf{n} = 0$ and (ii) $\mathbf{a} \cdot \mathbf{n} = d$.

7.28 Projection of a line on a plane

If P be the point of intersection of given line and plane and Q be the foot of the perpendicular from any point on the line to the plane then PQ is called the projection of given line on the given plane.

Image of line about a plane : Let line is $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$, plane is $a_2x + b_2y + c_2z + d = 0$.

Find point of intersection (say P) of line and plane. Find image (say Q) of point (x_1, y_1, z_1) about the plane. Line PQ is the reflected line.

Example: 55 The sine of angle between the straight line $\frac{x-2}{3} = \frac{y-3}{4} = \frac{z-4}{5}$ and the plane $2x - 2y + z = 5$ is

[Kurukshetra CEE 1995, 2001; DCE 2000]

- (a) $\frac{2\sqrt{3}}{5}$ (b) $\frac{\sqrt{2}}{10}$ (c) $\frac{4}{5\sqrt{2}}$ (d) $\frac{10}{6\sqrt{5}}$

Solution: (b) We know that $\sin \theta = \frac{al + bm + cn}{\sqrt{a^2 + b^2 + c^2} \sqrt{l^2 + m^2 + n^2}}$
 $\sin \theta = \frac{3(2) + 4(-2) + 5(1)}{\sqrt{9 + 16 + 25} \sqrt{4 + 4 + 1}} = \frac{3}{5\sqrt{2} \cdot 3}$
Hence, $\sin \theta = \frac{\sqrt{2}}{10}$

Example: 56 Value of k such that the line $\frac{x-1}{2} = \frac{y-1}{3} = \frac{z-k}{k}$ is perpendicular to normal to the plane $\mathbf{r}(2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}) = 0$ is

[Pb. CET 2001]

- (a) $-\frac{13}{4}$ (b) $-\frac{17}{4}$ (c) 4 (d) None of these

Solution: (a) We have, $\frac{x-1}{2} = \frac{y-1}{3} = \frac{z-k}{k}$
or vector form of equation of line is $\mathbf{r} = (\mathbf{i} + \mathbf{j} + k\mathbf{k}) + \lambda(2\mathbf{i} + 3\mathbf{j} + k\mathbf{k})$ i.e. $\mathbf{b} = 2\mathbf{i} + 3\mathbf{j} + k\mathbf{k}$ and normal to the plane, $\mathbf{n} = 2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}$.

Given that, $\mathbf{b} \cdot \mathbf{n} = 0$
 $\Rightarrow (2\mathbf{i} + 3\mathbf{j} + k\mathbf{k}) \cdot (2\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}) = 0$
 $\Rightarrow 4 + 9 + 4k = 0 \Rightarrow k = -13/4$.

Example: 57 The equation of line of intersection of the planes $4x + 4y - 5z = 12$, $8x + 12y - 13z = 32$ can be written as [MP PET 2004]

- (a) $\frac{x}{2} = \frac{y-1}{3} = \frac{z-2}{4}$ (b) $\frac{x}{2} = \frac{y}{3} = \frac{z-2}{4}$ (c) $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z}{4}$ (d) $\frac{x-1}{2} = \frac{y-2}{-3} = \frac{z}{4}$

Solution: (c) Let equation of line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$ (i)

We have $4x + 4y - 5z = 12$ (ii) and $8x + 12y - 13z = 32$ (iii)

Let $z = 0$. Now putting $z = 0$ in (ii) and (iii),

we get, $4x + 4y = 12$, $8x + 12y = 32$, on solving these equations, we get $x = 1, y = 2$.

Equation of line passing through $(1, 2, 0)$ is $\frac{x-1}{l} = \frac{y-2}{m} = \frac{z-0}{n}$

From equation (i) and (ii),

$$4l+4m-5n=0 \text{ and } 8l+12m-13n=0$$

$$\Rightarrow \frac{l}{8} = \frac{m}{12} = \frac{n}{16} \text{ i.e. } \frac{l}{2} = \frac{m}{3} = \frac{n}{4}. \text{ Hence, equation of line is } \frac{x-1}{2} = \frac{y-2}{3} = \frac{z}{4}.$$

Example: 58 The equation of the plane containing the two lines $\frac{x-1}{2} = \frac{y+1}{-1} = \frac{z}{3}$ and $\frac{x}{2} = \frac{y-2}{-1} = \frac{z+1}{-3}$ is [MP PET 2000]

- (a) $8x+y-5z-7=0$ (b) $8x+y+5z-7=0$ (c) $8x-y-5z-7=0$ (d) None of these

Solution: (a) Any plane through the first line may be written as

$$a(x-1)+b(y+1)+c(z)=0 \quad \dots(i)$$

$$\text{where, } 2a-b+3c=0 \quad \dots(ii)$$

It will pass through the second line, if the point $(0, 2, -1)$ on the second line also lies on (i)

$$\text{i.e. if } a(0-1)+b(2+1)+c(-1)=0, \text{ i.e., } -a+3b-c=0 \quad \dots(iii)$$

$$\text{Solving (ii) and (iii), we get } \frac{a}{-8} = \frac{b}{-1} = \frac{c}{5} \text{ i.e. } \frac{a}{8} = \frac{b}{1} = \frac{c}{-5}$$

$$\therefore \text{ Required plane is } 8(x-1)+1(y+1)-5(z)=0 \Rightarrow 8x+y-5z-7=0.$$

Example: 59 The plane which passes through the point $(3, 2, 0)$ and the line $\frac{x-3}{1} = \frac{y-6}{5} = \frac{z-4}{4}$ is [AIEEE 2002]

- (a) $x-y+z=1$ (b) $x+y+z=5$ (c) $x+2y-z=1$ (d) $2x-y+z=5$

Solution: (a) Any plane through the line $\frac{x-3}{1} = \frac{y-6}{5} = \frac{z-4}{4}$ is

$$a(x-3)+b(y-6)+c(z-4)=0 \quad \dots(i)$$

$$\text{where, } a+5b+4c=0 \quad \dots(ii)$$

Plane (i) passes through $(3, 2, 0)$, if

$$a(3-3)+b(2-6)+c(0-4)=0$$

$$-4b-4c=0 \text{ i.e. } b+c=0 \quad \dots(iii)$$

From equation (ii) and (iii), $a+b=0 \therefore a=-b=c$.

$$\therefore \text{ Required plane is } a(x-3)-a(y-6)+a(z-4)=0 \text{ i.e. } x-y+z-3+6-4=0 \text{ i.e. } x-y+z=1.$$

$$\text{Trick : } \begin{vmatrix} x-3 & y-6 & z-4 \\ 3-3 & 2-6 & 0-4 \\ 1 & 5 & 4 \end{vmatrix} = \begin{vmatrix} x-3 & y-6 & z-4 \\ 0 & -4 & -4 \\ 1 & 5 & 4 \end{vmatrix} \Rightarrow x-y+z=1.$$

Example: 60 The distance of point $(-1, -5, -10)$ from the point of intersection of the line $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$ and plane $x-y+z=5$ is [MP PET 2000]

- (a) 10 (b) 8 (c) 21 (d) 13

Solution: (d) Any point on the line $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12} = r$ is $(3r+2, 4r-1, 12r+2)$

This lies on $x-y+z=5$, then $3r+2-4r+1+12r+2=5$ i.e. $r=0$.

$$\therefore \text{ Point is } (2, -1, 2). \text{ Its distance from } (-1, -5, -10) \text{ is } \sqrt{9+16+144} = 13.$$

Example: 61 The value of k such that $\frac{x-4}{1} = \frac{y-2}{1} = \frac{z-k}{2}$ lies in the plane $2x-4y+z=7$ is [IIT Screening 2003]

- (a) 7 (b) -7 (c) No real value (d) 4

Solution: (a) Given, point $(4, 2, k)$ is on the line and it also passes through the plane $2x-4y+z=7 \Rightarrow 2(4)-4(2)+k=7 \Rightarrow k=7$.

Example: 62 The distance between the line $\mathbf{r} = (\mathbf{i} + \mathbf{j} + 2\mathbf{k}) + \lambda(2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k})$ and the plane $\mathbf{r} \cdot (2\mathbf{i} + \mathbf{j} - 3\mathbf{k}) = 5$ is

[Kurukshetra CEE 1996]

- (a) $\frac{5}{\sqrt{14}}$ (b) $\frac{6}{\sqrt{14}}$ (c) $\frac{7}{\sqrt{14}}$ (d) $\frac{8}{\sqrt{14}}$

Solution: (d) The given line is $\mathbf{r} = (\mathbf{i} + \mathbf{j} + 2\mathbf{k}) + \lambda(2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k})$

$$\mathbf{a} = \mathbf{i} + \mathbf{j} + 2\mathbf{k}, \mathbf{b} = 2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k}$$

Given plane, $\mathbf{r} \cdot (2\mathbf{i} + \mathbf{j} - 3\mathbf{k}) = 5 \Rightarrow \mathbf{r} \cdot \mathbf{n} = p$

$$\text{Since } \mathbf{b} \cdot \mathbf{n} = 4 + 5 - 9 = 0$$


\therefore The line is parallel to plane. Thus the distance between line and plane is equal to length of perpendicular from a point $\mathbf{a} = \mathbf{i} + \mathbf{j} + 2\mathbf{k}$ on line to given plane.

$$\text{Hence, required distance} = \left| \frac{(\mathbf{i} + \mathbf{j} + 2\mathbf{k}) \cdot (2\mathbf{i} + \mathbf{j} - 3\mathbf{k}) - 5}{\sqrt{4+1+9}} \right| = \left| \frac{2+1-6-5}{\sqrt{14}} \right| = \frac{8}{\sqrt{14}}.$$

Sphere

A sphere is the locus of a point which moves in space in such a way that its distance from a fixed point always remains constant.

The fixed point is called the centre and the constant distance is called the radius of the sphere.



7.29 General equation of sphere

The general equation of a sphere is $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ with centre $(-u, -v, -w)$

i.e. $-(1/2)$ coeff. of x , $-(1/2)$ coeff. of y , $-(1/2)$ coeff. of z and, radius $= \sqrt{u^2 + v^2 + w^2 - d}$

From the above equation, we note the following characteristics of the equation of a sphere :

- (i) It is a second degree equation in x, y, z ;
- (ii) The coefficients of x^2, y^2, z^2 are all equal;
- (iii) The terms containing the products xy, yz and zx are absent.

Note : \square The equation $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ represents,

- (i) A real sphere, if $u^2 + v^2 + w^2 - d > 0$.
- (ii) A point sphere, if $u^2 + v^2 + w^2 - d = 0$.
- (iii) An imaginary sphere, if $u^2 + v^2 + w^2 - d < 0$.

Important Tips

\Rightarrow If $u^2 + v^2 + w^2 - d < 0$, then the radius of sphere is imaginary, whereas the centre is real. Such a sphere is called "pseudo-sphere" or a "virtual sphere".

\Rightarrow The equation of the sphere contains four unknown constants u, v, w and d and therefore a sphere can be found to satisfy four conditions.

7.30 Equation in sphere in various forms

(1) Equation of sphere with given centre and radius

(i) **Cartesian form :** The equation of a sphere with centre (a, b, c) and radius R is

$$(x-a)^2 + (y-b)^2 + (z-c)^2 = R^2 \quad \dots\dots(i)$$

If the centre is at the origin, then equation (i) takes the form $x^2 + y^2 + z^2 = R^2$,

which is known as the standard form of the equation of the sphere.

(ii) **Vector form :** The equation of sphere with centre at $C(\mathbf{c})$ and radius ' a ' is $|\mathbf{r} - \mathbf{c}| = a$.

(2) Diameter form of the equation of a sphere

(i) **Cartesian form :** If (x_1, y_1, z_1) and (x_2, y_2, z_2) are the co-ordinates of the extremities of a diameter of a sphere, then its equation is $(x-x_1)(x-x_2) + (y-y_1)(y-y_2) + (z-z_1)(z-z_2) = 0$.

(ii) **Vector form** : If the position vectors of the extremities of a diameter of a sphere are \mathbf{a} and \mathbf{b} , then its equation is $(\mathbf{r} - \mathbf{a}) \cdot (\mathbf{r} - \mathbf{b}) = 0$ or $|\mathbf{r}|^2 - \mathbf{r} \cdot (\mathbf{a} + \mathbf{b}) + \mathbf{a} \cdot \mathbf{b} = 0$.

7.31 Section of a sphere by a plane

Consider a sphere intersected by a plane. The set of points common to both sphere and plane is called a plane section of a sphere. The plane section of a sphere is always a circle. The equations of the sphere and the plane taken together represent the plane section.

Let C be the centre of the sphere and M be the foot of the perpendicular from C on the plane. Then M is the centre of the circle and radius of the circle is given by


$$PM = \sqrt{CP^2 - CM^2}$$

The centre M of the circle is the point of intersection of the plane and line CM which passes through C and is perpendicular to the given plane.

Centre : The foot of the perpendicular from the centre of the sphere to the plane is the centre of the circle.

$$(\text{radius of circle})^2 = (\text{radius of sphere})^2 - (\text{perpendicular from centre of sphere on the plane})^2$$

Great circle : The section of a sphere by a plane through the centre of the sphere is a great circle. Its centre and radius are the same as those of the given sphere.



7.32 Condition of tangency of a plane to a sphere

A plane touches a given sphere if the perpendicular distance from the centre of the sphere to the plane is equal to the radius of the sphere.

(1) **Cartesian form** : The plane $lx + my + nz = p$ touches the sphere $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$, if $(ul + vm + wn - p)^2 = (l^2 + m^2 + n^2)(u^2 + v^2 + w^2 - d)$

(2) **Vector form** : The plane $\mathbf{r} \cdot \mathbf{n} = d$ touches the sphere $|\mathbf{r} - \mathbf{a}| = R$ if $\frac{|\mathbf{a} \cdot \mathbf{n} - d|}{|\mathbf{n}|} = R$.

Important Tips

☞ Two spheres S_1 and S_2 with centres C_1 and C_2 and radii r_1 and r_2 respectively

(i) Do not meet and lies farther apart iff $|C_1C_2| > r_1 + r_2$

(ii) Touch internally iff $|C_1C_2| = |r_1 - r_2|$

(iii) Touch externally iff $|C_1C_2| = r_1 + r_2$

(iv) Cut in a circle iff $|r_1 - r_2| < |C_1C_2| < r_1 + r_2$

(v) One lies within the other if $|C_1C_2| < |r_1 - r_2|$.

When two spheres touch each other the common tangent plane is $S_1 - S_2 = 0$ and when they cut in a circle, the plane of the circle is

$$S_1 - S_2 = 0 ; \text{coefficients of } x^2, y^2, z^2 \text{ being unity in both the cases.}$$

☞ Let p be the length of perpendicular drawn from the centre of the sphere $x^2 + y^2 + z^2 = r^2$ to the plane $Ax + By + Cz + D = 0$, then

(i) The plane cuts the sphere in a circle iff $p < r$ and in this case, the radius of circle is $\sqrt{r^2 - p^2}$.

(ii) The plane touches the sphere iff $p = r$.

(iii) The plane does not meet the sphere iff $p > r$.

Equation of concentric sphere : Any sphere concentric with the sphere $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ is $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + \lambda = 0$, where λ is some real which makes it a sphere.

7.33 Intersection of straight line and a sphere

Let the equations of the sphere and the straight line be $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ (i)

And $\frac{x - \alpha}{l} = \frac{y - \beta}{m} = \frac{z - \gamma}{n} = r$ (say)(ii)

Any point on the line (ii) is $(\alpha + lr, \beta + mr, \gamma + nr)$.

If this point lies on the sphere (i) then we have,

$$(\alpha + lr)^2 + (\beta + mr)^2 + (\gamma + nr)^2 + 2u(\alpha + lr) + 2v(\beta + mr) + 2w(\gamma + nr) + d = 0$$

$$\text{or, } r^2[l^2 + m^2 + n^2] + 2r[l(u + \alpha) + m(v + \beta) + n(w + \gamma)] + (\alpha^2 + \beta^2 + \gamma^2 + 2u\alpha + 2v\beta + 2w\gamma + d) = 0 \text{(iii)}$$

This is a quadratic equation in r and so gives two values of r and therefore the line (ii) meets the sphere (i) in two points which may be real, coincident and imaginary, according as root of (iii) are so.

Note : \square If l, m, n are the actual d.c.'s of the line, then $l^2 + m^2 + n^2 = 1$ and then the equation (iii) can be simplified.

7.34 Angle of intersection of two spheres

The angle of intersection of two spheres is the angle between the tangent planes to them at their point of intersection. As the radii of the spheres at this common point are normal to the tangent planes so this angle is also equal to the angle between the radii of the spheres at their point of intersection.

If the angle of intersection of two spheres is a right angle, the spheres are said to be orthogonal.

Condition for orthogonality of two spheres

Let the equation of the two spheres be

$$x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0 \text{(i)}$$

$$\text{and } x^2 + y^2 + z^2 + 2u'x + 2v'y + 2w'z + d' = 0 \text{(ii)}$$

If the sphere (i) and (ii) cut orthogonally, then $2uu' + 2vv' + 2ww' = d + d'$, which is the required condition.

Note : \square If the spheres $x^2 + y^2 + z^2 = a^2$ and $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ cut orthogonally, then $d = a^2$.

\square Two spheres of radii r_1 and r_2 cut orthogonally, then the radius of the common circle is $\frac{r_1 r_2}{\sqrt{r_1^2 + r_2^2}}$.

Example: 63

The centre of sphere passing through four points $(0, 0, 0)$, $(0, 2, 0)$, $(1, 0, 0)$ and $(0, 0, 4)$ is

[MP PET 2002]

- (a) $\left(\frac{1}{2}, 1, 2\right)$ (b) $\left(-\frac{1}{2}, 1, 2\right)$ (c) $\left(\frac{1}{2}, 1, -2\right)$ (d) $\left(1, \frac{1}{2}, 2\right)$

Solution: (a)

Let the equation of sphere be $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$

\therefore It passes through $(0, 0, 0)$, $\therefore d = 0$

Also, It passes through $(0, 2, 0)$ i.e., $v = -1$

Also, It passes through (1, 0, 0) i.e., $u = -1/2$

Also, it passes through (0, 0, 4) i.e., $w = -2$

\therefore Centre $(-u, -v, -w) = (1/2, 1, 1/2)$

Example: 64 The equation $|\mathbf{r}|^2 - \mathbf{r} \cdot (2\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}) - 10 = 0$ represents a [DCE 1998]

- (a) Plane (b) Sphere of radius 4 (c) Sphere of radius 3 (d) None of these

Solution: (b) The given equation is $|\mathbf{r}|^2 - \mathbf{r} \cdot (2\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}) - 10 = 0$

$$\Rightarrow x^2 + y^2 + z^2 - 2x - 4y + 2z - 10 = 0,$$

which is the equation of sphere, whose centre is (1, 2, -1) and radius $= \sqrt{1+4+1+10} = 4$.

Example: 65 The intersection of the spheres $x^2 + y^2 + z^2 + 7x - 2y - z = 13$ and $x^2 + y^2 + z^2 - 3x + 3y + 4z = 8$ is the same as the intersection of one of the sphere and the plane [AIEEE 2004]

- (a) $2x - y - z = 1$ (b) $x - 2y - z = 1$ (c) $x - y - 2z = 1$ (d) $x - y - z = 1$

Solution: (a) We have the spheres $x^2 + y^2 + z^2 + 7x - 2y - z - 13 = 0$ and $x^2 + y^2 + z^2 - 3x + 3y + 4z - 8 = 0$

Required plane is $S_1 - S_2 = 0$

$$\therefore (7x + 3x) - (2y + 3y) - (z + 4z) - 5 = 0$$

$$\text{i.e. } 10x - 5y + (-5z) - 5 = 0 \Rightarrow 2x - y - z = 1.$$

Example: 66 The radius of the circle in which the sphere $x^2 + y^2 + z^2 + 2x - 2y - 4z - 19 = 0$ is cut by the plane $x + 2y + 2z + 7 = 0$ is [AIEEE 2003]


- (a) 1 (b) 2 (c) 3 (d) 4

Solution: (c) For sphere $x^2 + y^2 + z^2 + 2x - 2y - 4z - 19 = 0$, Centre O is $(-1, 1, 2)$ and radius $= \sqrt{1+1+4+19} = 5$,

Now, OL = length of perpendicular from O to plane $x + 2y + 2z + 7 = 0$ is

$$= \frac{-1 + 2 + 4 + 7}{\sqrt{1+4+4}} = \frac{12}{3} = 4, \text{ i.e. } OL = 4.$$

$$\text{In } \triangle OLB, LB = \sqrt{OB^2 - OL^2} = \sqrt{25 - 16} = 3.$$



Example: 67 The radius of circular section of the sphere $|\mathbf{r}| = 5$ by the plane $\mathbf{r} \cdot (\mathbf{i} + \mathbf{j} + \mathbf{k}) = 4\sqrt{3}$ is [DCE 1999; AMU 1991]

- (a) 2 (b) 3 (c) 4 (d) 6

Solution: (b) Radius of the sphere = 5

Given plane is $x + y + z - 4\sqrt{3} = 0$

$$\text{Length of the perpendicular from the centre } (0, 0, 0) \text{ of the sphere to the plane} = \frac{4\sqrt{3}}{\sqrt{1+1+1}} = 4$$

$$\text{Hence, radius of circular section} = \sqrt{25 - 16} = 3.$$

Example: 68 The shortest distance from the plane $12x + 4y + 3z = 327$ to the sphere $x^2 + y^2 + z^2 + 4x - 2y - 6z = 155$ is [AIEEE 2003]

- (a) 26 (b) $11\frac{4}{13}$ (c) 13 (d) 39

Solution: (c) Centre of sphere is $(-2, 1, 3)$

$$\text{Radius of sphere is } \sqrt{4+1+9+155} = 13$$

$$\text{Distance of centre from plane} = \frac{-24 + 4 + 9 - 327}{\sqrt{144 + 16 + 9}} = \frac{338}{13}$$

$$\therefore \text{Plane cuts the sphere and hence } S.D. = \frac{338}{13} - 13 = \frac{169}{13} = 13.$$



Assignment

System of Co-ordinates

Basic Level

- From which of the following the distance of the point $(1,2,3)$ is $\sqrt{10}$
 (a) Origin (b) x -axis (c) y -axis (d) z -axis
- If $A(1,2,3); B(-1,-1,-1)$ be the points, then the distance AB is [MP PET 2001]
 (a) $\sqrt{5}$ (b) $\sqrt{21}$ (c) $\sqrt{29}$ (d) None of these
- Perpendicular distance of the point $(3,4,5)$ from the y -axis, is [MP PET 1994]
 (a) $\sqrt{34}$ (b) $\sqrt{41}$ (c) 4 (d) 5
- Distance between the points $(1,3,2)$ and $(2,1,3)$ is [MP PET 1988]
 (a) 12 (b) $\sqrt{12}$ (c) $\sqrt{6}$ (d) 6
- The shortest distance of the point (a,b,c) from the x -axis is [MP PET 1999; DCE 1999]
 (a) $\sqrt{(a^2 + b^2)}$ (b) $\sqrt{(b^2 + c^2)}$ (c) $\sqrt{(c^2 + a^2)}$ (d) $\sqrt{(a^2 + b^2 + c^2)}$
- Points $(1,1,1)$, $(-2,4,1)$, $(-1,5,5)$ and $(2,2,5)$ are the vertices of
 (a) Rectangle (b) Square (c) Parallelogram (d) Trapezium
- The triangle formed by the points $(0,7,10)$, $(-1,6,6)$ $(-4,9,6)$ is [Rajasthan PET 2001]
 (a) Equilateral (b) Isosceles (c) Right angled (d) Right angled isosceles
- The points $A(5,-1,1)$; $B(7,-4,7)$; $C(1,-6,10)$ and $D(-1,-3,4)$ are vertices of a [Rajasthan PET 2000]
 (a) Square (b) Rhombus (c) Rectangle (d) None of these
- The coordinates of a point which is equidistant from the points $(0,0,0)$, $(a,0,0)$, $(0,b,0)$ and $(0,0,c)$ are given by [MP PET 1993; Rajasthan PET 2003]
 (a) $\left(\frac{a}{2}, \frac{b}{2}, \frac{c}{2}\right)$ (b) $\left(-\frac{a}{2}, -\frac{b}{2}, \frac{c}{2}\right)$ (c) $\left(\frac{a}{2}, -\frac{b}{2}, -\frac{c}{2}\right)$ (d) $\left(-\frac{a}{2}, \frac{b}{2}, -\frac{c}{2}\right)$
- If $A(1,2,-1)$ and $B(-1,0,1)$ are given, then the coordinates of P which divides AB externally in the ratio $1:2$, are [MP PET 1997]
 (a) $\frac{1}{3}(1,4,-1)$ (b) $(3,4,-3)$ (c) $\frac{1}{3}(3,4,-3)$ (d) None of these
- The coordinates of the point which divides the join of the points $(2,-1,3)$ and $(4,3,1)$ in the ratio $3:4$ internally are given by [MP PET 1997]
 (a) $\frac{2}{7}, \frac{20}{7}, \frac{10}{7}$ (b) $\frac{15}{7}, \frac{20}{7}, \frac{3}{7}$ (c) $\frac{10}{7}, \frac{15}{7}, \frac{2}{7}$ (d) $\frac{20}{7}, \frac{5}{7}, \frac{15}{7}$
- Points $(-2,4,7)$, $(3,-6,-8)$ and $(1,-2,-2)$ are [AI CBSE 1982]

Three Dimensional Co-ordinate Geometry

- (a) Collinear (b) Vertices of an equilateral triangle
(c) Vertices of an isosceles triangle (d) None of these
13. Which of the following set of points are non-collinear [MP PET 1990]
(a) (1, -1, 1), (-1, 1, 1), (0, 0, 1) (b) (1, 2, 3), (3, 2, 1), (2, 2, 2)
(c) (-2, 4, -3), (4, -3, -2), (-3, -2, 4) (d) (2, 0, -1), (3, 2, -2), (5, 6, -4)
14. If the points (-1, 3, 2), (-4, 2, -2) and (5, 5, λ) are collinear, then $\lambda =$
(a) -10 (b) 5 (c) -5 (d) 10
15. The area of triangle whose vertices are (1, 2, 3), (2, 5, -1) and (-1, 1, 2) is [Kerala (Engg.) 2002]
(a) 150 sq. units (b) 145 sq. units (c) $\frac{\sqrt{155}}{2}$ sq. units (d) $\frac{155}{2}$ sq. units
16. Volume of a tetrahedron is K (area of one face) (length of perpendicular from the opposite vertex upon it), where K is
(a) $\frac{1}{2}$ (b) $\frac{1}{3}$ (c) $\frac{1}{4}$ (d) $\frac{1}{6}$
17. A point moves so that the sum of its distances from the points (4,0,0) and (-4,0,0) remains 10. The locus of the point is [MP PET 1988]
(a) $9x^2 - 25y^2 + 25z^2 = 225$ (b) $9x^2 + 25y^2 - 25z^2 = 225$
(c) $9x^2 + 25y^2 + 25z^2 = 225$ (d) $9x^2 + 25y^2 + 25z^2 + 225 = 0$
18. If the sum of the squares of the distances of a point from the three coordinate axes be 36, then its distance from the origin is
(a) 6 (b) $3\sqrt{2}$ (c) $2\sqrt{3}$ (d) None of these
19. All the points on the x -axis have [MP PET 1988]
(a) $x = 0$ (b) $y = 0$ (c) $x = 0, y = 0$ (d) $y = 0, z = 0$
20. The equations $|x| = p, |y| = p, |z| = p$ in xyz space represent [Orissa JEE 2002]
(a) Cube (b) Rhombus (c) Sphere of radius p (d) Point (p, p, p)
21. The orthocentre of the triangle with vertices (1,2,3), (2,3,1) and (3,1,2) is
(a) (1, 1, 1) (b) (2, 2, 2) (c) (6, 6, 6) (d) None of these
22. If $a + b + c = \lambda$, then circumcentre of the triangle with vertices (a, b, c) ; (b, c, a) and (c, a, b) is
(a) $(\lambda, \lambda, \lambda)$ (b) $(\lambda/2, \lambda/2, \lambda/2)$ (c) $(\lambda/3, \lambda/3, \lambda/3)$ (d) None of these
23. (-1,6,6), (-4,9,6) are two vertices of $\triangle ABC$. If its centroid be $(-5/3, 22/3, 22/3)$, then its third vertex is
(a) (0, 7, 10) (b) (7, 0, 10) (c) (10, 0, 7) (d) None of these
24. If points (2, 3, 4), (5, a , 6) and (7, 8, b) are collinear, then values of a and b are [AISSE 1989]
(a) $a = 6, b = \frac{-22}{3}$ (b) $a = 6, b = \frac{22}{3}$ (c) $a = \frac{22}{3}, b = 6$ (d) $a = \frac{-22}{3}, b = -6$

Direction cosines and Projection

Basic Level

356 Three Dimensional Co-ordinate

25. If a line makes angles of 30° and 45° with x -axis and y -axis, then the angle made by it with z -axis is
 (a) 45° (b) 60° (c) 120° (d) None of these
26. If a straight line in space is equally inclined to the coordinate axes, the cosine of its angle of inclination to any one of the axes is
 [MP PET 1992]
 (a) $\frac{1}{3}$ (b) $\frac{1}{2}$ (c) $\frac{1}{\sqrt{3}}$ (d) $\frac{1}{\sqrt{2}}$
27. If the length of a vector be 21 and direction ratios be 2, -3, 6, then its direction cosines are
 (a) $\frac{2}{21}, \frac{-1}{7}, \frac{2}{7}$ (b) $\frac{2}{7}, \frac{-3}{7}, \frac{6}{7}$ (c) $\frac{2}{7}, \frac{3}{7}, \frac{6}{7}$ (d) None of these
28. If O is the origin, $OP = 3$ with d.r.'s -1, 2, -2 then the co-ordinates of P are
 [Rajasthan PET 2000]
 (a) (-1, 2, -2) (b) (1, 2, 2) (c) $\left(-\frac{1}{9}, \frac{2}{9}, -\frac{2}{9}\right)$ (d) (3, 6, -9)
29. The numbers 3, 4, 5 can be
 (a) Direction cosines of a line (b) Direction ratios of a line in space
 (c) Coordinates of a point on the plane $y = 4, z = 0$ (d) Co-ordinates of a point on the plane $x + y - z = 0$
30. If l, m, n are the d.c.'s of a line, then
 (a) $l^2 + m^2 + n^2 = 0$ (b) $l^2 + m^2 + n^2 = 1$ (c) $l + m + n = 1$ (d) $l = m = n = 1$
31. If a line lies in the octant $OXYZ$ and it makes equal angles with the axes, then
 [MP PET 2001]
 (a) $l = m = n = \frac{1}{\sqrt{3}}$ (b) $l = m = n = \pm \frac{1}{\sqrt{3}}$ (c) $l = m = n = -\frac{1}{\sqrt{3}}$ (d) $l = m = n = \pm \frac{1}{\sqrt{2}}$
32. If a line makes equal angle with axes, then its direction ratios will be
 (a) 1, 2, 3 (b) 3, 1, 2 (c) 3, 2, 1 (d) 1, 1, 1
33. The coordinates of the point P are (x, y, z) and the direction cosines of the line OP , when O is the origin, are l, m, n . If $OP = r$, then
 (a) $l = x, m = y, n = z$ (b) $l = xr, m = yr, n = zr$ (c) $x = lr, y = mr, z = nr$ (d) None of these
34. The direction ratios of the diagonals of a cube which joins the origin to the opposite corner are (when the 3 concurrent edges of the cube are coordinate axes)
 [MP PET 1996]
 (a) $\frac{2}{\sqrt{3}}, \frac{2}{\sqrt{3}}, \frac{2}{\sqrt{3}}$ (b) -1, 1, -1 (c) 2, -2, 1 (d) 1, 2, 3
35. If the direction ratios of a line are 1, -3, 2, then the direction cosines of the line are
 [MP PET 1997]
 (a) $\frac{1}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{2}{\sqrt{14}}$ (b) $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$ (c) $\frac{-1}{\sqrt{14}}, \frac{3}{\sqrt{14}}, \frac{-2}{\sqrt{14}}$ (d) $\frac{-1}{\sqrt{14}}, \frac{-2}{\sqrt{14}}, \frac{-3}{\sqrt{14}}$
36. If a line make α, β, γ with the positive direction of x, y and z -axis respectively. Then $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma$ is
 [Orissa JEE 2002; MP PET 2002]
 (a) $1/2$ (b) $-1/2$ (c) -1 (d) 1
37. The direction-cosines of the line joining the points (4, 3, -5) and (-2, 1, -8) are
 [MP PET 2001; Kurukshetra CEE 1998]
 (a) $\left(\frac{6}{7}, \frac{2}{7}, \frac{3}{7}\right)$ (b) $\left(\frac{2}{7}, \frac{3}{7}, \frac{6}{7}\right)$ (c) $\left(\frac{6}{7}, \frac{3}{7}, \frac{2}{7}\right)$ (d) None of these
38. The direction ratios of the line joining the points (4, 3, -5) and (-2, 1, -8) are
 [AI CBSE 1984; MP PET 1988]

- (a) $\left(\frac{6}{7}, \frac{2}{7}, \frac{3}{7}\right)$ (b) 6, 2, 3 (c) 2, 4, -13 (d) None of these
39. The coordinates of a point P are (3, 12, 4) with respect to origin O , then the direction cosines of OP are [MP PET 1996]
- (a) 3, 12, 4 (b) $\frac{1}{4}, \frac{1}{3}, \frac{1}{2}$ (c) $\frac{3}{\sqrt{13}}, \frac{1}{\sqrt{13}}, \frac{2}{\sqrt{13}}$ (d) $\frac{3}{13}, \frac{12}{13}, \frac{4}{13}$
40. The direction cosines of a line segment AB are $\frac{-2}{\sqrt{17}}, \frac{3}{\sqrt{17}}, \frac{-2}{\sqrt{17}}$. If $AB = \sqrt{17}$ and the coordinates of A are (3, -6, 10), then the coordinates of B are
- (a) (1, -2, 4) (b) (2, 5, 8) (c) (-1, 3, -8) (d) (1, -3, 8)
41. If $\left(\frac{1}{2}, \frac{1}{3}, n\right)$ are the direction cosines of a line, then the value of n is [Kerala (Engg.) 2002]
- (a) $\frac{\sqrt{23}}{6}$ (b) $\frac{23}{6}$ (c) $\frac{2}{3}$ (d) $\frac{3}{2}$
42. If a line makes the angle α, β, γ with three dimensional coordinate axes respectively, then $\cos 2\alpha + \cos 2\beta + \cos 2\gamma =$ [MP PET 1994,95,99; Rajasthan PET 2003]
- (a) -2 (b) -1 (c) 1 (d) 2
43. A line makes angles of 45° and 60° with the positive axes of X and Y respectively. The angle made by the same line with the positive axis of Z , is [MP PET 1997]
- (a) 30° or 60° (b) 60° or 90° (c) 90° or 120° (d) 60° or 120°
44. If α, β, γ be the angles which a line makes with the positive direction of coordinate axes, then $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma =$ [Rajasthan PET 2000; AMU 2002; MP PET 1989,98,2000,03; Kerala (Engg.) 2001]
- (a) 2 (b) 1 (c) 3 (d) 0
45. A line makes angles α, β, γ with the coordinate axes. If $\alpha + \beta = 90^\circ$, then $\gamma =$
- (a) 0° (b) 90° (c) 180° (d) None of these
46. The coordinates of the points P and Q are (x_1, y_1, z_1) and (x_2, y_2, z_2) respectively, then the projection of the line PQ on the line whose direction cosines are l, m, n , will be
- (a) $(x_2 - x_1)l + (y_2 - y_1)m + (z_2 - z_1)n$ (b) $\left(\frac{x_2 - x_1}{l}\right) + \left(\frac{y_2 - y_1}{m}\right) + \left(\frac{z_2 - z_1}{n}\right)$
- (c) $\frac{x_1}{l} + \frac{y_1}{m} + \frac{z_1}{n}$ (d) $\frac{x_2}{l} + \frac{y_2}{m} + \frac{z_2}{n}$
47. The projection of the line segment joining the points (-1, 0, 3) and (2, 5, 1) on the line whose direction ratios are 6, 2, 3, is [AI CBSE 1985]
- (a) $10/7$ (b) $22/7$ (c) $18/7$ (d) None of these
48. The projection of any line on coordinate axes be respectively 3, 4, 5, then its length is [MP PET 1995; Rajasthan PET 2001]
- (a) 12 (b) 50 (c) $5\sqrt{2}$ (d) None of these
49. If θ is the angle between the lines AB and CD , then projection of line segment AB on line CD is [MP PET 1995]
- (a) $AB \sin \theta$ (b) $AB \cos \theta$ (c) $AB \tan \theta$ (d) $CD \cos \theta$

358 Three Dimensional Co-ordinate

50. The projections of a line on the co-ordinate axes are 4, 6, 12. The direction cosines of the line are
 (a) $\frac{2}{7}, \frac{3}{7}, \frac{6}{7}$ (b) 2, 3, 6 (c) $\frac{2}{11}, \frac{3}{11}, \frac{6}{11}$ (d) None of these
51. The projections of segment PQ on the coordinate planes are -9, 12, -8 respectively. The direction cosines of PQ are [Pb. CET 1998]
 (a) $\langle -\frac{9}{\sqrt{17}}, \frac{12}{\sqrt{17}}, \frac{-8}{\sqrt{17}} \rangle$ (b) $\langle -9, 12, -8 \rangle$
 (c) $\langle \frac{-9}{289}, \frac{12}{289}, \frac{-8}{289} \rangle$ (d) $\langle \frac{-9}{17}, \frac{12}{17}, \frac{-8}{17} \rangle$
52. The projections of a line segment on x, y, z axes are 12, 4, 3. The length and the direction cosines of the line segments are
 [Kerala (Engg.) 2000]
 (a) $13, \langle 12/13, 4/13, 3/13 \rangle$ (b) $19, \langle 12/19, 4/19, 3/19 \rangle$ (c) $11, \langle 12/11, 4/11, 3/11 \rangle$ (d) None of these
53. The coordinates of A and B be (1, 2, 3) and (7, 8, 7), then the projections of the line segment AB on the coordinate axes are
 (a) 6, 6, 4 (b) 4, 6, 4 (c) 3, 3, 2 (d) 2, 3, 2
54. A line segment (vector) has length 21 and direction ratios (2, -3, 6). If the line makes an obtuse angle with x -axis, the components of the line (vector) are
 (a) 6, -9, 18 (b) 2, -3, 6 (c) -18, 27, -54 (d) -6, 9, -18

Angle between Two Lines

Basic Level

55. The angle between the pair of lines with direction ratios (1, 1, 2) and $(\sqrt{3}-1, -\sqrt{3}-1, 4)$ is [MP PET 1997, 2000]
 (a) 30° (b) 45° (c) 60° (d) 90°
56. The angle between a line with direction ratios 2 : 2 : 1 and a line joining (3, 1, 4) to (7, 2, 12) is [DCE 2002]
 (a) $\cos^{-1}(2/3)$ (b) $\cos^{-1}(-2/3)$ (c) $\tan^{-1}(2/3)$ (d) None of these
57. The angle between the lines whose direction cosines are proportional to (1, 2, 1) and (2, -3, 6) is
 (a) $\cos^{-1}\left(\frac{2}{7\sqrt{6}}\right)$ (b) $\cos^{-1}\left(\frac{1}{7\sqrt{6}}\right)$ (c) $\cos^{-1}\left(\frac{3}{7\sqrt{6}}\right)$ (d) $\cos^{-1}\left(\frac{5}{7\sqrt{6}}\right)$
58. If the vertices of a triangle are $A(1, 4, 2)$, $B(-2, 1, 2)$, $C(2, -3, 4)$, then the angle B is equal to
 (a) $\cos^{-1}(1/\sqrt{3})$ (b) $\pi/2$ (c) $\cos^{-1}(\sqrt{6}/3)$ (d) $\cos^{-1}\sqrt{3}$
59. If the coordinates of the points P, Q, R, S be (1, 2, 3), (4, 5, 7), (-4, 3, -6) and (2, 0, 2) respectively, then
 (a) $PQ \parallel RS$ (b) $PQ \perp RS$ (c) $PQ = RS$ (d) None of these
60. If the coordinates of the points A, B, C, D be (1, 2, 3), (4, 5, 7), (-4, 3, -6) and (2, 9, 2) respectively, then the angle between the lines AB and CD is
 (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{3}$ (d) None of these
61. If the angle between the lines whose direction ratios are 2, -1, 2 and $a, 3, 5$ be 45° , then $a =$

Three Dimensional Co-ordinate Geometry

- (a) 1 (b) 2 (c) 3 (d) 4
62. If O be the origin and $P(2, 3, 4)$ and $Q(1, b, 1)$ be two points such that $OP \perp OQ$, then $b =$
 (a) 2 (b) -2 (c) No such real b exists (d) None of these
63. If d.r.'s of two straight lines are 5, -12, 13 and -3, 4, 5 then, angle between them is [Rajasthan PET 2001]
 (a) $\cos^{-1}\left(\frac{2}{65}\right)$ (b) $\cos^{-1}\left(\frac{1}{65}\right)$ (c) $\cos^{-1}\left(\frac{3}{65}\right)$ (d) $\frac{\pi}{3}$
64. If direction ratio of two lines are a_1, b_1, c_1 and a_2, b_2, c_2 then these lines are parallel if and only if
 (a) $a_1 = a_2, b_1 = b_2, c_1 = c_2$ (b) $a_1 a_2 + b_1 b_2 + c_1 c_2 = 0$ (c) $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ (d) None of these
65. If $A(k, 1, -1)$, $B(2k, 0, 2)$ and $C(2+2k, k, 1)$ be such that the line $AB \perp BC$, then the value of k will be
 (a) 1 (b) 2 (c) 3 (d) 0
66. $A(a, 7, 10)$, $B(-1, 6, 6)$ and $C(-4, 9, 6)$ are the vertices of a right angled isosceles triangle. If $\angle ABC = 90^\circ$, then $a =$
 (a) 0 (b) 2 (c) -1 (d) -3

Advance Level

67. The angle between two diagonals of a cube will be [MP PET 1996, 97, 2000; Rajasthan PET 2000, 02]
 (a) $\sin^{-1} \frac{1}{3}$ (b) $\cos^{-1} \frac{1}{3}$ (c) Constant (d) Variable
68. If a line makes angles $\alpha, \beta, \gamma, \delta$ with the four diagonals of a cube, then the value of $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta =$ [Rajasthan PET 2002]
 (a) 1 (b) $\frac{4}{3}$ (c) Constant (d) Variable
69. The angle between the lines whose direction cosines satisfy the equations $l+m+n=0, l^2+m^2-n^2=0$ is given by [MP PET 1993; Rajasthan PET 2001]
 (a) $\frac{2\pi}{3}$ (b) $\frac{\pi}{6}$ (c) $\frac{5\pi}{6}$ (d) $\frac{\pi}{3}$
70. If three mutually perpendicular lines have direction cosines $(l_1, m_1, n_1), (l_2, m_2, n_2)$, and (l_3, m_3, n_3) , then the line having direction cosines $l_1 + l_2 + l_3, m_1 + m_2 + m_3$ and $n_1 + n_2 + n_3$ make an angle ofwith each other
 (a) 0° (b) 30° (c) 60° (d) 90°
71. The straight lines whose direction cosines are given by $al+bm+cn=0, flm+gnl+hlm=0$ are perpendicular, if
 (a) $\frac{f}{a} + \frac{g}{b} + \frac{h}{c} = 0$ (b) $\sqrt{\frac{a}{f}} + \sqrt{\frac{b}{g}} + \sqrt{\frac{c}{h}} = 0$ (c) $\sqrt{af} = \sqrt{bg} = \sqrt{ch}$ (d) $\sqrt{\frac{a}{f}} = \sqrt{\frac{b}{g}} = \sqrt{\frac{c}{h}}$
72. The angle between the lines whose direction cosines are connected by the relations $l+m+n=0$ and $2lm+2nl-mn=0$, is
 (a) $\frac{\pi}{3}$ (b) $\frac{2\pi}{3}$ (c) π (d) None of these
73. $A(3, 2, 0)$, $B(5, 3, 2)$, $C(-9, 6, -3)$ are three points forming a triangle and AD is the bisector of the $\angle BAC$, then coordinates of D are

360 Three Dimensional Co-ordinate

- (a) $\left(\frac{17}{16}, \frac{57}{16}, \frac{28}{16}\right)$ (b) $\left(\frac{38}{16}, \frac{57}{16}, \frac{17}{16}\right)$ (c) $\left(\frac{38}{16}, \frac{17}{16}, \frac{57}{16}\right)$ (d) $\left(\frac{57}{16}, \frac{38}{16}, \frac{17}{16}\right)$
74. The direction cosines of two lines at right angles are $\langle l_1, m_1, n_1 \rangle$ and $\langle l_2, m_2, n_2 \rangle$. Then the d.c. of a line \perp to both the given lines are
 (a) $\langle m_1 n_2 - m_2 n_1, n_1 l_2 - n_2 l_1, l_1 m_2 - l_2 m_1 \rangle$ (b) $\langle l_1 + l_2, m_1 + m_2, n_1 + n_2 \rangle$
 (c) $\langle l_1 - l_2, m_1 - m_2, n_1 - n_2 \rangle$ (d) None of these
75. Three lines drawn from origin with direction cosines $l_1, m_1, n_1; l_2, m_2, n_2; l_3, m_3, n_3$ are coplanar iff $\begin{vmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{vmatrix} = 0$, since
 (a) All lines pass through origin (b) It is possible to find a line perpendicular to all these lines
 (c) Intersecting lines are coplanar (d) None of these
76. The direction cosines of a variable line in two adjacent positions are l, m, n and $l + \delta l, m + \delta m, n + \delta n$. If angle between these two positions is $\delta\theta$, where $\delta\theta$ is a small angle, then $\delta\theta^2$ is equal to
 (a) $\delta l^2 + \delta m^2 + \delta n^2$ (b) $\delta l + \delta m + \delta n$ (c) $\delta l \cdot \delta m + \delta m \cdot \delta n + \delta n \cdot \delta l$ (d) None of these
77. If direction cosines of two lines OA and OB are respectively proportional to 1, -2, -1 and 3, -2, 3 then direction cosine of line perpendicular to given both lines are
 (a) $\pm 4 / \sqrt{29}, \pm 3 / \sqrt{29}, \pm 2 / \sqrt{29}$, (b) $\pm 4 / \sqrt{29}, \pm 3 / \sqrt{29}, \mp 2 / \sqrt{29}$
 (c) $\pm 4 / \sqrt{29}, \pm 2 / \sqrt{29}, \pm 3 / \sqrt{29}$, (d) None of these
78. A mirror and a source of light are situated at the origin O and at a point on OX respectively. A ray of light from the source strikes the mirror and is reflected. If the d.r's of the normal to the plane are 1, -1, 1, then d.c's of the reflected ray are
 (a) $\frac{1}{3}, \frac{2}{3}, \frac{2}{3}$ (b) $-\frac{1}{3}, \frac{2}{3}, \frac{2}{3}$ (c) $-\frac{1}{3}, -\frac{2}{3}, -\frac{2}{3}$ (d) $-\frac{1}{3}, -\frac{2}{3}, \frac{2}{3}$

Straight Line

Basic Level

79. The equation of straight line passing through the point (a, b, c) and parallel to z-axis, is [MP PET 1995]
 (a) $\frac{x-a}{1} = \frac{y-b}{1} = \frac{z-c}{0}$ (b) $\frac{x-a}{0} = \frac{y-b}{1} = \frac{z-c}{1}$ (c) $\frac{x-a}{1} = \frac{y-b}{0} = \frac{z-c}{0}$ (d) $\frac{x-a}{0} = \frac{y-b}{0} = \frac{z-c}{1}$
80. Equation of x-axis is [MP PET 2002]
 (a) $\frac{x}{1} = \frac{y}{1} = \frac{z}{1}$ (b) $\frac{x}{0} = \frac{y}{1} = \frac{z}{1}$ (c) $\frac{x}{1} = \frac{y}{0} = \frac{z}{0}$ (d) $\frac{x}{0} = \frac{y}{0} = \frac{z}{1}$
81. The equation of straight line passing through the points (a, b, c) and $(a-b, b-c, c-a)$, is [MP PET 1994]
 (a) $\frac{x-a}{a-b} = \frac{y-b}{b-c} = \frac{z-c}{c-a}$ (b) $\frac{x-a}{b} = \frac{y-b}{c} = \frac{z-c}{a}$ (c) $\frac{x-a}{a} = \frac{y-b}{b} = \frac{z-c}{c}$ (d) $\frac{x-a}{2a-b} = \frac{y-b}{2b-c} = \frac{z-c}{2c-a}$
82. The equation of a line passing through the point $(-3, 2, -4)$ and equally inclined to the axes, are
 (a) $x-3 = y+2 = z-4$ (b) $x+3 = y-2 = z+4$ (c) $\frac{x+3}{1} = \frac{y-2}{2} = \frac{z+4}{3}$ (d) None of these
83. The straight line through (a, b, c) and parallel to x-axis are [DCE 1992]

- (a) $\frac{x-a}{1} = \frac{y-b}{0} = \frac{z-c}{0}$ (b) $\frac{x-a}{0} = \frac{y-b}{1} = \frac{z-c}{0}$ (c) $\frac{x-a}{0} = \frac{y-b}{0} = \frac{z-c}{1}$ (d) $\frac{x-a}{1} = \frac{y-b}{1} = \frac{z-c}{1}$
84. Equation of the line passing through the point (1, 2, 3) and parallel to the line $\frac{x-6}{12} = \frac{y-2}{4} = \frac{z+7}{5}$ is given by
- (a) $\frac{x+1}{12} = \frac{y+2}{4} = \frac{z+3}{5}$ (b) $\frac{x-1}{l} = \frac{y-2}{m} = \frac{z-3}{n}$, where $12l+4m+5n=0$
- (c) $\frac{x-1}{12} = \frac{y-2}{4} = \frac{z-3}{5}$ (d) None of these
85. Let G be the centroid of the triangle formed by the points (1, 2, 0), (2, 1, 1), (0, 0, 2). Then equation of the line OG is given by
- (a) $x=y=z$ (b) $\frac{x-1}{1} = \frac{y}{1} = \frac{z}{1}$ (c) $\frac{x-1}{1} = \frac{y-1}{1} = \frac{z-1}{0}$ (d) None of these
86. The direction cosines of the line $\frac{3x+1}{-3} = \frac{3y+2}{6} = \frac{z}{-1}$ are
- (a) $\left(\frac{1}{3}, \frac{2}{3}, 0\right)$ (b) $\left(-1, \frac{2}{3}, 1\right)$ (c) $\left(-\frac{1}{2}, 1, -\frac{1}{2}\right)$ (d) $\left(-\frac{1}{\sqrt{6}}, \frac{2}{\sqrt{6}}, -\frac{1}{\sqrt{6}}\right)$
87. The direction cosines of the line $x=y=z$ are [MP PET 1989]
- (a) $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$ (b) $\frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ (c) 1, 1, 1 (d) None of these
88. The direction ratio's of the line $x-y+z-5=0=x-3y-6$ are [MP PET 1999]
- (a) 3, 1, -2 (b) 2, -4, 1 (c) $\frac{3}{\sqrt{14}}, \frac{1}{\sqrt{14}}, \frac{-2}{\sqrt{14}}$ (d) $\frac{2}{\sqrt{41}}, \frac{-4}{\sqrt{41}}, \frac{1}{\sqrt{41}}$
89. The angle between two lines $\frac{x+1}{2} = \frac{y+3}{2} = \frac{z-4}{-1}$ and $\frac{x-4}{1} = \frac{y+4}{2} = \frac{z+1}{2}$ is [MP PET 1996]
- (a) $\cos^{-1}\left(\frac{1}{9}\right)$ (b) $\cos^{-1}\left(\frac{2}{9}\right)$ (c) $\cos^{-1}\left(\frac{3}{9}\right)$ (d) $\cos^{-1}\left(\frac{4}{9}\right)$
90. The angle between the lines $\frac{x+4}{1} = \frac{y-3}{2} = \frac{z+2}{3}$ and $\frac{x}{3} = \frac{y-1}{-2} = \frac{z}{1}$ is
- (a) $\sin^{-1}\left(\frac{1}{7}\right)$ (b) $\cos^{-1}\left(\frac{2}{7}\right)$ (c) $\cos^{-1}\left(\frac{1}{7}\right)$ (d) None of these
91. The angle between the lines $\frac{x}{1} = \frac{y}{0} = \frac{z}{-1}$ and $\frac{x}{3} = \frac{y}{4} = \frac{z}{5}$ is
- (a) $\cos^{-1} \frac{1}{5}$ (b) $\cos^{-1} \frac{1}{3}$ (c) $\cos^{-1} \frac{1}{2}$ (d) $\cos^{-1} \frac{1}{4}$
92. The value of λ for which the lines $\frac{x-1}{1} = \frac{y-2}{\lambda} = \frac{z+1}{-1}$ and $\frac{x+1}{-\lambda} = \frac{y+1}{2} = \frac{z-2}{1}$ are perpendicular to each other is
- (a) 0 (b) 1 (c) -1 (d) None of these
93. The angle between the straight lines $\frac{x+1}{2} = \frac{y-2}{5} = \frac{z+3}{4}$ and $\frac{x-1}{1} = \frac{y+2}{2} = \frac{z-3}{-3}$ is [MP PET 2000]
- (a) 45° (b) 30° (c) 60° (d) 90°
94. The angle between the lines $2x=3y=-z$ and $6x=-y=-4z$, is [MP PET 1994,99]
- (a) 0° (b) 30° (c) 45° (d) 90°

362 Three Dimensional Co-ordinate

95. The angle between the lines $x = 1, y = 2$ and $y = -1$ and $z = 0$ is [Kurukshetra CEE 1993]
 (a) 90° (b) 30° (c) 60° (d) 0°
96. The straight line $\frac{x-3}{3} = \frac{y-2}{1} = \frac{z-1}{0}$ is [Rajasthan PET 2002]
 (a) Parallel to x-axis (b) Parallel to y-axis (c) Parallel to z-axis (d) Perpendicular to z-axis
97. The lines $\frac{x-1}{2} = \frac{y-1}{3} = \frac{z-3}{0}$ and $\frac{x-2}{0} = \frac{y-3}{0} = \frac{z-4}{1}$ are
 (a) Parallel (b) Skew (c) Coincident (d) Perpendicular
98. The straight lines $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{3}$ and $\frac{x-1}{2} = \frac{y-2}{2} = \frac{z-3}{-2}$ are
 (a) Parallel lines (b) Intersecting at 60° (c) Skew lines (d) Intersecting at right angle
99. The angle between the lines $\frac{x-2}{3} = \frac{y+1}{-2}, z = 2$ and $\frac{x-1}{1} = \frac{2y+3}{3} = \frac{z+5}{2}$ is
 (a) $\pi/2$ (b) $\pi/3$ (c) $\pi/6$ (d) None of these
100. The lines $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ and $\frac{x-1}{-2} = \frac{y-2}{-4} = \frac{z-3}{-6}$ are [Kurukshetra CEE 2000]
 (a) Parallel (b) Intersecting (c) Skew (d) Coincident
101. The lines $\frac{x-1}{2} = \frac{y-2}{4} = \frac{z-3}{7}$ and $\frac{x-1}{4} = \frac{y-2}{5} = \frac{z-3}{7}$ are
 (a) Parallel (b) Intersecting (c) Skew (d) Perpendicular
102. Lines $\mathbf{r} = \mathbf{a}_1 + t\mathbf{b}_1$ and $\mathbf{r} = \mathbf{a}_2 + s\mathbf{b}_2$ are parallel iff [Kurukshetra CEE 1992]
 (a) \mathbf{b}_1 is parallel to $\mathbf{a}_2 - \mathbf{a}_1$ (b) \mathbf{b}_2 is parallel to $\mathbf{a}_2 - \mathbf{a}_1$
 (c) $\mathbf{b}_1 = \lambda\mathbf{b}_2$ for some real λ (d) None of these
103. The equation of the line passing through the points $a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$ and $b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$ [Rajasthan PET 2002]
 (a) $(a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) + t(b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k})$ (b) $(a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}) - t(b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k})$
 (c) $a_1(1-t)\mathbf{i} + a_2(1-t)\mathbf{j} + a_3(1-t)\mathbf{k} + (b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k})t$ (d) None of these
104. The vector equation of the line joining the points $\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ and $-2\mathbf{j} + 3\mathbf{k}$ is [MP PET 2003]
 (a) $\mathbf{r} = t(\mathbf{i} + \mathbf{j} + \mathbf{k})$ (b) $\mathbf{r} = t_1(\mathbf{i} - 2\mathbf{j} + \mathbf{k}) + t_2(3\mathbf{k} - 2\mathbf{j})$ (c) $\mathbf{r} = (\mathbf{i} - 2\mathbf{j} + \mathbf{k}) + t(2\mathbf{k} - \mathbf{i})$ (d) $\mathbf{r} = t(2\mathbf{k} - \mathbf{i})$
105. The acute angle between the line joining the points $(2, 1, -3), (-3, 1, 7)$ and a line parallel to $\frac{x-1}{3} = \frac{y}{4} = \frac{z+3}{5}$ through the point $(-1, 0, 4)$ is [MP PET 1998]
 (a) $\cos^{-1}\left(\frac{7}{5\sqrt{10}}\right)$ (b) $\cos^{-1}\left(\frac{1}{\sqrt{10}}\right)$ (c) $\cos^{-1}\left(\frac{3}{5\sqrt{10}}\right)$ (d) $\cos^{-1}\left(\frac{1}{5\sqrt{10}}\right)$
106. The shortest distance between the lines $\frac{x-3}{3} = \frac{y-8}{-1} = \frac{z-3}{1}$ and $\frac{x+3}{-3} = \frac{y+7}{2} = \frac{z-6}{4}$ is [MP PET 2002]
 (a) $\sqrt{30}$ (b) $2\sqrt{30}$ (c) $5\sqrt{30}$ (d) $3\sqrt{30}$
107. Shortest distance between lines $\frac{x-6}{1} = \frac{y-2}{-2} = \frac{z-2}{2}$ and $\frac{x+4}{3} = \frac{y}{-2} = \frac{z+1}{-2}$ is
 (a) 108 (b) 9 (c) 27 (d) None of these
108. The lines l_1 and l_2 intersect. The shortest distance between them is

Three Dimensional Co-ordinate Geometry

- (a) Positive (b) Zero (c) Negative (d) Infinity

109. The shortest distance between two straight lines given by $\frac{x-4}{1} = \frac{y+1}{2} = \frac{z-0}{-3}$ and $\frac{x-1}{1} = \frac{y+1}{4} = \frac{z-2}{-5}$ is [Pb. CET 2001]

- (a) $\frac{2}{\sqrt{5}}$ (b) $\frac{3}{\sqrt{5}}$ (c) $\frac{6}{\sqrt{5}}$ (d) None of these

110. The shortest distance between the lines $\mathbf{r} = (3\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}) + \mathbf{i}t$ and $\mathbf{r} = \mathbf{i} - \mathbf{j} + 2\mathbf{k} + \mathbf{j}s$ (t and s being parameters) is [AMU 1999]

- (a) $\sqrt{21}$ (b) $\sqrt{102}$ (c) 4 (d) 3

Advance Level

111. The equation of the line passing through the point (1, 2, -4) and perpendicular to the two lines $\frac{x-8}{2} = \frac{y+19}{-16} = \frac{z-10}{7}$ and $\frac{x-15}{-2} = \frac{y-29}{8} = \frac{z-5}{-5}$, will be [AI CBSE 1983]

- (a) $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z+4}{6}$ (b) $\frac{x-1}{-2} = \frac{y-2}{3} = \frac{z+4}{8}$ (c) $\frac{x-1}{3} = \frac{y-2}{2} = \frac{z+4}{8}$ (d) None of these

112. The equation of straight line $3x + 2y - z - 4 = 0$; $4x + y - 2z + 3 = 0$ in the symmetrical form is

- (a) $\frac{x-2}{3} = \frac{y-5}{2} = \frac{z}{5}$ (b) $\frac{x+2}{3} = \frac{y-5}{-2} = \frac{z}{5}$ (c) $\frac{x+2}{3} = \frac{y-5}{2} = \frac{z}{5}$ (d) None of these

113. The point of intersection of lines $\frac{x-4}{5} = \frac{y-1}{2} = \frac{z}{1}$ and $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ is [AISSE 1986]

- (a) (-1, -1, -1) (b) (-1, -1, 1) (c) (1, -1, -1) (d) (-1, 1, -1)

114. The length and foot of the perpendicular from the point (2, -1, 5) to the line $\frac{x-11}{10} = \frac{y+2}{-4} = \frac{z+8}{-11}$ are [DSSE 1987]

- (a) $\sqrt{14}$, (1, 2, -3) (b) $\sqrt{14}$, (1, -2, 3) (c) $\sqrt{14}$, (1, 2, 3) (d) None of these

115. The perpendicular distance of the point $(2, 4, -1)$ from the line $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}$ is [Kurukshetra CEE 1996]
 (a) 3 (b) 5 (c) 7 (d) 9
116. Distance of the point (x_1, y_1, z_1) from the line $\frac{x-x_2}{l} = \frac{y-y_2}{m} = \frac{z-z_2}{n}$, where l, m and n are the direction cosines of line is
 (a) $\sqrt{(x_1-x_2)^2 + (y_1-y_2)^2 + (z_1-z_2)^2 - [l(x_1-x_2) + m(y_1-y_2) + n(z_1-z_2)]^2}$
 (b) $\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2 + (z_2-z_1)^2}$
 (c) $\sqrt{(x_2-x_1)l + (y_2-y_1)m + (z_2-z_1)n}$
 (d) None of these
117. The length of the perpendicular from point $(1, 2, 3)$ to the line $\frac{x-6}{3} = \frac{y-7}{2} = \frac{z-7}{-2}$ is [MP PET 1997]
 (a) 5 (b) 6 (c) 7 (d) 8
118. The foot of the perpendicular from $(0, 2, 3)$ to the line $\frac{x+3}{5} = \frac{y-1}{2} = \frac{z+4}{3}$ is
 (a) $(-2, 3, 4)$ (b) $(2, -1, 3)$ (c) $(2, 3, -1)$ (d) $(3, 2, -1)$
119. The foot of the perpendicular from $(1, 2, 3)$ to the line joining the points $(6, 7, 7)$ and $(9, 9, 5)$ is
 (a) $(5, 3, 9)$ (b) $(3, 5, 9)$ (c) $(3, 9, 5)$ (d) $(3, 9, 9)$
120. If the equation of a line through a point \mathbf{a} and parallel to vector \mathbf{b} is $\mathbf{r} = \mathbf{a} + t\mathbf{b}$, where t is a parameter, then its perpendicular distance from the point \mathbf{c} is [MP PET 1998]
 (a) $|(\mathbf{c}-\mathbf{b}) \times \mathbf{a}| \div |\mathbf{a}|$ (b) $|(\mathbf{c}-\mathbf{a}) \times \mathbf{b}| \div |\mathbf{b}|$ (c) $|(\mathbf{a}-\mathbf{b}) \times \mathbf{c}| \div |\mathbf{c}|$ (d) $|(\mathbf{a}-\mathbf{b}) \times \mathbf{c}| \div |\mathbf{a} + \mathbf{c}|$
121. The distance of the point $B(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ from the line which is passing through $A(4\mathbf{i} + 2\mathbf{j} + 2\mathbf{k})$ and which is parallel to the vector $\vec{C} = 2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$ is [Roorkee 1993]
 (a) 10 (b) $\sqrt{10}$ (c) 100 (d) None of these

Plane

Basic Level

122. The ratio in which the line joining the points (a, b, c) and $(-a, -c, -b)$ is divided by the xy -plane is [MP PET 1994; Him.
 (a) $a:b$ (b) $b:c$ (c) $c:a$ (d) $c:b$
123. The ratio in which the line joining $(2, 4, 5)$ and $(3, 5, -4)$ is divided by the yz -plane is [MP PET 2002; Rajasthan PET 2002]
 (a) $2:3$ (b) $3:2$ (c) $-2:3$ (d) $4:-3$
124. xy -plane divides the line joining the points $(2, 4, 5)$ and $(-4, 3, -2)$ in the ratio [MP PET 1988]
 (a) $3:5$ (b) $5:2$ (c) $1:3$ (d) $3:4$
125. The coordinates of the point where the line through $P(3, 4, 1)$ and $Q(5, 1, 6)$ crosses the xy -plane are [MP PET 1997]
 (a) $\frac{3}{5}, \frac{13}{5}, \frac{23}{5}$ (b) $\frac{13}{5}, \frac{23}{5}, \frac{3}{5}$ (c) $\frac{13}{5}, \frac{23}{5}, 0$ (d) $\frac{13}{5}, 0, 0$
126. The plane XOZ divides the join of $(1, -1, 5)$ and $(2, 3, 4)$ in the ratio $\lambda:1$, then λ is [Pb. CET 1988]

364 Three Dimensional Co-ordinate

- (a) -3 (b) 3 (c) $-\frac{1}{3}$ (d) $\frac{1}{3}$
127. XOZ plane divides the join of (2, 3, 1) and (6, 7, 1) in the ratio [EAMCET 2003]
 (a) 3:7 (b) 2:7 (c) -3:7 (d) -2:7
128. The plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 3$ meets the coordinate axes in A, B, C. The centroid of the triangle ABC is
 (a) $\left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3}\right)$ (b) $\left(\frac{3}{a}, \frac{3}{b}, \frac{3}{c}\right)$ (c) $\left(\frac{1}{a}, \frac{1}{b}, \frac{1}{c}\right)$ (d) (a, b, c)
129. The ratio in which the plane $\mathbf{r} \cdot (\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}) = 17$ divides the line joining the points $-2\mathbf{i} + 4\mathbf{j} + 7\mathbf{k}$ and $3\mathbf{i} - 5\mathbf{j} + 8\mathbf{k}$ is [Kurukshetra CEE 1996; DCE 1999]
 (a) 1:5 (b) 1:10 (c) 3:5 (d) 3:10
130. If a plane cuts off intercepts $OA = a$, $OB = b$, $OC = c$ from the coordinate axes, then the area of the triangle ABC =
 (a) $\frac{1}{2} \sqrt{b^2c^2 + c^2a^2 + a^2b^2}$ (b) $\frac{1}{2}(bc + ca + ab)$
 (c) $\frac{1}{2}abc$ (d) $\frac{1}{2} \sqrt{(b-c)^2 + (c-a)^2 + (a-b)^2}$
131. The plane $\frac{x}{2} + \frac{y}{3} + \frac{z}{4} = 1$ cuts the axes in A, B, C, then the area of the $\triangle ABC$ is [MP PET 2000]
 (a) $\sqrt{29}$ (b) $\sqrt{41}$ (c) $\sqrt{61}$ (d) None of these
132. The volume of the tetrahedron included between the plane $2x - 3y + 4z - 12 = 0$ and the three coordinate planes is
 (a) $3\sqrt{(29)}$ (b) $6\sqrt{(29)}$ (c) 12 (d) None of these
133. A point located in space moves in such a way that sum of its distances from xy - and yz plane is equal to distance from zx plane, the locus of the point is
 (a) $x - y + z = 2$ (b) $x + y - z = 0$ (c) $x + y - z = 2$ (d) $x - y + z = 0$
134. The equation of a plane parallel to x - axis is [DCE 2001]
 (a) $ax + by + cz + d = 0$ (b) $ax + by + d = 0$ (c) $by + cz + d = 0$ (d) $ax + cz + d = 0$
135. In the space the equation $by + cz + d = 0$ represents a plane perpendicular to the plane [EAMCET 2002]
 (a) YOZ (b) Z=k (c) ZOX (d) XOY
136. The intercepts of the plane $5x - 3y + 6z = 60$ on the coordinate axes are [MP PET 2001]
 (a) (10, 20, -10) (b) (10, -20, 12) (c) (12, -20, 10) (d) (12, 20, -10)
137. The coordinates of the points A and B are (2, 3, 4) and (-2, 5, -4) respectively. If a point P moves, so that $PA^2 - PB^2 = k$ where k is constant, then the locus of P is
 (a) A line (b) A plane (c) A sphere (d) None of these
138. In a three dimensional xyz space the equation $x^2 - 5x + 6 = 0$ represents [Orissa JEE 2002]
 (a) Points (b) Plane (c) Curves (d) Pair of straight line
139. The equation of yz -plane is [MP PET 1988]
 (a) $x = 0$ (b) $y = 0$ (c) $z = 0$ (d) $x + y + z = 0$
140. The intercepts of the plane $2x - 3y + 4z = 12$ on the coordinate axes are given by

Three Dimensional Co-ordinate Geometry

- (a) 2, -3, 4 (b) 6, -4, -3 (c) 6, -4, 3 (d) 3, -2, 1.5
- 141.** The locus of the point (x, y, z) for which $z = k$, is
 (a) A plane parallel to xy plane at a distance k from it (b) A plane parallel to yz plane at a distance k from it
 (c) A plane parallel to zx plane at a distance k from it (d) A line parallel to z -axis at a distance k from it
- 142.** A point (x, y, z) moves parallel to x - axis. Which of the three variables x, y, z remains fixed
 (a) x (b) x and y (c) y and z (d) z and x
- 143.** If $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are three non-coplanar vectors, then the vector equation $\mathbf{r} = (1-p-q)\mathbf{a} + p\mathbf{b} + q\mathbf{c}$ represents a [EAMCET 2003]
 (a) Straight line (b) Plane
 (c) Plane passing through the origin (d) Sphere
- 144.** The direction cosines of the normal to the plane $3x + 4y + 12z = 52$ will be [MP PET 1997]
 (a) 3, 4, 12 (b) -3, -4, -12 (c) $\frac{3}{13}, \frac{4}{13}, \frac{12}{13}$ (d) $\frac{3}{\sqrt{13}}, \frac{4}{\sqrt{13}}, \frac{12}{\sqrt{13}}$
- 145.** The direction cosines of the normal to the plane $x + 2y - 3z + 4 = 0$ are [MP PET 1996]
 (a) $\frac{1}{\sqrt{14}}, -\frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$ (b) $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$ (c) $-\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$ (d) $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, -\frac{3}{\sqrt{14}}$
- 146.** Normal form of the plane $2x + 6y + 3z = 1$ is
 (a) $\frac{2}{7}x + \frac{6}{7}y + \frac{3}{7}z = 1$ (b) $\frac{2}{7}x + \frac{6}{7}y + \frac{3}{7}z = \frac{1}{7}$ (c) $\frac{2}{7}x + \frac{6}{7}y + \frac{3}{7}z = 0$ (d) None of these
- 147.** The equation of a plane which cuts equal intercepts of unit length on the axes, is [MP PET 1996]
 (a) $x + y + z = 0$ (b) $x + y + z = 1$ (c) $x + y - z = 1$ (d) $\frac{x}{a} + \frac{y}{a} + \frac{z}{a} = 1$
- 148.** The equation of the plane which is parallel to y - axis and cuts off intercepts of length 2 and 3 from x -axis and z -axis is
 (a) $3x + 2z = 1$ (b) $3x + 2z = 6$ (c) $2x + 3z = 6$ (d) $3x + 2z = 0$
- 149.** A planes π makes intercepts 3 and 4 respectively on z -axis and x -axis. If π is parallel to y - axis, then its equation is [EAMCET 2003]
 (a) $3x + 4z = 12$ (b) $3z + 4x = 12$ (c) $3y + 4z = 12$ (d) $3z + 4y = 12$
- 150.** The equation of the plane through the three points $(1, 1, 1)$, $(1, -1, 1)$, and $(-7, -3, -5)$, is [AISSE 1984]
 (a) $3x - 4z + 1 = 0$ (b) $3x - 4y + 1 = 0$ (c) $3x + 4y + 1 = 0$ (d) None of these
- 151.** The equation of the plane through $(1, 2, 3)$ and parallel to the plane $2x + 3y - 4z = 0$ is [MP PET 1990]
 (a) $2x + 3y + 4z = 4$ (b) $2x + 3y + 4z + 4 = 0$ (c) $2x - 3y + 4z + 4 = 0$ (d) $2x + 3y - 4z + 4 = 0$
- 152.** The equation of the plane through $(2, 3, 4)$ and parallel to the plane $x + 2y + 4z = 5$ is [Kurukshetra CEE 1999; MP PET 1999]
 (a) $x + 2y + 4z = 10$ (b) $x + 2y + 4z = 3$ (c) $x + y + 2z = 2$ (d) $x + 2y + 4z = 24$
- 153.** The equation of the plane passing through the points $(1, -3, -2)$ and perpendicular to planes $x + 2y + 2z = 5$ and $3x + 3y + 2z = 8$, is [AISSE 1987]
 (a) $2x - 4y + 3z - 8 = 0$ (b) $2x - 4y - 3z + 8 = 0$ (c) $2x + 4y + 3z + 8 = 0$ (d) None of theses
- 154.** The line drawn from $(4, -1, 2)$ to the point $(-3, 2, 3)$ meets a plane at right angles at the point $(-10, 5, 4)$, then the equation of plane is

366 Three Dimensional Co-ordinate

[DSSE 1985]

- (a) $7x - 3y - z + 89 = 0$ (b) $7x + 3y + z + 89 = 0$ (c) $7x - 3y + z + 89 = 0$ (d) None of these

155. $x + y + z + 2 = 0$ together with $x + y + z + 3 = 0$ represents in space [MP PET 1989]

- (a) A line (b) A point (c) A plane (d) None of these

156. The equation of the plane which contains the line of intersection of the planes $x + 2y + 3z - 4 = 0$ and $2x + y - z + 5 = 0$ and which is perpendicular to the plane $5x + 3y - 6z + 8 = 0$, is [DSSE 1987]

- (a) $33x + 50y + 45z - 41 = 0$ (b) $33x + 45y + 50z + 41 = 0$ (c) $45x + 45y + 50z - 41 = 0$ (d) $33x + 45y + 50z - 41 = 0$

157. The equation of the planes passing through the line of intersection of the planes $3x - y - 4z = 0$ and $x + 3y + 6 = 0$, whose distance from the origin is 1, are

- (a) $x - 2y - 2z - 3 = 0, 2x + y - 2z + 3 = 0$ (b) $x - 2y + 2z - 3 = 0, 2x + y + 2z + 3 = 0$
(c) $x + 2y - 2z - 3 = 0, 2x - y - 2z + 3 = 0$ (d) None of these

158. The equation of the plane which passes through the point (2, 1, 4) and parallel to the plane $2x + 3y + 5z + 6 = 0$ is

- (a) $2x + 3y + 5z + 27 = 0$ (b) $2x + 3y + 5z - 27 = 0$ (c) $2x + y + 4z - 27 = 0$ (d) $2x + y + 4z + 27 = 0$

159. The equation of a plane which passes through (2, -3, 1) and is normal to the line joining the points (3, 4, -1) and (2, -1, 5) is given by

- (a) $x + 5y - 6z + 19 = 0$ (b) $x - 5y + 6z - 19 = 0$ (c) $x + 5y + 6z + 19 = 0$ (d) $x - 5y - 6z - 19 = 0$

160. The coordinates of the point in which the line joining the points (3, 5, -7) and (-2, 1, 8) is intersected by the plane yz are given by

[MP PET 1993]

- (a) $\left(0, \frac{13}{5}, 2\right)$ (b) $\left(0, -\frac{13}{5}, -2\right)$ (c) $\left(0, -\frac{13}{5}, \frac{2}{5}\right)$ (d) $\left(0, \frac{13}{5}, \frac{2}{5}\right)$

161. If P be the point (2, 6, 3), then the equation of the plane through P at right angle to OP , O being the origin, is [MP PET

- (a) $2x + 6y + 3z = 7$ (b) $2x - 6y + 3z = 7$ (c) $2x + 6y - 3z = 49$ (d) $2x + 6y + 3z = 49$

162. The equation of the plane containing the line of intersection of the planes $2x - y = 0$ and $y - 3z = 0$ the perpendicular to the plane $4x + 5y - 3z - 8 = 0$ is

- (a) $28x - 17y + 9z = 0$ (b) $28x + 17y + 9z = 0$ (c) $28x - 17y - 9z = 0$ (d) $7x - 3y + z = 0$

163. The equation of the plane passing through (1, 1, 1) and (1, -1, -1) and perpendicular to $2x - y + z + 5 = 0$ is [EAMCET 2003]

- (a) $2x + 5y + z - 8 = 0$ (b) $x + y - z - 1 = 0$ (c) $2x + 5y + z + 4 = 0$ (d) $x - y + z - 1 = 0$

164. The equation of the plane through the intersection of the planes $x + y + z = 1$ and $2x + 3y - z + 4 = 0$ and parallel to x -axis is

[Orissa JEE 2003]

- (a) $y - 3z + 6 = 0$ (b) $3y - z + 6 = 0$ (c) $y + 3z + 6 = 0$ (d) $3y - 2z + 6 = 0$

165. If O is the origin and A is the point (a, b, c), then the equation of the plane through A and at right angles to OA is

- (a) $a(x-a) - b(y-b) - c(z-c) = 0$ (b) $a(x+a) + b(y+b) + c(z+c) = 0$
(c) $a(x-a) + b(y-b) + c(z-c) = 0$ (d) None of these

166. The equation of the plane through the point (1, 2, 3) and parallel to the plane $x + 2y + 5z = 0$ is [DCE 2002]

- (a) $(x-1) + 2(y-2) + 5(z-3) = 0$ (b) $x + 2y + 5z = 14$
(c) $x + 2y + 5z = 6$ (d) None of these

Three Dimensional Co-ordinate Geometry

- 167.** The equation of the plane passing through the intersection of the planes $x + y + z = 6$ and $2x + 3y + 4z + 5 = 0$ and the point $(1, 1, 1)$, is
- (a) $20x + 23y + 26z - 69 = 0$ (b) $20x + 23y + 26z + 69 = 0$
 (c) $23x + 20y + 26z + 69 = 0$ (d) None of these
- 168.** The equation of the plane passing through the intersection of the planes $x + 2y + 3z + 4 = 0$ and $4x + 3y + 2z + 1 = 0$ and the origin is
- [Kerala (Engg.) 2002]
- (a) $3x + 2y + z + 1 = 0$ (b) $3x + 2y + z = 0$ (c) $2x + 3y + z = 0$ (d) $x + y + z = 0$
- 169.** If the plane $x - 2y + 3z = 0$ is rotated through a right angle about its line of intersection with the plane $2x + 3y - 4z - 5 = 0$, then the equation of plane in its new position is
- (a) $28x - 17y + 9z = 0$ (b) $22x + 5y - 4z - 35 = 0$ (c) $25x + 17y - 52z - 25 = 0$ (d) $x + 35y - 10z - 70 = 0$
- 170.** The equation of the plane passing through the point $(-2, -2, 2)$ and containing the line joining the points $(1, 1, 1)$ and $(1, -1, 2)$ is
- (a) $x + 2y - 3z + 4 = 0$ (b) $3x - 4y + 1 = 0$ (c) $5x + 2y - 3z - 17 = 0$ (d) $x - 3y - 6z + 8 = 0$
- 171.** The equation of the plane containing the line $2x + z - 4 = 0, 2y + z = 0$ and passing through the point $(2, 1, -1)$ is [AMU 1990]
- (a) $x + y + z + 2 = 0$ (b) $x + y - z - 4 = 0$ (c) $x - y - z - 2 = 0$ (d) $x + y + z - 2 = 0$
- 172.** In three dimensional space, the equation $3y + 4z = 0$ represents
- [Kurukshetra CEE 1994]
- (a) A plane containing x -axis (b) A plane containing y -axis
 (c) A plane containing z -axis (d) A line with direction numbers 0, 3, 4
- 173.** Direction ratios of the normal to the plane passing through the point $(2, 1, 3)$ and the point of intersection of the planes $x + 2y + z = 3$ and $2x - y - z = 5$ are
- (a) 13, 6, 1 (b) 5, 7, 3 (c) 4, 3, 2 (d) None of these
- 174.** The plane of intersection of $x^2 + y^2 + z^2 + 2x + 2y + 2z + 2 = 0$ and $4x^2 + 4y^2 + 4z^2 + 4x + 4y + 4z - 1 = 0$ is [Pb. CET 1996]
- (a) $4x + 4y + 4z + 9 = 0$ (b) $x + y + z + 9 = 0$ (c) $4x + 4y + 4z + 1 = 0$ (d) They do not intersect
- 175.** If the planes $x + 2y + kz = 0$ and $2x + y - 2z = 0$ are at right angles, then the value of k is [MP PET 1999]
- (a) $-\frac{1}{2}$ (b) $\frac{1}{2}$ (c) -2 (d) 2
- 176.** The value of k for which the planes $3x - 6y - 2z = 7$ and $2x + y - kz = 5$ are perpendicular to each other, is [MP PET 1992]
- (a) 0 (b) 1 (c) 2 (d) 3
- 177.** If the given planes $ax + by + cz + d = 0$ and $a'x + b'y + c'z + d' = 0$ be mutually perpendicular, then [MP PET 1994]
- (a) $\frac{a}{a'} = \frac{b}{b'} = \frac{c}{c'}$ (b) $\frac{a}{a'} + \frac{b}{b'} + \frac{c}{c'} = 0$ (c) $aa' + bb' + cc' + dd' = 0$ (d) $aa' + bb' + cc' = 0$
- 178.** The angle between two planes is equal to
- (a) The angle between the tangents to them from any point
 (b) The angle between the normals to them from any point
 (c) The angle between the lines parallel to the planes from any point
 (d) None of these
- 179.** If the planes $3x - 2y + 2z + 17 = 0$ and $4x + 3y - kz = 25$ are mutually perpendicular, then $k =$ [MP PET 1995]
- (a) 3 (b) -3 (c) 9 (d) -6

368 Three Dimensional Co-ordinate

- 180.** The angle between the planes $2x - y + z = 6$ and $x + y + 2z = 7$ is [MP PET 1991,98,2000,01,03; Rajasthan PET 2001]
(a) 30° (b) 45° (c) 0° (d) 60°
- 181.** The angle between the planes $3x - 4y + 5z = 0$ and $2x - y - 2z = 5$ is [MP PET 1988; Kurukshetra CEE 2000]
(a) $\frac{\pi}{3}$ (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{6}$ (d) None of these
- 182.** If θ is the angle between the planes $2x - y + 2z = 3$, $6x - 2y + 3z = 5$, then $\cos \theta$ is equal to [Kerala (Engg.) 2001]
(a) $\frac{21}{20}$ (b) $\frac{11}{20}$ (c) $\frac{20}{21}$ (d) $\frac{12}{25}$
- 183.** The value of $aa' + bb' + cc'$ being negative, the origin will lie in the acute angle between the planes $ax + by + cz + d = 0$ and $a'x + b'y + c'z + d' = 0$, if [MP PET 2003]
(a) $a = a' = 0$ (b) d and d' are of same sign (c) d and d' are of opposite sign (d) $d = d' = 0$
- 184.** The equation of the plane which bisects the angle between the planes $3x - 6y + 2z + 5 = 0$ and $4x - 12y + 3z - 3 = 0$ which contains the origin is
(a) $33x - 13y + 32z + 45 = 0$ (b) $x - 3y + z - 5 = 0$ (c) $33x + 13y + 32z + 45 = 0$ (d) None of these
- 185.** The equation of the bisector of the obtuse angle between the planes $3x + 4y - 5z + 1 = 0$, $5x + 12y - 13z = 0$ is
(a) $11x + 4y - 3z = 0$ (b) $14x - 8y + 13 = 0$ (c) $x + y + z = 9$ (d) $13x - 7z + 18 = 0$
- 186.** The two points $(1, 1, 1)$ and $(-3, 0, 1)$ with respect to the plane $3x + 4y - 12z + 13 = 0$ lie on
(a) Opposite side (b) Same side (c) On the plane (d) None of these
- 187.** Distance between parallel planes $2x - 2y + z + 3 = 0$ and $4x - 4y + 2z + 5 = 0$ is [MP PET 1994, 95]
(a) $\frac{2}{3}$ (b) $\frac{1}{3}$ (c) $\frac{1}{6}$ (d) 2
- 188.** The distance between the planes $x + 2y + 3z + 7 = 0$ and $2x + 4y + 6z + 7 = 0$ is [MP PET 1991]
(a) $\frac{\sqrt{7}}{2\sqrt{2}}$ (b) $\frac{7}{2}$ (c) $\frac{\sqrt{7}}{2}$ (d) $\frac{7}{2\sqrt{2}}$
- 189.** Distance of the point $(2, 3, 4)$ from the plane $3x - 6y + 2z + 11 = 0$ is [MP PET 1990,96]
(a) 1 (b) 2 (c) 3 (d) 0
- 190.** The distance of the plane $6x - 3y + 2z - 14 = 0$ from the origin is [MP PET 2003]
(a) 2 (b) 1 (c) 14 (d) 8
- 191.** The distance of the point $(2, 3, -5)$ from the plane $x + 2y - 2z = 9$ is [MP PET 2001]
(a) 4 (b) 3 (c) 2 (d) 1
- 192.** If the points $(1, 1, k)$ and $(-3, 0, 1)$ be equidistant from the plane $3x + 4y - 12z + 13 = 0$, then $k =$
(a) 0 (b) 1 (c) 2 (d) None of these
- 193.** If the product of distances of the point $(1, 1, 1)$ from the origin and the plane $x - y + z + k = 0$ be 5, then $k =$
(a) -2 (b) -3 (c) 4 (d) 7
- 194.** If two planes intersect, then the shortest distance between the planes is [Kurukshetra CEE 1998]
(a) $\cos 0^\circ$ (b) $\cos 90^\circ$ (c) $\sin 90^\circ$ (d) 1
- 195.** The length of the perpendicular from the origin to the plane $3x + 4y + 12z = 52$ is [MP PET 2000]
(a) 3 (b) -4 (c) 5 (d) None of these

- 196.** If the length of perpendicular drawn from origin on a plane is 7 units and its direction ratios are $-3, 2, 6$, then that plane is [MP PET 1998]
- (a) $-3x + 2y + 6z - 7 = 0$ (b) $-3x + 2y + 6z - 49 = 0$ (c) $3x - 2y + 6z + 7 = 0$ (d) $-3x + 2y - 6z - 49 = 0$
- 197.** If a plane cuts off intercepts $-6, 3, 4$ from the coordinate axes, then the length of the perpendicular from origin to the plane is
- (a) $\frac{1}{\sqrt{61}}$ (b) $\frac{13}{\sqrt{61}}$ (c) $\frac{12}{\sqrt{29}}$ (d) $\frac{5}{\sqrt{41}}$
- 198.** If $A(-1, 2, 3)$, $B(1, 1, 1)$ and $C(2, -1, 3)$ are points on a plane. A unit normal vector to the plane ABC is [BIT Ranchi 1988]
- (a) $\pm\left(\frac{2\mathbf{i} + 2\mathbf{j} + \mathbf{k}}{3}\right)$ (b) $\pm\left(\frac{2\mathbf{i} - 2\mathbf{j} + \mathbf{k}}{3}\right)$ (c) $\pm\left(\frac{2\mathbf{i} - 2\mathbf{j} - \mathbf{k}}{3}\right)$ (d) $-\left(\frac{2\mathbf{i} + 2\mathbf{j} + \mathbf{k}}{3}\right)$
- 199.** If the position vectors of three points A, B and C are respectively $\mathbf{i} + \mathbf{j} + \mathbf{k}$, $2\mathbf{i} + 3\mathbf{j} - 4\mathbf{k}$ and $7\mathbf{i} + 4\mathbf{j} + 9\mathbf{k}$, then the unit vector to the plane containing the triangle ABC is [DCE 1999]
- (a) $31\mathbf{i} - 18\mathbf{j} - 9\mathbf{k}$ (b) $\frac{31\mathbf{i} - 38\mathbf{j} - 9\mathbf{k}}{\sqrt{2486}}$ (c) $\frac{31\mathbf{i} + 18\mathbf{j} + 9\mathbf{k}}{\sqrt{2486}}$ (d) None of these
- 200.** The projection of point (a, b, c) in yz plane are
- (a) $(0, b, c)$ (b) $(a, 0, c)$ (c) $(a, b, 0)$ (d) $(a, 0, 0)$

Advance Level

- 201.** A variable plane at a constant distance p from origin meets the coordinate axes in A, B, C . Through these points planes are drawn parallel to coordinate planes. Then locus of the point of intersection is
- (a) $\frac{1}{x^2} + \frac{1}{y^2} + \frac{1}{z^2} = \frac{1}{p^2}$ (b) $x^2 + y^2 + z^2 = p^2$ (c) $x + y + z = p$ (d) $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = p$
- 202.** A variable plane is at a constant distance p from the origin and meets the axes in A, B and C , then the locus of the centroid of the triangle ABC is
- (a) $x^{-2} + y^{-2} + z^{-2} = p^{-2}$ (b) $x^{-2} + y^{-2} + z^{-2} = 9p^{-2}$ (c) $x^{-2} + y^{-2} + z^{-2} = p^2$ (d) None of these
- 203.** The equation of the plane which bisects line joining $(2, 3, 4)$ and $(6, 7, 8)$ is [CET 1991, 93]
- (a) $x + y + z - 15 = 0$ (b) $x - y + z - 15 = 0$ (c) $x - y - z - 15 = 0$ (d) $x + y + z + 15 = 0$
- 204.** The equation of the plane which bisects the line joining the points $(-1, 2, 3)$ and $(3, -5, 6)$ at right angle, is
- (a) $4x - 7y - 3z = 8$ (b) $4x - 7y - 3z = 28$ (c) $4x - 7y + 3z = 28$ (d) $4x + 2y - 3z = 28$
- 205.** P is a fixed point (a, a, a) on a line through the origin equally inclined to the axes, then any plane through P perpendicular to OP , makes intercepts on the axes, the sum of whose reciprocals is equal to
- (a) a (b) $\frac{3}{2a}$ (c) $\frac{3a}{2}$ (d) None of these
- 206.** If from a point $P(a, b, c)$ perpendiculars PA and PB are drawn to yz and zx planes, then the equation of the plane OAB is
- (a) $bcx + cay + abz = 0$ (b) $bcx + cay - abz = 0$ (c) $bcx - cay + abz = 0$ (d) $-bcx + cay + abz = 0$
- 207.** If l_1, m_1, n_1 and l_2, m_2, n_2 are the direction ratios of two intersecting lines, then the direction ratios of lines through them and coplanar with them are given by
- (a) $l_1 + km_1, l_2 + km_2, l_3 + km_3$ (b) $kl_1l_2, km_1m_2, kn_1n_2$
- (c) $l_1 + kl_2, m_1 + km_2, n_1 + kn_2$ (d) $\frac{kl_1}{l_2}, \frac{km_1}{m_2}, \frac{kn_1}{n_2}, k$ being a number whatsoever
- 208.** The four points $(0, 4, 3)$, $(-1, -5, -3)$, $(-2, -2, 1)$ and $(1, 1, -1)$ lie in the plane
- (a) $4x + 3y + 2z - 9 = 0$ (b) $9x - 5y + 6z + 2 = 0$ (c) $3x + 4y + 7z - 5 = 0$ (d) None of these

370 Three Dimensional Co-ordinate

209. A plane meets the coordinate axes at A, B, C such that the centre of the triangle is $(3, 3, 3)$. The equation of the plane is

- (a) $x + y + z = 3$ (b) $x + y + z = 9$ (c) $3x + 3y + 3z = 1$ (d) $9x + 9y + 9z = 1$

210. Two system of rectangular axes have the same origin. If a plane cuts them at distance a, b, c and a', b', c' from the origin, then

[AIEEE 2003]

- (a) $\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + \frac{1}{a'^2} + \frac{1}{b'^2} + \frac{1}{c'^2} = 0$ (b) $\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + \frac{1}{a'^2} + \frac{1}{b'^2} - \frac{1}{c'^2} = 0$
 (c) $\frac{1}{a^2} - \frac{1}{b^2} - \frac{1}{c^2} + \frac{1}{a'^2} - \frac{1}{b'^2} - \frac{1}{c'^2} = 0$ (d) $\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} - \frac{1}{a'^2} - \frac{1}{b'^2} - \frac{1}{c'^2} = 0$

211. Which one of the following is the best condition for the plane $ax + by + cz + d = 0$ to intersect the x and y axes at equal angle

- (a) $|a| = |b|$ (b) $a = -b$ (c) $a = b$ (d) $a^2 + b^2 = 1$

212. If the equation $2x^2 - 2y^2 + 4z^2 + 6xz + 2yz + 3xy = 0$ represents a pair of planes, then the angle between the pair of planes is

- (a) $\cos^{-1}(4/9)$ (b) $\cos^{-1}(4/21)$ (c) $\cos^{-1}(4/17)$ (d) $\cos^{-1}(2/3)$

213. The points $A(-1, 3, 0), B(2, 2, 1)$ and $C(1, 1, 3)$ determine a plane. The distance from the plane to the point $D(5, 7, 8)$ is

[AMU 2001]

- (a) $\sqrt{66}$ (b) $\sqrt{71}$ (c) $\sqrt{73}$ (d) $\sqrt{76}$

214. The length and foot of the perpendicular from the point $(7, 14, 5)$ to the plane $2x + 4y - z = 2$, are [AISSE 1987]

- (a) $\sqrt{21}, (1, 2, 8)$ (b) $3\sqrt{21}, (3, 2, 8)$ (c) $21\sqrt{3}, (1, 2, 8)$ (d) $3\sqrt{21}, (1, 2, 8)$

215. The distance of the point $(1, 1, 1)$ from the plane passing through the points $(2, 1, 1), (1, 2, 1)$ and $(1, 1, 2)$ is [AISSE 1987]

- (a) $\frac{1}{\sqrt{3}}$ (b) 1 (c) $\sqrt{3}$ (d) None of these

216. Perpendicular is drawn from the point $(0, 3, 4)$ to the plane $2x - 2y + z = 10$. The coordinates of the foot of the perpendicular are

- (a) $(-8/3, 1/3, 16/3)$ (b) $(8/3, 1/3, 16/3)$ (c) $(8/3, -1/3, 16/3)$ (d) $(8/3, 1/3, -16/3)$

217. The equation of the plane containing the lines $\mathbf{r} - \mathbf{a} = t\mathbf{b}$ and $\mathbf{r} - \mathbf{b} = s\mathbf{a}$ is

- (a) $\mathbf{r} \cdot \mathbf{a} = \mathbf{a} \cdot \mathbf{b}$ (b) $[\mathbf{r} \mathbf{a} \mathbf{b}] = 0$ (c) $\mathbf{r} \cdot \mathbf{a} = \mathbf{r} \cdot \mathbf{b}$ (d) $\mathbf{r} \cdot \mathbf{b} = \mathbf{a} \cdot \mathbf{b}$

218. Let the points P, Q and R have position vectors $\mathbf{r}_1 = 3\mathbf{i} - 2\mathbf{j} - \mathbf{k}$; $\mathbf{r}_2 = \mathbf{i} + 3\mathbf{j} + 4\mathbf{k}$ and $\mathbf{r}_3 = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ relative to an origin O . The distance of P from the plane OQR is [Roorkee 1990]

- (a) 2 (b) 3 (c) 1 (d) 5

219. The projection of the point $(1, 3, 4)$ on the plane $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} + \mathbf{k}) + 3 = 0$ is

- (a) $(1, 3, 4)$ (b) $(-3, 5, 2)$ (c) $(-1, 4, 3)$ (d) None of these

220. If $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} + \mathbf{k}) + \frac{3}{2} = 0$ is the equation of plane and $\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}$ is a point, then a point equidistant from the plane on the opposite side is [AMU 1998]

- (a) $\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ (b) $3\mathbf{i} + \mathbf{j} + \mathbf{k}$ (c) $3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ (d) $3(\mathbf{i} + \mathbf{j} + \mathbf{k})$

221. If (p_1, q_1, r_1) be the image of (p, q, r) in the plane $ax + by + cz + d = 0$, then

- (a) $\frac{p_1 - p}{a} = \frac{q_1 - q}{b} = \frac{r_1 - r}{c}$ (b) $a(p + p_1) + b(q + q_1) + c(r + r_1) + 2d = 0$
 (c) Both (a) and (b) (d) None of these

Basic Level

222. The equation of the straight line passing through (1, 2, 3) and perpendicular to the plane $x + 2y - 5z + 9 = 0$ is [MP PET 1999]

- (a) $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{-5}$ (b) $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z+5}{3}$ (c) $\frac{x+1}{1} = \frac{y+2}{2} = \frac{z+3}{-5}$ (d) $\frac{x+1}{1} = \frac{y+2}{2} = \frac{z-5}{3}$

223. The equation of the perpendicular from the point (α, β, γ) to the plane $ax + by + cz + d = 0$ is [MP PET 2003]

- (a) $a(x-\alpha) + b(y-\beta) + c(z-\gamma) = 0$ (b) $\frac{x-\alpha}{a} = \frac{y-\beta}{b} = \frac{z-\gamma}{c}$
(c) $a(x-\alpha) + b(y-\beta) + c(z-\gamma) = abc$ (d) None of these

224. The equation of the plane passing through the points (3, 2, 2) and (1, 0, -1) and parallel to the line

$$\frac{x-1}{2} = \frac{y-1}{-2} = \frac{z-2}{3} \text{ is}$$

- (a) $4x - y - 2z + 6 = 0$ (b) $4x - y + 2z + 6 = 0$ (c) $4x - y - 2z - 6 = 0$ (d) None of these

225. The equation of the plane containing the line $\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$ and the point (0, 7, -7) is [Roorkee 1999]

- (a) $x + y + z = 1$ (b) $x + y + z = 2$ (c) $x + y + z = 0$ (d) None of these

226. The equation of plane through the line of intersection of planes $ax + by + cz + d = 0$, $a'x + b'y + c'z + d' = 0$ and parallel to the line $y = 0, z = 0$ is [Kurukshetra CEE 1998]

- (a) $(ab' - a'b)x + (bc' - b'c)y + (ad' - a'd)z = 0$ (b) $(ab' - a'b)x + (bc' - b'c)y + (ad' - a'd)z = 0$
(c) $(ab' - a'b)y + (ac' - a'c)z + (ad' - a'd)z = 0$ (d) None of these

227. The equation of the plane passing through the line $\frac{x-1}{5} = \frac{y+2}{6} = \frac{z-3}{4}$ and the point (4, 3, 7) is [MP PET 2001]

- (a) $4x + 8y + 7z = 41$ (b) $4x - 8y + 7z = 41$ (c) $4x - 8y - 7z = 41$ (d) $4x - 8y + 7z = 39$

228. The equation of the plane containing the line $2x - 5y + 2z = 6$, $2x + 3y - z = 5$ and parallel to the line $\frac{x}{1} = \frac{y}{-6} = \frac{z}{7}$ is

- (a) $6x + y - 10 = 0$ (b) $6x + y - 16 = 0$ (c) $12x + 2y - 1 = 0$ (d) $6x + y + 16 = 0$

229. The equation of the plane which is parallel to the line $\frac{x-4}{1} = \frac{y+3}{-4} = \frac{z+1}{7}$ and passes through the points (0, 0, 0) and (3, -1, 2), is [DSSE 1984]

- (a) $x + 19y + 11z = 0$ (b) $x - 19y - 11z = 0$ (c) $x - 19y + 11z = 0$ (d) None of these

230. Equation of a line passing through (1, -2, 3) and parallel to the plane $2x + 3y + z + 5 = 0$ is

- (a) $\frac{x-1}{-1} = \frac{y+2}{1} = \frac{z-3}{-1}$ (b) $\frac{x-1}{2} = \frac{y+2}{3} = \frac{z-3}{1}$ (c) $\frac{x+1}{-1} = \frac{y-2}{1} = \frac{z-3}{-1}$ (d) None of these

231. The equation of the plane through the line $3x - 4y + 5z = 10$, $2x + 2y - 3z = 4$ and parallel to the line $x = 2y = 3z$ is

- (a) $x - 20y + 27z = 14$ (b) $x + 4y + 27z = 14$ (c) $x - 20y + 3z = 14$ (d) None of these

372 Three Dimensional Co-ordinate

- 232.** The equation of the plane passing through the line $\frac{x-4}{1} = \frac{y-3}{1} = \frac{z-2}{2}$ and $\frac{x-3}{1} = \frac{y-2}{-4} = \frac{z}{5}$ is
 (a) $11x - y - 3z = 35$ (b) $11x + y - 3z = 35$ (c) $11x - y + 3z = 35$ (d) None of these
- 233.** The equation of the plane in which the lines $\frac{x-5}{4} = \frac{y-7}{4} = \frac{z+3}{-5}$ and $\frac{x-8}{7} = \frac{y-4}{1} = \frac{z-5}{3}$ lie, is [MP PET 2000]
 (a) $17x - 47y - 24z + 172 = 0$ (b) $17x + 47y - 24z + 172 = 0$
 (c) $17x + 47y + 24z + 172 = 0$ (d) $17x - 47y + 24z + 172 = 0$
- 234.** The equation of the line passing through (1, 2, 3) and parallel to the planes $x - y + 2z = 5$ and $3x + y + z = 6$, is [DSSE 1986]
 (a) $\frac{x-1}{-3} = \frac{y-2}{5} = \frac{z-3}{4}$ (b) $\frac{x-1}{-3} = \frac{y-2}{-5} = \frac{z-1}{4}$ (c) $\frac{x-1}{-3} = \frac{y-2}{-5} = \frac{z-1}{-4}$ (d) None of these
- 235.** The plane $x - 2y + z - 6 = 0$ and the line $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ are related as [Kurukshetra CEE 2001]
 (a) Parallel to the plane (b) Normal to the plane (c) Lies in the plane (d) None of these
- 236.** The condition that the line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$ lies in the plane $ax + by + cz + d = 0$ is
 (a) $ax_1 + by_1 + cz_1 + d = 0$ and $al + bm + cn \neq 0$ (b) $al + bm + cn = 0$ and $ax_1 + by_1 + cz_1 + d \neq 0$
 (c) $ax_1 + by_1 + cz_1 + d = 0$ and $al + bm + cn = 0$ (d) $ax_1 + by_1 + cz_1 = 0$ and $al + bm + cn = 0$
- 237.** $\mathbf{r} = \mathbf{i} + \mathbf{j} + \lambda(2\mathbf{i} + \mathbf{j} + 4\mathbf{k})$ and $\mathbf{r} \cdot (\mathbf{i} + 2\mathbf{j} - \mathbf{k}) = 3$ are the equation of line and plane respectively, then which of the following is true
 (a) The line is perpendicular to plane (b) The line lies in the plane
 (c) The line is parallel to plane but does not lie in plane (d) The line cuts the plane obliquely
- 238.** The line joining the points (3, 5, -7) and (-2, 1, 8) meets the yz -plane at point [Rajasthan PET 2003; MP PET 1993]
 (a) $\left(0, \frac{13}{5}, 2\right)$ (b) $\left(2, 0, \frac{13}{5}\right)$ (c) $\left(0, 2, \frac{13}{5}\right)$ (d) (2, 2, 0)
- 239.** Two lines which do not lie in the same plane are called
 (a) Parallel (b) Coincident (c) Intersecting (d) Skew
- 240.** The planes $x = cy + bz$, $y = az + cx$, $z = bx + ay$ pass through one line, if
 (a) $a + b + c = 0$ (b) $a + b + c = 1$ (c) $a^2 + b^2 + c^2 = 1$ (d) $a^2 + b^2 + c^2 + 2abc = 1$
- 241.** The line $\frac{x-3}{2} = \frac{y-4}{3} = \frac{z-5}{4}$ lies in the plane $4x + 4y - kz - d = 0$. The values of k and d are
 (a) 4, 8 (b) -5, -3 (c) 5, 3 (d) -4, -8
- 242.** If $4x + 4y - kz = 0$ is the equation of the plane through the origin that contains the line $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z}{4}$, then $k =$ [MP PET 1992]
 (a) 1 (b) 3 (c) 5 (d) 7
- 243.** If $\frac{x-1}{l} = \frac{y-2}{m} = \frac{z+1}{n}$ is the equation of the line through (1, 2, -1) and (-1, 0, 1); then (l, m, n) is [MP PET 1992]
 (a) (-1, 0, 1) (b) (1, 1, -1) (c) (1, 2, -1) (d) (0, 1, 0)
- 244.** Given the line $L: \frac{x-1}{3} = \frac{y+1}{2} = \frac{z-3}{-1}$ and plane $P: x - 2y - z = 0$. Then of the following assertions, the only one that is always true is
 (a) L is parallel to plane P (b) L is perpendicular to plane P (c) L lies in the plane P

Three Dimensional Co-ordinate Geometry

- 245.** The coordinates of the point where the line joining the points $(2, -3, 1)$, $(3, -4, -5)$ cuts the plane $2x + y + z = 7$ are
 (a) $(2, 1, 0)$ (b) $(3, 2, 5)$ (c) $(1, -2, 7)$ (d) None of these
- 246.** The point where the line $\frac{x-1}{2} = \frac{y-2}{-3} = \frac{z+3}{4}$ meets the plane $2x + 4y - z = 1$ is [DSSE 1981]
 (a) $(3, -1, 1)$ (b) $(3, 1, 1)$ (c) $(1, 1, 3)$ (d) $(1, 3, 1)$
- 247.** The coordinates of the point where the line $\frac{x-6}{-1} = \frac{y+1}{0} = \frac{z+3}{4}$ meets the plane $x + y - z = 3$ are [MP PET 1998]
 (a) $(2, 1, 0)$ (b) $(7, -1, -7)$ (c) $(1, 2, -6)$ (d) $(5, -1, 1)$
- 248.** The point of intersection of the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z+2}{3}$ and the plane $2x + 3y + z = 0$ is [MP PET 1989]
 (a) $(0, 1, -2)$ (b) $(1, 2, 3)$ (c) $(-1, 9, -25)$ (d) $\left(\frac{-1}{11}, \frac{9}{11}, \frac{-25}{11}\right)$
- 249.** If $p_1 = 0$ and $p_2 = 0$ be two non-parallel planes, then the equation $p_1 + \lambda p_2 = 0$, $\lambda \in R$ represents the family of all planes through the line of intersection of the planes $p_1 = 0$ and $p_2 = 0$, except the plane
 (a) $p_1 = 0$ (b) $p_2 = 0$ (c) $p_1 + p_2 = 0$ (d) $p_1 - p_2 = 0$
- 250.** The direction ratios of the normal to the plane passing through the points $(1, -2, 3)$, $(-1, 2, -1)$ and parallel to $\frac{x-2}{2} = \frac{y+1}{3} = \frac{z}{4}$ is [Tamilnadu (Engg.) 2002]
 (a) $(2, 3, 4)$ (b) $(4, 0, 7)$ (c) $(-2, 0, -1)$ (d) $(2, 0, -1)$
- 251.** The distance between the line $\frac{x-1}{3} = \frac{y+2}{-2} = \frac{z-1}{2}$ and the plane $2x + 2y - z = 6$ is
 (a) 9 units (b) 1 unit (c) 2 units (d) 3 units
- 252.** The distance of the point of intersection of the line $\frac{x-3}{1} = \frac{y-4}{2} = \frac{z-5}{2}$ and the plane $x + y + z = 17$ from the point $(3, 4, 5)$ is given by
 (a) 3 (b) $\frac{3}{2}$ (c) $\sqrt{3}$ (d) None of these

253. The distance of the point $(1, -2, 3)$ from the plane $x - y + z = 5$ measured parallel to the line $\frac{x}{2} = \frac{y}{3} = \frac{z}{-6}$, is [AI CBSE 1984]
- (a) 1 (b) $\frac{6}{7}$ (c) $\frac{7}{6}$ (d) None of these
254. If line $\frac{x-x_1}{l} = \frac{y-y_1}{m} = \frac{z-z_1}{n}$ is parallel to the plane $ax + by + cz + d = 0$, then [MNR 1995; MP PET 1995]
- (a) $\frac{a}{l} = \frac{b}{m} = \frac{c}{n}$ (b) $al + bm + cn = 0$ (c) $\frac{a}{l} + \frac{b}{m} + \frac{c}{n} = 0$ (d) None of these
255. The angle between the line $\frac{x-2}{a} = \frac{y-2}{b} = \frac{z-2}{c}$ and the plane $ax + by + cz + 6 = 0$ is
- (a) $\sin^{-1}\left(\frac{1}{\sqrt{a^2 + b^2 + c^2}}\right)$ (b) 45° (c) 60° (d) 90°
256. The angle between the line $\frac{x}{2} = \frac{y}{3} = \frac{z}{4}$ and the plane $3x + 2y - 3z = 4$ is [MP PET 2003]
- (a) 45° (b) 0° (c) $\cos^{-1}\left(\frac{24}{\sqrt{29}\sqrt{22}}\right)$ (d) 90°
257. The angle between the line $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z+3}{-2}$ and the plane $x + y + 4 = 0$, is [MP PET 1999]
- (a) 0° (b) 30° (c) 45° (d) 90°
258. The angle between the line $\frac{x+1}{3} = \frac{y-1}{2} = \frac{z-2}{4}$ and the plane $2x + y - 3z + 4 = 0$, is [AI CBSE 1981; Pb. CET 1997]
- (a) $\sin^{-1}\left(\frac{4}{\sqrt{406}}\right)$ (b) $\sin^{-1}\left(\frac{-4}{\sqrt{406}}\right)$ (c) $\sin^{-1}\left(\frac{4}{14\sqrt{29}}\right)$ (d) None of these

Advance Level

259. A straight line passes through the point $(2, -1, -1)$. It is parallel to the plane $4x + y + z + 2 = 0$ and is perpendicular to the line $x/1 = y/(-2) = (z-5)/1$. The equation of the straight line are
- (a) $(x-2)/4 = (y+1)/1 = (z+1)/1$ (b) $(x+2)/4 = (y-1)/1 = (z-1)/3$
- (c) $(x-2)/(-1) = (y+1)/1 = (z+1)/3$ (d) $(x+2)/(-1) = (y-1)/1 = (z-1)/3$
260. The equations of the projection of the line $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-3}{3}$ on the plane $x + y + z - 1 = 0$ are
- (a) $x + y + z - 1 = 0 = 2x - y - z + 3$ (b) $x + y - z - 1 = 0 = x + 2y - z - 3$
- (c) $2x - y + 3z - 1 = 0 = x + y + z + 1$ (d) $x + 2y - 3z = 0 = x + y + z + 1$
261. If a plane passes through the point $(1, 1, 1)$ and is perpendicular to the line $\frac{x-1}{3} = \frac{y-1}{0} = \frac{z-1}{4}$, then its perpendicular distance from the origin is [MP PET 1998]
- (a) $\frac{3}{4}$ (b) $\frac{4}{3}$ (c) $\frac{7}{5}$ (d) 1

374 Three Dimensional Co-ordinate Geometry

262. The line $\frac{x-2}{3} = \frac{y+1}{2} = \frac{z-1}{-1}$ intersects the curve $xy = c^2, z = 0$ if $c =$
- (a) ± 1 (b) $\pm 1/3$ (c) $\pm\sqrt{5}$ (d) None of these
263. The points on the line $\frac{x+1}{1} = \frac{y+3}{3} = \frac{z-2}{-2}$ distant $\sqrt{14}$ from the point in which the line meets the plane $3x + 4y + 5z - 5 = 0$ are
- (a) $(0, 0, 0), (2, -4, 6)$ (b) $(0, 0, 0), (3, -4, -5)$ (c) $(0, 0, 0), (2, 6, -4)$ (d) $(2, 6, -4), (3, -4, -5)$
264. The angle between the line $\mathbf{r} = (\mathbf{i} + 2\mathbf{j} - \mathbf{k}) + \lambda(\mathbf{i} - \mathbf{j} + \mathbf{k})$ and the normal to the plane $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} + \mathbf{k}) = 4$ is [MP PET 1997]
- (a) $\sin^{-1}\left(\frac{2\sqrt{2}}{3}\right)$ (b) $\cos^{-1}\left(\frac{2\sqrt{2}}{3}\right)$ (c) $\tan^{-1}\left(\frac{2\sqrt{2}}{3}\right)$ (d) $\cot^{-1}\left(\frac{2\sqrt{2}}{3}\right)$
265. Angle between the line $\mathbf{r} = (2\mathbf{i} - \mathbf{j} + \mathbf{k}) + \lambda(-\mathbf{i} + \mathbf{j} + \mathbf{k})$ and the plane $\mathbf{r} \cdot (3\mathbf{i} + 2\mathbf{j} - \mathbf{k}) = 4$ is [AMU 1993]
- (a) $\cos^{-1}\left(\frac{2}{\sqrt{42}}\right)$ (b) $\cos^{-1}\left(-\frac{2}{\sqrt{42}}\right)$ (c) $\sin^{-1}\left(\frac{2}{\sqrt{42}}\right)$ (d) $\sin^{-1}\left(-\frac{2}{\sqrt{42}}\right)$

Sphere

Basic Level

266. The ratio in which the sphere $x^2 + y^2 + z^2 = 504$ divides the line segment AB joining the points $A(12, -4, 8)$ and $(27, -9, 18)$ is given by
- (a) $2:3$ externally (b) $2:3$ internally (c) $1:2$ externally (d) None of these
267. The graph of the equation $y^2 + z^2 = 0$ in three dimensional space is
- (a) x -axis (b) z -axis (c) y -axis (d) yz -plane
268. A point moves so that the sum of the squares of its distances from two given points remains constant. The locus of the point is
- (a) A line (b) A plane (c) A sphere (d) None of these
269. The locus of the equation $x^2 + y^2 + z^2 + 1 = 0$ is
- (a) An empty set (b) A sphere (c) A degenerate set (d) A pair of planes
270. Let $(3, 4, -1)$ and $(-1, 2, 3)$ are the end points of a diameter of sphere. Then the radius of the sphere is equal to [Orissa JEE 2003]
- (a) 1 (b) 2 (c) 3 (d) 9
271. The number of spheres of radius ' a ' touching all the coordinate planes is
- (a) 4 (b) 8 (c) 1 (d) None of these
272. The equation of the sphere touching the three coordinate planes is [AMU 2002]
- (a) $x^2 + y^2 + z^2 + 2a(x + y + z) + 2a^2 = 0$ (b) $x^2 + y^2 + z^2 - 2a(x + y + z) + 2a^2 = 0$
- (c) $x^2 + y^2 + z^2 \pm 2a(x + y + z) + 2a^2 = 0$ (d) $x^2 + y^2 + z^2 \pm 2ax \pm 2ay \pm 2az + 2a^2 = 0$
273. Equation $ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy + 2ux + 2vy + 2wz + d = 0$ represent, a sphere, if [MP PET 1990]
- (a) $a = b = c$ (b) $f = g = h = 0$
- (c) $v = u = w$ (d) $a = b = c$ and $f = g = h = 0$
274. The centre of the sphere which passes through $(a, 0, 0), (0, b, 0), (0, 0, 0)$ is [AMU 1990]

- (a) $\left(\frac{a}{2}, 0, 0\right)$ (b) $\left(0, \frac{b}{2}, 0\right)$ (c) $\left(0, 0, \frac{c}{2}\right)$ (d) $\left(\frac{a}{2}, \frac{b}{2}, \frac{c}{2}\right)$
275. The equation $ax^2 + ay^2 + az^2 + 2ux + 2vy + 2wz + d = 0$, $a \neq 0$, represents a sphere if
 (a) $u^2 + v^2 + w^2 + ad \leq 0$ (b) $u^2 + v^2 + w^2 + ad \geq 0$ (c) $u^2 + v^2 + w^2 - ad \leq 0$ (d) $u^2 + v^2 + w^2 - ad \geq 0$
276. The radius of the sphere $x^2 + y^2 + z^2 - 6x + 8y - 10z + 1 = 0$ is [Kurukshetra CEE 1994]
 (a) 7 (b) 5 (c) 2 (d) 15
277. Centre of the sphere $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) + (z - z_1)(z - z_2) = 0$ is
 (a) (x_2, y_2, z_2) (b) $\left(\frac{x_1 - x_2}{2}, \frac{y_1 - y_2}{2}, \frac{z_1 - z_2}{2}\right)$ (c) $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2}\right)$ (d) (x_1, y_1, z_1)
278. The equation of the tangent plane at a point (x_1, y_1, z_1) on the sphere $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$ is
 (a) $xx_1 + yy_1 + zz_1 + ux + vy + wz + d = 0$ (b) $xx_1 + yy_1 + zz_1 + ux_1 + vy_1 + wz_1 + d = 0$
 (c) $xx_1 + yy_1 + zz_1 + u(x + x_1) + v(y + y_1) + w(z + z_1) + d = 0$ (d) None of these
279. If two spheres of radii r_1 and r_2 cut orthogonally, then the radius of the common circle is
 (a) $r_1 r_2$ (b) $\sqrt{(r_1^2 + r_2^2)}$ (c) $r_1 r_2 \sqrt{(r_1^2 + r_2^2)}$ (d) $\frac{r_1 r_2}{\sqrt{(r_1^2 + r_2^2)}}$
280. The equation of the sphere, concentric with the sphere $x^2 + y^2 + z^2 - 4x - 6y - 8z - 5 = 0$ and which passes through $(0, 1, 0)$, is [Pb. CET 1994]
 (a) $x^2 + y^2 + z^2 - 4x - 6y - 8z + 1 = 0$ (b) $x^2 + y^2 + z^2 - 4x - 6y - 8z + 5 = 0$
 (c) $x^2 + y^2 + z^2 - 4x - 6y - 5z + 2 = 0$ (d) $x^2 + y^2 + z^2 - 4x - 6y - 5z + 3 = 0$
281. The radius of the sphere which passes through the points $(0, 0, 0)$, $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$ is [AMU 1991]
 (a) $\frac{1}{2}$ (b) 1 (c) $\sqrt{3}$ (d) $\sqrt{3}/2$
282. The coordinates of the centre of the sphere $(x + 1)(x + 3) + (y - 2)(y - 4) + (z + 1)(z + 3) = 0$ are [AMU 1987]
 (a) $(1, -1, 1)$ (b) $(-1, 1, -1)$ (c) $(2, -3, 2)$ (d) $(-2, 3, -2)$
283. Equation of the sphere with centre $(1, -1, 1)$ and radius equal to that of sphere $2x^2 + 2y^2 + 2z^2 - 2x + 4y - 6z = 1$ is [DCE 1994]
 (a) $x^2 + y^2 + z^2 + 2x - 2y + 2z + 1 = 0$ (b) $x^2 + y^2 + z^2 - 2x + 2y - 2z - 1 = 0$
 (c) $x^2 + y^2 + z^2 - 2x + 2y - 2z + 1 = 0$ (d) None of these
284. The equation of the sphere concentric with the sphere $x^2 + y^2 + z^2 - 2x - 6y - 8z - 5 = 0$ and which passes through the origin is [Pb. CET 1990]
 (a) $x^2 + y^2 + z^2 - 2x - 6y - 8z = 0$ (b) $x^2 + y^2 + z^2 - 6y - 8z = 0$
 (c) $x^2 + y^2 + z^2 = 0$ (d) None of these
285. The equation of the sphere with centre at $(2, 3, -4)$ and touching the plane $2x + 6y - 3z + 15 = 0$ is
 (a) $x^2 + y^2 + z^2 - 4x - 6y + 8z - 20 = 0$ (b) $x^2 + y^2 + z^2 + 4x - 6y - 8z - 20 = 0$
 (c) $x^2 + y^2 + z^2 - 4x - 6y + 8z + 20 = 0$ (d) None of these
286. Spheres $x^2 + y^2 + z^2 + x + y + z - 1 = 0$ and $x^2 + y^2 + z^2 + x + y + z - 5 = 0$ [AMU 1991]

376 Three Dimensional Co-ordinate Geometry

- (a) Intersect in a plane (b) Intersect in five points (c) Do not intersect (d) None of these
287. If \mathbf{r} be position vector of any point on a sphere and \mathbf{a} and \mathbf{b} are respectively position vectors of the extremities of a diameter, then [AMU 1999]
- (a) $\mathbf{r} \cdot (\mathbf{a} - \mathbf{b}) = 0$ (b) $\mathbf{r} \cdot (\mathbf{r} - \mathbf{a}) = 0$ (c) $(\mathbf{r} + \mathbf{a}) \cdot (\mathbf{r} + \mathbf{b}) = 0$ (d) $(\mathbf{r} - \mathbf{a}) \cdot (\mathbf{r} - \mathbf{b}) = 0$
288. The centre of the sphere $\alpha \mathbf{r} - 2\mathbf{u} \cdot \mathbf{r} = \beta$, ($\alpha \neq 0$) is [AMU 1999]
- (a) $-\mathbf{u} / \alpha$ (b) \mathbf{u} / α (c) $\alpha \mathbf{u} / \beta$ (d) $\frac{\alpha + \beta}{\alpha} \mathbf{u}$
289. The spheres $\mathbf{r}^2 + 2\mathbf{u}_1 \cdot \mathbf{r} + 2\mathbf{d}_1 = 0$ and $\mathbf{r}^2 + 2\mathbf{u}_2 \cdot \mathbf{r} + 2\mathbf{d}_2 = 0$ cut orthogonally, if [AMU 1999]
- (a) $\mathbf{u}_1 \cdot \mathbf{u}_2 = 0$ (b) $\mathbf{u}_1 + \mathbf{u}_2 = 0$
 (c) $\mathbf{u}_1 \cdot \mathbf{u}_2 = \mathbf{d}_1 + \mathbf{d}_2$ (d) $(\mathbf{u}_1 - \mathbf{u}_2) \cdot (\mathbf{u}_1 + \mathbf{u}_2) = \mathbf{d}_1^2 + \mathbf{d}_2^2$

Advance level

290. If a sphere of constant radius k passes through the origin and meets the axis in A, B, C then the centroid of the triangle ABC lies on
- (a) $x^2 + y^2 + z^2 = k^2$ (b) $x^2 + y^2 + z^2 = 4k^2$ (c) $9(x^2 + y^2 + z^2) = 4k^2$ (d) $9(x^2 + y^2 + z^2) = k^2$
291. The smallest radius of the sphere passing through $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$ is [Pb. CET 1997,99; Kurukshetra CEE 1996]
- (a) $\sqrt{\frac{3}{5}}$ (b) $\sqrt{\frac{3}{8}}$ (c) $\sqrt{\frac{2}{3}}$ (d) $\sqrt{\frac{5}{12}}$
292. In order that bigger sphere (centre C_1 , radius R) may fully contain a smaller sphere (center C_2 , radius r), the correct relationship is [AMU 1991]
- (a) $C_1 C_2 < r + R$ (b) $C_1 C_2 < R - r$ (c) $C_1 C_2 < 2(R - r)$ (d) $C_1 C_2 < \frac{1}{2}(R + r)$
293. A sphere $x^2 + y^2 + z^2 = 9$ is cut by the plane $x + y + z = 3$. The radius of the circle so formed is
- (a) $\sqrt{6}$ (b) $\sqrt{3}$ (c) 3 (d) 6
294. The radius of the circle $x^2 + y^2 + z^2 - 2y - 4z = 11$, $x + 2y + 2z = 15$ is [AMU 1990,92]
- (a) 4 (b) $\sqrt{7}$ (c) 5 (d) 7
295. The line $\frac{x+1}{-1} = \frac{y-12}{5} = \frac{z-7}{2}$ cuts the surface $11x^2 - 5y^2 + z^2 = 0$ in the point
- (a) $(1, 1, 1)$ and $(1, 2, 3)$ (b) $(1, -1, 2)$ and $(1, 2, 4)$ (c) $(1, 2, 3)$ and $(2, -3, 1)$ (d) None of these
296. The equation of the sphere circumscribing the tetrahedron whose faces are $x = 0, y = 0, z = 0$ and $x/a + y/b + z/c = 1$ is
- (a) $x^2 + y^2 + z^2 = a^2 + b^2 + c^2$
 (b) $x^2 + y^2 + z^2 - ax - by - cz = 0$
 (c) $x^2 + y^2 + z^2 - 2ax - 2by - 2cz = 0$
 (d) None of these
297. A plane passes through a fixed point (a, b, c) . The locus of the foot of the perpendicular drawn to it from the origin is

(a) $x^2 + y^2 + z^2 + ax + by + cz = 0$

(b) $x^2 + y^2 + z^2 - ax - by - cz = 0$

(c) $x^2 + y^2 + z^2 + 2ax + 2by + 2cz = 0$

(d) $x^2 + y^2 + z^2 + 2ax - 2by - 2cz = 0$

298. The equation of the sphere passing through the point $(1, 3, -2)$ and the circle $y^2 + z^2 = 25$ and $x = 0$ is [DCE 1998]

(a) $x^2 + y^2 + z^2 + 11x + 25 = 0$

(b) $x^2 + y^2 + z^2 - 11x + 25 = 0$

(c) $x^2 + y^2 + z^2 + 11x - 25 = 0$

(d) $x^2 + y^2 + z^2 - 11x - 25 = 0$

299. Radius of the circle $\mathbf{r}^2 + \mathbf{r} \cdot (2\mathbf{i} - 2\mathbf{j} - 4\mathbf{k}) - 19 = 0$, $\mathbf{r} \cdot (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) + 8 = 0$ is [Kurukshetra CEE 1996, DCE 1997]

(a) 2

(b) 3

(c) 4

(d) 5


300. The shortest distance from the point $(1, 2, -1)$ to the surface of the sphere $x^2 + y^2 + z^2 = 24$ is [Pb. CET 1996]

(a) $3\sqrt{6}$

(b) $2\sqrt{6}$

(c) $\sqrt{6}$

(d) 2


Three Dimensional Co-ordinate Geometry
Assignment (Basic and Advance)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
c	c	a	c	b	b	d	b	a	b	d	a	c	d	c	b	c	b	d	a
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
b	c	a	b	d	c	b	a	b	b	a	d	c	a	a	d	a	b	d	d
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
a	b	d	a	b	a	b	c	b	a	d	a	a	d	c	a	a	b	d	d
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
d	b	b	c	c	a	b	b	d	a	a	b	b	a	b	a	b	d	d	c
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
b	b	a	c	a	d	a	a	d	c	a	b	d	d	a	d	d	d	a	a
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
b	c	c	c	a	d	b	b	c	c	a	b	a	c	c	a	c	c	b	b
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
b	d	c	b	c	d	c	d	d	a	c	c	d	c	a	c	b	b	a	c
141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
a	c	b	c	d	b	b	b	a	a	d	d	a	a	d	d	a	b	a	a
161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
d	a	b	a	c	a	a	b	b	d	d	a	a	a	d	a	d	b	a	d
181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
b	c	b	d	b	a	c	a	a	a	b	b	c	b	d	b	c	a	b	a
201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220
a	b	a	c	d	b	c	b	b	d	a	a	a	d	a	b	b	b	c	b
221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
c	a	b	d	c	c	b	b	b	a	a	d	a	a	a	c	b	a	d	d
241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260
c	c	b	c	c	a	d	d	b	d	d	a	a	b	d	b	c	b	c	a
261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280
c	c	c	a	d	a	a	c	a	c	b	d	d	d	d	a	c	c	d	b
281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300

d	d	b	a	a	c	d	d	c	c	c	b	a	b	c	b	b	c	c	c
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---