

8. Differentiation

- **Derivatives**

- Suppose f is a real-valued function and a is a point in its domain of definition. The derivative of f at a [denoted by $f'(a)$] is defined as

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}, \text{ provided the limit exists.}$$

Derivative of $f(x)$ at a is denoted by $f'(a)$.

- Suppose f is a real-valued function. The derivative of f { denoted by $f'(x)$ or $\frac{d}{dx}[f(x)]$ } is defined as

$$\frac{d}{dx}[f(x)] = f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \text{ provided the limit exists.}$$

This definition of derivative is called the first principle of derivative.

Example: Find the derivative of $f(x) = x^2 + 2x$ using first principle of derivative.

Solution: We know that $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$\begin{aligned} \therefore f'(x) &= \lim_{h \rightarrow 0} \frac{(x+h) + 2(x+h) - (x^2 + 2x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + h^2 + 2hx + 2x + 2h - x^2 - 2x}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^2 + 2hx + 2h}{h} \\ &= \lim_{h \rightarrow 0} (h + 2x + 2) \\ &= 0 + 2x + 2 = 2x + 2 \\ f'(x) &= 2x + 2 \end{aligned}$$

- **Derivatives of Polynomial Functions**

For the functions u and v (provided u' and v' are defined in a common domain),

- - $(u \pm v)' = u' \pm v'$
 - $(uv)' = u'v + uv'$ (Product rule)
 - $\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$ (Quotient rule)

- **Derivatives of Trigonometric Functions**

- $\frac{d}{dx}(x^n) = nx^{n-1}$ for any positive integer n
- $\frac{d}{dx}(a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0) = na_n x^{n-1} + (n-1)a_{n-1} x^{n-2} + \dots + a_1$

- $\frac{d}{dx}(\sin x) = \cos x$
- $\frac{d}{dx}(\cos x) = -\sin x$
- $\frac{d}{dx}(\tan x) = \sec^2 x$

Example: Find the derivative of the function $f(x) = (3x^2 + 4x + 1) \cdot \tan x$

Solution: We have,

$f(x) = 3x^2 + 4x + 1 \cdot \tan x$
Differentiating both sides with respect to x , $f'(x) = 3x^2 + 4x + 1 \cdot \frac{d}{dx} \tan x + \tan x \cdot \frac{d}{dx} (3x^2 + 4x + 1)$
 $f'(x) = 3x^2 + 4x + 1 \cdot \sec^2 x + \tan x \cdot 6x + 4 = 3x^2 + 4x + 1 \cdot \sec^2 x + 6x + 4 \tan x$

The derivatives of exponential functions are as follows:

- $\frac{d}{dx}(e^x) = e^x$
- $\frac{d}{dx}(e^{ax}) = ae^{ax}$

• **Mean value theorem:**

If $f: [a, b] \rightarrow \mathbf{R}$ is continuous on $[a, b]$ and differentiable on (a, b) , then there exists some $c \in (a, b)$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

Example: Verify Mean Value Theorem for the function:

$f(x) = 2x^2 - 17x + 30$ in the interval $\left[\frac{5}{2}, 6\right]$.

Solution:

$$f(x) = 2x^2 - 17x + 30$$

$$\therefore f'(x) = 4x - 17$$

The function $f(x)$ being a polynomial, is continuous on $\left[\frac{5}{2}, 6\right]$ and is differentiable on $\left(\frac{5}{2}, 6\right)$.

$$\text{Also, } f\left(\frac{5}{2}\right) = 2\left(\frac{5}{2}\right)^2 - 17\left(\frac{5}{2}\right) + 30 = 0$$

$$\text{and, } f(6) = 2(6)^2 - 17 \times 6 + 30 = 0$$

$$\therefore f\left(\frac{5}{2}\right) = f(6)$$


$$\frac{f(6) - f\left(\frac{5}{2}\right)}{6 - \frac{5}{2}} = 0$$

Now,

According to Mean Value Theorem (MVT), there exists $c \in \left(\frac{5}{2}, 6\right)$ such that $f'(c) = 0$.

$$\therefore 4c - 17 = 0$$

$$\Rightarrow c = \frac{17}{4} \in \left(\frac{5}{2}, 6\right)$$



Therefore, M.V.T is verified.