

MANIPAL

Engineering Entrance Exam

Solved Paper 2008

Chemistry

- Which of the following is diamagnetic ?
(a) H_2^+ (b) O_2
(c) Li_2 (d) He_2^+
- Which of the following can participate in linkage isomerism ?
(a) NO_2^-
(b) $\text{H}_2\ddot{\text{N}}\text{CH}_2\text{CH}_2\ddot{\text{N}}\text{H}_2$
(c) H_2O
(d) $:\text{NH}_3$
- By heating phenol with chloroform in alkali, it is converted into
(a) salicylic acid
(b) salicylaldehyde
(c) anisole
(d) phenyl benzoate
- Osmotic pressure observed when benzoic acid is dissolved in benzene is less than that expected from theoretical considerations. This is because
(a) benzoic acid is an organic solute
(b) benzoic acid has higher molar mass than benzene
(c) benzoic acid gets associated in benzene
(d) benzoic acid gets dissociated in benzene
- The formula mass of Mohr's salt is 392. The iron present in it is oxidised by KMnO_4 in acid medium. The equivalent mass of Mohr's salt is
(a) 392 (b) 31.6
(c) 278 (d) 156
- Solubility product of a salt AB is $1 \times 10^{-8} \text{ M}^2$ in a solution in which the concentration of A^+ ions is 10^{-3} M . The salt will precipitate when the concentration of B^- ions is kept
(a) between 10^{-8} M to 10^{-7} M
(b) between 10^{-7} M to 10^{-8} M
(c) $> 10^{-5} \text{ M}$
(d) $< 10^{-8} \text{ M}$
- The decomposition of a certain mass of CaCO_3 gave 11.2 dm^3 of CO_2 gas at STP. The mass of KOH required to completely neutralise the gas is
(a) 56 g (b) 28 g
(c) 42 g (d) 20 g
- The basicity of aniline is less than that of cyclohexylamine. This is due to
(a) $+R$ -effect of $-\text{NH}_2$ group
(b) $-I$ -effect of $-\text{NH}_2$ group
(c) $-R$ -effect of $-\text{NH}_2$ group
(d) hyperconjugation effect
- A distinctive and characteristic functional group of fats is
(a) a peptide group
(b) an ester group
(c) an alcoholic group
(d) a ketonic group
- Which of the following compound is expected to be optically active ?
(a) $(\text{CH}_3)_2\text{CHCHO}$
(b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$
(c) $\text{CH}_3\text{CH}_2\text{CHBrCHO}$
(d) $\text{CH}_3\text{CH}_2\text{CBr}_2\text{CHO}$
- Which cycloalkane has the lowest heat of combustion per CH_2 group ?
(a) Cyclopropane (b) Cyclobutane
(c) Cyclopentane (d) Cyclohexane
- The physical states of dispersing phase and dispersion medium in colloid like pesticide spray respectively, are
(a) gas, liquid (b) solid, gas
(c) liquid, solid (d) liquid, gas

13. Potassium dichromate is used
 - (a) in electroplating
 - (b) as a reducing agent
 - (c) it oxidises ferrous ions into ferric ions in acidic media as an oxidising agent
 - (d) as an insecticide
14. Which one of the following statements is incorrect for the sucrose ?
 - (a) It is obtained from cane sugar
 - (b) It is not reducing sugar
 - (c) On hydrolysis, it gives equal quantities of D-glucose and D-fructose
 - (d) It gives aspartame when it is heated at 210°C
15. Inductive effect involves
 - (a) displacement of σ -electrons
 - (b) delocalisation of π -electrons
 - (c) delocalisation of σ -electrons
 - (d) displacement of π -electrons
16. The atomic number of Ni and Cu are 28 and 29 respectively. The electronic configuration $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 3d^{10}$ represents
 - (a) Cu^+
 - (b) Cu^{2+}
 - (c) Ni^{2+}
 - (d) Ni
17. In which of the following complex ion, the central metal ion is in a state of sp^3d^2 hybridisation ?
 - (a) $[\text{CoF}_6]^{3-}$
 - (b) $[\text{Co}(\text{NH}_3)_6]^{3+}$
 - (c) $[\text{Fe}(\text{CN})_6]^{3-}$
 - (d) $[\text{Cr}(\text{NH}_3)_6]^{3+}$
18. The formation of $\text{O}_2^+[\text{PtF}_6]^-$ is the basis for the formation of xenon fluorides. This is because
 - (a) O_2 and Xe have comparable sizes
 - (b) Both O_2 and Xe are gases
 - (c) O_2 and Xe have comparable ionisation energies
 - (d) Both (a) and (c)
19. The density of a gas is 1.964 g dm^{-3} at 273 K and 76 cm Hg . The gas is
 - (a) CH_4
 - (b) C_2H_6
 - (c) CO_2
 - (d) Xe
20. ΔG° vs T plot in the Ellingham's diagram slopes downwards for the reactions
 - (a) $\text{Mg} + \frac{1}{2} \text{O}_2 \longrightarrow \text{MgO}$
 - (b) $2\text{Ag} + \frac{1}{2} \text{O}_2 \longrightarrow \text{Ag}_2\text{O}$
 - (c) $\text{CO} + \frac{1}{2} \text{O}_2 \longrightarrow \text{CO}_2$
 - (d) All of the above
21. When a mixture of calcium benzoate and calcium acetate is dry distilled, the resulting compound is
 - (a) acetophenone
 - (b) benzaldehyde
 - (c) benzophenone
 - (d) acetaldehyde
22. In a metallic crystal
 - (a) the valence electrons constitute a sea of mobile electrons
 - (b) the valence electrons are localised in between the kernels
 - (c) the valence electrons remain within the field of influence of their own kernels
 - (d) None of the above
23. Which of the following is correct, based on molecular orbital theory for peroxide ion ?
 - (a) Its bond order is one and it is paramagnetic
 - (b) Its bond order is two and it is diamagnetic
 - (c) Its bond order is one and it is diamagnetic
 - (d) Its bond order is two and it is paramagnetic
24. Insulin regulates the metabolism of
 - (a) minerals
 - (b) amino acids
 - (c) glucose
 - (d) vitamins
25. Which of the following electrolyte will have maximum flocculation value for $\text{Fe}(\text{OH})_3$ sol ?
 - (a) NaCl
 - (b) Na_2S
 - (c) $(\text{NH}_4)_3\text{PO}_4$
 - (d) K_2SO_4
26. The concentration of a reactant X decreases from 0.1 M to 0.005 M in 40 min . If the reaction follows first order kinetics, the rate of the reaction when the concentration of X is 0.01 M will be
 - (a) $1.73 \times 10^{-4} \text{ M min}^{-1}$
 - (b) $3.47 \times 10^{-4} \text{ M min}^{-1}$
 - (c) $3.47 \times 10^{-5} \text{ M min}^{-1}$
 - (d) $7.5 \times 10^{-4} \text{ M min}^{-1}$



27. At pH = 4, glycine exists as

- (a) $\text{H}_3\text{N}^+ - \text{CH}_2 - \text{COO}^-$
 (b) $\text{H}_3\text{N}^+ - \text{CH}_2 - \text{COOH}$
 (c) $\text{H}_2\text{N} - \text{CH}_2 - \text{COOH}$
 (d) $\text{H}_2\text{N} - \text{CH}_2 - \text{COO}^-$


28. Which of the following taking place in the blast furnace is endothermic ?

- (a) $\text{CaCO}_3 \longrightarrow \text{CaO} + \text{CO}_2$
 (b) $2\text{C} + \text{O}_2 \longrightarrow 2\text{CO}$
 (c) $\text{C} + \text{O}_2 \longrightarrow \text{CO}_2$
 (d) $\text{Fe}_2\text{O}_3 + 3\text{CO} \longrightarrow 2\text{Fe} + 3\text{CO}_2$

29. The emf E° of the following cells are :



emf of the following cell is




- (a) 0.64 V (b) 1.10 V
 (c) 1.56 V (d) -0.64 V

30. The formation of cyanohydrin from acetone is which type of reaction ?

- (a) Electrophilic substitution reaction
 (b) Electrophilic addition reaction
 (c) Nucleophilic addition reaction
 (d) Nucleophilic substitution reaction

31. Name the end product in the following series of reactions



- (a) CH_3OH (b) CH_4
 (c) $\text{CH}_3\text{COONH}_4$ (d) CH_3CN

32. The presence of unpaired electron in phosphorous atom is explained by which principle ?

- (a) Aufbau principle
 (b) Pauli's exclusion principle
 (c) Hund's rule
 (d) Heisenberg's principle

33. If a cricket ball having mass of 200 g is thrown with a speed of $3 \times 10^3 \text{ cm/s}$, then calculate the wavelength related to it.

- (a) $2.2 \times 10^{-27} \text{ cm}$
 (b) $1.104 \times 10^{-32} \text{ cm}$

(c) $1.104 \times 10^{-27} \text{ cm}$

(d) $1.104 \times 10^{-33} \text{ cm}$

34. Which type of stacking pattern is found in sodium chloride crystal lattice ?

- (a) $a-b-a-b$ (b) $a-a-a$
 (c) $a-b-c-a-b-c$ (d) None of these

35. Equivalent weight of a bivalent metal is 37.2. The molecular weight of its chloride is

- (a) 412.2 (b) 216
 (c) 145.4 (d) 108.2

36. Phenolphthalein is obtained by heating phthalic anhydride with conc. H_2SO_4 and

- (a) benzyl alcohol (b) benzene
 (c) phenol (d) benzoic acid

37. Freezing point of urea solution is -0.6°C . How much urea (m.wt. = 60 g/mol) will be required to dissolve in 3 kg water ?

($k_f = 1.5^\circ\text{C kg mol}^{-1}$)

- (a) 24 g (b) 36 g
 (c) 60 g (d) 72 g

38. If $K < 1.0$, what will be the value of ΔG° of the following ?

- (a) Zero (b) 1.0
 (c) Positive (d) Negative

39. The normality of a solution containing 32.5 g of $(\text{COOH})_2 \cdot 2\text{H}_2\text{O}$ per 0.5 L is

- (a) 10 N (b) 1 N
 (c) 2 N (d) 0.1 N

40. For the titration of KOH vs $(\text{COOH})_2 \cdot 2\text{H}_2\text{O}$, the suitable indicator is

- (a) methyl orange
 (b) phenolphthalein
 (c) methyl red
 (d) All can be used


41. The radius of Na^+ is 95 pm and that of Cl^- ion is 181 pm. The coordination number of Na^+ is

- (a) 8 (b) 6
 (c) 4 (d) unpredictable

42. In van der Waals' equation of state of the gas law, the constant 'b' is a measure of

- (a) intermolecular repulsion
 (b) intermolecular attraction
 (c) volume occupied by the molecules
 (d) intermolecular collisions per unit volume

43. Which reaction intermediate is formed during the condensation reaction between acetaldehyde and formaldehyde ?



44. 2, 2-dichloro propane on hydrolysis yields

- (a) acetone
 (b) 2, 2-propane diol
 (c) *iso*-propyl alcohol
 (d) acetaldehyde

45. Phenols are more acidic than alcohols because

- (a) phenoxide ion is stabilised by resonance
 (b) phenols are more soluble in polar solvents
 (c) phenoxide ions do not exhibit resonance
 (d) alcohols do not lose H atoms at all

46. Lemon gives sour taste because of

- (a) citric acid (b) tartaric acid
 (c) oxalic acid (d) acetic acid





47. When ammonium chloride is added to ammonia solution, the pH of the resulting solution will be

- (a) increased (b) seven
 (c) decreased (d) unchanged

48. Which of the following has highest second ionisation energy ?

- (a) Calcium (b) Chromium
 (c) Iron (d) Cobalt


49. The standard reduction potentials at 298 K for the following half-cell reactions are given below



Which one of the following is the strongest reducing agent ?

- (a) Zn(s) (b) Cr(s)
 (c) H₂(g) (d) Fe²⁺(aq)

50. Following reaction,






is an example of

- (a) elimination reaction
 (b) free radical substitution
 (c) nucleophilic substitution
 (d) electrophilic substitution

51. The unit of rate for a first order reaction is

- (a) L s⁻¹
 (b) mol⁻¹ L s⁻¹
 (c) mol L⁻¹ s⁻¹
 (d) mol s⁻¹

52. *Iso*-propyl amine with excess of acetyl chloride will give



53. C₂H₅CHO and (CH₃)₂CO can be distinguished by testing with

- (a) phenyl hydrazine
 (b) hydroxyl amine
 (c) Fehling solution
 (d) sodium bisulphite

54. Glucose molecule reacts with 'X' number of molecules of phenylhydrazine to yield osazone. The value of 'X' is

- (a) four (b) one
 (c) two (d) three

55. The oxidation number of chromium in CrO₅ is

- (a) + 3 (b) + 5
 (c) + 10 (d) + 6

56. Liquor ammonia bottles are opened only after cooling. This is because

- (a) it is a mild explosive
 (b) it generates high vapour pressure
 (c) both (a) and (b)
 (d) it is lachrymatory


57. What will be the proportion of moles of metal (Cu : Ni : Ag) at cathode according to the second law of Faraday ?

- (a) 1 : 2 : 1 (b) 2 : 2 : 1
 (c) 1 : 2 : 2 (d) 1 : 1 : 2

58. Which equation is true to calculate the energy of activation, if the rate of reaction is doubled by increasing temperature from T_1 K to T_2 K ?

(a) $\log_{10} \frac{k_1}{k_2} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$
 (b) $\log_{10} \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{1}{T_2} - \frac{1}{T_1} \right]$
 (c) $\log_{10} \frac{1}{2} = \frac{E_a}{2.303} \left[\frac{1}{T_2} - \frac{1}{T_1} \right]$
 (d) $\log_{10} 2 = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$


59. For a reversible reaction :



temperature at which the above reaction attains equilibrium is about

- (a) 400 K
 (b) 500 K
 (c) 273 K
 (d) 373 K

60. Which of the following gives aldol condensation reaction ?



Mathematics

1. The range of the function $f(x) = x^2 + \frac{1}{x^2 + 1}$

is

- (a) $[1, \infty)$ (b) $[2, \infty)$
 (c) $\left[\frac{3}{2}, \infty \right)$ (d) None of these

2. If $f(x) = \begin{cases} ax^2 + b, & b \neq 0, x \leq 1 \\ bx^2 + ax + c, & x > 1 \end{cases}$, then

$f(x)$ is continuous and differentiable at $x = 1$, if

- (a) $c = 0, a = 2b$ (b) $a = b, c \in R$
 (c) $a = b, c = 0$ (d) $a = b, c \neq 0$

3. If a circle passes through the point (1, 2) and cuts the circle $x^2 + y^2 = 4$ orthogonally, then

the equation of the locus of its centre is

- (a) $x^2 + y^2 - 3x - 8y + 1 = 0$
 (b) $x^2 + y^2 - 2x - 6y - 7 = 0$
 (c) $2x + 4y - 9 = 0$
 (d) $2x + 4y - 1 = 0$

4. If $\int f(x) \sin x \cos x \, dx$

$$= \frac{1}{2(b^2 - a^2)} \log[f(x)] + c,$$

then $f(x)$ is equal to

- (a) $\frac{1}{a^2 \sin^2 x + b^2 \cos^2 x}$
 (b) $\frac{1}{a^2 \sin^2 x - b^2 \cos^2 x}$
 (c) $\frac{1}{a^2 \cos^2 x - b^2 \sin^2 x}$
 (d) $\frac{1}{a^2 \cos^2 x + b^2 \sin^2 x}$

5. The points representing complex number z for which $|z - 3| = |z - 5|$ lie on the locus given by

- (a) an ellipse
 (b) a circle
 (c) a straight line
 (d) None of the above

6. The value of α , for which the equation $x^2 - (\sin \alpha - 2)x - (1 + \sin \alpha) = 0$ has roots whose sum of square is least, is

- (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{4}$
 (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{6}$

7. For $n \in N, 10^{n-2} \geq 81n$, if

- (a) $n > 5$ (b) $n \geq 5$
 (c) $n < 5$ (d) $n > 8$

8. The two consecutive terms in the expansion of $(3 + 2x)^{74}$ whose coefficients are equal, are
 (a) 11, 12 (b) 7, 8
 (c) 30, 31 (d) None of these
9. The value of $2.\overline{357}$ is
 (a) $\frac{2355}{999}$ (b) $\frac{2355}{1000}$
 (c) $\frac{2355}{1111}$ (d) None of these
10. Let $S_n = \frac{1}{1^3} + \frac{1+2}{1^3+2^3} + \dots + \frac{1+2+\dots+n}{1^3+2^3+\dots+n^3}$
 $n = 1, 2, 3, \dots$. Then, S_n is not greater than
 (a) $\frac{1}{2}$ (b) 1
 (c) 2 (d) 4
11. If $E(\theta) = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ and θ and ϕ differ by an odd multiple of $\frac{\pi}{2}$, then $E(\theta)E(\phi)$ is a
 (a) unit matrix
 (b) null matrix
 (c) diagonal matrix
 (d) None of the above
12. A parabola is drawn with its focus at (3, 4) and vertex at the focus of the parabola $y^2 - 12x - 4y + 4 = 0$. The equation of the parabola is
 (a) $y^2 - 8x - 6y + 25 = 0$
 (b) $y^2 - 6x + 8y - 25 = 0$
 (c) $x^2 - 6x - 8y + 25 = 0$
 (d) $x^2 + 6x - 8y - 25 = 0$
13. If p, p' denote the lengths of the perpendiculars from the focus and the centre of an ellipse with semi major axis of length a respectively on a tangent to the ellipse and r denotes the focal distance of the point, then
 (a) $ap' = rp + 1$ (b) $rp = ap'$
 (c) $ap = rp' + 1$ (d) $ap = rp'$
14. The equation of perpendicular bisectors of sides AB and AC of a $\triangle ABC$ are $x - y + 5 = 0$ and $x + 2y = 0$ respectively. If the coordinates of vertex A are (1, -2), the equation of BC is
 (a) $14x + 23y - 40 = 0$
 (b) $14x - 23y + 40 = 0$
 (c) $23x + 14y - 40 = 0$
 (d) $23x - 14y + 40 = 0$
15. If $\cos \theta = -\frac{\sqrt{3}}{2}$ and $\sin \alpha = -\frac{3}{5}$, where θ does not lie in the third quadrant, then $\frac{2 \tan \alpha + \sqrt{3} \tan \theta}{\cot^2 \theta + \cos \alpha}$ is equal to
 (a) $\frac{7}{22}$ (b) $\frac{5}{22}$ (c) $\frac{9}{22}$ (d) $\frac{22}{5}$
16. A parallelogram is constructed on the vectors $\vec{a} = 3\vec{\alpha} - \vec{\beta}$, $\vec{b} = \vec{\alpha} + 3\vec{\beta}$, if $|\vec{\alpha}| = |\vec{\beta}| = 2$ and angle between $\vec{\alpha}$ and $\vec{\beta}$ is $\frac{\pi}{3}$, then length of a diagonal of the parallelogram is
 (a) $4\sqrt{5}$ (b) $4\sqrt{3}$
 (c) $4\sqrt{17}$ (d) None of these
17. The value of c , so that for all real x , the vectors $c x \hat{i} - 6 \hat{j} + 3 \hat{k}$, $x \hat{i} + 2 \hat{j} + 2 c x \hat{k}$ make an obtuse angle, are
 (a) $c < 0$ (b) $0 < c < \frac{4}{3}$
 (c) $-\frac{4}{3} < c < 0$ (d) $c > 0$
18. The solution of the equation $y - x \frac{dy}{dx} = a \left(y^2 + \frac{dy}{dx} \right)$ is
 (a) $y = c(x + a)(1 - ay)$
 (b) $y = c(x + a)(1 + ay)$
 (c) $y = c(x - a)(1 + ay)$
 (d) None of the above
19. The order of the differential equation whose general solution is given by $y = (c_1 + c_2)\cos(x + c_3) - c_4 e^{x+c_5}$, where c_1, c_2, c_3, c_4, c_5 are arbitrary constants, is
 (a) 4 (b) 3 (c) 2 (d) 5
20. If $f(x) = x^3 + bx^2 + cx + d$ and $0 < b^2 < c$, then in $(-\infty, \infty)$
 (a) $f(x)$ is strictly increasing function
 (b) $f(x)$ has a local maxima
 (c) $f(x)$ strictly decreasing function
 (d) $f(x)$ is bounded

21. $\frac{d}{dx} \sin^{-1}(x\sqrt{1-x} + \sqrt{x}\sqrt{1-x^2})$
 (a) $-\frac{1}{2x\sqrt{1-x}} - \frac{1}{\sqrt{1-x^2}}$
 (b) $\frac{1}{2\sqrt{x}\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}}$
 (c) $\frac{1}{2\sqrt{x}\sqrt{1-x}} + \frac{1}{\sqrt{1-x^2}}$
 (d) $-\frac{1}{2\sqrt{x}\sqrt{1-x}} + \frac{1}{\sqrt{1-x^2}}$
22. $\int \frac{x \tan^{-1} x}{\sqrt{1+x^2}} dx$ is equal to
 (a) $\frac{x - \tan^{-1} x}{1-x^2} + c$ (b) $\frac{x + \tan^{-1} x}{\sqrt{1-x^2}} + c$
 (c) $\frac{x - \tan^{-1} x}{\sqrt{1+x^2}} + c$ (d) $\frac{x + \sqrt{1-x^2}}{\sqrt{1+x^2}} + c$
23. Let $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$, where $0 \leq \theta \leq 2\pi$. Then, the range of $|A|$ is
 (a) 0
 (b) $\{2, 4\}$
 (c) $[2, 4]$
 (d) None of the above
24. If $y = |\cos x| + |\sin x|$, then $\frac{dy}{dx}$ at $x = \frac{2\pi}{3}$ is
 (a) 0 (b) 1
 (c) $\frac{1-\sqrt{3}}{2}$ (d) $\frac{\sqrt{3}-1}{2}$
25. $\lim_{n \rightarrow \infty} \left(\frac{1}{1-n^2} + \frac{2}{1-n^2} + \dots + \frac{n}{1-n^2} \right)$ is equal to
 (a) 0 (b) $-\frac{1}{2}$
 (c) $\frac{1}{2}$ (d) None of these
26. Let $f(x) = \begin{cases} \sin x, & x \neq n\pi \\ 2, & x = n\pi \end{cases}$, where $n \in I$ and $g(x) = \begin{cases} x^2 + 1, & x \neq 2 \\ 3, & x = 2 \end{cases}$, then $\lim_{x \rightarrow 0} g[f(x)]$ is
 (a) 1 (b) 0
 (c) 3 (d) does not exist
27. The intercept made by the tangent to the curve $y = \int_0^x |t| dt$, which is parallel to the line $y = 2x$, on x-axis is equal to
 (a) 1 (b) -2
 (c) 2 (d) None of these
28. If $P = x^3 - \frac{1}{x^3}$ and $Q = x - \frac{1}{x}$, $x \in (0, \infty)$, then minimum value of $\frac{P}{Q^2}$ is
 (a) $2\sqrt{3}$ (b) $-2\sqrt{3}$
 (c) does not exist (d) None of these
29. Locus of the point which divides double ordinate of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ in the ratio 1:2 internally, is
 (a) $\frac{x^2}{a^2} - \frac{9y^2}{b^2} = \frac{1}{9}$ (b) $\frac{x^2}{a^2} + \frac{9y^2}{b^2} = 1$
 (c) $\frac{9y^2}{a^2} + \frac{9y^2}{b^2} = 1$ (d) None of these
30. From any point on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ tangents are drawn to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$. The area cut-off by the chord of contact on the asymptotes is equal to
 (a) $\frac{ab}{2}$ (b) ab
 (c) $2ab$ (d) $4ab$
31. The value of the sum of the series $3 \cdot {}^nC_0 - 8 \cdot {}^nC_1 + 13 \cdot {}^nC_2 - 18 \cdot {}^nC_3 + \dots$ upto $(n+1)$ terms is
 (a) 0 (b) 3^n
 (c) 5^n (d) None of these
32. If A is a skew-symmetric matrix, then trace of A is
 (a) 1 (b) -1
 (c) 0 (d) None of these
33. The arbitrary constant on which the value of the determinant

$$\begin{vmatrix} 1 & \alpha & \alpha^2 \\ \cos(p-d)a & \cos pa & \cos(p-d)a \\ \sin(p-d)a & \sin pa & \sin(p-d)a \end{vmatrix}$$

 does not depend, is
 (a) α (b) p (c) d (d) a

34. The sum of the first n terms of the series $\frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots$ is equal to
 (a) $2^n - n + 1$ (b) $1 - 2^{-n}$
 (c) $n + 2^{-n} - 1$ (d) $2^n - 1$
35. The base of a cliff is circular. From the extremities of a diameter of the base angles of elevation of the top of the cliff are 30° and 60° . If the height of the cliff be 500 m, then the diameter of the base of the cliff is
 (a) $\frac{2000}{\sqrt{3}}$ m (b) $\frac{1000}{\sqrt{3}}$ m
 (c) $\frac{2000}{\sqrt{2}}$ m (d) $1000\sqrt{3}$ m
36. The most general solutions of the equation $\sec x - 1 = (\sqrt{2} - 1) \tan x$ are given by
 (a) $n\pi + \frac{\pi}{8}$ (b) $2n\pi, 2n\pi + \frac{\pi}{4}$
 (c) $2n\pi$ (d) None of these
37. The maximum value of $\sin\left(x + \frac{\pi}{6}\right) + \cos\left(x + \frac{\pi}{6}\right)$ in the interval $\left[0, \frac{\pi}{2}\right]$ is attained at
 (a) $x = \frac{\pi}{12}$ (b) $x = \frac{\pi}{6}$
 (c) $x = \frac{\pi}{3}$ (d) $x = \frac{\pi}{2}$
38. If $z_r = \cos \frac{r\alpha}{n^2} + i \sin \frac{r\alpha}{n^2}$, where $r = 1, 2, 3, \dots, n$, then $\lim_{n \rightarrow \infty} z_1 z_2 \dots z_n$ is equal to
 (a) $\cos \alpha + i \sin \alpha$
 (b) $\cos\left(\frac{\alpha}{2}\right) - i \sin\left(\frac{\alpha}{2}\right)$
 (c) $e^{i\alpha/2}$
 (d) $\sqrt[3]{e^{i\alpha}}$
39. Negation of "Paris is in France and London is in England" is
 (a) Paris is in England and London is in France
 (b) Paris is not in France or London is not in England
 (c) Paris is in England or London is in France
 (d) None of the above
40. The area enclosed between the curves $y = x^3$ and $y = \sqrt{x}$ is
 (a) $\frac{5}{3}$ sq unit (b) $\frac{5}{4}$ sq unit
 (c) $\frac{5}{12}$ sq unit (d) $\frac{12}{5}$ sq unit
41. Find the equation of the bisector of the obtuse angle between the lines $3x - 4y + 7 = 0$ and $-12x - 5y + 2 = 0$,
 (a) $21x + 77y - 101 = 0$
 (b) $99x - 27y + 81 = 0$
 (c) $21x - 77y + 101 = 0$
 (d) None of the above
42. The equation of curve passing through the point $\left(1, \frac{\pi}{4}\right)$ and having slope of tangent at any point (x, y) as $\frac{y}{x} - \cos^2\left(\frac{y}{x}\right)$, is
 (a) $x = e^{1 + \tan\left(\frac{y}{x}\right)}$ (b) $x = e^{1 - \tan\left(\frac{y}{x}\right)}$
 (c) $x = e^{1 + \tan\left(\frac{x}{y}\right)}$ (d) $x = e^{1 - \tan\left(\frac{x}{y}\right)}$
43. If $P(n) : 2 + 4 + 6 + \dots + (2n), n \in N$, then $P(k) = k(k+1) + 2$ implies
 $P(k+1) = (k+1)(k+2) + 2$
 is true for all $k \in N$. So, statement $P(n) = n(n+1) + 2$ is true for
 (a) $n \geq 1$ (b) $n \geq 2$
 (c) $n \geq 3$ (d) None of these
44. The differential equation of all non-vertical lines in a plane is
 (a) $\frac{d^2y}{dx^2} = 0$ (b) $\frac{d^2x}{dy^2} = 0$
 (c) $\frac{dy}{dx} = 0$ (d) $\frac{dx}{dy} = 0$
45. The unit vector in ZOY plane and making angle 45° and 60° respectively with $\vec{a} = 2\hat{i} + 2\hat{j} - \hat{k}$ and $\vec{b} = 0\hat{i} + \hat{j} - \hat{k}$, is
 (a) $-\frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{k}$
 (b) $\frac{1}{\sqrt{2}}\hat{i} - \frac{1}{\sqrt{2}}\hat{k}$
 (c) $\frac{1}{3\sqrt{2}}\hat{i} + \frac{4}{3\sqrt{2}}\hat{j} + \frac{1}{3\sqrt{2}}\hat{k}$
 (d) None of the above

46. If $\int_2^e \left(\frac{1}{\log x} - \frac{1}{(\log x)^2} \right) dx = a + \frac{b}{\log 2}$, then
 (a) $a = e, b = -2$ (b) $a = e, b = 2$
 (c) $a = -e, b = 2$ (d) None of these
47. The circle $x^2 + y^2 - 4x - 4y + 4 = 0$ is inscribed in a triangle which has two of its sides along the coordinate axes. If the locus of the circumcentre of the triangle is $x + y - xy + k\sqrt{x^2 + y^2} = 0$, then the value of k is equal to
 (a) 2 (b) 1 (c) -2 (d) 3
48. The points of discontinuity of $\tan x$ are
 (a) $n\pi, n \in I$
 (b) $2n\pi, n \in I$
 (c) $(2n + 1)\frac{\pi}{2}, n \in I$
 (d) None of the above
49. The two curves $x^3 - 3xy^2 + 2 = 0$ and $3x^2y - y^3 - 2 = 0$
 (a) cut at right angles
 (b) touch each other
 (c) cut at an angle $\frac{\pi}{3}$
 (d) cut at an angle $\frac{\pi}{4}$
50. The period of the function $f(x) = \frac{\sin 8x \cos x - \sin 6x \cos 3x}{\cos 2x \cos x - \sin 3x \sin 4x}$ is
 (a) π (b) 2π
 (c) $\frac{\pi}{2}$ (d) None of these
51. The derivative of $f(\tan x)$ w.r.t. $g(\sec x)$ at $x = \frac{\pi}{4}$, where $f'(1) = 2$ and $g'(\sqrt{2}) = 4$, is
 (a) $\frac{1}{\sqrt{2}}$ (b) $\sqrt{2}$
 (c) 1 (d) None of these
52. If $a > 0, b > 0$ the maximum area of the triangle formed by the points $O(0, 0)$, $A(a \cos \theta, b \sin \theta)$ and $B(a \cos \theta, -b \sin \theta)$ is (in sq unit)
 (a) $\frac{ab}{2}$ when $\theta = \frac{\pi}{4}$
 (b) $\frac{3ab}{4}$ when $\theta = \frac{\pi}{4}$
 (c) $\frac{ab}{2}$ when $\theta = -\frac{\pi}{2}$
 (d) a^2b^2
53. If the two curves $y = a^x$ and $y = b^x$ intersect at an angle α , then $\tan \alpha$ equals
 (a) $\frac{\log a - \log b}{1 + \log a \log b}$ (b) $\frac{\log a + \log b}{1 - \log a \log b}$
 (c) $\frac{\log a - \log b}{1 - \log a \log b}$ (d) None of these
54. The number of roots of the equation $x - \frac{2}{x-1} = 1 - \frac{2}{x-1}$ is
 (a) 1 (b) 2
 (c) 0 (d) infinitely many
55. The vector $z = -4 + 5i$ is turned counterclockwise through an angle of 180° and stretched $1\frac{1}{2}$ times. The complex number corresponding to newly obtained vector is
 (a) $-6 + \frac{15}{2}i$ (b) $6 + \frac{15}{2}i$
 (c) $6 - \frac{15}{2}i$ (d) None of these
56. If $A = [a_{ij}]$ is a 4×4 matrix C_{ij} is the cofactor of the element a_{ij} in $|A|$, then the expression $a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} + a_{14}C_{14}$ is equal to
 (a) 0 (b) -1
 (c) 1 (d) $|A|$
57. If $A = \left\{ x : \frac{\pi}{6} \leq x \leq \frac{\pi}{3} \right\}$ and $f(x) = \cos x - x(1+x)$, then $f(A)$ is equal to
 (a) $\left[-\frac{\pi}{3}, -\frac{\pi}{6} \right]$
 (b) $\left[\frac{\pi}{6}, \frac{\pi}{3} \right]$
 (c) $\left[\frac{1}{2} - \frac{\pi}{3} \left(1 + \frac{\pi}{3} \right), \frac{\sqrt{3}}{2} - \frac{\pi}{6} \left(1 + \frac{\pi}{6} \right) \right]$
 (d) $\left[\frac{1}{2} + \frac{\pi}{3} \left(1 - \frac{\pi}{3} \right), \frac{\sqrt{3}}{2} + \frac{\pi}{6} \left(1 - \frac{\pi}{6} \right) \right]$


58. The contrapositive of $(p \vee q) \Rightarrow r$ is
 (a) $\sim r \Rightarrow (p \vee q)$
 (b) $r \Rightarrow (p \vee q)$
 (c) $\sim r \Rightarrow (\sim p \wedge \sim q)$
 (d) $p \Rightarrow (q \vee r)$
59. If the function $f(x) = ax^3 + bx^2 + 11x - 6$ satisfies the condition of Rolle's theorem in $[1, 3]$ and $f'\left(2 + \frac{1}{\sqrt{3}}\right) = 0$, then the values of a, b are respectively
 (a) $-1, 6$ (b) $-2, 1$
 (c) $1, -6$ (d) $-1, \frac{1}{2}$
60. If $f(x) = \cos(\log x)$, then

$$f\left(\frac{1}{x}\right)f\left(\frac{1}{y}\right) - \frac{1}{2}\left[f\left(\frac{x}{y}\right) + f(xy)\right]$$

 is equal to
 (a) $\cos(x - y)$ (b) $\log(x - y)$
 (c) $\cos(x + y)$ (d) None of these
61. If $\alpha = \sin^{-1} \frac{\sqrt{3}}{2} + \sin^{-1} \frac{1}{3}$ and $\beta = \cos^{-1} \frac{\sqrt{3}}{2} + \cos^{-1} \frac{1}{3}$, then
 (a) $\alpha > \beta$ (b) $\alpha = \beta$
 (c) $\alpha < \beta$ (d) $\alpha + \beta = 2\pi$
62. If $1 + \sin \theta + \sin^2 \theta + \dots \infty = 4 + 2\sqrt{3}, 0 < \theta < \pi$, $\theta \neq \frac{\pi}{2}$, then
 (a) $\theta = \frac{\pi}{3}$ (b) $\theta = \frac{\pi}{6}$
 (c) $\theta = \frac{\pi}{3}$ or $\frac{\pi}{6}$ (d) $\theta = \frac{\pi}{3}$ or $\frac{2\pi}{3}$
63. A round balloon of radius r subtends an angle α at the eye of the observer, while the angle of elevation of its centre is β . The height of the centre of balloon is
 (a) $r \operatorname{cosec} \alpha \sin \frac{\beta}{2}$ (b) $r \sin \alpha \operatorname{cosec} \frac{\beta}{2}$
 (c) $r \sin \frac{\alpha}{2} \operatorname{cosec} \beta$ (d) $r \operatorname{cosec} \frac{\alpha}{2} \sin \beta$
64. In $\triangle ABC$, $(a - b)^2 \cos^2 \frac{C}{2} + (a + b)^2 \sin^2 \frac{C}{2}$ is equal to
 (a) a^2 (b) b^2
 (c) c^2 (d) None of these

65. In a triangle $\left(1 - \frac{r_1}{r_2}\right)\left(1 - \frac{r_1}{r_3}\right) = 2$, then the triangle is
 (a) right angled (b) isosceles
 (c) equilateral (d) None of these
66. If $a + b + c = 0$, then the roots of the equation $4ax^2 + 3bx + 2c = 0$ are
 (a) equal (b) imaginary
 (c) real (d) None of these

67. The adjoining graph



- (a) connected
 (b) disconnected
 (c) Neither connected nor disconnected
 (d) None of the above
68. The solution of $\tan^{-1} x + 2 \cot^{-1} x = \frac{2\pi}{3}$ is
 (a) $-\frac{1}{\sqrt{3}}$ (b) $\frac{1}{\sqrt{3}}$
 (c) $-\sqrt{3}$ (d) $\sqrt{3}$
69. The conjugate of the complex number $\frac{(1+i)^2}{1-i}$ is
 (a) $1 - i$ (b) $1 + i$
 (c) $-1 + i$ (d) $-1 - i$
70. A graph G has ' m ' vertices of odd degree and ' n ' vertices of even degree. Then which of the following statements is necessarily true?
 (a) $m + n$ is an odd number
 (b) $m + n$ is an even number
 (c) $n + 1$ is an even number
 (d) $m + 1$ is an odd number
71. The value of $\sin \left[2 \cos^{-1} \frac{\sqrt{5}}{3} \right]$ is
 (a) $\frac{\sqrt{5}}{3}$ (b) $\frac{2\sqrt{5}}{3}$
 (c) $\frac{4\sqrt{5}}{9}$ (d) $\frac{2\sqrt{5}}{9}$

72. In the group (G, \otimes_{15}) , where $G = \{3, 6, 9, 12\}$, \otimes_{15} is multiplication modulo 15, the identity element is
 (a) 3 (b) 6 (c) 12 (d) 9
73. A group $(G, *)$ has 10 elements. The minimum number of elements of G , which are their own inverses is
 (a) 2 (b) 1 (c) 9 (d) 0
74. $\frac{3x^2 + 1}{x^2 - 6x + 8}$ is equal to
 (a) $3 + \frac{49}{2(x-4)} - \frac{13}{2(x-2)}$
 (b) $\frac{49}{2(x-4)} - \frac{13}{2(x-2)}$
 (c) $\frac{-49}{2(x-4)} + \frac{13}{2(x-2)}$
 (d) $\frac{49}{2(x-4)} + \frac{13}{2(x-2)}$
75. The orthocentre of the triangle with vertices $O(0, 0)$, $A\left(0, \frac{3}{2}\right)$, $B(-5, 0)$ is
 (a) $\left(\frac{5}{2}, \frac{3}{4}\right)$ (b) $\left(-\frac{5}{2}, \frac{3}{4}\right)$
 (c) $\left(-5, \frac{3}{2}\right)$ (d) $(0, 0)$
76. The range in which $y = -x^2 + 6x - 3$ is increasing, is
 (a) $x < 3$ (b) $x > 3$
 (c) $7 < x < 8$ (d) $5 < x < 6$
77. The area bounded by the curve $x = 4 - y^2$ and the y-axis is
 (a) 16 sq unit
 (b) 32 sq unit
 (c) $\frac{32}{3}$ sq unit
 (d) $\frac{16}{3}$ sq unit
78. The number of positive divisors of 252 is
 (a) 9 (b) 5 (c) 18 (d) 10
79. The remainder obtained when 5^{124} is divided by 124 is
 (a) 5 (b) 0
 (c) 2 (d) 1
80. Which of the following is not a group with respect to the given operation ?
 (a) The set of even integers including zero under addition
 (b) The set of odd integers under addition
 (c) $\{0\}$ under addition
 (d) $\{1, -1\}$ under multiplication

General English and Aptitude

Directions (Q. 1-4) : In each of the following questions, choose the most suitable alternative to fill in the blank.

- He is so of his own idea that he will not entertain any suggestion from others.
 (a) hopeful (b) enamoured
 (c) jealous (d) possessed
- Undoubtedly, English is the most spoken language in the world today.
 (a) broadly
 (b) widely
 (c) greatly
 (d) beautifully
- I will be leaving for Delhi tonight and to return by this weekend.
 (a) waiting (b) plan
 (c) going (d) making

- The vacancy by the dismissal of the superintendent is expected to be filled up by the promotion of a U.D.C.
 (a) made (b) created
 (c) caused (d) generated

Directions (Q. 5-8) : In each of the following questions, choose the alternative which best expresses the meaning of the word given in capital letters.

- HAGGLE
 (a) Postpone (b) Accept
 (c) Bargain (d) Reject
- ABSTRUSE
 (a) Awful (b) Irrelevant
 (c) Shallow (d) Profound
- PENCHANT
 (a) Like (b) Eagerness
 (c) Disability (d) Dislike

8. BARTER
(a) Deal (b) Return
(c) Lend (d) Exchange

Directions (Q. 9-12) : In each of the following questions, choose the alternative which can be substituted for the given words/sentence.

9. Bringing about gentle and painless death from incurable disease
(a) Gallows (b) Suicide
(c) Euphoria (d) Euthanasia
10. The act of killing one's wife
(a) Avicide
(b) Canicide
(c) Uxoricide
(d) Genocide
11. Stage between boyhood and youth
(a) Infancy (b) Adolescence
(c) Puberty (d) Maturity
12. Lack of enough blood
(a) Amnesia
(b) Insomnia
(c) Anaemia
(d) Allergy

Directions (Q. 13-16) : In each of the following questions, choose the alternative which best expresses the meaning of the given idiom/phrase.

13. To set the people by ears
(a) To box the people
(b) To insult and disgrace the people
(c) To punish heavily
(d) To excite people to a quarrel
14. To give chapter and verse for a thing
(a) To produce the proof of something
(b) To eulogise the qualities of a thing
(c) To make publicity of a thing
(d) To attach artificial value to a thing
15. Dog in the manger
(a) An undersized bull almost the shape of a dog
(b) A dog that has no kennel of its own
(c) A person who puts himself in difficulties on account of other people
(d) A person who prevents others from enjoying something useless to himself

16. To blow hot and cold
(a) Changing weather
(b) To be untrustworthy
(c) To change opinion often
(d) To be rich and poor frequently

Directions (Q. 17-20) : In each of the following questions, choose the alternative which is opposite in meaning to the word given in capital letters.

17. SAGACIOUS
(a) Casual (b) Cunning
(c) Foolish (d) False
18. EPILOGUE
(a) Conversation (b) Dialogue
(c) Dramatic (d) Prologue
19. AUSPICIOUS
(a) Spicy
(b) Unfavourable
(c) Conspicuous
(d) Condemnatory
20. ENGULFED
(a) Encircled (b) Groped
(c) Disfigured (d) Detached
21. Four of the following five are alike in a certain way and so form a group. Which is the one that does not belong to that group?
(a) Hill (b) Valley
(c) Dam (d) River
22. In a certain code 'CREAM' is written as 'NBDBQ'. How is 'BREAD' written in that code?
(a) EBFAQ
(b) EBDAQ
(c) BEDQA
(d) BEFQA
23. If black means white, white means red, red means yellow, yellow means blue, blue means green, green means purple and purple means orange, then what is the colour of lemon?
(a) Green (b) Purple
(c) Orange (d) Blue

Directions (Q. 24-25) : In the following questions, find the word which holds the same relation with the third word as there is between the first two word.

24. Hot : Oven :: Cold : ?

- (a) Ice cream
- (b) Air conditioner
- (c) Snow
- (d) Refrigerator

25. Push : Pull :: Throw : ?

- (a) Jump
- (b) Collect
- (c) Pick
- (d) Game

Directions (Q. 26-27) In each of the following questions, one letter or a set of letter is missing, you have to understand the pattern of the series and insert the appropriate letter?

26. R, M, ?, F, D, ?

- (a) C, B
- (b) J, H
- (c) H, C
- (d) I, C

27. - bcc - ac - aabb - ab - cc

- (a) aab ca
- (b) aba ca
- (c) ba cab
- (d) bca ca.

28. How many 9's are there in the following number series which are immediately preceded by 3 and followed by 6?

3 9 6 9 3 9 3 9 3 9 6 3 9 3 6 3 9 5 6 9 5 6 9 3 9 6 3 9

- (a) 0
- (b) 3
- (c) 2
- (d) 4

29. Some boys are sitting in a row, P is sitting fourteenth from the left and Q is seventh from the right. If there are four boys between P and Q, how many boys are there in the row?

- (a) 25
- (b) 23
- (c) 21
- (d) 19

30. Pointing to a photograph, a woman says, "this man's son's sister in my mother-in-law." How is the woman's husband related to the man in the photograph?

- (a) Grandson
- (b) Son
- (c) Son-in-law
- (d) Nephew.

31. The longest canal in the world is

- (a) Volga Baltic
- (b) Beloye-more Baltic
- (c) Suez Canal
- (d) Grand China Canal

32. The oldest Hindu epic is

- (a) Mahabhashya
- (b) Ramayan
- (c) Mahabharata
- (d) Ashtadhyayi

33. Who among the following is not associated with the Swaraj Party?

- (a) C.R. Das
- (b) M.L. Kelkar
- (c) Motilal Nehru
- (d) Mahatma Gandhi

34. Where is the 'City of palaces'?

- (a) London
- (b) Kolkata
- (c) Patiala
- (d) Lucknow

35. Our National Song is

- (a) Sare Jahan Se Achcha
- (b) Jana Gana Mana
- (c) Vande Mataram
- (d) All of the above

36. The Constitution of India was adopted by the Constituent Assembly on

- (a) Dec 11, 1946
- (b) Aug 15, 1957
- (c) Nov 26, 1949
- (d) Jan 26, 1949

37. Gandhiji's Dandi March started from

- (a) Bardoli
- (b) Ahmedabad
- (c) Surat
- (d) Bombay

38. Who was the viceroy of India at the time of formation of the Indian National Congress?

- (a) Lord Canning
- (b) Lord Dufferin
- (c) Lord Mayo
- (d) Lord Elgin

39. Which country was a major donor in financing the 'SAARC'?

- (a) Pakistan
- (b) Sri Lanka
- (c) India
- (d) Bangladesh

40. Where in the H. Q. of the European Economic Community?

- (a) Bonn
- (b) Rome
- (c) Brussels
- (d) Hague

Answers

➡ PHYSICS

1. (a)	2. (b)	3. (a)	4. (d)	5. (a)	6. (c)	7. (c)	8. (a)	9. (c)	10. (c)
11. (c)	12. (a)	13. (b)	14. (d)	15. (c)	16. (c)	17. (a)	18. (b)	19. (b)	20. (b)
21. (c)	22. (b)	23. (b)	24. (a)	25. (c)	26. (d)	27. (b)	28. (d)	29. (a)	30. (b)
31. (a)	32. (b)	33. (a)	34. (a)	35. (a)	36. (b)	37. (c)	38. (c)	39. (c)	40. (a)
41. (d)	42. (c)	43. (a)	44. (a)	45. (b)	46. (a)	47. (c)	48. (d)	49. (c)	50. (a)
51. (c)	52. (c)	53. (c)	54. (b)	55. (b)	56. (c)	57. (d)	58. (b)	59. (b)	60. (b)

➡ CHEMISTRY

1. (c)	2. (a)	3. (b)	4. (c)	5. (a)	6. (c)	7. (b)	8. (a)	9. (b)	10. (c)
11. (d)	12. (d)	13. (c)	14. (d)	15. (a)	16. (a)	17. (a)	18. (d)	19. (c)	20. (d)
21. (a)	22. (a)	23. (c)	24. (c)	25. (a)	26. (d)	27. (b)	28. (a)	29. (c)	30. (c)
31. (d)	32. (c)	33. (b)	34. (c)	35. (c)	36. (c)	37. (d)	38. (c)	39. (b)	40. (b)
41. (b)	42. (c)	43. (a)	44. (a)	45. (a)	46. (a)	47. (c)	48. (b)	49. (a)	50. (c)
51. (c)	52. (c)	53. (c)	54. (d)	55. (d)	56. (c)	57. (d)	58. (d)	59. (b)	60. (c)

➡ MATHEMATICS

1. (a)	2. (a)	3. (c)	4. (a)	5. (c)	6. (c)	7. (b)	8. (c)	9. (a)	10. (c)
11. (b)	12. (c)	13. (d)	14. (a)	15. (b)	16. (b)	17. (c)	18. (a)	19. (b)	20. (a)
21. (c)	22. (c)	23. (c)	24. (d)	25. (b)	26. (a)	27. (a)	28. (c)	29. (b)	30. (d)
31. (a)	32. (c)	33. (b)	34. (c)	35. (a)	36. (b)	37. (a)	38. (c)	39. (b)	40. (c)
41. (a)	42. (b)	43. (d)	44. (a)	45. (b)	46. (a)	47. (b)	48. (c)	49. (a)	50. (c)
51. (a)	52. (a)	53. (a)	54. (c)	55. (c)	56. (d)	57. (c)	58. (c)	59. (c)	60. (d)
61. (c)	62. (d)	63. (d)	64. (c)	65. (a)	66. (c)	67. (a)	68. (d)	69. (d)	70. (b)
71. (c)	72. (b)	73. (b)	74. (a)	75. (d)	76. (a)	77. (c)	78. (c)	79. (a)	80. (b)

➡ GENERAL ENGLISH AND APTITUDE

1. (b)	2. (b)	3. (b)	4. (c)	5. (c)	6. (d)	7. (a)	8. (d)	9. (d)	10. (c)
11. (b)	12. (c)	13. (d)	14. (a)	15. (d)	16. (c)	17. (c)	18. (d)	19. (b)	20. (d)
21. (c)	22. (b)	23. (d)	24. (d)	25. (b)	26. (d)	27. (c)	28. (b)	29. (a)	30. (a)
31. (b)	32. (b)	33. (d)	34. (b)	35. (c)	36. (c)	37. (b)	38. (b)	39. (b)	40. (c)

Hints & Explanations

Physics

1. Surface tension (T) of a liquid is equal to the work (W) required to increase the surface area (ΔA) of the liquid film by unity at constant temperature.

$$W = T \Delta A = \text{surface energy}$$

Also, volume of big drop = $27 \times$ volume of small drop

$$\text{ie, } V' = 27V$$

where V' is volume of big drop of diameter D and V the volume of small drop of diameter d .

$$\therefore \frac{4}{3} \pi \left(\frac{D}{2} \right)^3 = 27 \times \frac{4}{3} \pi \left(\frac{d}{2} \right)^3$$

$$\Rightarrow \frac{D}{2} = 3 \times \frac{d}{2}$$

$$\Rightarrow d = \frac{D}{3}$$

$$\text{Radius of small drop, } r = \frac{d}{2} = \frac{D}{6}$$

$$\therefore \text{Change in surface energy} = T (A_2 - A_1)$$

$$= T [27 \cdot 4\pi r^2 - 4\pi R^2]$$

$$= T 4\pi \left[27 \left(\frac{D}{6} \right)^2 - \left(\frac{D}{2} \right)^2 \right]$$

$$= 4\pi T \left[\frac{3D^2}{4} - \frac{D^2}{4} \right] = 2\pi D^2 T$$

2. From Kepler's third law of planetary motion. The square of the period of revolution (T) of any planet around the sun is directly proportional to the cube of the semi major axis (R) of its elliptical orbit.

$$\text{ie, } T^2 \propto R^3$$

$$\text{Given, } T_p = 27T_e$$

$$\frac{T_e^2}{T_p^2} = \frac{R_e^3}{R_p^3}$$

$$\frac{T_e^2}{T_p^2} = \frac{R_e^3}{R_p^3}$$


$$\frac{T_e^2}{(27T_e)^2} = \frac{R_e^3}{R_p^3}$$

$$\frac{R_p}{R_e} = (27)^{2/3}$$

$$\frac{R_p}{R_e} = 3^2$$

$$R_p = 9R_e$$

3. The net force acting on a unit mass placed at O due to three equal masses M at vertices A, B and C is the gravitational field intensity at point O . The gravitational force on the particle placed at the point of intersection of three medians.




$$\vec{F}_1 + \vec{F}_2 + \vec{F}_3 = 0$$

Since, the resultant of \vec{F}_1 and \vec{F}_2 is equal and opposite to \vec{F}_3 .

4. In a turbulent flow, the velocity of the liquid, in contact with the walls of the tube is equal to critical velocity.
5. Potential due to charge (q) at point (r) is given by

$$V = \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{r}$$




Since, charge Q is rotated in a circle of radius r , hence its potential remains same at all points on the path, hence $\Delta V = 0$.

Also, work done $= q\Delta V$

where q is charge and $\Delta V = 0$.

\therefore Work done $= 0$.

6. The circuit diagram of the given situation is shown. Since, it is a closed mesh, the voltage is



Given, $V' = 0.5$ volt, $I = 0.1$ A, $R = 20\Omega$

$\therefore V = 0.5 + 0.1 \times 20$

$V = 2.5$ volt

7. A dipole placed in an external electric field is acted upon by a torque which tends to align the dipole in the direction of the field. Therefore, work must be done to change the orientation of the dipole against the torque. If dipole be rotated from an initial orientation $\theta = \theta_1$ to final orientation $\theta = \theta_2$, the total work required is

$$W = \int_{\theta_1}^{\theta_2} pE \sin \theta d\theta$$

$$W = pE [-\cos \theta]_{\theta_1}^{\theta_2}$$

where p is dipole moment and E the electric field.

In first case,

$$W = pE (1 - \cos 60^\circ)$$

$$W = pE \left(1 - \frac{1}{2}\right) = \frac{pE}{2}$$


$$\Rightarrow pE = 2W$$

In second case,

$$W_2 = pE (1 - \cos 180^\circ)$$

$$W_2 = 2W (1 + 1) = 4W$$

8. Magnetic moment or magnetic dipole moment is a measure of the strength of a magnetic source. It is given by



$$\mu = IA \quad \dots(i)$$

where I is current and A the area.

The effective current (I) is $I = \frac{e}{T}$

where e is electron charge and T the time period.

Also, $\nu = \frac{1}{T}$ = frequency

and area $A = \pi r^2$, where r is radius of circular path.


Putting these values in Eq. (i), we get

$$\mu = e\nu \cdot \pi r^2$$

Note : Magnetic moment can also be expressed terms of angular momentum (L)

$$\mu = IA = -\frac{2}{2M_e} L, \text{ where } M_e \text{ is mass of electron.}$$

9. From Coulomb's law, the force of attraction between two charged particles (q), kept at distance r apart is



when two identical spheres are brought in contact, charge on them is equalised, hence total charge on C is equally shared when brought in contact with sphere B having a charge q . Therefore, charge on B and C is $\frac{q}{2}$.

From Coulomb's law, the force on C is

$$F_C = \frac{q \times q/2}{4\pi\epsilon_0(r/2)^2} - \frac{(q/2)(q/2)}{4\pi\epsilon_0(r/2)^2}$$

$$= \frac{qq}{4\pi\epsilon_0 r^2} (2 - 1) = F$$

Note : The force will be opposite because A and B spheres will repel the third sphere.

10. The standard equation of wave is

$$y = a \sin(\omega t - kx) \quad \dots(i)$$

where a is amplitude, ω the angular velocity and x the displacement at instant t .

Given equation is

$$y = 0.1 \sin(100\pi t - kx) \quad \dots(ii)$$


Comparing Eq. (i) with Eq. (ii), we get

$$\omega = 100\pi$$

$$\therefore \text{Wave number} = \frac{\omega}{v} = \frac{100\pi}{100} = \pi \text{ m}^{-1}$$

11. At constant pressure, the volume of a given mass of a gas is directly proportional to its absolute temperature (T).

ie, $\frac{V}{T} = \text{constant.}$



This is another form of Charles' law. Hence, variation of volume with temperature is as follows.

Hence, correct graph will be (c).

12. Optical rotation or optical activity is the rotation of linearly polarized light as it travels through certain materials. It occurs in solutions of chiral molecules (eg, sugar), solids with rotated crystal planes (eg, quartz) and spin polarized gases of atoms or molecules. Any linear polarization of light can be written as an equal combination of right hand (RHC) and left hand (LHC) circularly polarized light.

$$E_{\theta_D} = E_{\text{RHC}} + e^{i2\theta_D} E_{\text{LHC}}$$

where E is the electric field of light.

13. The average power or simply power is the average amount of work done or energy transferred per unit time.

The instantaneous power is then the limiting value of the average power as the time interval Δt approaches zero.

$$P = \lim_{\Delta t \rightarrow 0} \frac{\Delta W}{\Delta t}$$

$$\therefore W = \int P dt$$

Given, $P = 3t^2 - 2t + 1$


$$\therefore W = \int_2^4 (3t^2 - 2t + 1) dt$$

Using $\int x^n dx = \frac{x^{n+1}}{n+1}$, we have

$$W = [t^3 - t^2 + t]_2^4 = 56 - 12 + 2$$

$$\Rightarrow W = 46 \text{ J}$$

14. If a constant force \vec{F} is applied on a body for a short interval of time Δt , then the impulse of this force is $F \times \Delta t$.



Since, impulse = change in momentum (Δp)

$$\therefore F \times \Delta t = \Delta p$$

$$\Rightarrow F = \frac{\Delta p}{\Delta t}$$


Change in x-direction

$$F = \frac{m [30 - (-15 \sin 30^\circ)]}{0.01}$$

$$F = \frac{0.1 \times 37.5}{0.01} = 375 \text{ N}$$

Note : Unit of impulse is same as that of momentum.

15. From Coulomb's law, the force of attraction/repulsion acting between two stationary point charges (q_1, q_2) separated by a distance (r) is



$$F = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2}$$

Taking the net force, we have

$$F_{\text{net}} = 2F \cos \theta = 2 \left[\frac{1}{4\pi\epsilon_0} \frac{Qqx}{\left(x^2 + \frac{d^2}{4}\right)^{3/2}} \right]$$

For maximum,

$$\frac{dF_{\text{net}}}{dx} = 0$$


$$\therefore \left(x^2 + \frac{d^2}{4}\right)^{3/2} - \frac{3}{2}x \left(x^2 + \frac{d^2}{4}\right)^{1/2} (2x) = 0$$

$$\therefore \left(x^2 + \frac{d^2}{4}\right)^{1/2} \left(x^2 + \frac{d^2}{4} - 3x^2\right) = 0$$

$$\text{or } 2x^2 = \frac{d^2}{4}$$

$$\Rightarrow x = \frac{d}{2\sqrt{2}}$$

16. Let x be the equivalent resistance of entire network between A and B. Hence, we have



$R_{AB} = 1 + \text{resistance of parallel combination of } 1\Omega \text{ and } x\Omega$

$$\therefore R_{AB} = 1 + \frac{x}{1+x}$$


$$\therefore x = 1 + \frac{x}{1+x}$$

$$\Rightarrow x + x^2 = 1 + x + x$$

$$\Rightarrow x^2 - x - 1 = 0$$

$$\Rightarrow x = \frac{1 + \sqrt{1+4}}{2} = \frac{1 + \sqrt{5}}{2} \Omega$$

17. For a circular coil of radius a carrying a current i , the magnetic field at point P , distance x from coil is given by



$$B = \frac{\mu_0 i a^2}{2(a^2 + x^2)^{3/2}} \text{ NA}^{-1} \text{ m}^{-1} \quad \dots(i)$$

At the centre of coil $x = 0$

$$\therefore B' = \frac{\mu_0 i}{2a} \text{ NA}^{-1} \text{ m}^{-1} \quad \dots(ii)$$

$$\text{Given, } B = \frac{1}{8} B'$$

$$\therefore \frac{\mu_0 i a^2}{2(a^2 + x^2)^{3/2}} = \frac{1}{8} \left(\frac{\mu_0 i}{2a} \right)$$

$$\Rightarrow \frac{a^2}{(a^2 + x^2)^{3/2}} = \frac{1}{8a}$$

$$\Rightarrow 8a^3 = (a^2 + x^2)^{3/2}$$


$$\Rightarrow a^2 + x^2 = 4a^2$$

$$\Rightarrow x = \sqrt{3} \cdot a$$

Given, $a = R$

$$\therefore x = \sqrt{3} R$$

18. Let S be the light source. If light falls on the surface at critical angle C , it grazes along the surface as shown.



$$\sin C = \frac{1}{n} = \frac{3}{5}$$

From $\triangle QSR$, we have

$$\tan C = \frac{QR}{QS} = \frac{r}{4}$$

$$\Rightarrow \frac{3}{4} = \frac{r}{4}$$

$$\Rightarrow r = 3$$

Hence, diameter $= 2r = 2 \times 3 = 6 \text{ m}$

19. Critical angle (θ_c) is the angle of incidence (i) in the denser medium for which the angle of refraction in the rarer medium is 90° . For total internal reflection to take place

$$i > \theta_c$$

Taking sine on both sides, we get


$$\sin i > \sin \theta_c$$

[as angle i at both faces will be 45°]

$$\Rightarrow \frac{1}{\sqrt{2}} > \frac{1}{\mu}$$

$$\Rightarrow \mu > \sqrt{2}$$

20. When an object is placed in front of such a lens, the rays are first of all refracted from the convex surface, then reflected from the polished plane surface and again refracted from convex surface.



Let f_i, f_m be focal lengths of convex surface and mirror (plane polished surface) respectively, then effective focal length is

$$\frac{1}{F} = \frac{1}{f_i} + \frac{1}{f_m} + \frac{1}{f_i} = \frac{2}{f_i} + \frac{1}{f_m}$$

$$\text{Since, } f_m = \frac{R}{2} = \infty$$

$$\therefore \frac{1}{F} = \frac{2}{f_i}$$

From lens formula

$$\frac{1}{f_i} = (\mu - 1) \left(\frac{1}{R} \right)$$

$$\therefore \frac{1}{F} = \frac{2(\mu - 1)}{R}$$

$$\Rightarrow F = \frac{R}{2(\mu - 1)}$$

$$\text{or } R_{\text{eq}} = 2F = \frac{R}{(\mu - 1)}$$

21. From Kirchhoff's second law

$$V = \sum ir \quad (\text{for closed mesh})$$

where V is potential difference, i the current and r the resistance.

$$\therefore E + E = Ir + Ir = 2Ir$$

$$\text{or } I = \frac{E}{r} \quad \dots(i)$$

$$V_x - V_y = E - Ir$$

Putting the value of I from Eq. (i), we get

$$V_x - V_y = E - \frac{E}{r} \times V = 0.$$

22. Induced emf is given by

$$e = -\frac{d\phi}{dt}$$

If the radius of loop is r at a time t , then the instantaneous magnetic flux is given by

$$\phi = \pi r^2 B$$

$$\therefore e = -\frac{d}{dt} (\pi r^2 B)$$

$$e = -\pi B \left(\frac{2r}{dt} \frac{dr}{dt} \right)$$

$$e = -2\pi B r \frac{dr}{dt}$$

$$\text{Numerically, } e = 2\pi B r \left(\frac{dr}{dt} \right)$$

23. The minimum energy needed to ionise an atom is called ionisation energy. The potential difference through which an electron should be accelerated to acquire this much energy is called ionisation potential.

$$(E_2)_H - (E_1)_H = 10.2 \text{ eV}$$

$$\text{or } \frac{(E_1)_H}{4} - (E_1)_H = 10.2 \text{ eV}$$

$$\therefore (E_1)_H = -13.6 \text{ eV}$$

Hence, ionisation potential energy is

$$= (E_\infty)_H - (E_1)_H = 13.6 \text{ eV}$$

$$\therefore \text{Ionisation potential} = 13.6 \text{ V}$$

24. Since, train (source) is moving towards pedestrain (observer), the perceived frequency will be higher than the original.

$$f' = f \left(\frac{v + v_o}{v - v_s} \right)$$

Here, $v_o = 0$ (as observer is stationary)

$v_s = 25 \text{ m/s}$ (velocity of source)

$v = 350 \text{ m/s}$ (velocity of sound)

and $f = 1 \text{ kHz}$ (original frequency)

$$\text{Hence, } f' = 1000 \left(\frac{350 + 0}{350 - 25} \right)$$

$$= 1000 \times \frac{350}{325}$$

$$= 1077 \text{ Hz}$$

25. For two coherent sources, the resultant intensity is given by

$$I = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \phi$$

For maximum intensity, $\cos \phi = +1$

$$\therefore I_{\max} = I_1 + I_2 + 2\sqrt{I_1 I_2}$$

$$= (\sqrt{I_1} + \sqrt{I_2})^2$$

For minimum intensity, $\cos \phi = -1$

$$\therefore I_{\min} = I_1 + I_2 - 2\sqrt{I_1 I_2}$$

$$= (\sqrt{I_1} - \sqrt{I_2})^2$$

$$\text{Hence, } I_{\max} = (\sqrt{9I} + \sqrt{I})^2$$


$$= (3\sqrt{I} + \sqrt{I})^2 = 16I$$

$$\text{and } I_{\min} = (\sqrt{9I} - \sqrt{I})^2$$

$$= (3\sqrt{I} - \sqrt{I})^2 = 4I$$

Note : In an interference pattern, maximum intensity is obtained for constructive interference and minimum intensity is obtained for destructive interference.

26. An optical fibre is a device based on total internal reflection by which a light signal can be transferred from one place to the other with a negligible loss of energy.



It consists of a very long and thin fibre of quartz glass (refractive index = 1.7). Each fibre is of thickness nearly 10^{-4} cm . The fibre is coated all around by a thin layer of a material (refractive index = 1.5) rarer than the quartz glass.

When a light ray is incident at one end A of the fibre making a small angle of incidence. It suffers refraction from air to quartz and strikes the fibre-layer interface at an angle of incidence greater than the critical angle. It therefore, suffers total internal reflection and strikes its opposite interface. At this interface, it again suffers total internal reflection. Thus, the ray reaches the other end B of the fibre.

27. Bernoulli's theorem for unit mass of liquid

$$\frac{p}{\rho} + \frac{1}{2} v^2 = \text{constant}$$

As the liquid starts flowing, its pressure energy decreases

$$\frac{1}{2} v^2 = \frac{p_1 - p_2}{\rho} \Rightarrow \frac{1}{2} v^2 = \frac{3.5 \times 10^5 - 3 \times 10^5}{10^3}$$

$$\Rightarrow v^2 = \frac{2 \times 0.5 \times 10^5}{10^3} \Rightarrow v^2 = 100$$

$$\Rightarrow v = 10 \text{ m/s}$$

28. $a_1 v_1 = a_2 v_2$

$$\Rightarrow 4.20 \times 5.18 = 7.60 \times v_2$$

$$\Rightarrow v_2 = 2.86 \text{ m/s}$$

29. de-Broglie wavelength of a particle is given by

$$\lambda = \frac{h}{mv} \quad \dots(i)$$

where h is Planck's constant.

If kinetic energy of particle of mass m is ν , then

$$K = \frac{1}{2} mv^2$$

$$\Rightarrow v = \sqrt{\frac{2K}{m}} \quad \dots(ii)$$

Combining Eqs. (i) and (ii), we get

$$\lambda = \frac{h}{m\sqrt{\frac{2K}{m}}} = \frac{h}{\sqrt{2mK}} \quad \dots(iii)$$

Given : $m = 9.1 \times 10^{-31}$ kg,

$$K = 10 \text{ keV} = 10 \times 10^3 \times 1.6 \times 10^{-19} \text{ J}$$

$$h = 6.6 \times 10^{-34} \text{ J-s}$$

Substituting the above values in Eq. (iii), we get

$$\begin{aligned} \lambda &= \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 9.1 \times 10^{-31} \times 10 \times 10^3 \times 1.6 \times 10^{-19}}} \\ &= 1.22 \times 10^{-11} \\ &\approx 0.12 \text{ \AA} \end{aligned}$$

Note : If an electron is accelerated through a potential difference of V volt, then Eq. (iii) takes the form

$$\lambda = \frac{h}{\sqrt{2meV}}$$

After putting the numerical values for electrons, we get

$$\lambda = \sqrt{\frac{150}{V}} \text{ \AA}$$

30. The given equations of waves be written as

$$y_1 = 0.25 \sin(310t) \quad \dots(i)$$

$$\text{and } y_2 = 0.25 \sin(316t) \quad \dots(ii)$$

Comparing Eqs. (i) and (ii) with the standard wave equation, written as

$$y = a \sin(\omega t) \quad \dots(iii)$$

We have,

$$\omega_1 = 310$$

$$\Rightarrow f_1 = \frac{310}{2\pi} \text{ unit}$$

$$\text{and } \omega_2 = 316$$

$$\Rightarrow f_2 = \frac{316}{2\pi} \text{ unit}$$

Hence, beat frequency

$$\begin{aligned} &= f_2 - f_1 \\ &= \frac{316}{2\pi} - \frac{310}{2\pi} \\ &= \frac{3}{\pi} \text{ unit} \end{aligned}$$

31. The activity or decay rate R of a radioactive substance is the number of decays per second.

$$\therefore R = \lambda N$$

$$\text{or } R = \lambda N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$$

$$\text{or } R = R_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$$

where $R_0 = \lambda N_0$ is the activity of the radioactive substance at time $t = 0$.

According to question

$$\frac{R}{R_0} = 1 - \frac{75}{100} = 25\%$$

$$\therefore \frac{25}{100} = \left(\frac{1}{2}\right)^{t/T_{1/2}}$$


$$\text{or } \left(\frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^{t/T_{1/2}}$$

$$\text{or } \frac{t}{T_{1/2}} = 2$$

$$\therefore t = 2T_{1/2} = 2 \times 3.20 = 6.40 \text{ h}$$

$$\text{or } t \approx 6.38 \text{ h}$$

32. Initially, the capacitance of capacitor



$$C = \frac{\epsilon_0 A}{d}$$

$$\therefore \frac{\epsilon_0 A}{d} = 1 \mu\text{F} \quad \dots(i)$$

When it is filled with dielectrics of dielectric constants K_1 and K_2 as shown, then there are two capacitors connected in parallel. So,

$$C' = \frac{K_1 \epsilon_0 (A/2)}{d} + \frac{K_2 \epsilon_0 (A/2)}{d}$$

(as area becomes half)


$$C' = \frac{4\epsilon_0 A}{2d} + \frac{6\epsilon_0 A}{2d}$$

$$= \frac{2\epsilon_0 A}{d} + \frac{3\epsilon_0 A}{d}$$

Using Eq. (i), we obtain

$$C' = 2 \times 1 + 3 \times 1 = 5 \mu\text{F}$$

33. The given combination can be represented as



All the gates in the combination are NAND gates.

Output of gate-1, $Y_1 = \overline{A \cdot A}$

Output of gate-2, $Y_2 = \overline{B \cdot B}$

Output of gate-3, $Y = \overline{Y_1 \cdot Y_2}$

$$\therefore Y = \overline{\overline{A \cdot A} \cdot \overline{B \cdot B}} \quad \dots(i)$$

By Demorgan's theorem, we have

$$\overline{A \cdot A} = \overline{A}$$

$$\text{and } \overline{B \cdot B} = \overline{B}$$

Therefore, Eq. (i) becomes,

$$Y = \overline{\overline{A} \cdot \overline{B}}$$

Again from Demorgan's theorem

$$\overline{\overline{A} \cdot \overline{B}} = \overline{\overline{A}} + \overline{\overline{B}}$$

$$\therefore Y = \overline{\overline{A}} + \overline{\overline{B}} = A + B$$

(as $\overline{\overline{A}} = A$)

Hence, the combination behaves as OR gate.

34. In a junction transistor, both the electron and hole play role, hence they are called the bi-polar devices or the bi-polar transistors and they are abbreviated as BJT in short form.


There are three parts in a transistor; namely emitter, base and collector. In the emitter part of the transistor the doping is more and it is less in collector part.

The doping is very less in the base part. So, emitter current is actually the sum of base and collector current.


$$\text{i.e., } I_E = I_B + I_C$$

Hence, maximum current flows in emitter region.

Note : Figure shown, represents the transistor available in the market. For identifying the electrodes, the collector terminal is generally kept far separated as compared to the remaining two electrodes or a red dot is etched near the collector terminal.



35. The energy band scheme of semiconductors is shown here.



In semiconductors, valence band and conduction band are separated by an energy gap called the forbidden energy gap. It is very small. At room temperature some electrons in valence band acquire thermal energy. This energy is more than forbidden energy gap E_g , thus they jump into the conduction band and leaves their vacancy in the valence band which act as holes.

Hence, at room temperature valence band is partially empty and conduction band is partially filled.

36. When coil is open, there is no current in it, hence no flux associated with it, i.e., $\phi = 0$.

Also, we know that flux linked with the coil is directly proportional to the current in the coil, i.e., $\phi \propto i$

$$\text{or } \phi = Li$$

where L is proportionality constant known as self-inductance.

$$\therefore L = \frac{\phi}{i} = 0$$

Again since $i = 0$, hence, $R = \infty$.

37. If a current i is passed in the circuit and it is changed with a rate $\frac{di}{dt}$, the induced emf e produced in the circuit is directly proportional to the rate of change of current.

Thus,

$$e \propto \frac{di}{dt}$$

When the proportionality constant is removed, the constant L comes here.

$$\therefore e = -L \frac{di}{dt} \quad \dots(i)$$


The minus sign here is a reflection of Lenz's law.

Here, $di = (2 - 10)A = -8A$, $dt = 0.1 \text{ s}$, $e = 3.28 \text{ V}$.

$$\therefore 3.28 = -\frac{L(-8)}{0.1}$$

$$\therefore L = \frac{3.28 \times 0.1}{8} = 0.04 \text{ H}$$

38. When rod is bent in the form of square, then each side has resistance of $\frac{1}{4} \Omega$. As shown R_1, R_2 and R_3 are connected in series, so their equivalent resistance



$$R' = R_1 + R_2 + R_3 \\ = \frac{1}{4} + \frac{1}{4} + \frac{1}{4} = \frac{3}{4} \Omega$$

Now, R' and R_4 are connected in parallel, so equivalent resistance of the circuit is

$$R = \frac{R' \times R_4}{R' + R_4} \\ = \frac{(3/4)(1/4)}{(3/4) + (1/4)} \\ = \frac{(3/16)}{1} = \frac{3}{16} \Omega$$

$$39. \tan \phi = \frac{\omega L - \frac{1}{\omega C}}{R}$$

ϕ being the angle by which the current leads the voltage.

Given, $\phi = 45^\circ$

$$\therefore \tan 45^\circ = \frac{\omega L - \frac{1}{\omega C}}{R}$$

$$\Rightarrow 1 = \frac{\omega L - \frac{1}{\omega C}}{R}$$


$$\Rightarrow R = \omega L - \frac{1}{\omega C}$$

$$\Rightarrow C = \frac{1}{\omega(\omega L - R)} \\ = \frac{1}{2\pi f(2\pi fL - R)}$$

Note : In series resonant L-C-R circuit, $\frac{1}{\omega CR}$ is greater than unity.

40. The imaginary surface joining the points of same potential in an electric field is called the equipotential surface, i.e., the potential difference between any two points on an equipotential surface is zero. Hence, if a charge is moved on an equipotential surface from one point to the other, no work is needed to be done. But this is possible only when the charge is moved perpendicular to the electric field (i.e., perpendicular to the lines of force). It means that the electric lines of force at

each point of an equipotential surface are normal to the surface.



Hence, the angle between electric field and equipotential surface is 90° .

Alternative : Potential gradient along equipotential surface is zero.

$$\text{ie, } E \cos \theta = -\frac{dV}{dr} = 0$$

$$\therefore \theta = 90^\circ$$

41. According to Newton's second law of motion, force acting on a body is equal to the rate of change of momentum during impact.

$$F = \frac{\Delta p}{\Delta t}$$

Also, $F = ma$

$$\therefore ma = \frac{p_2 - p_1}{\Delta t}$$

$$\text{or } a = \frac{mv_2 - (-mv_1)}{m \Delta t}$$

$$\text{or } a = \frac{v_2 + v_1}{\Delta t}$$

$$\therefore a = \frac{\sqrt{2 \times 10 \times 20} + \sqrt{2 \times 10 \times 5}}{0.02}$$

$$\text{or } a = \frac{20 + 10}{0.02} = 1500 \text{ m/s}^2$$

42. Volume remains constant after coalescing.

Thus,

$$\frac{4}{3} \pi R^3 = 2 \times \frac{4}{3} \pi r^3$$

where R is radius of bigger drop and r is radius of each smaller drop.

$$\therefore R = 2^{1/3} r$$

Now, surface energy per unit surface area is the surface tension.

So, surface energy, $W = T \Delta A$

$$\text{or } W = 4\pi R^2 T$$

Therefore, surface energy of bigger drop

$$W_1 = 4\pi (2^{1/3} r)^2 T$$

$$= (2^{2/3}) 4\pi r^2 T$$

Surface energy of smaller drop

$$W_2 = 4\pi r^2 T$$

Hence, required ratio

$$\frac{W_1}{W_2} = 2^{2/3} : 1$$

43. In fission process, when a parent nucleus breaks into daughter products, then some mass is lost in the form of energy. Thus, mass of fission products < mass of parent nucleus

$$\Rightarrow \frac{\text{Mass of fission products}}{\text{Mass of parent nucleus}} < 1$$

44. Width of central maximum is given by

$$w = \frac{2f\lambda}{a} \quad \dots(i)$$

where f is focal length of lens, a is width of slit and λ is wavelength of light used.

From Eq. (i), it is clear that fringe width

$$w \propto \lambda$$

So, when blue light is used in the experiment instead of red light, the fringes will become narrower.

45. If object in a denser medium is seen from a rarer medium then image of object will appear at a lesser distance. The distance between object and its image, called as normal shift is given by

$$x = t \left[1 - \frac{1}{\mu} \right]$$

$$\text{Here, } t = 6 \text{ cm, } \mu = 1.5$$

$$\therefore x = 6 \left[1 - \frac{1}{1.5} \right]$$

$$= 6 \left[\frac{0.5}{1.5} \right]$$

$$= 2 \text{ cm}$$

46. When plates of capacitor are separated by a dielectric medium of dielectric constant K , its capacity

$$C_m = \frac{K\epsilon_0 A}{d} = KC_0$$

ie, $C_m = KC_0$

Here, $C_0 = C$

$\therefore C_m = KC$

Now, two capacitors of capacities KC and C are in series, their effective capacitance


$$\frac{1}{C'} = \frac{1}{KC} + \frac{1}{C}$$

or $\frac{1}{C'} = \frac{1+K}{KC}$

$\therefore C' = \frac{KC}{K+1}$

47. When a person of mass m is placed on a weighing machine which is placed in a lift, then actual weight of the person is mg .

This acts on a weighing machine which offers a reaction R given by the reading of weighing machine. This reaction exerted by the surface of contact on the person is the apparent weight of the person.



When lift is accelerating upwards at the rate of a , then reaction

$$R - mg = ma$$

$\therefore R = m(g + a)$

Hence, apparent weight is greater than actual weight.

48. Due to high resistance of voltmeter, connected in series, the effective resistance of circuit will increase and hence the current in circuit will decrease. Due to which the ammeter and voltmeter will not be damaged.

49. $T = 100 \mu\text{s} = 10^{-4} \text{ s}$

$$\lambda = \frac{0.6931}{T} = \frac{0.6931}{10^{-4}} = 0.6931 \times 10^4 \text{ s}^{-1}$$

Number of atoms in 215 mg

$$= \frac{6.023 \times 10^{23}}{215} \times 215 \times 10^{-3}$$

$\therefore N = 6.023 \times 10^{20}$

Activity, $\frac{dN}{dt} = \lambda N$
 $= 0.6931 \times 10^4 \times 6.023 \times 10^{20}$
 $= 4.17 \times 10^{24} \text{ Bq}$

50. For maximum wavelength of Balmer series

$$\frac{1}{\lambda_{\max}} = R \left(\frac{1}{2^2} - \frac{1}{3^2} \right) = \frac{R \times 5}{36} \quad \dots(i)$$

For minimum wavelength of Balmer series,

$$\frac{1}{\lambda_{\min}} = R \left(\frac{1}{2^2} - \frac{1}{\infty} \right) = \frac{R}{4} \quad \dots(ii)$$

From Eqs. (i) and (ii), we have

$\therefore \frac{\lambda_{\min}}{\lambda_{\max}} = \frac{R \times 5}{36} \times \frac{4}{R} = \frac{5}{9}$

51. When the ball is released from the top of tower then ratio of distances covered by the ball in first, second and third second

$$h_I : h_{II} : h_{III} = 1 : 3 : 5 \text{ [because } h_n \propto (2n-1)]$$

\therefore Ratio of work done

$$mgh_I : mgh_{II} : mgh_{III} = 1 : 3 : 5$$

52. $m_1 v_1 - m_2 v_2 = (m_1 + m_2)v$

$$\Rightarrow 2 \times 3 - 1 \times 4 = (2 + 1)v$$

$$\Rightarrow v = \frac{2}{3} \text{ m/s}$$

53. Total energy radiated from a body

$$Q = A\epsilon\sigma T^4 t$$

$$\Rightarrow Q \propto AT^4 \propto r^2 T^4 \quad (\because A = 4\pi r^2)$$

$$\Rightarrow \frac{Q_p}{Q_q} = \left(\frac{r_p}{r_q} \right)^2 \left(\frac{T_p}{T_q} \right)^4$$

$$= \left(\frac{8}{2} \right)^2 \left[\frac{(273 + 127)}{(273 + 527)} \right]^4 = 1$$

54. According to Newton's law of cooling t_1 will be less than t_2 .

55. If an electron and a photon propagates in the form of waves having the same wavelength, it implies that they have same momentum. This is according to de-Broglie equation

$$p \propto \frac{1}{\lambda}$$

$$\begin{aligned} 56. \lambda_0 &= \frac{h_C}{W_0} = \frac{12400}{4} \\ &= 3100 \text{ \AA} \\ &= 310 \text{ nm} \end{aligned}$$

57. Threshold wavelength

$$\lambda_0 = \frac{12375}{2.1} = 5892.8 \text{ \AA}$$

58. Electric flux


$$\phi = \frac{\text{pulse power}}{\text{area}} = \frac{10^{12}}{10^{-4}} = 10^{16} \text{ W/cm}^2$$

59. A laser device produces amplification in the ultraviolet or visible region.



60. $\frac{Q}{t} \propto \frac{r^2}{l}$, from the given options, option (b) has higher value of $\frac{r^2}{l}$.

Chemistry


1. Molecules or ions having no unpaired electrons are diamagnetic, eg,



2. NO_2^- can participate in linkage isomerism because it may be bonded to metal either through nitrogen or through oxygen.




3. **Reimer-Tiemann reaction** : In this reaction phenol reacts with chloroform and alkali to form salicylaldehyde.



4. Benzoic acid in benzene exist as a dimer. So, number of molecules decreases and hence, osmotic pressure decreases.

5. Mohr's salt is $(\text{NH}_4)_2\text{SO}_4 \cdot \text{FeSO}_4 \cdot 6\text{H}_2\text{O}$. The equation is



Total change in oxidation number of iron

$$= (+3) - (+2) = +1$$

So, equivalent wt. of Mohr's salt

$$\begin{aligned} &= \frac{\text{mol. wt. of Mohr's salt}}{\text{change in oxidation number}} \\ &= \frac{392}{1} = 392 \end{aligned}$$

6. A salt is precipitated only when the product of ionic concentration is more than its solubility product.

$$K_{sp} = 1 \times 10^{-8}$$


$$[A^+] = 10^{-3} \text{ M}$$

$$[B^-] = \frac{1 \times 10^{-8}}{10^{-3}} = 10^{-5} \text{ M}$$

So, AB will be precipitated only when the concentration of B^- is more than 10^{-5} M .

7. Weight of 11.2 dm^3 of CO_2 gas at STP


$$= 44 / 2 = 22 \text{ g}$$




\therefore KOH required for complete neutralisation of CO_2 (44 g) = 56 g

\therefore KOH required for neutralisation of 22 g of $\text{CO}_2 = \frac{56}{44} \times 22 = 28$ g

8. $-\text{NH}_2$ has +R effect, it donates electrons to the benzene ring. As a result, the lone pair of electrons on the N-atom get delocalized over the benzene ring and thus less readily available for protonation. Hence, aniline is a weaker base than cyclohexylamine.



9. Fats are also known as triglycerides. These triglycerides are the triesters of fatty acid with glycerol. So, the characteristic feature of fat is ester group.
10. Compounds having asymmetric C-atom (C) is optically active, eg,





The C-atom whose four valencies are satisfied by four different groups is asymmetric C-atom.


11. In cyclopropane the angle strain is maximum. Hence, it is highly strained molecule and consequently most unstable. The angle strain in cyclobutane is less than cyclopropane. Hence, cyclobutane is more stable. This stability increases upto 6 membered rings, then decreases from 7 to 11 membered rings and from the 12 membered rings onwards attain the stability of 6 membered ring. Heat of combustion is a method of measuring chemical stability. Hence, cyclohexane has the lowest heat of combustion.

12. The physical states of dispersing phase and dispersion medium in colloid like pesticide spray are liquid and gas respectively.

13. Potassium dichromate is an oxidising agent it oxidises ferrous ions into ferric ions in acid medium.




14. Sucrose is obtained from cane sugar. On hydrolysis, it gives equimolar mixture of D (+) glucose and D (-) fructose.







In sucrose, glucose is present in the pyranose form and fructose in the furanose form. Further since $\text{C}_1 - \alpha$ of glucose is connected to $\text{C}_2 - \beta$ of fructose, therefore sucrose is a non-reducing sugar.

15. During inductive effect shifting of σ electrons takes place due to which partial charges are developed on the atoms.





16. Electronic configuration of




So, the given configuration is of Cu^+ .

17. (i) Electronic configuration of Co^{3+} ion




F^- is a weak ligand. It cannot pair up electrons within d -subshell and forms outer orbital octahedral complex.


(ii) $[Co(NH_3)_6]^{3+}$ ion



(iii) $[Fe(CN)_6]^{3-}$ ion



(iv) $[Cr(NH_3)_6]^{3+}$ ion



NH_3 and CN^- are strong ligands. So, they form inner orbital complex.

18. The first ionisation energy of xenon is quite close to that of oxygen and the molecular diameter of xenon and oxygen are almost identical.

Based on the above facts it is suggested that as oxygen combines with PtF_6 , so xenon should also form similar compound with PtF_6 .

19. We know that,

$$pV = nRT$$

or
$$pV = \frac{w}{M} RT$$

or
$$M = \frac{w}{V} \frac{RT}{p}$$

or
$$M = d \frac{RT}{p}$$

$$d = 1.964 \text{ g/dm}^3 = 1.964 \times 10^{-3} \text{ g/cc.}$$

$$p = 76 \text{ cm} = 1 \text{ atm}$$

$$R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$= 82.1 \text{ cc atm K}^{-1} \text{ mol}^{-1}$$



$$T = 273 \text{ K}$$

$$M = \frac{1.964 \times 10^{-3} \times 82.1 \times 273}{1} = 44.$$



The molecular weight of CO_2 is 44.

So, the gas is CO_2 .

20.




[where, M = highly reactive metal]




ΔG° vs T plot in the Ellingham's diagram slopes downwards for the reaction (I) and reaction (II). For reaction (III) ΔG° vs T in the diagram slopes upwards.

21.



22. According to electron sea model, in a metallic crystal a lattice of positive ions, called kernels, remains immersed in a sea of mobile valence electrons. These electrons move freely within the boundaries of metallic crystal.

23. The electronic configuration of peroxide ion (O_2^{2-}) is as :



$$\text{Bond order} = \frac{8 - 6}{2} = 1$$

Since, all the electrons in molecular orbitals are paired, it is diamagnetic.

24. Insulin regulates the metabolism of carbohydrates (glucose).

25. Flocculation value $\propto \frac{1}{\text{coagulating power}}$

Fe(OH)_3 is a positively charged sol.



To coagulate Fe(OH)_3 , negatively charged electrolyte is used and greater the value of negative charge, coagulating power will be strong. Among the given electrolytes, NaCl has lowest coagulating power, so its flocculation value will be maximum.

$$\begin{aligned} 26. \quad k &= \frac{2.303}{t} \log \frac{A_0}{A} \\ &= \frac{2.303}{40} \log \frac{0.1}{0.005} \\ &= \frac{2.303}{40} \log 20 = 0.075 \end{aligned}$$

Rate of reaction when concentration of X is 0.01 M will be

$$0.075 \times 0.01 = 7.5 \times 10^{-4} \text{ M min}^{-1}$$

27. At pH = 4, an amphoteric Zwitter ion structure changes into cation when an acid is added to it.



28. Bond breaking processes or decomposition processes are endothermic process.

$$29. \quad E_{\text{Cu}^{2+}/\text{Cu}}^\circ - E_{\text{Ag}^+/\text{Ag}}^\circ = -0.46 \text{ V} \quad \dots (i)$$


$$E_{\text{Cu}^{2+}/\text{Cu}}^\circ - E_{\text{Zn}^{2+}/\text{Zn}}^\circ = 1.10 \text{ V} \quad \dots (ii)$$

On subtracting Eq (i) from equation (ii), we get




$$E_{\text{Ag}^+/\text{Ag}}^\circ - E_{\text{Zn}^{2+}/\text{Zn}}^\circ = +1.56 \text{ V}$$

30. Ketone undergoes nucleophilic addition reaction because if nucleophiles attack first followed by electrophiles, anion is formed as intermediate and if electrophiles attack first

followed by nucleophiles, cation, formed as intermediate. The stability of anion is more than cation, so nucleophiles attack first followed by electrophiles.




So, probable mechanism will be



32. The presence of unpaired electron in phosphorus atom is explained by Hund's rule, which states that "Pairing of electrons in *d* and *f*-orbitals can't occur until each orbital of a given subshell contains one electron or is singly occupied"

$${}_{15}\text{P} : 1s^2, 2s^2, 2p^6, 3s^2, 3p_x^1, 3p_y^1, 3p_z^1$$



33. Given, $m = 200 \text{ g}$, $v = 3 \times 10^3 \text{ cm/s}$,

$$h = 6.624 \times 10^{-27} \text{ erg s}, \lambda = ?$$

$$\lambda = \frac{h}{mv}$$

$$= \frac{6.624 \times 10^{-27}}{200 \times 3 \times 10^3}$$

$$\therefore \lambda = 1.104 \times 10^{-32} \text{ cm}$$

34. Cubic close stacking is $a-b-c-a-b-c \dots$ pattern of stacking spheres. NaCl has face centred cubic arrangement which is also called cubic close stacking arrangement.


35. Equivalent weight of bivalent metal = 37.2

$$\therefore \text{Atomic weight of metal} = 37.2 \times 2 = 74.4$$

\therefore Formula of its chloride is MCl_2

Hence, molecular weight of MCl_2

$$= 74.4 + 2 \times 35.5 = 145.4$$



37.
$$\frac{\Delta T_f \times 1000 \times w}{M \times W} = \frac{k_f \times 1000 \times w}{M \times W}$$

$$w = \frac{\Delta T_f \times M \times W}{k_f \times 1000}$$

$$\Delta T_f = 0 - (-0.6) = 0.6^\circ \text{C}$$

$$M = 60 \text{ g/mol}$$

$$k_f = 1.5^\circ \text{C kg mol}^{-1}$$

$$W = 3 \times 10^3 \text{ g}$$

$$w = \frac{0.6 \times 60 \times 3 \times 10^3}{1.5 \times 1000}$$

$$= 72 \text{ g}$$

38. If $K < 1$, $\Delta G^\circ > 0$, hence the value of ΔG° is positive if $K < 1$.

39. Molecular weight of $(\text{COOH})_2 \cdot 2\text{H}_2\text{O} = 126 \text{ g}$

$$\therefore \text{Equivalent weight} = \frac{126}{2} = 63$$

[\therefore basicity = 2]

Weight of $(\text{COOH})_2 \cdot 2\text{H}_2\text{O}$ in solution

$$= 32.5 \text{ g}$$

$$\text{Normality} = \frac{\text{number of gram equivalent}}{\text{volume (in litre)}}$$

$$= \frac{32.5 / 63}{0.5}$$

$$= 1 \text{ N}$$

40. In acidic medium phenolphthalein is colourless and in alkaline medium it is pink. Therefore, in the titration of strong base (KOH) vs weak acid (oxalic acid) phenolphthalein is used.

41.
$$\frac{r_+}{r_-} = \frac{95}{181} = 0.524$$

\therefore The radius ratio lies between 0.414 to 0.732.



\therefore The coordination number of Na^+ is 6.




42. van der Waals' equation for one mole of a gas is

$$\left(p + \frac{a}{V^2} \right) (V - b) = RT$$


where, b = volume correction, which arises due to finite size of molecules.

43. The α -hydrogen atoms of acetaldehyde due to $-E$ effect of >C=O group is slightly acidic in nature. In crossed aldol condensation between formaldehyde and acetaldehyde, in the first step OH^- ion (from the base added) abstracts one of these acidic α -hydrogen to form carbanion or enolate ion which is stabilised by resonance.





45. Phenol is more acidic than alcohol because phenoxide ion is stabilised by resonance.



phenoxide ion

resonance structure of phenoxide ion of phenol


46. Citric acid is found in lemon, therefore lemon gives sour taste.
47. Due to common ion effect, rate of ionisation of NH_4OH decreases, so lower $[\text{OH}^-]$ is obtained. Hence, pH value decreases.
48. The electronic configuration of ${}_{24}\text{Cr}$ is $=[\text{Ar}]_{18} 3d^5, 4s^1$ Chromium after loss of one electron gain stable configuration due to presence of half filled d orbitals, therefore its second ionisation enthalpy is highest.
49. The metals having higher negative value of standard reduction potential are placed above hydrogen in electrochemical series. The metals placed above hydrogen has a great tendency to donate electrons or to undergo oxidation. The metals having great power to undergo oxidation are strongest reducing agent. Zn has highest negative value of standard reduction potential. Therefore, it is the strongest reducing agent.
50. $(\text{CH}_3)_3\text{CBr} + \text{H}_2\text{O} \longrightarrow (\text{CH}_3)_3\text{C} - \text{OH} + \text{HBr}$

Br is substituted by $-\text{OH}^-$ (nucleophile)





$\text{S}_\text{N}1$ (unimolecular nucleophilic substitution reaction)

51. Rate of a reaction = $\frac{\text{change in concentration}}{\text{time}}$

\therefore units of rate = $\frac{\text{mol / L}}{\text{s}} = \text{mol L}^{-1} \text{s}^{-1}$




53. Fehling solution is only reduced by aldehydes but not by ketones, therefore it is used to distinguish between aldehyde ($\text{C}_2\text{H}_5\text{CHO}$) and ketone ($(\text{CH}_3)_2\text{CO}$).



\therefore Three molecules of phenyl hydrazine are utilised in the formation of glucosazone.

55. The structure of CrO_5 is as



The structure shows that four oxygen atom in CrO_5 are involve in the formation of peroxide linkage (OS of oxygen = - 1) while one is present as oxide (OS of oxygen = - 2). Therefore,

CrO_5

$$\begin{array}{ccccccc} x & + & 1 \times & (-2) & + & 4 & (-1) = 0 \\ \text{for Cr} & & \text{for O} & & \text{for (O-O)} & & \\ & & & & x - 2 - 4 = 0 & & \\ & & & & x = + 6 & & \end{array}$$

56. Liquor ammonia bottles are opened only after cooling because it has high vapour pressure and it is mild explosive.
57. According to Faraday's second law of electrolysis "when the same quantity of electricity is passed through solutions of different electrolytes connected in series, the weights of substances produced at the electrodes are directly proportional to their equivalent weights".

If M stands for atomic weight of substances

$$\text{Equivalent weight of Cu} = \frac{M}{2}$$

$$\text{Equivalent weight of Ni} = \frac{M}{2}$$

$$\text{Equivalent weight of Ag} = \frac{M}{1}$$

So, the proportion of moles of metals

$$\frac{M}{2} : \frac{M}{2} : \frac{M}{1}$$

$$\therefore 1 : 1 : 2$$

58. According to question, the rate of reaction is doubled on increasing temperature from T_1 K to T_2 K.

At T_1 , $k_1 = k$ (say)

and T_2 , $k_2 = 2k$

On applying,

$$\log_{10} \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$\log_{10} \frac{2k}{k} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$\log_{10} 2 = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

59. $X(g) + 3Y(g) \rightleftharpoons 2Z(g)$
 $\Delta S^\circ = 2S^\circ(Z) - \{S^\circ(X) + 3S^\circ(Y)\}$
 $= 2 \times 50 - \{60 + 3 \times 40\}$
 $= 100 - 180 = -80 \text{ J K}^{-1} \text{ mol}^{-1}$

$$\Delta G = \Delta H - T\Delta S$$


At equilibrium,

$$\Delta G = 0$$

$$\therefore \Delta H = T\Delta S$$

$$\begin{aligned} T &= \frac{\Delta H}{\Delta S} \\ &= \frac{-40 \times 10^3 \text{ J}}{-80 \text{ JK}^{-1} \text{ mol}^{-1}} \\ &= 500 \text{ K} \end{aligned}$$

60. Those aldehydes or ketones which have α -hydrogen atoms undergo aldol condensation reaction.



Mathematics

$$\begin{aligned}
 1. \text{ Given, } f(x) &= x^2 + \frac{1}{x^2 + 1} \\
 \therefore f(x) &= x^2 + \left(\frac{1}{x^2 + 1} - 1 \right) + 1 \\
 &= (x^2 + 1) - \left(\frac{x^2}{x^2 + 1} \right) \\
 &= 1 + x^2 \left(1 - \frac{1}{x^2 + 1} \right) \\
 &\geq 1, \forall x \in R
 \end{aligned}$$

Hence, range of $f(x)$ is $[1, \infty)$.

$$\begin{aligned}
 2. \text{ Given,} \\
 f(x) &= \begin{cases} ax^2 + b, & b \neq 0, x \leq 1 \\ bx^2 + ax + c, & x > 1 \end{cases} \\
 \Rightarrow f'(x) &= \begin{cases} 2ax, & b \neq 0, x \leq 1 \\ 2bx + a, & x > 1 \end{cases}
 \end{aligned}$$

Since, $f(x)$ is continuous at $x = 1$

$$\begin{aligned}
 \therefore \lim_{x \rightarrow 1^-} f(x) &= \lim_{x \rightarrow 1^+} f(x) \\
 \Rightarrow a + b &= b + a + c \\
 \Rightarrow c &= 0
 \end{aligned}$$

Also, $f(x)$ is differentiable at $x = 1$.

$$\therefore (\text{LHD at } x = 1) = (\text{RHD at } x = 1)$$

$$\begin{aligned}
 \Rightarrow 2a &= 2b(1) + a \\
 \Rightarrow a &= 2b
 \end{aligned}$$

3. Let the equation of the circle be

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

Since, this passes through $(1, 2)$.

$$\therefore 1^2 + 2^2 + 2g(1) + 2f(2) + c = 0$$

$$\Rightarrow 5 + 2g + 4f + c = 0 \quad \dots (i)$$

Also, the circle $x^2 + y^2 = 4$ intersects the circle $x^2 + y^2 + 2gx + 2fy + c = 0$ orthogonally.

$$\therefore 2(g \cdot 0 + f \cdot 0) = c - 4$$

$$\Rightarrow c = 4$$

On putting the value of c in Eq. (i), we get

$$2g + 4f + 9 = 0$$

Hence, the locus of centre $(-g, -f)$ is

$$-2x - 4y + 9 = 0$$

$$\text{or } 2x + 4y - 9 = 0$$

4. Given,

$$\begin{aligned}
 \int f(x) \sin x \cos x \, dx &= \frac{1}{2(b^2 - a^2)} \\
 &\quad \log[f(x)] + c
 \end{aligned}$$

On differentiating both sides, we get

$$f(x) \sin x \cos x = \frac{d}{dx} \left(\frac{\log[f(x)]}{2(b^2 - a^2)} + c \right)$$

$$\Rightarrow f(x) \sin x \cos x = \frac{1}{2(b^2 - a^2)} \cdot \frac{1}{f(x)} f'(x)$$

$$\Rightarrow 2(b^2 - a^2) \sin x \cos x = \frac{f'(x)}{[f(x)]^2}$$

On integrating both sides, we get

$$\begin{aligned}
 \int (2b^2 \sin x \cos x - 2a^2 \sin x \cos x) \, dx \\
 = \int \frac{f'(x)}{[f(x)]^2} \, dx
 \end{aligned}$$

$$\Rightarrow -b^2 \cos^2 x - a^2 \sin^2 x = -\frac{1}{f(x)}$$

$$\therefore f(x) = \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x}$$

5. Given, $|z - 3| = |z - 5|$

On squaring both sides, we get

$$(z - 3)(\bar{z} - 3) = (z - 5)(\bar{z} - 5)$$

$$\Rightarrow z\bar{z} - 3\bar{z} - 3z + 9 = z\bar{z} - 5\bar{z} - 5z + 25$$

$$\Rightarrow 2\bar{z} + 2z = 16 \Rightarrow z + \bar{z} = 8$$

$$\Rightarrow 2x = 8 \Rightarrow x = 4$$

(putting $z = x + iy$)

Hence, locus of z is a straight line parallel to y -axis.

6. Let the roots of the equation be p, q .

$$\text{Let } S = p^2 + q^2$$

$$= (p + q)^2 - 2pq \quad \dots (i)$$

Given equation is

$$x^2 - (\sin \alpha - 2)x - (1 + \sin \alpha) = 0$$

$$\therefore p + q = (\sin \alpha - 2), pq = -(1 + \sin \alpha)$$

From Eq. (i),

$$\begin{aligned} S &= (\sin \alpha - 2)^2 + 2(1 + \sin \alpha) \\ &= \sin^2 \alpha - 4 \sin \alpha + 4 + 2 + 2 \sin \alpha \\ &= \sin^2 \alpha - 2 \sin \alpha + 6 \end{aligned}$$

$$\Rightarrow S = (\sin \alpha - 1)^2 + 5$$

This is least when $\sin \alpha - 1 = 0$

$$\Rightarrow \alpha = \frac{\pi}{2}$$

$$7. \text{ Let } P(n) : 10^{n-2} \geq 81n$$

$$\text{For } n = 4, \quad 10^2 \geq 81 \times 4$$

$$\text{For } n = 5, \quad 10^3 \geq 81 \times 5$$

Hence, by mathematical induction for $n \geq 5$, the proposition is true.

$$8. \text{ General term of } (3 + 2x)^{74} \text{ is}$$

$$T_{r+1} = {}^{74}C_r (3)^{74-r} 2^r x^r$$

Let two consecutive terms are T_{r+1} th and T_{r+2} th terms.

According to the given condition,

$$\text{Coefficient of } T_{r+1} = \text{Coefficient of } T_{r+2}$$

$$\Rightarrow {}^{74}C_r 3^{74-r} 2^r = {}^{74}C_{r+1} 3^{74-(r+1)} 2^{r+1}$$

$$\Rightarrow \frac{{}^{74}C_{r+1}}{{}^{74}C_r} = \frac{3}{2}$$

$$\Rightarrow \frac{74-r}{r+1} = \frac{3}{2}$$

$$\Rightarrow 148 - 2r = 3r + 3$$

$$\Rightarrow r = 29$$

Hence, two consecutive terms are 30 and 31.

$$9. \text{ Now, } 2.\overline{357} = 2 + 0.357357357\dots$$

$$\Rightarrow 2.\overline{357} = 2 + 0.357 + 0.000357 + \dots$$

$$\Rightarrow 2.\overline{357} = 2 + \frac{357}{10^3} + \frac{357}{10^6} + \dots$$

$$\Rightarrow 2.\overline{357} = 2 + \frac{357}{1 - \frac{1}{10^3}}$$

$$= 2 + \frac{357}{999}$$

$$\Rightarrow 2.\overline{357} = \frac{2355}{999}$$

10. Given,

$$S_n = \frac{1}{1^3} + \frac{1+2}{1^3+2^3}$$

$$+ \dots + \frac{1+2+3+\dots+n}{1^3+2^3+3^3+\dots+n^3}$$

$$\text{Now, } T_n = \frac{1+2+3+\dots+n}{1^3+2^3+3^3+\dots+n^3} = \frac{\Sigma n}{\Sigma n^3}$$

$$= \frac{n(n+1)/2}{\{n(n+1)/2\}^2}$$

$$= \frac{2}{n(n+1)}$$

$$= 2 \left(\frac{1}{n} - \frac{1}{n+1} \right)$$

\therefore

$$S_n = T_1 + T_2 + \dots + T_n$$

$$= 2 \left(\frac{1}{1} - \frac{1}{2} \right) + 2 \left(\frac{1}{2} - \frac{1}{3} \right) + \dots$$

$$+ 2 \left(\frac{1}{n} - \frac{1}{n+1} \right)$$

$$= 2 \left(1 - \frac{1}{n+1} \right)$$

$$= 2 - \frac{2}{n+1} \leq 2 \quad \left(\because \frac{2}{n+1} \leq 1 \right)$$

$$11. \text{ Given, } E(\theta) = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$$

$$E(\theta)E(\phi) = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$$

$$\times \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \theta \cos^2 \phi + \cos \theta \sin \theta \cos \phi \sin \phi & \\ \cos \theta \sin \theta \cos^2 \phi + \sin^2 \theta \cos \phi \sin \phi & \\ \cos^2 \theta \cos \phi \sin \phi + \cos \theta \sin \theta \sin^2 \phi & \\ \cos \theta \sin \theta \cos \phi \sin \phi + \sin^2 \theta \sin^2 \phi & \end{bmatrix}$$

$$\begin{aligned}
&= \begin{bmatrix} \cos \theta \cos \phi \cos(\theta - \phi) \\ \cos \phi \sin \theta \cos(\theta - \phi) \\ \cos \theta \sin \phi \cos(\theta - \phi) \\ \sin \theta \sin \phi \cos(\theta - \phi) \end{bmatrix} \\
&= \begin{bmatrix} \cos \theta \cos \phi \cos(2n+1)\frac{\pi}{2} \\ \cos \phi \sin \theta \cos(2n+1)\frac{\pi}{2} \\ \cos \theta \sin \phi \cos(2n+1)\frac{\pi}{2} \\ \sin \theta \sin \phi \cos(2n+1)\frac{\pi}{2} \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \quad \begin{bmatrix} \therefore \cos(2n+1)\frac{\pi}{2} = 0 \end{bmatrix}
\end{aligned}$$

12. Given equation can be rewritten as

$$(y-2)^2 = 12x$$

Here, vertex and foci are (0, 2) and (3, 2).

\therefore Vertex of the required parabola is (3, 2) and focus is (3, 4). The axis of symmetry is $x = 3$ and latusrectum = $4 \cdot 2 = 8$

Hence, required equation is

$$(x-3)^2 = 8(y-2)$$

$$\text{or } x^2 - 6x - 8y + 25 = 0$$

13. Tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at

$(a \cos \theta, b \sin \theta)$ is

$$\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1 \quad \dots (i)$$

$\therefore p$ = perpendicular distance from focus $(ae, 0)$ to the line (i)

$$\begin{aligned}
&= \frac{\left| \frac{ae}{a} \cos \theta + 0 - 1 \right|}{\sqrt{\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2}}} \\
&= \frac{1 - e \cos \theta}{\sqrt{\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2}}} \quad \dots (i)
\end{aligned}$$

Also, p' = perpendicular distance from centre $(0, 0)$ to the line (i)

$$= \frac{1}{\sqrt{\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2}}} \quad \dots (ii)$$

Again, $r = SP = a(1 - e \cos \theta)$


$$\therefore ap = \frac{a - ae \cos \theta}{\sqrt{\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2}}} = rp'$$

14. Let $B(x_1, y_1)$ and $C(x_2, y_2)$ be two vertices and $P\left(\frac{x_1+1}{2}, \frac{y_1-2}{2}\right)$ lies on perpendicular

bisector $x - y + 5 = 0$

$$\therefore \frac{x_1+1}{2} - \frac{y_1-2}{2} = -5$$

$$\Rightarrow x_1 - y_1 = -13 \quad \dots (i)$$



Also, PN is perpendicular to AB

$$\therefore \frac{y_1+2}{x_1-1} \times 1 = -1$$

$$\Rightarrow y_1 + 2 = -x_1 + 1$$

$$\Rightarrow x_1 + y_1 = -1 \quad \dots (ii)$$

On solving Eqs. (i) and (ii), we get

$$x_1 = -7, y_1 = 6$$

\therefore The coordinates of B are $(-7, 6)$. Similarly, the coordinates of C are $\left(\frac{11}{5}, \frac{2}{5}\right)$.

Hence, the equation of BC is

$$y - 6 = \frac{\frac{2}{5} - 6}{\frac{11}{5} + 7} (x + 7)$$

$$\Rightarrow y - 6 = \frac{-14}{23}(x + 7)$$

$$\Rightarrow 14x + 23y - 40 = 0$$

15. Given, $\cos \theta = -\frac{\sqrt{3}}{2} < 0$ and θ does not lie in third quadrant.

$\therefore \theta$ must be lying in 2nd quadrant.

$$\Rightarrow \tan \theta = -\frac{1}{\sqrt{3}}$$

$$\text{and } \cot \theta = -\sqrt{3} \quad \dots (i)$$


Also, α lies in 3rd quadrant and $\sin \alpha = -\frac{3}{5}$

$$\therefore \tan \alpha = \frac{3}{4} \text{ and } \cos \alpha = -\frac{4}{5} \quad \dots (ii)$$

$$\begin{aligned} \therefore \frac{2 \tan \alpha + \sqrt{3} \tan \theta}{\cot^2 \theta + \cos \alpha} &= \frac{2 \cdot \frac{3}{4} - \sqrt{3} \cdot \frac{1}{\sqrt{3}}}{3 - \frac{4}{5}} \\ &= \frac{\frac{3}{2} - 1}{3 - \frac{4}{5}} = \frac{5}{22} \end{aligned}$$

16. Diagonal $\vec{AC} = \vec{a} + \vec{b}$

$$\Rightarrow |\vec{AC}| = |\vec{a} + \vec{b}|$$



$$\Rightarrow |\vec{AC}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b}$$

$$\begin{aligned} \Rightarrow |\vec{AC}|^2 &= \{ |3\vec{\alpha} - \vec{\beta}|^2 + |\vec{\alpha} + 3\vec{\beta}|^2 \\ &\quad + 2(3\vec{\alpha} - \vec{\beta}) \cdot (\vec{\alpha} + 3\vec{\beta}) \} \\ &= 9\vec{\alpha}^2 + \vec{\beta}^2 - 6\vec{\alpha} \cdot \vec{\beta} + \vec{\alpha}^2 + 9\vec{\beta}^2 + 6\vec{\alpha} \cdot \vec{\beta} \\ &\quad + 6\vec{\alpha}^2 - 6\vec{\beta}^2 + 16\vec{\alpha} \cdot \vec{\beta} \end{aligned}$$

$$\Rightarrow |\vec{AC}|^2 = 16\vec{\alpha}^2 + 4\vec{\beta}^2 + 16\vec{\alpha} \cdot \vec{\beta}$$

$$\Rightarrow |\vec{AC}|^2 = 64 + 16 + 16|\vec{\alpha}||\vec{\beta}|\cos \frac{\pi}{3}$$

$$|\vec{AC}|^2 = 80 + 16 \times 4 \times \frac{1}{2} = 112$$

$$\Rightarrow |\vec{AC}| = \sqrt{112} = 4\sqrt{7}$$

Other diagonal is $|\vec{BD}| = |\vec{a} - \vec{b}|$

$$\begin{aligned} \Rightarrow |\vec{BD}|^2 &= |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}| \cdot |\vec{b}| \\ &= 64 + 16 - 16 \times 4 \times \frac{1}{2} \\ &= 80 - 32 \\ &= 48 \end{aligned}$$

$$\Rightarrow |\vec{BD}| = \sqrt{48} = 4\sqrt{3}$$

17. For an obtuse angle

$$(cx\hat{i} - 6\hat{j} + 3\hat{k}) \cdot (x\hat{i} + 2\hat{j} + 2cx\hat{k}) < 0$$

$$\Rightarrow cx^2 - 12 + 6cx < 0$$

$$\Rightarrow cx^2 + 6cx - 12 < 0$$

We know that, if

$$ax^2 + bx + c > 0 \text{ or } < 0, \forall x$$

$$\text{Then, } b^2 - 4ac < 0$$

$$\therefore (6c)^2 - 4c(-12) < 0$$

$$\Rightarrow 3c^2 + 4c < 0$$

$$\Rightarrow 3c\left(c + \frac{4}{3}\right) < 0$$

$$\Rightarrow -\frac{4}{3} < c < 0$$

18. Given,

$$y - x \frac{dy}{dx} = a \left(y^2 + \frac{dy}{dx} \right)$$

$$\Rightarrow \frac{dy}{dx}(a + x) = y - ay^2$$

$$\Rightarrow \int \frac{dy}{y(1 - ay)} = \int \frac{dx}{a + x}$$

$$\Rightarrow \int \left(\frac{1}{y} + \frac{a}{1 - ay} \right) dy = \int \frac{dx}{a + x}$$

$$\Rightarrow \log y - \log(1 - ay) = \log(a + x) + \log c$$

$$\Rightarrow \log y = \log(1 - ay)(a + x)c$$

$$\Rightarrow y = c(1 - ay)(a + x)$$

19. Given,

$$y = (c_1 + c_2) \cos(x + c_3) - c_4 e^{x+c_5}$$

$$\Rightarrow y = (c_1 \cos c_3 + c_2 \cos c_3) \cos x - (c_1 \sin c_3 + c_2 \sin c_3) \sin x - c_4 e^{c_5} e^x$$

$$\Rightarrow y = A \cos x - B \sin x + C e^x$$

$$\text{where, } A = c_1 \cos c_3 + c_2 \cos c_3$$

$$B = c_1 \sin c_3 + c_2 \sin c_3$$

$$\text{and } C = -c_4 e^{c_5}$$

Which is an equation containing three arbitrary constant.

Hence, the order of the differential equation is 3.

20. Given, $f(x) = x^3 + bx^2 + cx + d$

$$\Rightarrow f'(x) = 3x^2 + 2bx + c$$

(As we know, if $ax^2 + bx + c > 0$ for all $x \Rightarrow a > 0$ and $D < 0$)

$$\text{Now, } D = 4b^2 - 12c = 4(b^2 - c) - 8c$$

(where $b^2 - c < 0$ and $c > 0$)

$$\therefore D = (-ve) \text{ ie, } D < 0$$

$$\Rightarrow f'(x) = 3x^2 + 2bx + c > 0$$

for all $x \in (-\infty, \infty)$.

(as $D < 0$ and $a > 0$)

Hence, $f(x)$ is strictly increasing function.

21. $\frac{d}{dx} \{\sin^{-1}(x\sqrt{1-x} + \sqrt{x}\sqrt{1-x^2})\}$

$$= \frac{d}{dx} \sin^{-1} \{(x\sqrt{1-(\sqrt{x})^2} + \sqrt{x}\sqrt{1-x^2})\}$$

$$\left[\because \frac{\sin^{-1} A + \sin^{-1} B}{= \sin^{-1}(A\sqrt{1-B^2} + B\sqrt{1-A^2})} \right]$$

$$\Rightarrow \frac{d}{dx} \{\sin^{-1}(x\sqrt{1-(\sqrt{x})^2} + \sqrt{x}\sqrt{1-x^2})\}$$

$$= \frac{d}{dx} (\sin^{-1} x + \sin^{-1} \sqrt{x})$$

$$= \frac{1}{\sqrt{1-x^2}} + \frac{1}{2\sqrt{x}\sqrt{1-x}}$$

22. Let $I = \int \frac{x \tan^{-1} x}{(1+x^2)^{3/2}} dx$

$$\text{Put } x = \tan \theta \Rightarrow dx = \sec^2 \theta d\theta$$

$$\therefore I = \int \frac{\tan \theta \tan^{-1}(\tan \theta)}{(1 + \tan^2 \theta)^{3/2}} \sec^2 \theta d\theta$$

$$\Rightarrow I = \int \frac{\theta \tan \theta \sec^2 \theta}{\sec^3 \theta} d\theta$$

$$\Rightarrow I = \int \theta \sin \theta d\theta$$

$$\Rightarrow I = \theta(-\cos \theta) + \int \cos \theta d\theta$$

$$\Rightarrow I = -\theta \cos \theta + \sin \theta + c$$

$$\Rightarrow I = -\tan^{-1} x \frac{1}{\sqrt{1+x^2}} + \frac{x}{\sqrt{1+x^2}} + c$$

$$\Rightarrow I = \frac{(x - \tan^{-1} x)}{\sqrt{1+x^2}} + c$$

$$23. |A| = \begin{vmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{vmatrix}$$

$$\Rightarrow |A| = 2(1 + \sin^2 \theta)$$

Now, $0 \leq \sin^2 \theta \leq 1$ for all $\theta \in [0, 2\pi]$.

$$\Rightarrow 2 \leq 2 + 2 \sin^2 \theta \leq 4 \text{ for all } \theta \in [0, 2\pi].$$

\therefore The range of $|A|$ is $[2, 4]$.

24. In the neighbourhood for $x = \frac{2\pi}{3}$, we have

$$\cos x < 0 \text{ and } \sin x > 0$$

$$\therefore y = -\cos x + \sin x$$

$$\Rightarrow \frac{dy}{dx} = \sin x + \cos x$$

$$\Rightarrow \left(\frac{dy}{dx} \right)_{\left(x = \frac{2\pi}{3} \right)} = \sin \frac{2\pi}{3} + \cos \frac{2\pi}{3}$$

$$= \frac{\sqrt{3} - 1}{2}$$

$$25. \lim_{n \rightarrow \infty} \left(\frac{1}{1-n^2} + \frac{2}{1-n^2} + \dots + \frac{n}{1-n^2} \right)$$

$$= \lim_{n \rightarrow \infty} \left(\frac{1+2+\dots+n}{1-n^2} \right)$$

$$= \lim_{n \rightarrow \infty} \frac{n(n+1)}{2(1-n^2)}$$

$$= \lim_{n \rightarrow \infty} \frac{\left(1 + \frac{1}{n}\right)}{2\left(\frac{1}{n^2} - 1\right)} = -\frac{1}{2}$$

$$26. \quad g[f(x)] = \begin{cases} [f(x)]^2 + 1, & f(x) \neq 2 \\ 3, & f(x) = 2 \end{cases}$$

$$\Rightarrow g[f(x)] = \begin{cases} \sin^2 x + 1, & x \neq n\pi \\ 3, & x = n\pi \end{cases}$$

$$\therefore \text{RHL} = \lim_{h \rightarrow 0} g[f(0+h)]$$

$$= \lim_{h \rightarrow 0} (\sin^2 h + 1) = 1$$

$$\text{and LHL} = \lim_{h \rightarrow 0} g[f(0-h)]$$

$$= \lim_{h \rightarrow 0} (\sin^2 h + 1) = 1$$

$$\therefore \lim_{x \rightarrow 0} g[f(x)] = 1$$

$$27. \text{ Given, } y = \int_0^x |t| dt$$

On differentiating w.r.t. x , we get

$$\frac{dy}{dx} = |x|$$

Since, the tangent is parallel to the line $y = 2x$

$$\therefore |x| = 2 \Rightarrow x = \pm 2$$

$$\text{When } x = 2, y = \int_0^2 |t| dt = \int_0^2 t dt$$

$$= \left[\frac{t^2}{2} \right]_0^2 = 2$$

When $x = -2$,

$$y = \int_0^{-2} |t| dt = \int_{-2}^0 t dt$$

$$= \left[\frac{t^2}{2} \right]_{-2}^0 = -2$$

Then, tangent at $(2, 2)$ is

$$y - 2 = 2(x - 2)$$

$$\Rightarrow 2x - y - 2 = 0$$

For x -intercept, put $y = 0$, then $x = 1$

Also, tangent at $(-2, -2)$ is

$$y + 2 = 2(x + 2)$$

$$\Rightarrow 2x - y + 2 = 0$$

For x -intercept, put $y = 0$, then $x = -1$.

28. Given,

$$P = x^3 - \frac{1}{x^3}, Q = x - \frac{1}{x}$$

$$\therefore \frac{P}{Q^2} = \frac{\left(x - \frac{1}{x}\right)\left(x^2 + 1 + \frac{1}{x^2}\right)}{\left(x - \frac{1}{x}\right)^2}$$


$$= \frac{\left(x - \frac{1}{x}\right)^2 + 3}{\left(x - \frac{1}{x}\right)}$$

$$= \left(x - \frac{1}{x}\right) + \frac{3}{\left(x - \frac{1}{x}\right)}$$

Clearly, the minimum does not exist.

29. Let $P(a \cos \theta, b \sin \theta)$, $Q(a \cos \theta, -b \sin \theta)$

Given, $PR : RQ = 1 : 2$



Let a point $R(h, k)$ divides the line joining the points P and Q internally in the ratio $1 : 2$.

$$\therefore h = a \cos \theta \Rightarrow \cos \theta = \frac{h}{a} \quad \dots (i)$$

$$\text{and } k = \frac{b}{3} \sin \theta \Rightarrow \sin \theta = \frac{3k}{b} \quad \dots (ii)$$

On squaring and adding Eqs. (i) and (ii), we get

$$\frac{h^2}{a^2} + \frac{9k^2}{b^2} = 1$$

Hence, locus of R is

$$\frac{x^2}{a^2} + \frac{9y^2}{b^2} = 1$$

30. Let $P(x_1, y_1)$ be a point on the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \text{ then } \frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} = 1$$

The chord of contact of tangents from P to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2 \text{ is } \frac{x x_1}{a^2} - \frac{y y_1}{b^2} = 2 \quad \dots (i)$$

The equation of the asymptotes are

$$\frac{x}{a} - \frac{y}{b} = 0 \text{ and } \frac{x}{a} + \frac{y}{b} = 0$$

The point of intersection of Eq. (i) with the two asymptotes are given by

$$X_1 = \frac{2a}{\frac{x_1}{a} - \frac{y_1}{b}}, Y_1 = \frac{2b}{\frac{x_1}{a} - \frac{y_1}{b}}$$

$$\text{and } X_2 = \frac{2a}{\frac{x_1}{a} + \frac{y_1}{b}}, Y_2 = \frac{-2b}{\frac{x_1}{a} + \frac{y_1}{b}}$$

\therefore Area of triangle

$$= \frac{1}{2} (X_1 Y_2 - X_2 Y_1)$$

$$= \frac{1}{2} \left(\frac{4ab \times 2}{\frac{x_1^2}{a^2} - \frac{y_1^2}{b^2}} \right) = 4ab$$

31. Let S denotes the sum of the series. The general term of the given series is $T_r = (-1)^r (3 + 5r)^n C_r$ (nth term of AP)

$$\begin{aligned} \therefore S &= \sum_{r=0}^n (-1)^r (3 + 5r)^n C_r \\ &= 3 \sum_{r=0}^n (-1)^r {}^n C_r + 5 \sum_{r=0}^n (-1)^r r {}^n C_r \end{aligned}$$

$$\begin{aligned} &= 3(C_0 - C_1 + C_2 - C_3 + C_4 - \dots \\ &\quad + (-1)^n \cdot C_n) + 5(-C_1 + 2C_2 - 3C_3 \\ &\quad + 4C_4 - \dots + (-1)^n \cdot n \cdot C_n) \end{aligned}$$

$$\Rightarrow S = 0 + 0 = 0$$

32. Since, diagonal element of a skew-symmetric matrix are all zero.

$$\therefore \text{tr}(A) = \sum_{i=1}^n a_{ii} = 0$$

$$33. \text{ Let } \Delta = \begin{vmatrix} 1 & \alpha & \alpha^2 \\ \cos(p-d)a & \cos pa & \cos(p-d)a \\ \sin(p-d)a & \sin pa & \sin(p-d)a \end{vmatrix}$$

Applying $C_3 \rightarrow C_3 - C_1$, we get

$$\begin{aligned} \Rightarrow \Delta &= \begin{vmatrix} 1 & \alpha & \alpha^2 - 1 \\ \cos(p-d)a & \cos pa & 0 \\ \sin(p-d)a & \sin pa & 0 \end{vmatrix} \\ &= (\alpha^2 - 1) \{ -\cos pa \sin(p-d)a \\ &\quad + \sin pa \cos(p-d)a \} \\ &= (\alpha^2 - 1) \sin \{ -(p-d)a + pa \} \end{aligned}$$

$$\Rightarrow \Delta = (\alpha^2 - 1) \sin da$$

which is independent of p.


34. Let S_n be the sum of first n terms of the series

$$\begin{aligned} &\frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots \\ \therefore S_n &= \left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{4}\right) + \left(1 - \frac{1}{8}\right) + \dots \\ &\quad + \left(1 - \frac{1}{2^n}\right) \end{aligned}$$

$$\begin{aligned} &= n - \frac{1}{2} \left(\frac{1 - \frac{1}{2^n}}{1 - \frac{1}{2}} \right) \\ &= n - 1 + \frac{1}{2^n} = n - 1 + 2^{-n} \end{aligned}$$

35. In $\triangle AEC$,

$$\begin{aligned} \tan 60^\circ &= \frac{500}{d_1} \\ \Rightarrow d_1 &= \frac{500}{\sqrt{3}} \text{ m} \quad \dots (i) \end{aligned}$$



and in $\triangle BEC$,

$$\tan 30^\circ = \frac{500}{d_2}$$

$$\Rightarrow d_2 = 500\sqrt{3} \text{ m} \quad \dots \text{ (ii)}$$

\therefore Required diameter,

$$\begin{aligned} AB &= d_1 + d_2 \\ &= \frac{500}{\sqrt{3}} + 500\sqrt{3} = \frac{2000}{\sqrt{3}} \text{ m} \end{aligned}$$

36. Given,

$$\sec x - 1 = (\sqrt{2} - 1) \tan x$$

$$\Rightarrow 1 - \cos x = (\sqrt{2} - 1) \sin x$$

$$\Rightarrow \sin \frac{x}{2} \left(\sin \frac{x}{2} - (\sqrt{2} - 1) \cos \frac{x}{2} \right) = 0$$

$$\Rightarrow \sin \frac{x}{2} = 0$$

$$\text{or } \tan \frac{x}{2} = \sqrt{2} - 1 = \tan \frac{45^\circ}{2}$$

$$\Rightarrow \frac{x}{2} = n\pi \quad \text{or} \quad \frac{x}{2} = n\pi + \frac{\pi}{8}$$

$$\Rightarrow x = 2n\pi, 2n\pi + \frac{\pi}{4}$$

37. Now, $\cos\left(x + \frac{\pi}{6}\right) + \sin\left(x + \frac{\pi}{6}\right)$

$$= \sqrt{2} \left[\frac{1}{\sqrt{2}} \cos\left(x + \frac{\pi}{6}\right) + \frac{1}{\sqrt{2}} \sin\left(x + \frac{\pi}{6}\right) \right]$$

$$= \sqrt{2} \cos\left(x + \frac{\pi}{6} - \frac{\pi}{4}\right)$$

$$= \sqrt{2} \cos\left(x - \frac{\pi}{12}\right)$$

For maximum value $x = \frac{\pi}{12}$

38. Given,

$$z_r = \cos \frac{r\alpha}{n^2} + i \sin \frac{r\alpha}{n^2}$$

(where $r = 1, 2, 3, \dots, n$)

$$z_1 = \cos \frac{\alpha}{n^2} + i \sin \frac{\alpha}{n^2}$$

$$z_2 = \cos \frac{2\alpha}{n^2} + i \sin \frac{2\alpha}{n^2}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$z_n = \cos \frac{n\alpha}{n^2} + i \sin \frac{n\alpha}{n^2}$$

$$\therefore \lim_{n \rightarrow \infty} (z_1 z_2 \dots z_n) = \lim_{n \rightarrow \infty} \left(\cos \frac{\alpha}{n^2} + i \sin \frac{\alpha}{n^2} \right)$$

$$\times \left(\cos \frac{2\alpha}{n^2} + i \sin \frac{2\alpha}{n^2} \right) \dots \left(\cos \frac{n\alpha}{n^2} + i \sin \frac{n\alpha}{n^2} \right)$$

$$= \lim_{n \rightarrow \infty} \left[\cos \left\{ \frac{\alpha}{n^2} (1 + 2 + 3 + \dots + n) \right\} + i \sin \right.$$

$$\left. \left\{ \frac{\alpha}{n^2} (1 + 2 + 3 + \dots + n) \right\} \right]$$

$$= \lim_{n \rightarrow \infty} \left[\cos \frac{\alpha n(n+1)}{2n^2} + i \sin \frac{\alpha n(n+1)}{2n^2} \right]$$

$$= \cos \frac{\alpha}{2} + i \sin \frac{\alpha}{2} = e^{\frac{i\alpha}{2}}$$

39. Let p : Paris is in France


and q : London is in England

Given, $p \wedge q$

Its negation is $\sim (p \wedge q) = \sim p \vee \sim q$

Hence, Paris is not in France or London is not in England.

40. Solving the given curves $y = \sqrt{x}$ or $y^2 = x$ and $y = x^3$



we get, the points of intersection (0, 0) and (1, 1).

$$\begin{aligned}\therefore \text{Required area} &= \int_0^1 (\sqrt{x} - x^3) dx \\ &= \left[\frac{x^{3/2}}{3/2} - \frac{x^4}{4} \right]_0^1 \\ &= \frac{5}{12} \text{ sq unit}\end{aligned}$$

41. Given equations of lines are $3x - 4y + 7 = 0$ and $-12x - 5y + 2 = 0$

$$\begin{aligned}a_1a_2 + b_1b_2 &= 3 \times (-12) + (-4)(-5) \\ &= -36 + 20 = -16\end{aligned}$$

$$\Rightarrow a_1a_2 + b_1b_2 < 0$$

\therefore Obtuse angle bisector is

$$\frac{3x - 4y + 7}{\sqrt{3^2 + (-4)^2}} = -\frac{-12x - 5y + 2}{\sqrt{(-12)^2 + (-5)^2}}$$

$$\Rightarrow 13(3x - 4y + 7) = -5(-12x - 5y + 2)$$

$$\Rightarrow 21x + 77y - 101 = 0$$

42. Given, $\frac{dy}{dx} = \frac{y}{x} - \cos^2\left(\frac{y}{x}\right)$

$$\Rightarrow \frac{x dy - y dx}{x^2} = -\left(\cos^2 \frac{y}{x}\right) dx$$

$$\Rightarrow \sec^2\left(\frac{y}{x}\right) \left(\frac{x dy - y dx}{x^2}\right) = -\frac{dx}{x}$$

$$\Rightarrow \sec^2 \frac{y}{x} \cdot d\left(\frac{y}{x}\right) = -\frac{dx}{x}$$

$$\Rightarrow \tan \frac{y}{x} = -\log x + c$$

$$\text{When } x = 1, y = \frac{\pi}{4},$$

$$c = 1$$

$$\therefore \tan\left(\frac{y}{x}\right) = 1 - \log x$$

$$\Rightarrow \log x = 1 - \tan\left(\frac{y}{x}\right)$$

$$\Rightarrow x = e^{1 - \tan\left(\frac{y}{x}\right)}$$

43. Here, $P(1) = 2$ and from the equation

$$P(k) = k(k+1) + 2$$

$$\Rightarrow P(1) = 4$$

So, $P(1)$ is not true.

Hence, mathematical induction is not applicable.

44. The general equation of all non-vertical lines in a plane is $ax + by = 1$, where $b \neq 0$.

$$\text{On differentiating, we get } a + b \frac{dy}{dx} = 0$$

On again differentiating, we get

$$b \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow \frac{d^2y}{dx^2} = 0$$

45. Let the required vector be $\vec{c} = x \hat{i} + y \hat{k}$

As it is a unit vector, therefore

$$|\vec{c}| = 1 \Rightarrow x^2 + y^2 = 1 \quad \dots(i)$$

\vec{a} and \vec{c} are inclined at the angle 45° .

$$\therefore \cos 45^\circ = \frac{2x - y}{\sqrt{4 + 4 + 1}}$$

$$\Rightarrow 2x - y = \frac{3}{\sqrt{2}} \quad \dots(ii)$$

\vec{b} and \vec{c} are inclined at an angle 60° .

$$\therefore -\frac{y}{\sqrt{2}} = \cos 60^\circ \Rightarrow y = -\frac{1}{\sqrt{2}} \quad \dots(iii)$$

From Eqs. (ii) and (iii), we get $x = \frac{1}{\sqrt{2}}$

Hence, the required vector is $\frac{1}{\sqrt{2}} \hat{i} - \frac{1}{\sqrt{2}} \hat{k}$.

46. Given,

$$\int_2^e \left(\frac{1}{\log x} - \frac{1}{(\log x)^2} \right) dx = a + \frac{b}{\log 2}$$

$$\text{Put } \log x = z \Rightarrow x = e^z \Rightarrow dx = e^z dz$$

$$\therefore \int_2^e \left(\frac{1}{\log x} - \frac{1}{(\log x)^2} \right) dx$$

$$= \int_{\log 2}^1 \left(\frac{1}{z} - \frac{1}{z^2} \right) e^z dz$$

$$= \int_{\log 2}^1 e^z \left(\frac{1}{z} + d \left(\frac{1}{z} \right) \right) dz$$

$$= \left[e^z \cdot \frac{1}{z} \right]_{\log 2}^1 = e - \frac{2}{\log 2}$$

$$\therefore a = e \quad \text{and} \quad b = -2$$


47. Let the equation of AB be $\frac{x}{a} + \frac{y}{b} = 1$.

Since, the line AB touches the circle

$$x^2 + y^2 - 4x - 4y + 4 = 0$$

$$\therefore \frac{\left| \frac{2}{a} + \frac{2}{b} - 1 \right|}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} = 2$$

[Since, O (0, 0) and C (2, 2) lie on the same side of AB, therefore $\frac{2}{a} + \frac{2}{b} - 1 < 0$]



$$\Rightarrow \frac{-(2b + 2a - ab)}{\sqrt{a^2 + b^2}} = 2$$

$$\Rightarrow 2a + 2b - ab + 2\sqrt{a^2 + b^2} = 0 \quad \dots(i)$$

Since, $\triangle OAB$ is a right angled triangle. So, its circumcentre is the mid point of AB.

$$\therefore h = \frac{a}{2} \quad \text{and} \quad k = \frac{b}{2} \quad \dots(ii)$$

$$\Rightarrow a = 2h \quad \text{and} \quad b = 2k$$

From Eqs. (i) and (ii), we get

$$4h + 4k - 4hk + 2\sqrt{4h^2 + 4k^2} = 0$$

$$\Rightarrow h + k - hk + \sqrt{h^2 + k^2} = 0$$

So, the locus of P(h, k) is

$$x + y - xy + \sqrt{x^2 + y^2} = 0$$

But the locus of the circumcentre is given to be

$$x + y - xy + k\sqrt{x^2 + y^2} = 0$$

$$\therefore k = 1$$

48. Let $f(x) = \tan x$

The point of discontinuity of $f(x)$ are those points where $\tan x$ is infinite.

ie, $\tan x = \tan \infty$

$$\Rightarrow x = (2n + 1) \frac{\pi}{2}, n \in I$$

49. Given curves are

$$x^3 - 3xy^2 + 2 = 0$$

$$\text{and} \quad 3x^2y - y^3 - 2 = 0$$

On differentiating w.r.t. x, we get

$$\left(\frac{dy}{dx} \right)_{c_1} = \frac{x^2 - y^2}{2xy}$$

$$\text{and} \quad \left(\frac{dy}{dx} \right)_{c_2} = \frac{-2xy}{x^2 - y^2}$$

$$\text{Now,} \quad \left(\frac{dy}{dx} \right)_{c_1} \times \left(\frac{dy}{dx} \right)_{c_2} = -1$$

Hence, the two curves cut at right angles.

50. Given,

$$f(x) = \frac{2 \sin 8x \cos x - 2 \sin 6x \cos 3x}{2 \cos 2x \cos x - 2 \sin 3x \sin 4x}$$

$$= \frac{(\sin 9x + \sin 7x) - (\sin 9x + \sin 3x)}{(\cos 3x + \cos x) + (\cos 7x - \cos x)}$$

$$= \frac{\sin 7x - \sin 3x}{\cos 7x + \cos 3x} = \frac{2 \sin 2x \cos 5x}{2 \cos 5x \cos 2x}$$

$$= \tan 2x$$

$$\therefore \text{Period of } f(x) = \frac{\pi}{2}$$

51. Let $y = f(\tan x)$ and $u = g(\sec x)$

On differentiating w.r.t. x, we get

$$\frac{dy}{dx} = f'(\tan x) \sec^2 x$$

$$\text{and} \quad \frac{du}{dx} = g'(\sec x) \cdot \sec x \tan x$$

$$\therefore \frac{dy}{du} = \frac{dy/dx}{du/dx} = \frac{f'(\tan x) \sec^2 x}{g'(\sec x) \sec x \tan x}$$

$$\begin{aligned} \therefore \left(\frac{dy}{du} \right)_{x=\pi/4} &= \frac{f' \left(\tan \frac{\pi}{4} \right)}{g' \left(\sec \frac{\pi}{4} \right) \sin \frac{\pi}{4}} \\ &= \frac{f'(1) \cdot \sqrt{2}}{g'(\sqrt{2})} = \frac{2 \cdot \sqrt{2}}{4} = \frac{1}{\sqrt{2}} \end{aligned}$$

$$52. \text{ Area of } \Delta = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ a \cos \theta & b \sin \theta & 1 \\ a \cos \theta & -b \sin \theta & 1 \end{vmatrix}$$

$$\begin{aligned} \Rightarrow \Delta &= \frac{1}{2} |1(-ab \sin \theta \cos \theta \\ &\quad - ab \sin \theta \cos \theta)| \\ &= \frac{ab \sin 2\theta}{2} \end{aligned}$$

Since, maximum value of $\sin 2\theta$ is 1, when $\theta = \frac{\pi}{4}$

$$\therefore \Delta_{\max} = \frac{ab}{2}$$

53. The angle between the two curves is the angle between their tangents at their point of contact.

\therefore The point of intersection of two curves is $a^x = b^x$

For distinct values of a and b , the equality is true, if $x = 0$

Now, if $x = 0$, then $y = a^0 = 1$

\therefore The two curves intersect at $(0, 1)$.

Let m_1 and m_2 be the slope of the tangent to the curve $y = a^x$ and $y = b^x$ respectively at $P(0, 1)$.

$$\therefore m_1 = \left(\frac{dy}{dx} \right)_P = (a^x \log a)_{(0,1)} = \log a$$

$$m_2 = \left(\frac{dy}{dx} \right)_P = (b^x \log b)_{(0,1)} = \log b$$

If α be the angle of intersection of the tangents.

$$\text{Then, } \tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$\Rightarrow \tan \alpha = \left| \frac{\log a - \log b}{1 + \log a \log b} \right|$$

$$54. \text{ Given, } x - \frac{2}{x-1} = 1 - \frac{2}{x-1}$$

For domain of equation $x \neq 1$

But here $x = 1$

Which is not possible.

Hence, no solution exist.

55. We know that, when we multiply a complex number by i , it rotates the vector for z by an angle of $\frac{\pi}{2}$ in the anticlockwise direction.

\therefore Multiplying by i^2 will rotate the vector for z by π .

$$\therefore z' = i^2 (-4 + 5i) = -(-4 + 5i)$$

Stretching $\frac{3}{2}$ times

$$\begin{aligned} \therefore z'' &= \frac{3}{2} (4 - 5i) \\ &= 6 - \frac{15}{2}i \end{aligned}$$

56. By definition, Any row of a matrix/determinant A when multiplied by its corresponding cofactors results in the value of $|A|$.

57. Since, $f(x)$ is a continuous decreasing function on $\left[\frac{\pi}{6}, \frac{\pi}{3} \right]$.

$\therefore f(x)$ attains every value between $\left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ its minimum value

$$\text{ie, } f\left(\frac{\pi}{3}\right) = \frac{1}{2} - \frac{\pi}{3} \left(1 + \frac{\pi}{3}\right)$$

and maximum value is

$$f\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} - \frac{\pi}{6} \left(1 + \frac{\pi}{6}\right)$$

58. Given, $(p \vee q) \Rightarrow r$

$$\equiv \sim r \Rightarrow \sim (p \vee q)$$

$$\equiv \sim r \Rightarrow (\sim p \wedge \sim q)$$

59. By Rolle's theorem,

$$f(1) = f(3)$$

$$\Rightarrow a + b + 11 - 6 = 27a + 9b + 33 - 6$$

$$\Rightarrow 26a + 8b + 22 = 0$$

$$\Rightarrow 13a + 4b + 11 = 0 \quad \dots(i)$$

$$\text{Now, } f'(x) = 3ax^2 + 2bx + 11$$

$$\Rightarrow f'\left(2 + \frac{1}{\sqrt{3}}\right)$$

$$= 3a\left(2 + \frac{1}{\sqrt{3}}\right)^2 + 2b\left(2 + \frac{1}{\sqrt{3}}\right) + 11$$

$$\Rightarrow 0 = 3a\left(4 + \frac{1}{3} + \frac{4}{\sqrt{3}}\right) + 4b + \frac{2b}{\sqrt{3}} + 11$$

$$\Rightarrow 13a + 4b + \frac{12a}{\sqrt{3}} + \frac{2b}{\sqrt{3}} + 11 = 0$$

$$\Rightarrow -11 + \frac{12a}{\sqrt{3}} + \frac{2b}{\sqrt{3}} + 11 = 0$$

[from Eq. (i)]

$$\Rightarrow 6a + b = 0 \quad \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$a = 1, b = -6$$

60. Now, $f\left(\frac{1}{x}\right)f\left(\frac{1}{y}\right) - \frac{1}{2}\left[f\left(\frac{x}{y}\right) + f(xy)\right]$

$$= \cos\left(\log \frac{1}{x}\right)\cos\left(\log \frac{1}{y}\right)$$

$$- \frac{1}{2}\left[\cos\left(\log \frac{x}{y}\right) + \cos[\log(xy)]\right]$$

$$= \cos(-\log x)\cos(-\log y)$$

$$- \frac{1}{2}[\cos(\log x - \log y) + \cos(\log x + \log y)]$$

$$= \cos(\log x)\cos(\log y)$$

$$- \frac{1}{2}[2\cos(\log x)\cos(\log y)]$$

$$= \cos(\log x)\cos(\log y)$$

$$- \cos(\log x)\cos(\log y) = 0$$

61. Now, $\alpha + \beta = \sin^{-1} \frac{\sqrt{3}}{2} + \cos^{-1} \frac{\sqrt{3}}{2}$
 $+ \sin^{-1} \frac{1}{3} + \cos^{-1} \frac{1}{3}$

$$= \frac{\pi}{2} + \frac{\pi}{2} = \pi$$

Also, $\alpha = \frac{\pi}{3} + \sin^{-1} \frac{1}{3} < \frac{\pi}{3} + \sin^{-1} \frac{1}{2}$

As $\sin \theta$ is increasing in $\left[0, \frac{\pi}{2}\right]$.

$$\therefore \alpha < \frac{\pi}{3} + \frac{\pi}{6} = \frac{\pi}{2}$$

$$\Rightarrow \beta > \frac{\pi}{2} > \alpha$$

$$\Rightarrow \alpha < \beta$$

62. Given,

$$1 + \sin \theta + \sin^2 \theta + \dots \infty = 4 + 2\sqrt{3}$$

$$\Rightarrow \frac{1}{1 - \sin \theta} = 4 + 2\sqrt{3} \quad (\because 0 < \sin \theta < 1)$$

$$\Rightarrow 1 - \sin \theta = \frac{1}{4 + 2\sqrt{3}} \times \frac{4 - 2\sqrt{3}}{4 - 2\sqrt{3}}$$


$$\Rightarrow 1 - \sin \theta = \frac{4 - 2\sqrt{3}}{16 - 12} = 1 - \frac{\sqrt{3}}{2}$$

$$\Rightarrow \sin \theta = \frac{\sqrt{3}}{2} = \sin \frac{\pi}{3}$$

$$\theta = \frac{\pi}{3}$$

or $\theta = \pi - \frac{\pi}{3} = \frac{2\pi}{3}$

63. In ΔAPC , $\sin(\angle PAC) = \frac{CP}{AC}$



$$\Rightarrow AC = \frac{r}{\sin \frac{\alpha}{2}} = r \operatorname{cosec} \frac{\alpha}{2} \quad \dots(i)$$

Again in ΔABC ,

$$\sin \beta = \frac{BC}{AC}$$

$$\Rightarrow BC = AC \sin \beta$$

$$\Rightarrow H = r \sin \beta \operatorname{cosec} \left(\frac{\alpha}{2} \right)$$

[from Eq. (i)]

$$\begin{aligned} 64. (a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2} \\ = (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2} \\ + (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2} \\ = a^2 + b^2 + 2ab \left(\sin^2 \frac{C}{2} - \cos^2 \frac{C}{2} \right) \\ = a^2 + b^2 - 2ab \cos C \\ = a^2 + b^2 - (a^2 + b^2 - c^2) \\ = c^2 \end{aligned}$$

$$65. \text{ Given, } \left(1 - \frac{r_1}{r_2} \right) \left(1 - \frac{r_1}{r_3} \right) = 2$$

$$\therefore \left(1 - \frac{s-b}{s-a} \right) \left(1 - \frac{s-c}{s-a} \right) = 2$$

$$\Rightarrow \frac{(b-a)(c-a)}{(s-a)^2} = 2$$

$$\Rightarrow a^2 = b^2 + c^2$$

Hence, triangle is a right angled triangle.

$$66. \text{ Given, } a + b + c = 0$$

Let the roots of the equation $4ax^2 + 3bx + 2c = 0$ are α and β .

$$\text{Now, } D = b^2 - 4ac$$

$$= 9b^2 - 4(4a)(2c)$$

$$= 9b^2 - 32ac$$

$$= 9(a+c)^2 - 32ac$$

Hence, roots are real.

67. The given graph is a connected graph.

$$68. \text{ Given, } \tan^{-1} x + 2 \cot^{-1} x = \frac{2\pi}{3}$$

$$\Rightarrow \tan^{-1} x + 2 \tan^{-1} \frac{1}{x} = \frac{2\pi}{3}$$

$$\Rightarrow \tan^{-1} x + \tan^{-1} \left(\frac{2 \left(\frac{1}{x} \right)}{1 - \left(\frac{1}{x} \right)^2} \right) = \frac{2\pi}{3}$$

$$\left[\because 2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2} \right]$$

$$\Rightarrow \tan^{-1} x + \tan^{-1} \left(\frac{2x}{x^2-1} \right) = \frac{2\pi}{3}$$

$$\Rightarrow \tan^{-1} \left(\frac{x + \frac{2x}{x^2-1}}{1 - \frac{2x^2}{x^2-1}} \right) = \frac{2\pi}{3}$$

$$\Rightarrow \frac{x^3 - x + 2x}{x^2 - 1 - 2x^2} = \tan \left(\frac{2\pi}{3} \right)$$

$$\Rightarrow \frac{x^3 + x}{-1 - x^2} = \tan \left(\frac{2\pi}{3} \right)$$

$$\Rightarrow \frac{x(x^2+1)}{-1(x^2+1)} = -\sqrt{3}$$

$$\Rightarrow x = \sqrt{3}$$

$$69. \text{ Given complex number is } \frac{(1+i)^2}{1-i}$$

$$= \frac{(1+i^2+2i)}{1-i} \times \frac{1+i}{1+i}$$

$$= \frac{2i(1+i)}{1-i^2}$$

$$= \frac{2i+2i^2}{1+1}$$

$$= \frac{2i-2}{2}$$

$$= i-1$$

\therefore Required conjugate is $-i-1$.

70. Since, the number of vertices of odd degree is always even and number of even degree is always even. So, $m + n$ is an even number.

71. We have, $\sin \left[2 \cos^{-1} \frac{\sqrt{5}}{3} \right]$

$$= \sin \left[\cos^{-1} \left(2 \cdot \left(\frac{\sqrt{5}}{3} \right)^2 - 1 \right) \right]$$

$$[\because 2 \cos^{-1} x = \cos^{-1} (2x^2 - 1)]$$

$$= \sin \left[\cos^{-1} \left(\frac{1}{9} \right) \right]$$

$$= \sin \left[\sin^{-1} \sqrt{1 - \left(\frac{1}{9} \right)^2} \right]$$

$$[\because \cos^{-1} x = \sin^{-1} (\sqrt{1 - x^2})]$$

$$= \sin \left[\sin^{-1} \sqrt{\frac{80}{81}} \right]$$

$$= \frac{\sqrt{80}}{9}$$

$$= \frac{4\sqrt{5}}{9}$$

72. Given, $G = \{3, 6, 9, 12\}$

The composition table is

\times_{15}	3	6	9	12
3	9	3	12	6
6	3	6	9	12
9	12	9	6	3
12	6	12	3	9

From the table 6 is the identity element.

73. Since, identity element is its own inverse. So, minimum number of element is 1.

74. We have, $\frac{3x^2 + 1}{x^2 - 6x + 8}$

On dividing, we get

$$\frac{3x^2 + 1}{x^2 - 6x + 8} = 3 + \frac{18x - 23}{x^2 - 6x + 8} \quad \dots(i)$$

Now, $\frac{18x - 23}{(x - 2)(x - 4)} = \frac{A}{x - 2} + \frac{B}{x - 4}$

$$\Rightarrow 18x - 23 = A(x - 4) + B(x - 2)$$

$$\Rightarrow 18x - 23 = (A + B)x - 4A - 2B$$

Equating the coefficient of x and constant term, we get

$$A + B = 18$$

$$-4A - 2B = -23$$

On solving these equations, we get

$$A = -\frac{13}{2}, B = \frac{49}{2}$$

$$\therefore \frac{18x - 23}{(x - 2)(x - 4)} = -\frac{13}{2(x - 2)} + \frac{49}{2(x - 4)}$$

Then, from Eq. (i), we get


$$\frac{3x^2 + 1}{x^2 - 6x + 8} = 3 - \frac{13}{2(x - 2)} + \frac{49}{2(x - 4)}$$

75. ΔAOB is the given triangle.

$$\text{Slope of } AB = \frac{\frac{3}{2} - 0}{0 + 5} = \frac{3}{10}$$

$$\text{Slope of } BO = \frac{0 - 0}{0 + 5} = 0$$

The equation of line passing through A and perpendicular to BO is $y - 0 = -0 \left(x - \frac{3}{2} \right)$



$$\Rightarrow y = 0 \quad \dots(i)$$

and equation of line passing through O and perpendicular to AB is $y - 0 = -\frac{10}{3}(x - 0)$

$$\Rightarrow y = -\frac{10}{3}x \quad \dots(ii)$$

The intersection point of Eqs. (i) and (ii) is $(0, 0)$, which is the required orthocentre.

76. Given, $y = -x^2 + 6x - 3$


$$\Rightarrow \frac{dy}{dx} = -2x + 6 = y'$$

y is increasing, if $y' > 0$

$$\Rightarrow -2x + 6 > 0 \Rightarrow -2x > -6$$

$$\Rightarrow 2x < 6 \Rightarrow x < 3$$

77. Given, curve $x = 4 - y^2$ and y -axis.



The required area is $ABCOA$

$$= 2 \times \text{area of } ABOA$$

$$= 2 \times \int_0^4 \sqrt{4-x} \, dx$$

$$= 2 \left[-\frac{(4-x)^{3/2}}{3/2} \right]_0^4$$

$$\begin{aligned} &= 2 \left[-\frac{2}{3} \times 0 + \frac{2}{3} (4)^{3/2} \right] \\ &= \frac{4}{3} \times 8 \\ &= \frac{32}{3} \text{ sq unit} \end{aligned}$$

78. We know that, if $a = p_1^{\alpha_1} \cdot p_2^{\alpha_2} \dots$

Then, the total number of positive divisors of a is

$$T(a) = (\alpha_1 + 1)(\alpha_2 + 1) \dots$$

$$\text{Given, } 252 = 2^2 \times 3^2 \times 7^1$$

$$\text{Here, } \alpha_1 = 2, \alpha_2 = 2, \alpha_3 = 1$$

$$\begin{aligned} \therefore T(a) &= (2+1)(2+1)(1+1) \\ &= 3 \cdot 3 \cdot 2 \\ &= 18 \end{aligned}$$

79. We have, $5^{124} = (5^3)^{41} \cdot 5$

$$\text{Now, } 5^3 \equiv 1 \pmod{124}$$

$$\therefore (5^3)^{41} \equiv 1 \pmod{124}$$

$$(5^3)^{41} \cdot 5 \equiv 1 \cdot 5 \pmod{124}$$

$$\Rightarrow 5^{124} \equiv 5 \pmod{124}$$

80. The set of odd integers under addition is not a group. Since, addition of two odd numbers is always an even number. *ie*, closure property does not hold.