

MIND MAP : LEARNING MADE SIMPLE

CHAPTER - 7

Integrals

It is the inverse of differentiation. Let, $\frac{d}{dx} F(x) = f(x)$. Then $\int f(x) dx = F(x) + c$, 'c': constant of integral. These integrals are called indefinite or general integrals.

Properties of indefinite integrals are

$$(i) \int [f(x) + g(x)] dx = \int f(x) dx + \int g(x) dx, \quad (ii) \int kf(x) dx = k \int f(x) dx,$$

For eg : $\int (3x^2 + 2x) dx = x^3 + x^2 + c$ where k is real.

The method in which we change the variable to some other variable is called the method of substitution

$$\int \tan x dx = \log |\sec x| + c \quad \int \cot x dx = \log |\sin x| + c$$

$$\int \sec x dx = \log |\sec x + \tan x| + c \quad \int \csc x dx = \log |\csc x - \cot x| + c.$$

$$(i) \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + c \quad (ii) \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + c$$

$$(iii) \int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + c \quad (iv) \int \frac{dx}{\sqrt{x^2 - a^2}} = \log |x + \sqrt{x^2 - a^2}| + c$$

$$(v) \int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + c \quad (vi) \int \frac{dx}{\sqrt{x^2 + a^2}} = \log |x + \sqrt{x^2 + a^2}| + c.$$

$$\int f_1(x) f_2(x) dx = f_1(x) \int f_2(x) dx - \int \left[\frac{d}{dx} f_1(x) \right] \int f_2(x) dx$$

$$(i) \int \sqrt{x^2 - a^2} dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log |x + \sqrt{x^2 - a^2}| + c.$$

$$(ii) \int \sqrt{x^2 + a^2} dx = \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2 + a^2}| + c.$$

$$(iii) \int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + c.$$

Let the area function be defined by $A(x) = \int_a^x f(x) dx \forall x \geq a$, where f is continuous on $[a, b]$ then $A'(x) = f(x) \forall x \in [a, b]$.

Let f be a continuous function of x defined on $[a, b]$ and let F be another function such that $\frac{d}{dx} F(x) = f(x) \forall x \in \text{domain of } f_1$ then $\int_a^b f(x) dx = [F(x) + c]_a^b = F(b) - F(a)$. This is called the definite integral of f over the range $[a, b]$, where a and b are called the limits of integration, a being the lower limit and b be the upper limit.

$$\int_{-\pi/4}^{\pi/4} \sin^2 x dx$$

$$= 2 \int_0^{\pi/4} \sin^2 x dx$$

$$= 2 \int_0^{\pi/4} \left(\frac{1 - \cos 2x}{2} \right) dx$$

$$= \int_0^{\pi/4} (1 - \cos 2x) dx$$

$$= \left[x - \frac{\sin 2x}{2} \right]_0^{\pi/4}$$

$$= \frac{\pi}{4} - \frac{1}{2}$$

$$(i) \int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1 \text{ like, } \int dx = x + c$$

$$(ii) \int \cos x dx = \sin x + c \quad (iii) \int \sin x dx = -\cos x + c$$

$$(iv) \int \sec^2 x dx = \tan x + c \quad (v) \int \csc^2 x dx = -\cot x + c$$

$$(vi) \int \sec x \tan x dx = \sec x + c \quad (vii) \int \csc x \cot x dx = -\csc x + c$$

$$(viii) \int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x + c \quad (ix) \int \frac{dx}{\sqrt{1-x^2}} = -\cos^{-1} x + c$$

$$(x) \int \frac{dx}{1+x^2} = \tan^{-1} x + c \quad (xi) \int \frac{dx}{1+x^2} = -\cot^{-1} x + c$$

$$(xii) \int e^x dx = e^x + c \quad (xiii) \int a^x dx = \frac{a^x}{\log a} + c$$

$$(xiv) \int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x + c \quad (xv) \int \frac{dx}{x\sqrt{x^2-1}} = -\csc^{-1} x + c$$

$$(xvi) \int \frac{1}{x} dx = \log |x| + c$$

A rational function of the form $\frac{P(x)}{Q(x)} (Q(x) \neq 0) = T(x) + \frac{P_1(x)}{Q(x)}$, $P_1(x)$

has degree less than that of Q(x). We can integrate $\frac{P_1(x)}{Q(x)}$ by expressing

it in the following forms -

$$(i) \frac{px+q}{(x-a)(x-b)} = \frac{A}{x-a} + \frac{B}{x-b}, a \neq b.$$

$$(ii) \frac{px+q}{(x+a)^2} = \frac{A}{x-a} + \frac{B}{(x-a)^2}$$

$$(iii) \frac{px^2+qx+r}{(x-a)(x-b)(x-c)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$$

$$(iv) \frac{px^2+qx+r}{(x-a)^2(x-b)} = \frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b} \quad (v) \frac{px^2+qx+r}{(x-a)(x^2+bx+c)} = \frac{A}{x-a} + \frac{Bx+C}{x^2+bx+c}$$