

Chapter : 28. THE PLANE

Exercise : 28A

Question: 1

Find the equation

Solution:

(i) A(2, 2, -1), B(3, 4, 2) and C(7, 0, 6)

Given Points :

$$A = (2, 2, -1)$$

$$B = (3, 4, 2)$$

$$C = (7, 0, 6)$$

To Find : Equation of plane passing through points A, B & C

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Vector :

If A and B be two points with position vectors $\vec{a}$ & $\vec{b}$ respectively, where

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{AB} = \vec{b} - \vec{a}$$

$$= (b_1 - a_1)\hat{i} + (b_2 - a_2)\hat{j} + (b_3 - a_3)\hat{k}$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

4) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

5) Equation of Plane :

If A = (a_1, a_2, a_3) , B = (b_1, b_2, b_3) , C = (c_1, c_2, c_3) are three non-collinear points,

Then, the vector equation of the plane passing through these points is

$$\vec{r} \cdot (\vec{AB} \times \vec{AC}) = \vec{a} \cdot (\vec{AB} \times \vec{AC})$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given points,

$$A = (2, 2, -1)$$

$$B = (3, 4, 2)$$

$$C = (7, 0, 6)$$

Position vectors are given by,

$$\vec{a} = 2\hat{i} + 2\hat{j} - \hat{k}$$

$$\vec{b} = 3\hat{i} + 4\hat{j} + 2\hat{k}$$

$$\vec{c} = 7\hat{i} + 6\hat{k}$$

Now, vectors $\vec{AB}$ & $\vec{AC}$ are

$$\vec{AB} = \vec{b} - \vec{a}$$

$$= (3 - 2)\hat{i} + (4 - 2)\hat{j} + (2 + 1)\hat{k}$$

$$\therefore \vec{AB} = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{AC} = \vec{c} - \vec{a}$$

$$= (7 - 2)\hat{i} + (0 - 2)\hat{j} + (6 + 1)\hat{k}$$

$$\therefore \vec{AC} = 5\hat{i} - 2\hat{j} + 7\hat{k}$$

Therefore,

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 5 & -2 & 7 \end{vmatrix}$$

$$= \hat{i}(2 \times 7 - (-2) \times 3) - \hat{j}(1 \times 7 - 5 \times 3) + \hat{k}(1 \times (-2) - 5 \times 2)$$

$$= 20\hat{i} + 8\hat{j} - 12\hat{k}$$

Now,

$$\vec{a} \cdot (\vec{AB} \times \vec{AC}) = (2 \times 20) + (2 \times 8) + ((-1) \times (-12))$$

$$= 40 + 16 + 12$$

$$= 68$$

$$\therefore \vec{a} \cdot (\vec{AB} \times \vec{AC}) = 68 \dots\dots\dots \text{eq(1)}$$

And

$$\vec{r} \cdot (\vec{AB} \times \vec{AC}) = (x \times 20) + (y \times 8) + (z \times (-12))$$

$$= 20x + 8y - 12z$$

$$\therefore \vec{r} \cdot (\vec{AB} \times \vec{AC}) = 20x + 8y - 12z \dots\dots\dots \text{eq(2)}$$

Vector equation of the plane passing through points A, B & C is

$$\vec{r} \cdot (\vec{AB} \times \vec{AC}) = \vec{a} \cdot (\vec{AB} \times \vec{AC})$$

From eq(1) and eq(2)

$$20x + 8y - 12z = 68$$

This is $5x + 2y - 3z = 17$ vector equation of required plane.

(ii) Given Points :

$$A = (0, -1, -1)$$

$$B = (4, 5, 1)$$

$$C = (3, 9, 4)$$

To Find : Equation of plane passing through points A, B & C

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Vector :

If A and B be two points with position vectors $\vec{a}$ & $\vec{b}$ respectively, where

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{AB} = \vec{b} - \vec{a}$$

$$= (b_1 - a_1)\hat{i} + (b_2 - a_2)\hat{j} + (b_3 - a_3)\hat{k}$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

4) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

5) Equation of Plane :

If A = (a₁, a₂, a₃), B = (b₁, b₂, b₃), C = (c₁, c₂, c₃) are three non-collinear points,

Then, vector equation of the plane passing through these points is

$$\vec{r} \cdot (\vec{AB} \times \vec{AC}) = \vec{a} \cdot (\vec{AB} \times \vec{AC})$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given points,

$$A = (0, -1, -1)$$

$$B = (4, 5, 1)$$

$$C = (3, 9, 4)$$

Position vectors are given by,

$$\vec{a} = -\hat{j} - \hat{k}$$

$$\vec{b} = 4\hat{i} + 5\hat{j} + \hat{k}$$

$$\vec{c} = 3\hat{i} + 9\hat{j} + 4\hat{k}$$

Now, vectors $\vec{AB}$ & $\vec{AC}$ are

$$\vec{AB} = \vec{b} - \vec{a}$$

$$= (4 - 0)\hat{i} + (5 + 1)\hat{j} + (1 + 1)\hat{k}$$

$$\therefore \vec{AB} = 4\hat{i} + 6\hat{j} + 2\hat{k}$$

$$\vec{AC} = \vec{c} - \vec{a}$$

$$= (3 - 0)\hat{i} + (9 + 1)\hat{j} + (4 + 1)\hat{k}$$

$$\therefore \vec{AC} = 3\hat{i} + 10\hat{j} + 5\hat{k}$$

Therefore,

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 6 & 2 \\ 3 & 10 & 5 \end{vmatrix}$$

$$= \hat{i}(6 \times 5 - 10 \times 2) - \hat{j}(4 \times 5 - 2 \times 3) + \hat{k}(4 \times 10 - 3 \times 6)$$

$$= 10\hat{i} - 14\hat{j} + 22\hat{k}$$

Now,

$$\vec{a} \cdot (\vec{AB} \times \vec{AC}) = (0 \times 10) + ((-1) \times (-14)) + ((-1) \times 22)$$

$$= 0 + 14 - 22$$

$$= -8$$

$$\therefore \vec{a} \cdot (\vec{AB} \times \vec{AC}) = -8 \dots\dots\dots \text{eq(1)}$$

And

$$\vec{r} \cdot (\vec{AB} \times \vec{AC}) = (x \times 10) + (y \times (-14)) + (z \times 22)$$

$$= 10x - 14y + 22z$$

$$\therefore \vec{r} \cdot (\vec{AB} \times \vec{AC}) = 10x - 14y + 22z \dots\dots\dots \text{eq(2)}$$

Vector equation of plane passing through points A, B & C is

$$\vec{r} \cdot (\overline{AB} \times \overline{AC}) = \vec{a} \cdot (\overline{AB} \times \overline{AC})$$

From eq(1) and eq(2)

$$10x - 14y + 22z = -8$$

This is $5x - 7y + 11z = -4$ vector equation of required plane

(iii) Given Points :

$$A = (-2, 6, -6)$$

$$B = (-3, 10, 9)$$

$$C = (-5, 0, -6)$$

To Find : Equation of plane passing through points A, B & C

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Vector :

If A and B be two points with position vectors $\vec{a}$ & $\vec{b}$ respectively, where

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\overline{AB} = \vec{b} - \vec{a}$$

$$= (b_1 - a_1)\hat{i} + (b_2 - a_2)\hat{j} + (b_3 - a_3)\hat{k}$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

4) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

5) Equation of Plane :

If A = (a_1, a_2, a_3) , B = (b_1, b_2, b_3) , C = (c_1, c_2, c_3) are three non-collinear points,

Then, vector equation of the plane passing through these points is

$$\vec{r} \cdot (\overline{AB} \times \overline{AC}) = \vec{a} \cdot (\overline{AB} \times \overline{AC})$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given points,

$$A = (-2, 6, -6)$$

$$B = (-3, 10, 9)$$

$$C = (-5, 0, -6)$$

Position vectors are given by,

$$\vec{a} = -2\hat{i} + 6\hat{j} - 6\hat{k}$$

$$\vec{b} = -3\hat{i} + 10\hat{j} + 9\hat{k}$$

$$\vec{c} = -5\hat{i} - 6\hat{k}$$

Now, vectors $\overline{AB}$ & $\overline{AC}$ are

$$\overline{AB} = \vec{b} - \vec{a}$$

$$= (-3 + 2)\hat{i} + (10 - 6)\hat{j} + (9 + 6)\hat{k}$$

$$\therefore \overline{AB} = -\hat{i} + 4\hat{j} + 15\hat{k}$$

$$\overline{AC} = \vec{c} - \vec{a}$$

$$= (-5 + 2)\hat{i} + (0 - 6)\hat{j} + (-6 + 6)\hat{k}$$

$$\therefore \overline{AC} = -3\hat{i} - 6\hat{j} + 0\hat{k}$$

Therefore,

$$\overline{AB} \times \overline{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 4 & 15 \\ -3 & -6 & 0 \end{vmatrix}$$

$$= \hat{i}(4 \times 0 - (-6) \times 15) - \hat{j}((-1) \times 0 - (-3) \times 15) + \hat{k}((-1) \times (-6) - (-3) \times 4)$$

$$= 90\hat{i} - 45\hat{j} + 18\hat{k}$$

Now,

$$\vec{a} \cdot (\overline{AB} \times \overline{AC}) = ((-2) \times 90) + (6 \times (-45)) + ((-6) \times 18)$$

$$= -180 - 270 - 108$$

$$= -558$$

$$\therefore \vec{a} \cdot (\overline{AB} \times \overline{AC}) = -558 \dots\dots\dots \text{eq(1)}$$

And

$$\vec{r} \cdot (\overline{AB} \times \overline{AC}) = (x \times 90) + (y \times (-45)) + (z \times 18)$$

$$= 90x - 45y + 18z$$

$$\therefore \vec{r} \cdot (\overline{AB} \times \overline{AC}) = 90x - 45y + 18z \dots\dots\dots \text{eq(2)}$$

Vector equation of plane passing through points A, B & C is

$$\vec{r} \cdot (\overline{AB} \times \overline{AC}) = \vec{a} \cdot (\overline{AB} \times \overline{AC})$$

From eq(1) and eq(2)

$$90x - 45y + 18z = -558$$

This is $10x - 5y + 2z = -62$ vector equation of required plane

Question: 2

Show that the four

Solution:

Given Points :

$$A = (3, 2, -5)$$

$$B = (-1, 4, -3)$$

$$C = (-3, 8, -5)$$

$$D = (-3, 2, 1)$$

To Prove : Points A, B, C & D are coplanar.

To Find : Equation of plane passing through points A, B, C & D.

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Equation of line

If A and B are two points having position vectors $\vec{a}$ & $\vec{b}$ then equation of line passing through two points is given by,

$$\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

4) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

5) Coplanarity of two lines :

If two lines $\vec{r}_1 = \vec{a} + \lambda\vec{b}$ & $\vec{r}_2 = \vec{c} + \mu\vec{d}$ are coplanar then

$$\vec{a} \cdot (\vec{b} \times \vec{d}) = \vec{c} \cdot (\vec{b} \times \vec{d})$$

6) Equation of plane :

If two lines $\vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$ & $\vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$ are coplanar then equation of the plane containing them is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given points,

$$A = (3, 2, -5)$$

$$B = (-1, 4, -3)$$

$$C = (-3, 8, -5)$$

$$D = (-3, 2, 1)$$

Position vectors are given by,

$$\vec{a} = 3\hat{i} + 2\hat{j} - 5\hat{k}$$

$$\vec{b} = -1\hat{i} + 4\hat{j} - 3\hat{k}$$

$$\vec{c} = -3\hat{i} + 8\hat{j} - 5\hat{k}$$

$$\vec{d} = -3\hat{i} + 2\hat{j} + \hat{k}$$

Equation of line passing through points A & B is

$$\vec{r}_1 = \vec{a} + \lambda(\vec{b} - \vec{a})$$

$$\vec{b} - \vec{a} = (-1 - 3)\hat{i} + (4 - 2)\hat{j} + (-3 + 5)\hat{k}$$

$$= -4\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\therefore \vec{r}_1 = (3\hat{i} + 2\hat{j} - 5\hat{k}) + \lambda(-4\hat{i} + 2\hat{j} + 2\hat{k})$$

$$\text{Let, } \vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$$

Where,

$$\vec{a}_1 = 3\hat{i} + 2\hat{j} - 5\hat{k} \text{ \& } \vec{b}_1 = -4\hat{i} + 2\hat{j} + 2\hat{k}$$

And the equation of the line passing through points C & D is

$$\vec{r}_2 = \vec{c} + \mu(\vec{d} - \vec{c})$$

$$\vec{d} - \vec{c} = (-3 + 3)\hat{i} + (2 - 8)\hat{j} + (1 + 5)\hat{k}$$

$$= -6\hat{j} + 6\hat{k}$$

$$\therefore \vec{r}_2 = (-3\hat{i} + 8\hat{j} - 5\hat{k}) + \lambda(-6\hat{j} + 6\hat{k})$$

$$\text{Let, } \vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$$

Where,

$$\vec{a}_2 = -3\hat{i} + 8\hat{j} - 5\hat{k} \text{ \& } \vec{b}_2 = -6\hat{j} + 6\hat{k}$$

Now,

$$\vec{b_1} \times \vec{b_2} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -4 & 2 & 2 \\ 0 & -6 & 6 \end{vmatrix}$$

$$= \hat{i}(12 + 12) - \hat{j}(-24 - 0) + \hat{k}(24 + 0)$$

$$\therefore (\vec{b_1} \times \vec{b_2}) = 24\hat{i} + 24\hat{j} + 24\hat{k}$$

Therefore,

$$\vec{a_1} \cdot (\vec{b_1} \times \vec{b_2}) = (3 \times 24) + (2 \times 24) + ((-5) \times 24)$$

$$= 72 + 48 - 120$$

$$= 0$$

$$\therefore \vec{a_1} \cdot (\vec{b_1} \times \vec{b_2}) = 0 \dots\dots\dots \text{eq(1)}$$

And

$$\vec{a_2} \cdot (\vec{b_1} \times \vec{b_2}) = ((-3) \times 24) + (8 \times 24) + ((-5) \times 24)$$

$$= -72 + 192 - 120$$

$$= 0$$

$$\therefore \vec{a_2} \cdot (\vec{b_1} \times \vec{b_2}) = 0 \dots\dots\dots \text{eq(2)}$$

From eq(1) and eq(2)

$$\vec{a_1} \cdot (\vec{b_1} \times \vec{b_2}) = \vec{a_2} \cdot (\vec{b_1} \times \vec{b_2})$$

Hence lines $\vec{r_1}$ & $\vec{r_2}$ are coplanar

Therefore, points A, B, C & D are also coplanar.

As lines $\vec{r_1}$ & $\vec{r_2}$ are coplanar therefore equation of the plane passing through two lines containing four given points is

$$\vec{r} \cdot (\vec{b_1} \times \vec{b_2}) = \vec{a_1} \cdot (\vec{b_1} \times \vec{b_2})$$

Now,

$$\vec{r} \cdot (\vec{b_1} \times \vec{b_2}) = (x \times 24) + (y \times 24) + (z \times 24)$$

$$= 24x + 24y + 24z$$

From eq(1)

$$\vec{a_1} \cdot (\vec{b_1} \times \vec{b_2}) = 0$$

Therefore, equation of required plane is

$$24x + 24y + 24z = 0$$

$$x + y + z = 0$$

Question: 3

Show that the four

Solution:

Given Points :

$$A = (0, -1, 0)$$

$$B = (2, 1, -1)$$

$$C = (1, 1, 1)$$

$$D = (3, 3, 0)$$

To Prove : Points A, B, C & D are coplanar.

To Find : Equation of plane passing through points A, B, C & D.

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Equation of line

If A and B are two points having position vectors $\vec{a}$ & $\vec{b}$ then equation of line passing through two points is given by,

$$\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

4) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

5) Coplanarity of two lines :

If two lines $\vec{r}_1 = \vec{a} + \lambda\vec{b}$ & $\vec{r}_2 = \vec{c} + \mu\vec{d}$ are coplanar then

$$\vec{a} \cdot (\vec{b} \times \vec{d}) = \vec{c} \cdot (\vec{b} \times \vec{d})$$

6) Equation of plane :

If two lines $\vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$ & $\vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$ are coplanar then equation of the plane containing them is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given points,

$$A = (0, -1, 0)$$

$$B = (2, 1, -1)$$

$$C = (1, 1, 1)$$

$$D = (3, 3, 0)$$

Position vectors are given by,

$$\vec{a} = -\hat{j}$$

$$\vec{b} = 2\hat{i} + \hat{j} - \hat{k}$$

$$\vec{c} = \hat{i} + \hat{j} + \hat{k}$$

$$\vec{d} = 3\hat{i} + 3\hat{j}$$

Equation of line passing through points A & D is

$$\vec{r}_1 = \vec{a} + \lambda(\vec{d} - \vec{a})$$

$$\vec{d} - \vec{a} = (3 - 0)\hat{i} + (3 + 1)\hat{j} + (0 - 0)\hat{k}$$

$$= 3\hat{i} + 4\hat{j}$$

$$\therefore \vec{r}_1 = (-\hat{j}) + \lambda(3\hat{i} + 4\hat{j})$$

$$\text{Let, } \vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$$

Where,

$$\vec{a}_1 = -\hat{j} \text{ \& } \vec{b}_1 = 3\hat{i} + 4\hat{j}$$

And equation of line passing through points B & C is

$$\vec{r}_2 = \vec{b} + \mu(\vec{c} - \vec{b})$$

$$\vec{c} - \vec{b} = (1 - 2)\hat{i} + (1 - 1)\hat{j} + (1 + 1)\hat{k}$$

$$= -\hat{i} + 0\hat{j} + 2\hat{k}$$

$$\therefore \vec{r}_2 = (2\hat{i} + \hat{j} - \hat{k}) + \lambda(-\hat{i} + 2\hat{k})$$

$$\text{Let, } \vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$$

Where,

$$\vec{a}_2 = 2\hat{i} + \hat{j} - \hat{k} \text{ \& } \vec{b}_2 = -\hat{i} + 2\hat{k}$$

Now,

$$\begin{aligned}\bar{b}_1 \times \bar{b}_2 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 4 & 0 \\ -1 & 0 & 2 \end{vmatrix} \\ &= \hat{i}(8-0) - \hat{j}(6-0) + \hat{k}(0+4) \\ \therefore (\bar{b}_1 \times \bar{b}_2) &= 8\hat{i} - 6\hat{j} + 4\hat{k}\end{aligned}$$

Therefore,

$$\begin{aligned}\bar{a}_1 \cdot (\bar{b}_1 \times \bar{b}_2) &= (0 \times 8) + ((-1) \times (-6)) + (0 \times 4) \\ &= 0 + 6 + 0 \\ &= 6 \\ \therefore \bar{a}_1 \cdot (\bar{b}_1 \times \bar{b}_2) &= 6 \dots\dots\dots \text{eq(1)}\end{aligned}$$

And

$$\begin{aligned}\bar{a}_2 \cdot (\bar{b}_1 \times \bar{b}_2) &= (2 \times 8) + (1 \times (-6)) + ((-1) \times 4) \\ &= 16 - 6 - 4 \\ &= 6 \\ \therefore \bar{a}_2 \cdot (\bar{b}_1 \times \bar{b}_2) &= 6 \dots\dots\dots \text{eq(2)}\end{aligned}$$

From eq(1) and eq(2)

$$\bar{a}_1 \cdot (\bar{b}_1 \times \bar{b}_2) = \bar{a}_2 \cdot (\bar{b}_1 \times \bar{b}_2)$$

Hence lines $\bar{r}_1$ & $\bar{r}_2$ are coplanar

Therefore, points A, B, C & D are also coplanar.

As lines $\bar{r}_1$ & $\bar{r}_2$ are coplanar therefore equation of the plane passing through two lines containing four given points is

$$\bar{r} \cdot (\bar{b}_1 \times \bar{b}_2) = \bar{a}_1 \cdot (\bar{b}_1 \times \bar{b}_2)$$

Now,

$$\begin{aligned}\bar{r} \cdot (\bar{b}_1 \times \bar{b}_2) &= (x \times 8) + (y \times (-6)) + (z \times 4) \\ &= 8x - 6y + 4z\end{aligned}$$

From eq(1)

$$\bar{a}_1 \cdot (\bar{b}_1 \times \bar{b}_2) = 6$$

Therefore, equation of required plane is

$$8x - 6y + 4z = 6$$

$$4x - 3y + 2z = 3$$

Question: 4

Write the equatio

Solution:

Given :

X - intercept, a = 2

Y - intercept, b = - 4

Z - intercept, c = 5

To Find : Equation of plane

Formula :

If a, b & c are the intercepts made by plane on X, Y & Z axes respectively, then equation of the plane is given by,

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

$$\therefore \frac{x}{2} + \frac{y}{-4} + \frac{z}{5} = 1$$

Multiplying above equation throughout by 40,

$$\therefore \frac{40x}{2} + \frac{40y}{-4} + \frac{40z}{5} = 40$$

$$20x - 10y + 8z = 40$$

$$10x - 5y + 4z = 20$$

This the equation of the required plane.

Question: 5

Reduce the equati

Solution:

Given :

Equation of plane : $4x - 3y + 2z = 12$

To Find :

- 1) Equation of plane in intercept form
- 2) Intercepts made by the plane with the co-ordinate axes.

Formula :

$$\text{If } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

is the equation of a plane in intercept form then intercept made by it with co-ordinate axes are

X-intercept = a

Y-intercept = b

Z-intercept = c

Given the equation of plane:

$$4x - 3y + 2z = 12$$

Dividing the above equation throughout by 12

$$\therefore \frac{4x}{12} + \frac{-3y}{12} + \frac{2z}{12} = 1$$

$$\therefore \frac{x}{3} + \frac{y}{-4} + \frac{z}{6} = 1$$

This is the equation of a plane in intercept form.

Comparing the above equation with

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

We get,

$$a = 3$$

$$b = -4$$

$$c = 6$$

Therefore, intercepts made by plane with co-ordinate axes are

X-intercept = 3

Y-intercept = -4

Z-intercept = 6

Question: 6

Find the equation

Solution:

Equation of the plane making a, b & c intercepts with X, Y & Z axes respectively is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

But, the plane makes equal intercepts on the co-ordinate axes

Therefore, $a = b = c$

Therefore the equation of the plane is

$$\frac{x}{a} + \frac{y}{a} + \frac{z}{a} = 1$$

$$x + y + z = a$$

As plane passes through the point (2, -3, 7),

Substituting $x = 2$, $y = -3$ & $z = 7$

$$2 - 3 + 7 = a$$

Therefore, $a = 6$

Hence, required equation of plane is

$$x + y + z = 6$$

Question: 7

A plane meets the

Solution:

Given :

X-intercept = A

Y-intercept = B

Z-intercept = C

Centroid of $\Delta ABC = (1, -2, 3)$

To Find : Equation of a plane

Formulae :

1) Centroid Formula :

For ΔABC if co-ordinates of A, B & C are

A = (x_1, x_2, x_3)

B = (y_1, y_2, y_3)

C = (z_1, z_2, z_3)

Then co-ordinates of the centroid of ΔABC are

$$G = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$$

2) Equation of plane :

Equation of the plane making a, b & c intercepts with X, Y & Z axes respectively is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

As the plane makes intercepts at points A, B & C on X, Y & Z axes respectively, let co-ordinates of A, B, C be

A = $(a, 0, 0)$

B = $(0, b, 0)$

C = $(0, 0, c)$

By centroid formula,

The centroid of ΔABC is given by

$$G = \left(\frac{a + 0 + 0}{3}, \frac{0 + b + 0}{3}, \frac{0 + 0 + c}{3} \right)$$

$$G = \left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3} \right)$$

But, Centroid of $\Delta ABC = (1, -2, 3)$ given

$$\therefore \frac{a}{3} = 1, \frac{b}{3} = -2, \frac{c}{3} = 3$$

Therefore, a = 3, b = -6, c = 9

Therefore,

X-intercept = a = 3

Y-intercept = b = -6

Z-intercept = c = 9

Therefore, equation of required plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

$$\therefore \frac{x}{3} + \frac{y}{-6} + \frac{z}{9} = 1$$

Question: 8

Find the Cartesian

Solution:

Given :

A = $(1, 2, 3)$

Direction ratios of perpendicular vector = $(2, 3, -4)$

To Find : Equation of a plane

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

If a plane is passing through point A, then the equation of a plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{a}$ = *position vector of A*

$\vec{n}$ = *vector perpendicular to the plane*

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For point A = (1, 2, 3), position vector is

$$\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$$

Vector perpendicular to the plane with direction ratios (2, 3, -4) is

$$\vec{n} = 2\hat{i} + 3\hat{j} - 4\hat{k}$$

$$\text{Now, } \vec{a} \cdot \vec{n} = (1 \times 2) + (2 \times 3) + (3 \times (-4))$$

$$= 2 + 6 - 12$$

$$= -4$$

Equation of the plane passing through point A and perpendicular to vector $\vec{n}$ is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore \vec{r} \cdot (2\hat{i} + 3\hat{j} - 4\hat{k}) = -4$$

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore \vec{r} \cdot (2\hat{i} + 3\hat{j} - 4\hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + 3\hat{j} - 4\hat{k})$$

$$= 2x + 3y - 4z$$

Therefore, equation of the plane is

$$2x + 3y - 4z = -4$$

Or

$$2x + 3y - 4z + 4 = 0$$

Question: 9

If O is the origin

Solution:

Given :

$$P = (1, 2, -3)$$

$$O = (0, 0, 0)$$

$$\vec{n} = \overrightarrow{OP}$$

To Find : Equation of a plane

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Vector :

If A and B be two points with position vectors $\vec{a}$ & $\vec{b}$ respectively, where

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{AB} = \vec{b} - \vec{a}$$

$$= (b_1 - a_1)\hat{i} + (b_2 - a_2)\hat{j} + (b_3 - a_3)\hat{k}$$

3) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

4) Equation of plane :

If a plane is passing through point A, then the equation of a plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{a}$ = *position vector of A*

$\vec{n}$ = *vector perpendicular to the plane*

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For points,

$$P = (1, 2, -3)$$

$$O = (0, 0, 0)$$

Position vectors are

$$\vec{p} = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{o} = 0\hat{i} + 0\hat{j} + 0\hat{k}$$

Vector

$$\vec{OP} = \vec{p} - \vec{o}$$

$$= (1 - 0)\hat{i} + (2 - 0)\hat{j} + (3 - 0)\hat{k}$$

$$\therefore \vec{OP} = \hat{i} + 2\hat{j} + 3\hat{k}$$

Now,

$$\vec{p} \cdot \vec{OP} = (1 \times 1) + (2 \times 2) + (3 \times 3)$$

$$= 1 + 4 + 9$$

$$= 14$$

And

$$\vec{r} \cdot \vec{OP} = (x \times 1) + (y \times 2) + (z \times 3)$$

$$= x + 2y + 3z$$

Equation of the plane passing through point A and perpendicular to the vector $\vec{n}$ is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\text{But, } \vec{n} = \vec{OP}$$

Therefore, the equation of the plane is

$$\vec{r} \cdot \vec{OP} = \vec{p} \cdot \vec{OP}$$

$$x + 2y + 3z = 14$$

$$x + 2y + 3z - 14 = 0$$

Exercise : 28B

Question: 1

Find the vector a

Solution:

Given :

$$d = 5$$

$$\hat{n} = \hat{k}$$

To Find : Equation of a plane

Formulae :

1) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

2) Equation of plane :

Equation of plane which is at a distance of 5 units from the origin and having $\hat{n}$ as a unit vector normal to it is

$$\vec{r} \cdot \hat{n} = d$$

$$\text{Where, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given d = 5 and $\hat{n} = \hat{k}$,

Equation of plane is

$$\vec{r} \cdot \hat{n} = d$$

$$\therefore \vec{r} \cdot \hat{k} = 5$$

This is a vector equation of the plane

Now,

$$\vec{r} \cdot \hat{k} = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot \hat{k}$$

$$= (x \times 0) + (y \times 0) + (z \times 1)$$

$$= z$$

$$\therefore \vec{r} \cdot \hat{k} = z$$

Therefore, the equation of the plane is

This is - the Cartesian $z = 5$ equation of the plane.

Question: 2

Find the vector a

Solution:

Given :

$$d = 7$$

$$\vec{n} = 3\hat{i} + 5\hat{j} - 6\hat{k}$$

To Find : Equation of plane

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

Equation of plane which is at a distance of 5 units from the origin and having $\hat{n}$ as a unit vector normal to it is

$$\vec{r} \cdot \hat{n} = d$$

Where, $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

For given normal vector

$$\vec{n} = 3\hat{i} + 5\hat{j} - 6\hat{k}$$

Unit vector normal to the plane is

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\therefore \hat{n} = \frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{3^2 + 5^2 + (-6)^2}}$$

$$\therefore \hat{n} = \frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{9 + 25 + 36}}$$

$$\therefore \hat{n} = \frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{70}}$$

Equation of the plane is

$$\vec{r} \cdot \hat{n} = d$$

$$\therefore \vec{r} \cdot \left(\frac{3\hat{i} + 5\hat{j} - 6\hat{k}}{\sqrt{70}} \right) = 7$$

$$\therefore \vec{r} \cdot (3\hat{i} + 5\hat{j} - 6\hat{k}) = 7\sqrt{70}$$

This is a vector equation of the plane.

Now,

$$\vec{r} \cdot (3\hat{i} + 5\hat{j} - 6\hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + 5\hat{j} - 6\hat{k})$$

$$= (x \times 3) + (y \times 5) + (z \times (-6))$$

$$= 3x + 5y - 6z$$

Therefore equation of the plane is

$$3x + 5y - 6z = 7\sqrt{70}$$

This is the Cartesian equation of the plane.

Question: 3

Find the vector a

Solution:

Given :

$$d = \frac{6}{\sqrt{29}}$$

$$\vec{n} = 2\hat{i} - 3\hat{j} + 4\hat{k}$$

To Find : Equation of a plane

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then the unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

Equation of plane which is at a distance of 5 units from the origin and having $\hat{n}$ as a unit vector normal to it is

$$\vec{r} \cdot \hat{n} = d$$

$$\text{Where, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given normal vector

$$\vec{n} = 2\hat{i} - 3\hat{j} + 4\hat{k}$$

Unit vector normal to the plane is

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\therefore \hat{n} = \frac{2\hat{i} - 3\hat{j} + 4\hat{k}}{\sqrt{2^2 + (-3)^2 + 4^2}}$$

$$\therefore \hat{n} = \frac{2\hat{i} - 3\hat{j} + 4\hat{k}}{\sqrt{4 + 9 + 16}}$$

$$\therefore \hat{n} = \frac{2\hat{i} - 3\hat{j} + 4\hat{k}}{\sqrt{29}}$$

Equation of the plane is

$$\vec{r} \cdot \hat{n} = d$$

$$\therefore \vec{r} \cdot \left(\frac{2\hat{i} - 3\hat{j} + 4\hat{k}}{\sqrt{29}} \right) = \frac{6}{\sqrt{29}}$$

$$\therefore \vec{r} \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 6$$

This is a vector equation of the plane.

Now,

$$\vec{r} \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} - 3\hat{j} + 4\hat{k})$$

$$= (x \times 2) + (y \times (-3)) + (z \times 4)$$

$$= 2x - 3y + 4z$$

Therefore equation of the plane is

$$2x - 3y + 4z = 6$$

This is the Cartesian equation of the plane.

Question: 4

Find the vector a

Solution:

Given :

$$d = 6$$

direction ratios of $\vec{n}$ are (2, -1, -2)

$$\therefore \vec{n} = 2\hat{i} - \hat{j} - 2\hat{k}$$

To Find : Equation of plane

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then the unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

Equation of plane which is at a distance of 5 units from the origin and having $\hat{n}$ as a unit vector normal to it is

$$\vec{r} \cdot \hat{n} = d$$

$$\text{Where, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For given normal vector

$$\vec{n} = 2\hat{i} - \hat{j} - 2\hat{k}$$

Unit vector normal to the plane is

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\therefore \hat{n} = \frac{2\hat{i} - \hat{j} - 2\hat{k}}{\sqrt{2^2 + (-1)^2 + (-2)^2}}$$

$$\therefore \hat{n} = \frac{2\hat{i} - \hat{j} - 2\hat{k}}{\sqrt{4 + 1 + 4}}$$

$$\therefore \hat{n} = \frac{2\hat{i} - \hat{j} - 2\hat{k}}{\sqrt{9}}$$

$$\therefore \hat{n} = \frac{2\hat{i} - \hat{j} - 2\hat{k}}{3}$$

Equation of the plane is

$$\vec{r} \cdot \hat{n} = d$$

$$\therefore \vec{r} \cdot \left(\frac{2\hat{i} - \hat{j} - 2\hat{k}}{3} \right) = 6$$

$$\therefore \vec{r} \cdot (2\hat{i} - \hat{j} - 2\hat{k}) = 18$$

This is vector equation of the plane.

Now,

$$\vec{r} \cdot (2\hat{i} - \hat{j} - 2\hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} - \hat{j} - 2\hat{k})$$

$$= (x \times 2) + (y \times (-1)) + (z \times (-2))$$

$$= 2x - y - 2z$$

Therefore equation of the plane is

$$2x - y - 2z = 18$$

This is Cartesian equation of the plane.

Question: 5

Find the vector,

Solution:

Given :

$$A = (1, 4, 6)$$

$$\vec{n} = \hat{i} - 2\hat{j} + \hat{k}$$

To Find : Equation of plane.

Formulae :

1) Position Vector :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

Equation of plane passing through point A and having $\vec{n}$ as a unit vector normal to it is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\text{Where, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Position vector of point A = (1, 4, 6) is

$$\vec{a} = \hat{i} + 4\hat{j} + 6\hat{k}$$

Now,

$$\vec{a} \cdot \vec{n} = (\hat{i} + 4\hat{j} + 6\hat{k}) \cdot (\hat{i} - 2\hat{j} + \hat{k})$$

$$= (1 \times 1) + (4 \times (-2)) + (6 \times 1)$$

$$= 1 - 8 + 6$$

$$= -1$$

Equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore \vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = -1$$

This is vector equation of the plane.

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Therefore

$$\vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - 2\hat{j} + \hat{k})$$

$$= (x \times 1) + (y \times (-2)) + (z \times 1)$$

$$= x - 2y + z$$

Therefore equation of the plane is

$$x - 2y + z = -1$$

This is Cartesian equation of the plane.

Question: 6

Find the length of

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (3\hat{i} - 12\hat{j} - 4\hat{k}) + 39 = 0$$

To Find :

i) Length of perpendicular = d

ii) Unit normal vector = $\hat{n}$

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Length of perpendicular :

The length of the perpendicular from the origin to the plane

$\vec{r} \cdot \vec{n} = p$ is given by,

$$d = \frac{p}{|\vec{n}|}$$

Given the equation of the plane is

$$\vec{r} \cdot (3\hat{i} - 12\hat{j} - 4\hat{k}) + 39 = 0$$

$$\therefore \vec{r} \cdot (3\hat{i} - 12\hat{j} - 4\hat{k}) = -39$$

$$\therefore \vec{r} \cdot (-3\hat{i} + 12\hat{j} + 4\hat{k}) = 39$$

Comparing the above equation with

$$\vec{r} \cdot \vec{n} = p$$

We get,

$$\vec{n} = -3\hat{i} + 12\hat{j} + 4\hat{k} \text{ \& } p = 39$$

Therefore,

$$\begin{aligned}
 |\vec{n}| &= \sqrt{(-3)^2 + 12^2 + 4^2} \\
 &= \sqrt{9 + 144 + 16} \\
 &= \sqrt{169} \\
 &= 13
 \end{aligned}$$

The length of the perpendicular from the origin to the given plane is

$$\begin{aligned}
 d &= \frac{p}{|\vec{n}|} \\
 \therefore d &= \frac{39}{13} \\
 \therefore d &= 3
 \end{aligned}$$

Vector normal to the plane is

$$\vec{n} = -3\hat{i} + 12\hat{j} + 4\hat{k}$$

Therefore, the unit vector normal to the plane is

$$\begin{aligned}
 \hat{n} &= \frac{\vec{n}}{|\vec{n}|} \\
 \therefore \hat{n} &= \frac{-3\hat{i} + 12\hat{j} + 4\hat{k}}{13} \\
 \therefore \hat{n} &= \frac{-3\hat{i}}{13} + \frac{12\hat{j}}{13} + \frac{4\hat{k}}{13}
 \end{aligned}$$

Question: 7

Find the Cartesia

Solution:

Given :

Vector equation of the plane is

$$\vec{r} \cdot (3\hat{i} + 5\hat{j} - 9\hat{k}) = 8$$

To Find : Cartesian equation of the given plane.

Formulae :

1) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

Given the equation of the plane is

$$\vec{r} \cdot (3\hat{i} + 5\hat{j} - 9\hat{k}) = 8$$

Here,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\begin{aligned}
 \therefore \vec{r} \cdot (3\hat{i} + 5\hat{j} - 9\hat{k}) &= (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + 5\hat{j} - 9\hat{k}) \\
 &= (x \times 3) + (y \times 5) + (z \times (-9)) \\
 &= 3x + 5y - 9z
 \end{aligned}$$

Therefore equation of the plane is

$$3x + 5y - 9z = 8$$

This is the Cartesian equation of the given plane.

Question: 8

Find the vector e

Solution:

Given :

Cartesian equation of the plane is

$$5x - 7y + 2z + 4 = 0$$

To Find : Vector equation of the given plane.

Formulae :

1) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

Given the equation of the plane is

$$5x - 7y + 2z + 4 = 0$$

$$\Rightarrow 5x - 7y + 2z = -4$$

The term $(5x - 7y + 2z)$ can be written as

$$(5x - 7y + 2z) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (5\hat{i} - 7\hat{j} + 2\hat{k})$$

$$\text{But } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore (5x - 7y + 2z) = \vec{r} \cdot (5\hat{i} - 7\hat{j} + 2\hat{k})$$

Therefore the equation of the plane is

$$\vec{r} \cdot (5\hat{i} - 7\hat{j} + 2\hat{k}) = -4$$

or

$$\vec{r} \cdot (-5\hat{i} + 7\hat{j} - 2\hat{k}) = 4$$

This is Vector equation of the given plane.

Question: 9

Find a unit vector

Solution:

Given :

$$\text{Equation of plane : } x - 2y + 2z = 6$$

To Find : unit normal vector = $\hat{n}$

Formula :

Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then the unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

From the given equation of a plane

$$x - 2y + 2z = 6$$

direction ratios of vector normal to the plane are (1, -2, 2).

Therefore, the equation of normal vector is

$$\vec{n} = \hat{i} - 2\hat{j} + 2\hat{k}$$

Therefore unit normal vector is given by

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\therefore \hat{n} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{1^2 + (-2)^2 + 2^2}}$$

$$\therefore \hat{n} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{1 + 4 + 4}}$$

$$\therefore \hat{n} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{9}}$$

$$\therefore \hat{n} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3}$$

$$\therefore \hat{n} = \frac{\hat{i}}{3} - \frac{2\hat{j}}{3} + \frac{2\hat{k}}{3}$$

Question: 10

Find the direction

Solution:

Given :

Equation of plane : $3x - 6y + 2z = 7$

To Find : Direction cosines of the normal, i.e. $l, m \& n$

Formula :

1) Direction cosines :

If a, b & c are direction ratios of the vector, then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

For the given equation of a plane

$$3x - 6y + 2z = 7$$

Direction ratios of normal vector are (3, -6, 2)

$$\begin{aligned}\sqrt{a^2 + b^2 + c^2} &= \sqrt{3^2 + (-6)^2 + 2^2} \\ &= \sqrt{9 + 36 + 4} \\ &= \sqrt{49} \\ &= \pm 7\end{aligned}$$

Therefore, direction cosines are

$$\begin{aligned}l &= \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \pm \frac{3}{7} \\ m &= \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \mp \frac{6}{7} \\ n &= \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \pm \frac{2}{7} \\ (l, m, n) &= \pm \left(\frac{3}{7}, -\frac{6}{7}, \frac{2}{7} \right)\end{aligned}$$

Question: 11

For each of the f

Solution:

(i) $2x + 3y - z = 5$

Given :

Equation of plane : $2x + 3y - z = 5$

To Find :

Direction cosines of the normal i.e. $l, m \& n$

Distance of the plane from the origin = d

Formulae :

1) Direction cosines :

If a, b & c are direction ratios of the vector then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

2) The distance of the plane from the origin :

Distance of the plane from the origin is given by,

$$d = \frac{p}{|\vec{n}|}$$

For the given equation of plane

$$2x + 3y - z = 5$$

Direction ratios of normal vector are (2, 3, -1)

Therefore, equation of normal vector is

$$\vec{n} = 2\hat{i} + 3\hat{j} - \hat{k}$$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{2^2 + 3^2 + (-1)^2}$$

$$= \sqrt{4 + 9 + 1}$$

$$= \sqrt{14}$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{2}{\sqrt{14}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{3}{\sqrt{14}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{-1}{\sqrt{14}}$$

$$(l, m, n) = \left(\frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}, \frac{-1}{\sqrt{14}} \right)$$

Now, the distance of the plane from the origin is

$$d = \frac{p}{|\vec{n}|}$$

$$\therefore d = \frac{5}{\sqrt{14}}$$

(ii) Given :

Equation of plane : $z = 3$

To Find :

Direction cosines of the normal, i.e. l, m & n

The distance of the plane from the origin = d

Formulae :

3) Direction cosines :

If a, b & c are direction ratios of the vector, then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

4) The distance of the plane from the origin :

Distance of the plane from the origin is given by,

$$d = \frac{p}{|\vec{n}|}$$

For the given equation of a plane

$$z = 3$$

Direction ratios of normal vector are (0, 0, 1)

Therefore, equation of normal vector is

$$\vec{n} = \hat{k}$$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{0^2 + 0^2 + 1^2}$$

$$= \sqrt{1}$$

$$= 1$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{1} = 0$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{1} = 0$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{1}{1} = 1$$

$$(l, m, n) = (0, 0, 1)$$

Now, the distance of the plane from the origin is

$$d = \frac{p}{|\vec{n}|}$$

$$\therefore d = \frac{3}{1}$$

$$\therefore d = 3$$

(iii) Given :

Equation of plane : $3y + 5 = 0$

To Find :

Direction cosines of the normal, i.e. l, m & n

The distance of the plane from the origin = d

Formulae :

1) Direction cosines :

If a, b & c are direction ratios of the vector, then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

2) Distance of the plane from the origin :

Distance of the plane from the origin is given by,

$$d = \frac{p}{|\vec{n}|}$$

For the given equation of a plane

$$3y + 5 = 0$$

$$\Rightarrow -3y = 5$$

Direction ratios of normal vector are (0, -3, 0)

Therefore, equation of normal vector is

$$\vec{n} = -3\hat{j}$$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{0^2 + (-3)^2 + 0^2}$$

$$= \sqrt{9}$$

$$= 3$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{3} = 0$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{-3}{3} = -1$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{3} = 0$$

$$(l, m, n) = (0, -1, 0)$$

Now, distance of the plane from the origin is

$$d = \frac{p}{|\vec{n}|}$$

$$\therefore d = \frac{5}{3}$$

Question: 12

Find the vector a

Solution:

Given :

$$A = (2, -1, 1)$$

Direction ratios of perpendicular vector = (4, 2, -3)

To Find : Equation of a plane

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

If a plane is passing through point A, then the equation of a plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{a}$ = *position vector of A*

$\vec{n}$ = *vector perpendicular to the plane*

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For point A = (2, -1, 1), position vector is

$$\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$$

Vector perpendicular to the plane with direction ratios (4, 2, -3) is

$$\vec{n} = 4\hat{i} + 2\hat{j} - 3\hat{k}$$

$$\text{Now, } \vec{a} \cdot \vec{n} = (2 \times 4) + ((-1) \times 2) + (1 \times (-3))$$

$$= 8 - 2 - 3$$

$$= 3$$

Equation of the plane passing through point A and perpendicular to vector $\vec{n}$ is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore \vec{r} \cdot (4\hat{i} + 2\hat{j} - 3\hat{k}) = 3$$

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore \vec{r} \cdot (4\hat{i} + 2\hat{j} - 3\hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (4\hat{i} + 2\hat{j} - 3\hat{k})$$

$$= 4x + 2y - 3z$$

Therefore, the equation of the plane is

$$4x + 2y - 3z = 3$$

Or

$$4x + 2y - 3z - 3 = 0$$

Question: 13

Find the coordina

Solution:

$$(i) 2x + 3y + 4z - 12 = 0$$

Given :

$$\text{Equation of plane : } 2x + 3y + 4z + 12 = 0$$

To Find :

coordinates of the foot of the perpendicular

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

From the given equation of the plane

$$2x + 3y + 4z - 12 = 0$$

$$\Rightarrow 2x + 3y + 4z = 12$$

Direction ratios of the vector normal to the plane are (2, 3, 4)

Let, P = (x, y, z) be the foot of perpendicular drawn from origin to the plane.

Therefore perpendicular drawn is $\overline{OP}$.

$$\therefore \overline{OP} = x\hat{i} + y\hat{j} + z\hat{k}$$

Let direction ratios of $\overline{OP}$ are (x, y, z)

As normal vector and $\overline{OP}$ are parallel

$$\therefore \frac{x}{2} = \frac{y}{3} = \frac{z}{4} = k(\text{say})$$

$$\Rightarrow x = 2k, y = 3k, z = 4k$$

As point P lies on the plane, we can write

$$2(2k) + 3(3k) + 4(4k) = 12$$

$$\Rightarrow 4k + 9k + 16k = 12$$

$$\Rightarrow 29k = 12$$

$$\therefore k = \frac{12}{29}$$

$$\therefore x = 2k = \frac{24}{29},$$

$$y = 3k = \frac{36}{29}$$

$$z = 4k = \frac{48}{29}$$

Therefore co-ordinates of the foot of perpendicular are

$$P(x, y, z) = \left(\frac{24}{29}, \frac{36}{29}, \frac{48}{29}\right)$$

$$P = \left(\frac{24}{29}, \frac{36}{29}, \frac{48}{29}\right)$$

(ii) Given :

$$\text{Equation of plane : } 5y + 8 = 0$$

To Find :

coordinates of the foot of the perpendicular

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

From the given equation of the plane

$$5y + 8 = 0$$

$$\Rightarrow 5y = -8$$

Direction ratios of the vector normal to the plane are (0, 5, 0)

Let, P = (x, y, z) be the foot of perpendicular drawn from origin to the plane.

Therefore perpendicular drawn is $\overline{OP}$.

$$\therefore \overline{OP} = x\hat{i} + y\hat{j} + z\hat{k}$$

Let direction ratios of $\overline{OP}$ are (x, y, z)

As normal vector and $\overline{OP}$ are parallel

$$\therefore \frac{0}{x} = \frac{5}{y} = \frac{0}{z} = \frac{1}{k}(\text{say})$$

$$\Rightarrow x = 0, y = 5k, z = 0$$

As point P lies on the plane, we can write

$$5(5k) = -8$$

$$\Rightarrow 25k = -8$$

$$\therefore k = \frac{-8}{25}$$

$$\therefore x = 0,$$

$$y = 5k = 5 \times \frac{-8}{25} = \frac{-8}{5}$$

$$z = 0$$

Therefore co-ordinates of the foot of perpendicular are

$$P(x, y, z) = \left(0, \frac{-8}{5}, 0\right)$$

$$P = \left(0, \frac{-8}{5}, 0\right)$$

Question: 14

Find the length a

Solution:

Given :

Equation of plane : $3x - y - z = 7$

$$A = (2, 3, 7)$$

To Find :

- i) Length of perpendicular = d
- ii) coordinates of the foot of the perpendicular

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Length of perpendicular :

The length of the perpendicular from point A with position vector $\vec{a}$ to the plane is given by,

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

Given equation of the plane is

$$3x - y - z = 7 \dots\dots\dots \text{eq(1)}$$

Therefore direction ratios of normal vector of the plane are

$$(3, -1, -1)$$

Therefore normal vector of the plane is

$$\vec{n} = 3\hat{i} - \hat{j} - \hat{k}$$

$$\therefore |\vec{n}| = \sqrt{3^2 + (-1)^2 + (-1)^2}$$

$$= \sqrt{9 + 1 + 1}$$

$$= \sqrt{11}$$

From eq(1), p = 7

Given point A = (2, 3, 7)

Position vector of A is

$$\vec{a} = 2\hat{i} + 3\hat{j} + 7\hat{k}$$

Now,

$$\vec{a} \cdot \vec{n} = (2\hat{i} + 3\hat{j} + 7\hat{k}) \cdot (3\hat{i} - \hat{j} - \hat{k})$$

$$= (2 \times 3) + (3 \times (-1)) + (7 \times (-1))$$

$$= 6 - 3 - 7$$

$$= -4$$

Length of the perpendicular from point A to the plane is

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

$$\therefore d = \frac{|-4 - 7|}{\sqrt{11}}$$

$$\therefore d = \frac{11}{\sqrt{11}}$$

$$\therefore d = \sqrt{11}$$

Let P be the foot of perpendicular drawn from point A to the given plane,

Let P = (x, y, z)

$$\vec{AP} = (x - 2)\hat{i} + (y - 3)\hat{j} + (z - 7)\hat{k}$$

As normal vector and $\vec{AP}$ are parallel

$$\therefore \frac{x - 2}{3} = \frac{y - 3}{-1} = \frac{z - 7}{-1} = k(\text{say})$$

$$\Rightarrow x = 3k + 2, y = -k + 3, z = -k + 7$$

As point P lies on the plane, we can write

$$3(3k + 2) - (-k + 3) - (-k + 7) = 7$$

$$\Rightarrow 9k + 6 + k - 3 + k - 7 = 7$$

$$\Rightarrow 11k = 11$$

$$\therefore k = 1$$

$$\therefore x = 3k + 2 = 5,$$

$$y = -k + 3 = 2$$

$$z = -k + 7 = 6$$

Therefore co-ordinates of the foot of perpendicular are

$$P(x, y, z) = (5, 2, 6)$$

$$P = (5, 2, 6)$$

Question: 15

Find the length a

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (2\hat{i} - 2\hat{j} + 4\hat{k}) + 5 = 0$$

$$A = (1, 1, 2)$$

To Find :

i) Length of perpendicular = d

ii) coordinates of the foot of the perpendicular

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Length of perpendicular :

The length of the perpendicular from point A with position vector $\vec{a}$ to the plane is given by,

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

Given equation of the plane is

$$\vec{r} \cdot (2\hat{i} - 2\hat{j} + 4\hat{k}) + 5 = 0 \dots\dots\dots \text{eq(1)}$$

$$\therefore \vec{r} \cdot (2\hat{i} - 2\hat{j} + 4\hat{k}) = -5$$

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Therefore equation of plane is

$$2x - 2y + 4z = -5 \dots\dots\dots \text{eq(2)}$$

From eq(1) normal vector of the plane is

$$\vec{n} = 2\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{2^2 + (-2)^2 + 4^2}$$

$$= \sqrt{4 + 4 + 16}$$

$$= \sqrt{24}$$

From eq(1), p = -5

Given point A = (1, 1, 2)

Position vector of A is

$$\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$$

Now,

$$\vec{a} \cdot \vec{n} = (\hat{i} + \hat{j} + 2\hat{k}) \cdot (2\hat{i} - 2\hat{j} + 4\hat{k})$$

$$= (1 \times 2) + (1 \times (-2)) + (2 \times 4)$$

$$= 2 - 2 + 8$$

$$= 8$$

Length of the perpendicular from point A to the plane is

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

$$\therefore d = \frac{|8 + 5|}{\sqrt{24}}$$

$$\therefore d = \frac{13}{\sqrt{24}}$$

$$\therefore d = \frac{13\sqrt{6}}{\sqrt{24} \cdot \sqrt{6}}$$

$$\therefore d = \frac{13\sqrt{6}}{\sqrt{144}}$$

$$\therefore d = \frac{13\sqrt{6}}{12}$$

Let P be the foot of perpendicular drawn from point A to the given plane,

Let P = (x, y, z)

$$\vec{AP} = (x - 1)\hat{i} + (y - 1)\hat{j} + (z - 2)\hat{k}$$

As normal vector and $\vec{AP}$ are parallel

$$\therefore \frac{x - 1}{2} = \frac{y - 1}{-2} = \frac{z - 2}{4} = k(\text{say})$$

$$\Rightarrow x = 2k + 1, y = -2k + 1, z = 4k + 2$$

As point P lies on the plane, we can write

$$2(2k + 1) - 2(-2k + 1) + 4(4k + 2) = -5$$

$$= 4k + 2 + 4k - 2 + 16k + 8 = -5$$

$$\Rightarrow 24k = -13$$

$$\therefore k = \frac{-13}{24}$$

$$\therefore x = 2\left(\frac{-13}{24}\right) + 1 = \frac{-1}{12},$$

$$y = -2\left(\frac{-13}{24}\right) + 1 = \frac{25}{12}$$

$$z = 4\left(\frac{-13}{24}\right) + 2 = \frac{-1}{6}$$

Therefore co-ordinates of the foot of perpendicular are

$$P(x, y, z) = \left(\frac{-1}{12}, \frac{25}{12}, \frac{-1}{6}\right)$$

$$P \equiv \left(\frac{-1}{12}, \frac{25}{12}, \frac{-1}{6}\right)$$

Question: 16

From the point P(

Solution:

Given :

Equation of plane : $2x + y - 2z + 3 = 0$

P = (1, 2, 4)

To Find :

- i) Equation of perpendicular
- ii) Length of perpendicular = d
- iii) coordinates of the foot of the perpendicular

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

Where, $|\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

2) Length of perpendicular :

The length of the perpendicular from point A with position vector $\vec{a}$ to the plane is given by,

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

Given equation of the plane is

$$2x + y - 2z + 3 = 0$$

$$\Rightarrow 2x + y - 2z = -3 \dots\dots\dots \text{eq(1)}$$

From eq(1) direction ratios of normal vector of the plane are

(2, 1, -2)

Therefore, equation of normal vector is

$$\vec{n} = 2\hat{i} + \hat{j} - 2\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{2^2 + 1^2 + (-2)^2}$$

$$= \sqrt{4 + 1 + 4}$$

$$= \sqrt{9}$$

$$= 3$$

From eq(1), p = -3

Given point P = (1, 2, 4)

Position vector of A is

$$\vec{p} = \hat{i} + 2\hat{j} + 4\hat{k}$$

$$\text{Here, } \vec{a} = \vec{p}$$

Now,

$$\begin{aligned}\therefore \vec{a} \cdot \vec{n} &= (\hat{i} + 2\hat{j} + 4\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k}) \\ &= (1 \times 2) + (2 \times 1) + (4 \times (-2)) \\ &= 2 + 2 - 8 \\ &= -4\end{aligned}$$

Length of the perpendicular from point A to the plane is

$$\begin{aligned}d &= \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|} \\ \therefore d &= \frac{|-4 + 3|}{3} \\ \therefore d &= \frac{1}{3}\end{aligned}$$

Let Q be the foot of perpendicular drawn from point P to the given plane,

$$\text{Let } Q = (x, y, z)$$

$$\overrightarrow{PQ} = (x - 1)\hat{i} + (y - 2)\hat{j} + (z - 4)\hat{k}$$

As normal vector and $\overrightarrow{PQ}$ are parallel, we can write,

$$\therefore \frac{x - 1}{2} = \frac{y - 2}{1} = \frac{z - 4}{-2}$$

This is the equation of perpendicular.

$$\therefore \frac{x - 1}{2} = \frac{y - 2}{1} = \frac{z - 4}{-2} = k(\text{say})$$

$$\Rightarrow x = 2k + 1, y = k + 2, z = -2k + 4$$

As point Q lies on the plane, we can write

$$\begin{aligned}2(2k + 1) + (k + 2) - 2(-2k + 4) &= -3 \\ \Rightarrow 4k + 2 + k + 2 + 4k - 8 &= -3 \\ \Rightarrow 9k &= 1\end{aligned}$$

$$\therefore k = \frac{1}{9}$$

$$\therefore x = 2\left(\frac{1}{9}\right) + 1 = \frac{11}{9},$$

$$y = \frac{1}{9} + 2 = \frac{19}{9}$$

$$z = -2\left(\frac{1}{9}\right) + 4 = \frac{34}{9}$$

Therefore co-ordinates of the foot of perpendicular are

$$Q(x, y, z) = \left(\frac{11}{9}, \frac{19}{9}, \frac{34}{9}\right)$$

$$Q \equiv \left(\frac{11}{9}, \frac{19}{9}, \frac{34}{9}\right)$$

Question: 17

Find the coordina

Solution:

Given :

$$\text{Equation of plane : } 2x - y + z + 1 = 0$$

$$P = (3, 2, 1)$$

To Find :

- i) Length of perpendicular = d
- ii) coordinates of the foot of the perpendicular
- iii) Image of point P in the plane.

Formulae :

1) Unit Vector :

$$\text{Let } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k} \text{ be any vector}$$

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Length of perpendicular :

The length of the perpendicular from point A with position vector $\vec{a}$ to the plane is given by,

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

Given equation of the plane is

$$2x - y + z + 1 = 0$$

$$= 2x - y + z = -1 \dots\dots\dots \text{eq(1)}$$

From eq(1) direction ratios of normal vector of the plane are

$$(2, -1, 1)$$

Therefore, equation of normal vector is

$$\vec{n} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\therefore |\vec{n}| = \sqrt{2^2 + (-1)^2 + 1^2}$$

$$= \sqrt{4 + 1 + 1}$$

$$= \sqrt{6}$$

From eq(1), $p = -1$

Given point P = (3, 2, 1)

Position vector of A is

$$\vec{p} = 3\hat{i} + 2\hat{j} + \hat{k}$$

Here, $\vec{a} = \vec{p}$

Now,

$$\therefore \vec{a} \cdot \vec{n} = (3\hat{i} + 2\hat{j} + \hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k})$$

$$= (3 \times 2) + (2 \times (-1)) + (1 \times 1)$$

$$= 6 - 2 + 1$$

$$= 5$$

Length of the perpendicular from point A to the plane is

$$d = \frac{|\vec{a} \cdot \vec{n} - p|}{|\vec{n}|}$$

$$\therefore d = \frac{|5 + 1|}{\sqrt{6}}$$

$$\therefore d = \frac{6}{\sqrt{6}}$$

$$\therefore d = \sqrt{6}$$

Let Q be the foot of perpendicular drawn from point P to the given plane,

Let Q = (x, y, z)

$$\overrightarrow{PQ} = (x - 3)\hat{i} + (y - 2)\hat{j} + (z - 1)\hat{k}$$

As normal vector and $\overrightarrow{PQ}$ are parallel, we can write,

$$\therefore \frac{x - 3}{2} = \frac{y - 2}{-1} = \frac{z - 1}{1} = k(\text{say})$$

$$\Rightarrow x = 2k + 3, y = -k + 2, z = k + 1$$

As point A lies on the plane, we can write

$$2(2k + 3) - (-k + 2) + (k + 1) = -1$$

$$= 4k + 6 + k - 2 + k + 1 = -1$$

$$= 6k = -6$$

$$\therefore k = -1$$

$$\therefore x = 2(-1) + 3 = 1,$$

$$y = -(-1) + 2 = 3$$

$$z = (-1) + 1 = 0$$

Therefore, co-ordinates of the foot of perpendicular are

$$Q(x, y, z) = (1, 3, 0)$$

$$Q \equiv (1, 3, 0)$$

Let, R(a, b, c) be image of point P in the given plane.

Therefore, the power of points P and R in the given plane will be equal and opposite.

$$2a - b + c + 1 = -(2(3) - 2 + 1 + 1)$$

$$= 2a - b + c + 1 = -6$$

$$= 2a - b + c = -7 \dots\dots\dots \text{eq(2)}$$

$$\text{Now, } \overrightarrow{PR} = (a-3)\hat{i} + (b-2)\hat{j} + (c-1)\hat{k}$$

As $\overrightarrow{PR}$ & $\vec{n}$ are parallel

$$\therefore \frac{a-3}{2} = \frac{b-2}{-1} = \frac{c-1}{1} = k(\text{say})$$

$$= a = 2k+3, b = -k+2, c = k+1$$

substituting a, b, c in eq(2)

$$2(2k+3) - (-k+2) + (k+1) = -7$$

$$= 4k + 6 + k - 2 + k + 1 = -7$$

$$= 6k = -12$$

$$\therefore k = -2$$

$$\therefore a = 2(-2) + 3 = -1,$$

$$b = -(-2) + 2 = 4$$

$$c = (-2) + 1 = -1$$

Therefore, co-ordinates of the image of P are

$$R(a, b, c) = (-1, 4, -1)$$

$$R \equiv (-1, 4, -1)$$

Question: 18

Find the coordina

Solution:

Given :

$$\text{Equation of plane : } 2x - y + z + 3 = 0$$

$$P = (1, 3, 4)$$

To Find : Image of point P in the plane.

Note :

If two vectors with direction ratios (a_1, a_2, a_3) & (b_1, b_2, b_3) are parallel then

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

Given equation of the plane is

$$2x - y + z + 3 = 0$$

$$= 2x - y + z = -3 \dots\dots\dots \text{eq(1)}$$

From eq(1) direction ratios of normal vector of the plane are

$$(2, -1, 1)$$

Therefore, equation of normal vector is

$$\vec{n} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\text{Given point is } P = (1, 3, 4)$$

Let, R(a, b, c) be image of point P in the given plane.

Therefore, the power of points P and R in the given plane will be equal and opposite.

$$= 2a - b + c + 3 = - (2(1) - 3 + 4 + 3)$$

$$= 2a - b + c + 3 = -6$$

$$= 2a - b + c = -9 \dots\dots\dots \text{eq(2)}$$

$$\text{Now, } \overrightarrow{PR} = (a-1)\hat{i} + (b-3)\hat{j} + (c-4)\hat{k}$$

As $\overrightarrow{PR}$ & $\vec{n}$ are parallel

$$\therefore \frac{a-1}{2} = \frac{b-3}{-1} = \frac{c-4}{1} = k(\text{say})$$

$$= a = 2k+1, b = -k+3, c = k+4$$

substituting a, b, c in eq(2)

$$2(2k+1) - (-k+3) + (k+4) = -9$$

$$= 4k + 2 + k - 3 + k + 4 = -9$$

$$= 6k = -12$$

$$\therefore k = -2$$

$$\therefore a = 2(-2) + 1 = -3,$$

$$b = -(-2) + 3 = 5$$

$$c = (-2) + 4 = 2$$

Therefore, co-ordinates of the image of P are

$$R(a, b, c) = (-3, 5, 2)$$

Question: 19

Find the point wh

Solution:

Given :

$$\text{Equation of plane : } 2x + 4y - z = 1$$

Equation of line :

$$\frac{x-1}{2} = \frac{y-2}{-3} = \frac{z+3}{4}$$

To Find : Point of intersection of line and plane.

Let P(a, b, c) be point of intersection of plane and line.

As point P lies on the line, we can write,

$$\frac{a-1}{2} = \frac{b-2}{-3} = \frac{c+3}{4} = k(\text{say})$$

$$= a = 2k+1, b = -3k+2, c = 4k-3 \dots\dots\dots (1)$$

Also point P lies on the plane

$$2a + 4b - c = 1$$

$$= 2(2k+1) + 4(-3k+2) - (4k-3) = 1 \dots\dots \text{from (1)}$$

$$= 4k + 2 - 12k + 8 - 4k + 3 = 1$$

$$= -12k = -12$$

$$= k = 1$$

$$\therefore a = 2(1) + 1 = 3,$$

$$b = -3(1) + 2 = -1$$

$$c = 4(1) - 3 = 1$$

Therefore, co-ordinates of point of intersection of given line and plane are

$$P \equiv (3, -1, 1)$$

Question: 20

Find the coordina

Solution:

Given :

$$\text{Equation of plane : } 2x + y + z = 7$$

Points :

$$A = (3, -4, -5)$$

$$B = (2, -3, 1)$$

To Find : Point of intersection of line and plane.

Formula :

Equation of line passing through A = (x₁, y₁, z₁) &

B = (x₂, y₂, z₂) is

$$\frac{x - x_1}{x_1 - x_2} = \frac{y - y_1}{y_1 - y_2} = \frac{z - z_1}{z_1 - z_2}$$

Equation of line passing through A = (3, -4, -5) & B = (2, -3, 1) is

$$\frac{x - 3}{3 - 2} = \frac{y + 4}{-4 + 3} = \frac{z + 5}{-5 - 1}$$

$$\therefore \frac{x - 3}{1} = \frac{y + 4}{-1} = \frac{z + 5}{-6}$$

Let P(a, b, c) be point of intersection of plane and line.

As point P lies on the line, we can write,

$$\frac{a - 3}{1} = \frac{b + 4}{-1} = \frac{c + 5}{-6} = k(\text{say})$$

$$\Rightarrow a = k + 3, b = -k - 4, c = -6k - 5 \dots\dots\dots(1)$$

Also point P lies on the plane

$$2a + b + c = 7$$

$$\Rightarrow 2(k + 3) + (-k - 4) + (-6k - 5) = 7 \dots\dots\text{from (1)}$$

$$\Rightarrow 2k + 6 - k - 4 - 6k - 5 = 7$$

$$\Rightarrow -5k = 10$$

$$\Rightarrow k = -2$$

$$\therefore a = (-2) + 3 = 1,$$

$$b = -(-2) - 4 = -2$$

$$c = -6(-2) - 5 = 7$$

Therefore, co-ordinates of point of intersection of given line and plane are

$$P \equiv (1, -2, 7)$$

Question: 21

Find the distance

Solution:

Given :

$$\text{Equation of plane : } 3x + 2y + 2z + 5 = 0$$

Equation of line :

$$\frac{x + 3}{3} = \frac{y - 2}{6} = \frac{z}{2}$$

$$\text{Point : } P = (2, 3, 4)$$

To Find : Distance of point P from the given plane parallel to the given line.

Formula :

1) Equation of line :

Equation of line passing through A = (x₁, y₁, z₁) & having direction ratios (a, b, c) is

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$$

2) Distance formula :

The distance between two points A = (a₁, a₂, a₃) & B = (b₁, b₂, b₃) is

$$d = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$$

For the given line,

$$\frac{x + 3}{3} = \frac{y - 2}{6} = \frac{z}{2}$$

Direction ratios are (a, b, c) = (3, 6, 2)

Let Q be the point on the plane such that $\overline{PQ}$ is parallel to the given line.

Therefore direction ratios of given line and line PQ will be same.

Therefore equation of line PQ with point P = (2, 3, 4) and with direction ratios (3, 6, 2) is

$$\frac{x-2}{3} = \frac{y-3}{6} = \frac{z-4}{2}$$

Let co-ordinates of Q be (u, v, w)

As point Q lies on the line PQ, we can write,

$$\frac{u-2}{3} = \frac{v-3}{6} = \frac{w-4}{2} = k(\text{say})$$

$$\Rightarrow u = 3k+2, v = 6k+3, w = 2k+4 \dots\dots\dots(1)$$

Also point Q lies on the plane

$$3u + 2v + 2w = -5$$

$$\Rightarrow 3(3k+2) + 2(6k+3) + 2(2k+4) = -5 \dots\dots\text{from (1)}$$

$$\Rightarrow 9k + 6 + 12k + 6 + 4k + 8 = -5$$

$$\Rightarrow 25k = -25$$

$$\Rightarrow k = -1$$

$$\therefore u = 3(-1) + 2 = -1,$$

$$v = 6(-1) + 3 = -3$$

$$w = 2(-1) + 4 = 2$$

Therefore, co-ordinates of point Q are

$$Q = (-1, -3, 2)$$

Now distance between points P and Q by distance formula is

$$\begin{aligned} d &= \sqrt{(2+1)^2 + (3+3)^2 + (4-2)^2} \\ &= \sqrt{(3)^2 + (6)^2 + (2)^2} \\ &= \sqrt{9 + 36 + 4} \\ &= \sqrt{49} \\ &= 7 \end{aligned}$$

Therefore distance of point P from the given plane measured parallel to the given line is

$$d = 7 \text{ units}$$

Question: 22

Find the distance

Solution:

Given :

$$\text{Equation of plane : } x + 2y - z = 1$$

Equation of line :

$$\frac{x+1}{3} = \frac{y+1}{2} = \frac{z}{3}$$

$$\text{Point : } P = (0, -3, 2)$$

To Find : Distance of point P from the given plane parallel to the given line.

Formula :

1) Equation of line :

Equation of line passing through A = (x₁, y₁, z₁) & having direction ratios (a, b, c) is

$$\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$$

2) Distance formula :

The distance between two points A = (a₁, a₂, a₃) & B = (b₁, b₂, b₃) is

$$d = \sqrt{(a_1-b_1)^2 + (a_2-b_2)^2 + (a_3-b_3)^2}$$

For the given line,

$$\frac{x+1}{3} = \frac{y+1}{2} = \frac{z}{3}$$

Direction ratios are (a, b, c) = (3, 2, 3)

Let Q be the point on the plane such that $\overline{PQ}$ is parallel to the given line.

Therefore direction ratios of given line and line PQ will be same.

Therefore equation of line PQ with point P = (0, -3, 2) and with direction ratios (3, 2, 3) is

$$\frac{x-0}{3} = \frac{y+3}{2} = \frac{z-2}{3}$$

Let co-ordinates of Q be (u, v, w)

As point Q lies on the line PQ, we can write,

$$\frac{u}{3} = \frac{v+3}{2} = \frac{w-2}{3} = k(\text{say})$$

$$\Rightarrow u = 3k, v = 2k-3, w = 3k+2 \dots\dots\dots(1)$$

Also point Q lies on the plane

$$u + 2v - w = 1$$

$$\Rightarrow (3k) + 2(2k-3) - (3k+2) = 1 \dots\dots\text{from (1)}$$

$$\Rightarrow 3k + 4k - 6 - 3k - 2 = 1$$

$$\Rightarrow 4k = 9$$

$$\Rightarrow k = \frac{9}{4}$$

$$\therefore u = 3\left(\frac{9}{4}\right) = \frac{27}{4},$$

$$v = 2\left(\frac{9}{4}\right) - 3 = \frac{6}{4}$$

$$w = 3\left(\frac{9}{4}\right) + 2 = \frac{35}{4}$$

Therefore, co-ordinates of point Q are

$$Q \equiv \left(\frac{27}{4}, \frac{6}{4}, \frac{35}{4}\right)$$

Now distance between points P and Q by distance formula is

$$\begin{aligned} d &= \sqrt{\left(0 - \frac{27}{4}\right)^2 + \left(-3 - \frac{6}{4}\right)^2 + \left(2 - \frac{35}{4}\right)^2} \\ &= \sqrt{\left(\frac{-27}{4}\right)^2 + \left(\frac{-18}{4}\right)^2 + \left(\frac{-27}{4}\right)^2} \\ &= \sqrt{45.5625 + 20.25 + 45.5625} \\ &= \sqrt{111.375} \\ &= 10.55 \end{aligned}$$

Therefore distance of point P from the given plane measured parallel to the given line is

$$d = 10.55 \text{ units}$$

Question: 23

Find the equation

Solution:

Given :

$$\text{Equation of plane : } x + y - z = 8$$

Equation of line :

$$\frac{x-1}{3} = \frac{y}{2} = \frac{z+1}{7}$$

$$\text{Point : } P = (4, 6, 2)$$

To Find : Equation of line.

Formula :

Equation of line passing through A = (x₁, y₁, z₁) &

B = (x₂, y₂, z₂) is

$$\frac{x-x_1}{x_1-x_2} = \frac{y-y_1}{y_1-y_2} = \frac{z-z_1}{z_1-z_2}$$

let Q (a, b, c) be point of intersection of plane and line.

As point Q lies on the line, we can write,

$$\frac{a-1}{3} = \frac{b}{2} = \frac{c+1}{7} = k(\text{say})$$

$$\Rightarrow a = 3k+1, b = 2k, c = 7k-1$$

Also point Q lies on the plane,

$$a + b - c = 8$$

$$\Rightarrow (3k+1) + (2k) - (7k-1) = 8$$

$$\Rightarrow 3k + 1 + 2k - 7k + 1 = 8$$

$$\Rightarrow -2k = 6$$

$$\Rightarrow k = -3$$

$$\therefore a = 3(-3) + 1 = -8,$$

$$b = -2(-3) = -6$$

$$c = 7(-3) - 1 = -22$$

Therefore, co-ordinates of point of intersection of given line and plane are Q = (-8, -6, -22)

Now, equation of line passing through P(4,6,2) and

Q(-8, -6, -22) is

$$\frac{x-4}{4+8} = \frac{y-6}{6+6} = \frac{z-2}{2+22}$$

$$\therefore \frac{x-4}{12} = \frac{y-6}{12} = \frac{z-2}{24}$$

$$\therefore \frac{x-4}{1} = \frac{y-6}{1} = \frac{z-2}{2}$$

This is the equation of required line

Question: 24

Show that the dis

Solution:

Given :

Equation of plane : $x - y + z = 5$

Equation of line :

$$\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$$

Point : P = (-1, -5, -10)

To Prove : Distance of point P from the given plane parallel to the given line is 13 units.

Formula :

1) Equation of line :

Equation of line passing through A = (x₁, y₁, z₁) & having direction ratios (a, b, c) is

$$\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$$

2) Distance formula :

The distance between two points A = (a₁, a₂, a₃) & B = (b₁, b₂, b₃) is

$$d = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$$

For the given line,

$$\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$$

Direction ratios are (a, b, c) = (3, 4, 12)

Let Q be the point on the plane such that $\overline{PQ}$ is parallel to the given line.

Therefore direction ratios of given line and line PQ will be same.

Therefore equation of line PQ with point P = (-1, -5, -10) and with direction ratios (3, 4, 12) is

$$\frac{x+1}{3} = \frac{y+5}{4} = \frac{z+10}{12}$$

Let co-ordinates of Q be (u, v, w)

As point Q lies on the line PQ, we can write,

$$\frac{u+1}{3} = \frac{v+5}{4} = \frac{w+10}{12} = k(\text{say})$$

$$\Rightarrow u = 3k-1, v = 4k-5, w = 12k-10 \dots\dots\dots(1)$$

Also point Q lies on the plane

$$u - v + w = 5$$

$$\Rightarrow (3k-1) - (4k-5) + (12k-10) = 5 \dots\dots\text{from (1)}$$

$$\Rightarrow 3k - 1 - 4k + 5 + 12k - 10 = 5$$

$$\Rightarrow 11k = 11$$

$$\Rightarrow k = 1$$

$$\therefore u = 3(1) - 1 = 2,$$

$$v = 4(1) - 5 = -1$$

$$w = 12(1) - 10 = 2$$

Therefore, co-ordinates of point Q are

$$Q \equiv (2, -1, 2)$$

Now distance between points P and Q by distance formula is

$$d = \sqrt{(-1-2)^2 + (-5+1)^2 + (-10-2)^2}$$

$$= \sqrt{(-3)^2 + (-4)^2 + (-12)^2}$$

$$= \sqrt{9 + 16 + 144}$$

$$= \sqrt{169}$$

$$= 13$$

Therefore distance of point P from the given plane measured parallel to the given line is

$$d = 13 \text{ units}$$

Hence proved.

Question: 25

Find the distance

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

Equation of line :

$$\vec{r} = (2\hat{i} - \hat{j} + 2\hat{k}) + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k})$$

$$\text{Point : P} = (-1, -5, -10)$$

To Find : Distance of point P from the given plane parallel to the given line.

Formula :

1) Equation of line :

Equation of line passing through A = (x₁, y₁, z₁) & having direction ratios (a, b, c) is

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$$

2) Distance formula :

The distance between two points A = (a₁, a₂, a₃) & B = (b₁, b₂, b₃) is

$$d = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$$

for the given plane,

$$\vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

$$\text{Here, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

$$\Rightarrow x - y + z = 5 \dots\dots\dots\text{eq(1)}$$

For the given line,

$$\vec{r} = (2\hat{i} - \hat{j} + 2\hat{k}) + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k})$$

$$\text{Here, } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore (3\hat{i} + 4\hat{j} + 2\hat{k})\lambda = (x\hat{i} + y\hat{j} + z\hat{k}) - (2\hat{i} - \hat{j} + 2\hat{k})$$

$$\therefore 3\lambda\hat{i} + 4\lambda\hat{j} + 2\lambda\hat{k} = (x-2)\hat{i} + (y+1)\hat{j} + (z-2)\hat{k}$$

Comparing coefficients of $\hat{i}, \hat{j}$ & $\hat{k}$

$$= 3\lambda = (x - 2), 4\lambda = (y + 1) \text{ \& } 2\lambda = (z - 2)$$

$$\Rightarrow \lambda = \frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{2} \dots\dots\dots \text{eq(2)}$$

Direction ratios for above line are (a, b, c) = (3, 4, 2)

Let Q be the point on the plane such that $\overline{PQ}$ is parallel to the given line.

Therefore direction ratios of given line and line PQ will be same.

Therefore equation of line PQ with point P = (-1, -5, -10) and with direction ratios (3, 4, 2) is

$$\frac{x+1}{3} = \frac{y+5}{4} = \frac{z+10}{2}$$

Let co-ordinates of Q be (u, v, w)

As point Q lies on the line PQ, we can write,

$$\frac{u+1}{3} = \frac{v+5}{4} = \frac{w+10}{2} = k(\text{say})$$

$$\Rightarrow u = 3k-1, v = 4k-5, w = 2k-10 \dots\dots\dots (3)$$

Also point Q lies on the given plane

Therefore from eq(1), we can write,

$$u - v + w = 5$$

$$\Rightarrow (3k-1) - (4k-5) + (2k-10) = 5 \dots\dots \text{from (3)}$$

$$\Rightarrow 3k - 1 - 4k + 5 + 2k - 10 = 5$$

$$\Rightarrow k = 11$$

$$\Rightarrow k = 11$$

$$\therefore u = 3(11) - 1 = 32,$$

$$v = 4(11) - 5 = 39$$

$$w = 2(11) - 10 = 12$$

Therefore, co-ordinates of point Q are

$$Q \equiv (32, 39, 12)$$

Now the distance between points P and Q by distance formula is

$$d = \sqrt{(-1 - 32)^2 + (-5 - 39)^2 + (-10 - 12)^2}$$

$$= \sqrt{(-33)^2 + (-44)^2 + (-22)^2}$$

$$= \sqrt{1089 + 1936 + 484}$$

$$= \sqrt{3509}$$

$$= 59.24$$

Therefore distance of point P from the given plane measured parallel to the given line is

$$d = 59.24 \text{ units}$$

Question: 26

Prove that the no

Solution:

Given :

Equations of plane are :

$$4x + 11y + 2z + 3 = 0$$

$$3x - 2y + 5z = 8$$

To Prove : $\overline{n_1}$ & $\overline{n_2}$ are perpendicular.

Formula :

1) Dot Product :

If $\overline{a}$ & $\overline{b}$ are two vectors

$$\overline{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\overline{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

Note :

Direction ratios of the plane given by

$$ax + by + cz = d$$

are (a, b, c).

For plane

$$4x + 11y + 2z + 3 = 0$$

direction ratios of normal vector are (4, 11, 2)

therefore, equation of normal vector is

$$\vec{n}_1 = 4\hat{i} + 11\hat{j} + 2\hat{k}$$

And for plane

$$3x - 2y + 5z = 8$$

direction ratios of the normal vector are (3, -2, 5)

therefore, the equation of normal vector is

$$\vec{n}_2 = 3\hat{i} - 2\hat{j} + 5\hat{k}$$

Now,

$$\vec{n}_1 \cdot \vec{n}_2 = (4\hat{i} + 11\hat{j} + 2\hat{k}) \cdot (3\hat{i} - 2\hat{j} + 5\hat{k})$$

$$= (4 \times 3) + (11 \times (-2)) + (2 \times 5)$$

$$= 12 - 22 + 10$$

$$= 0$$

$$\therefore \vec{n}_1 \cdot \vec{n}_2 = 0$$

Therefore, normals to the given planes are perpendicular.

Question: 27

Show that the line

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 7$$

Equation of a line :

$$\vec{r} = (2\hat{i} - 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} - \hat{j} + 4\hat{k})$$

To Prove : Given line is parallel to the given plane.

Comparing given plane i.e.

$$\vec{r} \cdot (\hat{i} + 5\hat{j} + \hat{k}) = 7$$

with $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$, we get,

$$\vec{n} = \hat{i} + 5\hat{j} + \hat{k}$$

This is the vector perpendicular to the given plane.

Now, comparing the given the equation of line i.e.

$$\vec{r} = (2\hat{i} - 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} - \hat{j} + 4\hat{k})$$

with $\vec{r} = \vec{a} + \lambda\vec{b}$, we get,

$$\vec{b} = \hat{i} - \hat{j} + 4\hat{k}$$

Now,

$$\vec{n} \cdot \vec{b} = (\hat{i} + 5\hat{j} + \hat{k}) \cdot (\hat{i} - \hat{j} + 4\hat{k})$$

$$= (1 \times 1) + (5 \times (-1)) + (1 \times 4)$$

$$= 1 - 5 + 4$$

$$= 0$$

$$\therefore \vec{n} \cdot \vec{b} = 0$$

Therefore, a vector normal to the plane is perpendicular to the vector parallel to the line.

Hence, the given line is parallel to the given plane.

Question: 28

Find the equation

Solution:

Given :

$$d = 3\sqrt{3}$$

$$\alpha = \beta = \gamma$$

To Find : Equation of plane

Formulae :

1) Distance of plane from the origin :

If $\bar{n} = a\hat{i} + b\hat{j} + c\hat{k}$ is the vector normal to the plane, then distance of the plane from the origin is

$$d = \frac{p}{|\bar{n}|}$$

$$\text{Where, } |\bar{n}| = \sqrt{a^2 + b^2 + c^2}$$

$$\mathbf{2) } l^2 + m^2 + n^2 = 1$$

$$\text{Where } l = \cos \alpha, m = \cos \beta, n = \cos \gamma$$

3) Equation of plane :

If $\bar{n} = a\hat{i} + b\hat{j} + c\hat{k}$ is the vector normal to the plane, then equation of the plane is

$$\bar{r} \cdot \bar{n} = p$$

$$\text{As } \alpha = \beta = \gamma$$

$$\therefore \cos \alpha = \cos \beta = \cos \gamma$$

$$\Rightarrow l = m = n$$

$$l^2 + m^2 + n^2 = 1$$

$$\therefore 3l^2 = 1$$

$$\therefore l = \frac{1}{\sqrt{3}}$$

Therefore equation of normal vector of the plane having direction cosines l, m, n is

$$\bar{n} = l\hat{i} + m\hat{j} + n\hat{k}$$

$$\therefore \bar{n} = \frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k}$$

$$\therefore |\bar{n}| = \sqrt{\left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{1}{\sqrt{3}}\right)^2 + \left(\frac{1}{\sqrt{3}}\right)^2}$$

$$= \sqrt{\frac{1}{3} + \frac{1}{3} + \frac{1}{3}}$$

$$= \sqrt{1}$$

$$= 1$$

Now,

distance of the plane from the origin is

$$d = \frac{p}{|\bar{n}|}$$

$$\therefore 3\sqrt{3} = \frac{p}{1}$$

$$\therefore p = 3\sqrt{3}$$

Therefore equation of required plane is

$$\bar{r} \cdot \bar{n} = p$$

$$\therefore (x\hat{i} + y\hat{j} + z\hat{k}) \cdot \left(\frac{1}{\sqrt{3}}\hat{i} + \frac{1}{\sqrt{3}}\hat{j} + \frac{1}{\sqrt{3}}\hat{k}\right) = 3\sqrt{3}$$

$$\therefore \frac{x}{\sqrt{3}} + \frac{y}{\sqrt{3}} + \frac{z}{\sqrt{3}} = 3\sqrt{3}$$

$$\therefore x + y + z = 3\sqrt{3} \cdot \sqrt{3}$$

$$\therefore x + y + z = 9$$

This is the required equation of the plane.

Question: 29

A vector

Solution:

Given :

$$|\vec{n}| = 8$$

$$\alpha = 45^\circ$$

$$\beta = 60^\circ$$

$$P = (\sqrt{2}, -1, 1)$$

To Find : Equation of plane

Formulae :

$$1) l^2 + m^2 + n^2 = 1$$

$$\text{Where } l = \cos \alpha, m = \cos \beta, n = \cos \gamma$$

2) Equation of plane :

If $\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$ is the vector normal to the plane, then equation of the plane is

$$\vec{r} \cdot \vec{n} = p$$

$$\text{As } \alpha = 45^\circ \text{ \& } \beta = 60^\circ$$

$$\therefore l = \cos \alpha = \cos 45^\circ = \frac{1}{\sqrt{2}} \text{ and}$$

$$m = \cos \beta = \cos 60^\circ = \frac{1}{2}$$

$$\text{But, } l^2 + m^2 + n^2 = 1$$

$$\therefore \left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{1}{2}\right)^2 + n^2 = 1$$

$$\therefore \frac{1}{2} + \frac{1}{4} + n^2 = 1$$

$$\therefore n^2 = 1 - \frac{3}{4}$$

$$\therefore n^2 = \frac{1}{4}$$

$$\therefore n = \frac{1}{2}$$

Therefore direction cosines of the normal vector of the plane are (l, m, n)

Hence direction ratios are (kl, km, kn)

Therefore the equation of normal vector is

$$\vec{n} = kl\hat{i} + km\hat{j} + kn\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{(kl)^2 + (km)^2 + (kn)^2}$$

$$\therefore |\vec{n}| = \sqrt{\left(\frac{k}{\sqrt{2}}\right)^2 + \left(\frac{k}{2}\right)^2 + \left(\frac{k}{2}\right)^2}$$

$$\therefore 8 = \sqrt{\frac{k^2}{2} + \frac{k^2}{4} + \frac{k^2}{4}}$$

$$\therefore 8 = \sqrt{k^2}$$

$$\therefore k = 8$$

$$\vec{n} = \left(\frac{8}{\sqrt{2}}\right)\hat{i} + \left(\frac{8}{2}\right)\hat{j} + \left(\frac{8}{2}\right)\hat{k}$$

$$\therefore \vec{n} = 4\sqrt{2}\hat{i} + 4\hat{j} + 4\hat{k}$$

Now, equation of the plane is

$$\vec{r} \cdot \vec{n} = p$$

$$\therefore \vec{r} \cdot (4\sqrt{2}\hat{i} + 4\hat{j} + 4\hat{k}) = p \dots\dots\dots eq(1)$$

$$\text{But } \vec{r} = (x\hat{i} + y\hat{j} + z\hat{k})$$

$$\therefore (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (4\sqrt{2}\hat{i} + 4\hat{j} + 4\hat{k}) = p$$

$$=4\sqrt{2}x + 4y + 4z = p$$

As point P ($\sqrt{2}$, -1, 1) lies on the plane by substituting it in above equation,

$$4\sqrt{2}(\sqrt{2}) + 4(-1) + 4(1) = p$$

$$=8 - 4 + 4 = p$$

$$\Rightarrow P = 8$$

From eq(1)

$$\therefore \vec{r} \cdot (4\sqrt{2}\hat{i} + 4\hat{j} + 4\hat{k}) = 8$$

Dividing throughout by 4

$$\therefore \vec{r} \cdot (\sqrt{2}\hat{i} + \hat{j} + \hat{k}) = 2$$

This is the equation of required plane.

Question: 30

Find the vector e

Solution:

Given :

$$\vec{a} = 2\hat{i} - 3\hat{j} - 5\hat{k}$$

$$\text{Equation of plane : } \vec{r} \cdot (6\hat{i} - 3\hat{j} + 5\hat{k}) = -2$$

To Find :

Equation of line

Point of intersection

Formula :

Equation of line passing through point A with position vector $\vec{a}$ and parallel to vector $\vec{b}$ is

$$\vec{r} = \vec{a} + \lambda \vec{b}$$

$$\text{Where, } \vec{r} = (x\hat{i} + y\hat{j} + z\hat{k})$$

From the given equation of the plane

$$\vec{r} \cdot (6\hat{i} - 3\hat{j} + 5\hat{k}) = -2 \dots\dots\dots \text{eq(1)}$$

The normal vector of the plane is

$$\vec{n} = 6\hat{i} - 3\hat{j} + 5\hat{k}$$

As the given line is perpendicular to the plane therefore $\vec{n}$ will be parallel to the line.

$$\therefore \vec{n} = \vec{b}$$

Now, the equation of the line passing through $\vec{a} = (2\hat{i} - 3\hat{j} - 5\hat{k})$ and parallel to $\vec{b} = (6\hat{i} - 3\hat{j} + 5\hat{k})$ is

$$\vec{r} = \vec{a} + \lambda \vec{b}$$

$$\therefore \vec{r} = (2\hat{i} - 3\hat{j} - 5\hat{k}) + \lambda(6\hat{i} - 3\hat{j} + 5\hat{k})$$

$$\dots\dots\dots \text{eq(2)}$$

This is the required equation line.

Substituting $\vec{r} = (x\hat{i} + y\hat{j} + z\hat{k})$ in eq(1)

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (6\hat{i} - 3\hat{j} + 5\hat{k}) = -2$$

$$=6x - 3y + 5z = -2 \dots\dots\dots \text{eq(3)}$$

Also substituting $\vec{r} = (x\hat{i} + y\hat{j} + z\hat{k})$ in eq(2)

$$(x\hat{i} + y\hat{j} + z\hat{k}) = (2\hat{i} - 3\hat{j} - 5\hat{k}) + \lambda(6\hat{i} - 3\hat{j} + 5\hat{k})$$

$$\therefore (6\hat{i} - 3\hat{j} + 5\hat{k})\lambda = (x\hat{i} + y\hat{j} + z\hat{k}) - (2\hat{i} - 3\hat{j} - 5\hat{k})$$

$$\therefore 6\lambda\hat{i} - 3\lambda\hat{j} + 5\lambda\hat{k} = (x - 2)\hat{i} + (y + 3)\hat{j} + (z + 5)\hat{k}$$

Comparing coefficients of $\hat{i}, \hat{j}$ & $\hat{k}$

$$=6\lambda = (x - 2), -3\lambda = (y + 3) \text{ \& } 5\lambda = (z + 5)$$

$$\lambda = \frac{x-2}{6} = \frac{y+3}{-3} = \frac{z+5}{5} \dots\dots\dots \text{eq(4)}$$

Let Q(a, b, c) be the point of intersection of given line and plane

As point Q lies on the given line.

Therefore from eq(4)

$$\frac{a-2}{6} = \frac{b+3}{-3} = \frac{c+5}{5} = k(\text{say})$$

$$\Rightarrow a = 6k+2, b = -3k-3, c = 5k-5$$

Also point Q lies on the plane.

Therefore from eq(3)

$$6a - 3b + 5c = -2$$

$$\Rightarrow 6(6k+2) - 3(-3k-3) + 5(5k-5) = -2$$

$$\Rightarrow 36k + 12 + 9k + 9 + 25k - 25 = -2$$

$$\Rightarrow 70k = 2$$

$$\Rightarrow k = \frac{1}{35}$$

$$\therefore a = 6\left(\frac{1}{35}\right) + 2 = \frac{76}{35}$$

$$b = -3\left(\frac{1}{35}\right) - 3 = \frac{-108}{35}$$

$$c = 5\left(\frac{1}{35}\right) - 5 = \frac{-170}{35} = \frac{-34}{7}$$

Therefore co-ordinates of the point of intersection of line and plane are

$$Q \equiv \left(\frac{76}{35}, \frac{-108}{35}, \frac{-34}{7}\right)$$

Exercise : 28C

Question: 1

Find the distance

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

Plane r. $(3i - 4j + 12k) = 9$ can be written in cartesian form as

$$3x - 4y + 12z = 9$$

$$3x - 4y + 12z - 9 = 0$$

$$\text{Point} = (2i - j - 4k)$$

Which can be also written as

$$\text{Point} = (2, -1, -4)$$

$$\text{Distance} = \frac{|(2 \times 3) + (-1 \times -4) + (-4 \times 12) + (-9)|}{\sqrt{(3)^2 + (-4)^2 + 12^2}}$$

$$= \frac{|6 + 4 - 48 - 9|}{\sqrt{9 + 16 + 144}}$$

$$= \frac{|-47|}{\sqrt{169}}$$

$$= \frac{47}{13} \text{ units}$$

Question: 2

Find the distance

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

Plane r. $(i + j + k) + 17 = 0$ can be written in cartesian form as

$$x + y + z + 17 = 0$$

$$\text{Point} = (i + 2j + 5k)$$

Which can be also written as

$$\text{Point} = (1, 2, 5)$$

$$\begin{aligned}\text{Distance} &= \frac{|(1 \times 1) + (2 \times 1) + (5 \times 1) + (17)|}{\sqrt{(1)^2 + (1)^2 + 1^2}} \\&= \frac{|1 + 2 + 5 + 17|}{\sqrt{1 + 1 + 1}} \\&= \frac{|25|}{\sqrt{3}} \\&= \frac{25\sqrt{3}}{3} \text{ units}\end{aligned}$$

Question: 3

Find the di

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

Plane $r. (2i - 5j + 3k) = 13$ can be written in cartesian form as

$$2x - 5y + 3z = 13$$

$$2x - 5y + 3z - 13 = 0$$

$$\text{Point} = (3, 4, 5)$$

$$\begin{aligned}\text{Distance} &= \frac{|(3 \times 2) + (4 \times -5) + (5 \times 3) - (13)|}{\sqrt{(2)^2 + (-5)^2 + 3^2}} \\&= \frac{|6 - 20 + 15 - 13|}{\sqrt{4 + 25 + 9}} \\&= \frac{|-12|}{\sqrt{38}} \\&= \frac{12\sqrt{38}}{38} = \frac{6\sqrt{38}}{19} \text{ units}\end{aligned}$$

Question: 4

Find the di

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

Plane $r. (2i - 2j + 4k) + 5 = 0$ can be written in cartesian form as

$$2x - 2y + 4z + 5 = 0$$

$$\text{Point} = (1, 1, 2)$$

$$\begin{aligned}\text{Distance} &= \frac{|(1 \times 2) + (1 \times -2) + (2 \times 4) + (5)|}{\sqrt{(2)^2 + (-2)^2 + (4)^2}} \\&= \frac{|2 - 2 + 8 + 5|}{\sqrt{4 + 4 + 16}} \\&= \frac{|13|}{\sqrt{24}} \\&= \frac{13}{2\sqrt{6}} = \frac{13\sqrt{6}}{12} \text{ units}\end{aligned}$$

Question: 5

Find the di

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

$$2x + y + 2z + 5 = 0$$

$$\text{Point} = (2, 1, 0)$$

$$\text{Distance} = \frac{|(2 \times 2) + (1 \times 1) + (0 \times 2) + (5)|}{\sqrt{(2)^2 + (1)^2 + (2)^2}}$$

$$= \frac{|4 + 1 + 0 + 5|}{\sqrt{4 + 1 + 4}}$$

$$= \frac{|10|}{\sqrt{9}}$$

$$= \frac{10}{3} \text{ units}$$

Question: 6

Find the di

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

$$x - 2y + 4z = 9$$

$$x - 2y + 4z - 9 = 0$$

$$\text{Point} = (2, 1, -1)$$

$$\text{Distance} = \frac{|(2 \times 1) + (1 \times -2) + (-1 \times 4) - (9)|}{\sqrt{(1)^2 + (-2)^2 + (4)^2}}$$

$$= \frac{|2 - 2 - 4 - 9|}{\sqrt{1 + 4 + 16}}$$

$$= \frac{|-13|}{\sqrt{21}}$$

$$= \frac{13}{\sqrt{21}} = \frac{13\sqrt{21}}{21} \text{ units}$$

Question: 7

Show that t

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

First Plane $r \cdot (i + 2j - 2k) = 5$ can be written in cartesian form as

$$x + 2y - 2z = 5$$

$$x + 2y - 2z - 5 = 0$$

$$\text{Point} = (1, 2, 1)$$

$$\text{Distance for first plane} = \frac{|(1 \times 1) + (2 \times 2) + (1 \times -2) - (5)|}{\sqrt{(1)^2 + (2)^2 + (-2)^2}}$$

$$= \frac{|1 + 4 - 2 - 5|}{\sqrt{1 + 4 + 4}}$$

$$= \frac{|-2|}{\sqrt{9}}$$

$$= \frac{2}{3} \text{ units}$$

Second Plane $r \cdot (2i - 2j + k) + 3 = 0$ can be written in cartesian form as

$$2x - 2y + z + 3 = 0$$

$$\text{Point} = (1, 2, 1)$$

$$\text{Distance for second plane} = \frac{|(1 \times 2) + (2 \times -2) + (1 \times 1) + (3)|}{\sqrt{(2)^2 + (-2)^2 + (1)^2}}$$

$$\begin{aligned}
&= \frac{|2 - 4 + 1 + 3|}{\sqrt{4 + 4 + 1}} \\
&= \frac{|2|}{\sqrt{9}} \\
&= \frac{2}{3} \text{ units}
\end{aligned}$$

Hence proved.

Question: 8

Show that t

Solution:

$$\text{Formula : Distance} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

where (x_1, y_1, z_1) is point from which distance is to be calculated

Therefore ,

$$\text{Plane} = 3x + 4y - 12z + 13 = 0$$

$$\text{First Point} = (-3, 0, 1)$$

$$\text{Distance for first point} = \frac{|(-3 \times 3) + (0 \times 4) + (1 \times -12) + (13)|}{\sqrt{(3)^2 + (4)^2 + (-12)^2}}$$

$$= \frac{|-9 + 0 - 12 + 13|}{\sqrt{9 + 16 + 144}}$$

$$= \frac{|-8|}{\sqrt{169}}$$

$$= \frac{8}{13} \text{ units}$$

$$\text{Plane} = 3x + 4y - 12z + 13 = 0$$

$$\text{Second Point} = (1, 1, 1)$$

$$\text{Distance for first point} = \frac{|(1 \times 3) + (1 \times 4) + (1 \times -12) + (13)|}{\sqrt{(3)^2 + (4)^2 + (-12)^2}}$$

$$= \frac{|3 + 4 - 12 + 13|}{\sqrt{9 + 16 + 144}}$$

$$= \frac{|8|}{\sqrt{169}}$$

$$= \frac{8}{13} \text{ units}$$

Hence proved.

Question: 9

Find the di

Solution:

Formula : The distance between two parallel planes, say

$$\text{Plane 1: } ax + by + cz + d_1 = 0 \text{ \&}$$

Plane 2: $ax + by + cz + d_2 = 0$ is given by the formula

$$\text{Distance} = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

where (d_1, d_2) are constants of the planes

Therefore ,

$$\text{First Plane } 2x + 3y + 4 = 4$$

$$2x + 3y + 4 - 4 = 0 \text{ (1)}$$

$$\text{Second plane } 4x + 6y + 8z = 12$$

$$4x + 6y + 8z - 12 = 0$$

$$2(2x + 3y + 4z - 6) = 0$$

$$2x + 3y + 4z - 6 = 0 \text{ (2)}$$

Using equation (1) and (2)

$$\begin{aligned}
 \text{Distance between both planes} &= \frac{|-6 - (-4)|}{\sqrt{(2)^2 + (3)^2 + (4)^2}} \\
 &= \frac{|-6 + 4|}{\sqrt{4 + 9 + 16}} \\
 &= \frac{|-2|}{\sqrt{29}} \\
 &= \frac{2}{\sqrt{29}} = \frac{2\sqrt{29}}{29} \text{ units}
 \end{aligned}$$

Question: 10

Find the di

Solution:

Formula : The distance between two parallel planes, say

Plane 1: $ax + by + cz + d_1 = 0$ &

Palne 2: $ax + by + cz + d_2 = 0$ is given by the formula

$$\text{Distance} = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

where (d_1, d_2) are costants of the planes

Therefore ,

First Plane $x + 2y - 2z + 4 = 0$ (1)

Second plane $x + 2y - 2z - 8 = 0$ (2)

Using equation (1) and (2)

$$\begin{aligned}
 \text{Distance between both planes} &= \frac{|-8 - (4)|}{\sqrt{(1)^2 + (2)^2 + (2)^2}} \\
 &= \frac{|-12|}{\sqrt{1 + 4 + 4}} \\
 &= \frac{12}{\sqrt{9}} \\
 &= \frac{12}{3} = 4 \text{ units}
 \end{aligned}$$

Question: 11

Find the eq

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel , then their normal vectors are same

Therefore ,

Parallel Plane $x - 2y + 2z - 3 = 0$

Normal vector = $(i - 2j + 2k)$

$\therefore$ Normal vector of required plane = $(i - 2j + 2k)$

Equation of required planes $r \cdot (i - 2j + 2k) = d$

In cartesian form $x - 2y + 2z = d$

It should be at unit distance from point $(1,1,1)$

$$\begin{aligned}
 \text{Distance} &= \frac{|(1 \times 1) + (1 \times -2) + (1 \times 2) - (d)|}{\sqrt{(1)^2 + (-2)^2 + (2)^2}} \\
 &= \frac{|1 - 2 + 2 - d|}{\sqrt{1 + 4 + 4}} \\
 &= \frac{|1 - d|}{\sqrt{9}} \\
 1 &= \frac{\pm(1 - d)}{3}
 \end{aligned}$$

$$3 = \pm(1-d)$$

$$\text{For } + \text{ sign} = > 3 = 1 - d = > d = -2$$

$$\text{For } - \text{ sign} = > 3 = -1 + d = > d = 4$$

Therefore equations of planes are :-

$$\text{For } d = -2 \text{ For } d = 4$$

$$x - 2y + 2z = d \quad x - 2y + 2z = d$$

$$x - 2y + 2z = -2 \quad x - 2y + 2z = 4$$

$$x - 2y + 2z + 2 = 0 \quad x - 2y + 2z - 4 = 0$$

$$\text{Required planes} = x - 2y + 2z + 2 = 0$$

$$x - 2y + 2z - 4 = 0$$

Question: 12

Find the eq

Solution:

$$\text{Formula : Plane} = r \cdot (n) = d$$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

The distance between two parallel planes, say

$$\text{Plane 1: } ax + by + cz + d_1 = 0 \text{ \&}$$

Plane 2: $ax + by + cz + d_2 = 0$ is given by the formula

$$\text{Distance} = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

If two planes are parallel, then their normal vectors are same

Therefore,

$$\text{Parallel Plane } 2x - 3y + 5z + 7 = 0$$

$$\text{Normal vector} = (2i - 3j + 5k)$$

$$\therefore \text{Normal vector of required plane} = (2i - 3j + 5k)$$

$$\text{Equation of required plane } r \cdot (2i - 3j + 5k) = d$$

$$\text{In cartesian form } 2x - 3y + 5z = d$$

Plane passes through point $(3, 4, -1)$ therefore it will satisfy it.

$$2(3) - 3(4) + 5(-1) = d$$

$$6 - 12 - 5 = d$$

$$d = -11$$

$$\text{Equation of required plane } 2x - 3y + 5z = -11$$

$$2x - 3y + 5z + 11 = 0$$

Therefore,

$$\text{First Plane } 2x - 3y + 5z + 7 = 0 \text{ (1)}$$

$$\text{Second plane } 2x - 3y + 5z + 11 = 0 \text{ (2)}$$

Using equation (1) and (2)

$$\text{Distance between both planes} = \frac{|11 - (7)|}{\sqrt{(2)^2 + (-3)^2 + (5)^2}}$$

$$= \frac{|4|}{\sqrt{4 + 9 + 25}}$$

$$= \frac{4}{\sqrt{38}}$$

$$= \frac{4\sqrt{38}}{38} = \frac{2\sqrt{38}}{19} \text{ units}$$

Question: 13

Find the eq

Solution:

Formula : The equation of mid parallel plane is , say

Plane 1: $ax + by + cz + d_1 = 0$ &

Plane 2: $ax + by + cz + d_2 = 0$ is given by the formula

$$\text{Mid parallel plane} = ax + by + cz + \frac{(d_1 + d_2)}{2} = 0$$

where (d_1, d_2) are constants of the planes

Therefore ,

First Plane $2x - 3y + 6z + 21 = 0$ (1)

Second plane $2x - 3y + 6z - 14 = 0$ (2)

Using equation (1) and (2)

$$\text{Mid parallel plane} = 2x - 3y + 6z + \frac{21-14}{2} = 0$$

$$4x - 6y + 12z + 7 = 0$$

Exercise : 28D

Question: 1

Show that t

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel , then their normal vectors are either same or proportional to each other

Therefore ,

Plane 1 : $-2x - y + 6z = 5$

Normal vector (Plane 1) = $(2i - j + 6k)$ (1)

Plane 2 : $-5x - 2.5y + 15z = 12$

Normal vector (Plane 2) = $(5i - 2.5j + 15k)$ (2)

Multiply equation(1) by 5 and equation(2) by 2

Normal vector (Plane 1) = $5(2i - j + 6k)$

$$= 10i - 5j + 30k$$

Normal vector (Plane 2) = $2(5i - 2.5j + 15k)$

$$= 10i - 5j + 30k$$

Since, both normal vectors are same .Therefore both planes are parallel

Question: 2

Find the ve

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel , then their normal vectors are same.

Therefore ,

Parallel Plane $r \cdot (2i - 3j + 5k) + 5 = 0$

Normal vector = $(2i - 3j + 5k)$

$\therefore$ Normal vector of required plane = $(2i - 3j + 5k)$

Equation of required plane $r \cdot (2i - 3j + 5k) = d$

In cartesian form $2x - 3y + 5z = d$

Plane passes through point $(3, 4, -1)$ therefore it will satisfy it.

$$2(3) - 3(4) + 5(-1) = d$$

$$6 - 12 - 5 = d$$

$$d = -11$$

$$\text{Equation of required plane } r \cdot (2i - 3j + 5k) = -11$$

$$r \cdot (2i - 3j + 5k) + 11 = 0$$

Question: 3

Find the ve

Solution:

$$\text{Formula : Plane} = r \cdot (n) = d$$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel, then their normal vectors are same.

Therefore,

$$\text{Parallel Plane } r \cdot (i + j + k) = 2$$

$$\text{Normal vector} = (i + j + k)$$

$$\therefore \text{Normal vector of required plane} = (i + j + k)$$

$$\text{Equation of required plane } r \cdot (i + j + k) = d$$

$$\text{In cartesian form } x + y + z = d$$

Plane passes through point (a, b, c) therefore it will satisfy it.

$$(a) + (b) + (c) = d$$

$$d = a + b + c$$

$$\text{Equation of required plane } r \cdot (i + j + k) = a + b + c$$

Question: 4

Find the ve

Solution:

$$\text{Formula : Plane} = r \cdot (n) = d$$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel, then their normal vectors are same.

Therefore,

$$\text{Parallel Plane } r \cdot (2i - j + 2k) = 5$$

$$\text{Normal vector} = (2i - j + 2k)$$

$$\therefore \text{Normal vector of required plane} = (2i - j + 2k)$$

$$\text{Equation of required plane } r \cdot (2i - j + 2k) = d$$

$$\text{In cartesian form } 2x - y + 2z = d$$

Plane passes through point $(1, 1, 1)$ therefore it will satisfy it.

$$2(1) - (1) + 2(1) = d$$

$$d = 2 - 1 + 2 = 3$$

$$\text{Equation of required plane } r \cdot (2i - j + 2k) = 3$$

Question: 5

Find the eq

Solution:

$$\text{Formula : Plane} = r \cdot (n) = d$$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel, then their normal vectors are same.

Therefore,

$$\text{Parallel Plane } 2x - y + 3z + 7 = 0$$

Normal vector = $(2i - j + 3k)$

$\therefore$ Normal vector of required plane = $(2i - j + 3k)$

Equation of required plane $r \cdot (2i - j + 3k) = d$

In cartesian form $2x - y + 3z = d$

Plane passes through point $(1, 4, -2)$ therefore it will satisfy it.

$$2(1) - (4) + 3(-2) = d$$

$$d = 2 - 4 - 6 = -8$$

Equation of required plane $2x - y + 3z = -8$

$$2x - y + 3z + 8 = 0$$

Question: 6

Find the eq

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel , then their normal vectors are same.

Therefore ,

Parallel Plane $2x - 3y + 7z + 13 = 0$

Normal vector = $(2i - 3j + 7k)$

$\therefore$ Normal vector of required plane = $(2i - 3j + 7k)$

Equation of required plane $r \cdot (2i - 3j + 7k) = d$

In cartesian form $2x - 3y + 7z = d$

Plane passes through point $(0, 0, 0)$ therefore it will satisfy it.

$$2(0) - (0) + 3(0) = d$$

$$d = 0$$

Equation of required plane $2x - 3y + 7z = 0$

Question: 7

Find the eq

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel , then their normal vectors are same.

Therefore ,

Parallel Plane $3x - 5y + 4z = 11$

Normal vector = $(3i - 5j + 4k)$

$\therefore$ Normal vector of required plane = $(3i - 5j + 4k)$

Equation of required plane $r \cdot (3i - 5j + 4k) = d$

In cartesian form $3x - 5y + 4z = d$

Plane passes through point $(-1, 0, 7)$ therefore it will satisfy it.

$$3(-1) - 5(0) + 4(7) = d$$

$$d = -3 + 28 = 25$$

Equation of required plane $3x - 5y + 4z = 25$

Question: 8

Find the eq

Solution:

Formula : Plane = $r \cdot (n) = d$

Where r = any random point

n = normal vector of plane

d = distance of plane from origin

If two planes are parallel, then their normal vectors are same

Therefore,

Parallel Plane $x - 2y + 2z - 3 = 0$

Normal vector = $(i - 2j + 2k)$

$\therefore$ Normal vector of required plane = $(i - 2j + 2k)$

Equation of required planes $r \cdot (i - 2j + 2k) = d$

In cartesian form $x - 2y + 2z = d$

It should be at unit distance from point $(1, 2, 3)$

$$\text{Distance} = \frac{|(1 \times 1) + (2 \times -2) + (3 \times 2) - (d)|}{\sqrt{(1)^2 + (-2)^2 + (2)^2}}$$

$$= \frac{|1 - 4 + 6 - d|}{\sqrt{1 + 4 + 4}}$$

$$= \frac{|3 - d|}{\sqrt{9}}$$

$$1 = \frac{\pm(3 - d)}{3}$$

$$3 = \pm(3 - d)$$

$$\text{For } + \text{ sign} \Rightarrow 3 = 3 - d \Rightarrow d = 0$$

$$\text{For } - \text{ sign} \Rightarrow 3 = -3 + d \Rightarrow d = 6$$

Therefore equations of planes are :-

For $d = 0$ For $d = 6$

$$x - 2y + 2z = d \quad x - 2y + 2z = d$$

$$x - 2y + 2z = 0 \quad x - 2y + 2z = 6$$

$$\text{Required planes} = x - 2y + 2z = 0$$

$$x - 2y + 2z - 6 = 0$$

Question: 9

Find the di

Solution:

Formula : The distance between two parallel planes, say

$$\text{Plane 1: } ax + by + cz + d_1 = 0 \text{ \&}$$

$$\text{Plane 2: } ax + by + cz + d_2 = 0 \text{ is given by the formula}$$

$$\text{Distance} = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

where (d_1, d_2) are constants of the planes

Therefore,

$$\text{First Plane } x + 2y + 3z + 7 = 0$$

$$2(x + 2y + 3z + 7) = 0$$

$$2x + 4y + 6z + 14 = 0 \dots\dots (1)$$

$$\text{Second plane } 2x + 4y + 6z + 7 = 0 \dots\dots (2)$$

Using equation (1) and (2)

$$\text{Distance between both planes} = \frac{|7 - (14)|}{\sqrt{(2)^2 + (4)^2 + (6)^2}}$$

$$= \frac{|-7|}{\sqrt{4 + 16 + 36}}$$

$$= \frac{|-7|}{\sqrt{56}}$$

$$= \frac{7}{\sqrt{56}} \text{ units}$$

Question: 1

Find the equation

Solution:

Equation of plane through the line of intersection of planes in Cartesian form is

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes,

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$x + y + z - 6 + \lambda(2x + 2y + 4z + 5) = 0$$

$$(1 + 2\lambda)x + (1 + 2\lambda)y + (1 + 4\lambda)z - 6 + 5\lambda = 0 \quad (2)$$

Now plane passes through (1,1,1) then it must satisfy the plane equation,

$$(1 + 2\lambda).1 + (1 + 2\lambda).1 + (1 + 4\lambda).1 - 6 + 5\lambda = 0$$

$$1 + 2\lambda + 1 + 2\lambda + 1 + 4\lambda - 6 + 5\lambda = 0$$

$$3 + 8\lambda - 6 + 5\lambda = 0$$

$$13\lambda = 3$$

$$\lambda = \frac{3}{13}$$

Putting in equation (2)

$$\left(1 + 2 \cdot \frac{3}{13}\right)x + \left(1 + 2 \cdot \frac{3}{13}\right)y + \left(1 + 4 \cdot \frac{3}{13}\right)z - 6 + 5 \cdot \frac{3}{13} = 0$$

$$\left(\frac{13+6}{13}\right)x + \left(\frac{13+6}{13}\right)y + \left(\frac{13+12}{13}\right)z + \frac{-78+15}{13} = 0$$

$$19x + 19y + 25z - 63 = 0$$

So, the required equation of plane is $19x + 19y + 25z = 63$.

Question: 2

Find the equation

Solution:

Equation of plane through the line of intersection of planes in Cartesian form is

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes,

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$x - 3y + z + 6 + \lambda(x + 2y + 3z + 5) = 0$$

$$(1 + \lambda)x + (-3 + 2\lambda)y + (1 + 3\lambda)z + 6 + 5\lambda = 0 \quad (2)$$

Now plane passes through (0,0,0) then it must satisfy the plane equation,

$$(1 + \lambda).0 + (-3 + 2\lambda).0 + (1 + 3\lambda).0 + 6 + 5\lambda = 0$$

$$5\lambda = -6$$

$$\lambda = \frac{-6}{5}$$

Putting in equation (2)

$$\left(1 + \frac{-6}{5}\right)x + \left(-3 + 2 \cdot \frac{-6}{5}\right)y + \left(1 + 3 \cdot \frac{-6}{5}\right)z + 6 + 5 \cdot \frac{-6}{5} = 0$$

$$\left(\frac{5+(-6)}{5}\right)x + \left(\frac{-15-12}{5}\right)y + \left(\frac{5+(-18)}{5}\right)z + \frac{30+(-30)}{5} = 0$$

$$-x - 27y - 13z = 0$$

$$x + 27y + 13z = 0$$

So, required equation of plane is $x + 27y + 13z = 0$.

Question: 3

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes,

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$2x + 3y - z + 1 + \lambda(x + y - 2z + 3) = 0$$

$$(2 + \lambda)x + (3 + \lambda)y + (-1 - 2\lambda)z + 1 + 3\lambda = 0 \quad (2)$$

Now as the plane $3x - y - 2z - 4 = 0$ is perpendicular to the given plane,

$$\text{For } \theta = 90^\circ, \cos 90^\circ = 0$$

$$A_1A_2 + B_1B_2 + C_1C_2 = 0 \quad (3)$$

On comparing with standard equations in Cartesian form,

$$A_1 = 2 + \lambda, B_1 = 3 + \lambda, C_1 = -1 - 2\lambda \text{ and } A_2 = 3, B_2 = -1, C_2 = -2$$

Putting values in equation (3), we have

$$(2 + \lambda).3 + (3 + \lambda).(-1) + (-1 - 2\lambda).(-2) = 0$$

$$6 + 3\lambda - 3 - \lambda + 2 + 4\lambda = 0$$

$$5 + 6\lambda = 0$$

$$\lambda = \frac{-5}{6}$$

Putting in equation (2)

$$\left(2 + \frac{-5}{6}\right)x + \left(3 + \frac{-5}{6}\right)y + \left(-1 - 2 \cdot \frac{-5}{6}\right)z + 1 + 3 \cdot \frac{-5}{6} = 0$$

$$\left(\frac{12 - 5}{6}\right)x + \left(\frac{18 - 5}{6}\right)y + \left(\frac{-6 + 10}{6}\right)z + \frac{6 - 15}{6} = 0$$

$$7x + 13y + 4z - 9 = 0$$

$$7x + 13y + 4z = 9$$

So, required equation of plane is $7x + 13y + 4z = 9$.

Question: 4

Find the equation

Solution:

Equation of plane through the line of intersection of planes in Cartesian form is

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes,

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$2x - y + \lambda(3z - y) = 0$$

$$2x + (-1 - \lambda)y + 3\lambda z = 0 \quad (2)$$

Now as the plane is perpendicular to the given plane,

$$\text{For } \theta = 90^\circ, \cos 90^\circ = 0$$

$$A_1A_2 + B_1B_2 + C_1C_2 = 0 \quad (3)$$

On comparing with standard equations in Cartesian form,

$$A_1 = 2, B_1 = -1 - \lambda, C_1 = 3\lambda \text{ and } A_2 = 4, B_2 = 5, C_2 = -3$$

Putting values in equation (3),

$$2.4 + (-1 - \lambda).5 + 3\lambda.(-3) = 0$$

$$8 - 5 - 5\lambda - 9\lambda = 0$$

$$-14\lambda = -3$$

$$\lambda = \frac{3}{14}$$

Putting in equation (2)

$$2x + \left(-1 - \frac{3}{14}\right)y + 3\left(\frac{3}{14}\right)z = 0$$

$$2x + \left(\frac{-14-3}{14}\right)y + \frac{9}{14}z = 0$$

$$28x - 17y + 9z = 0$$

So, required equation of plane is $28x - 17y + 9z = 0$.

Question: 5

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes,

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$x - 2y + z - 1 + \lambda(2x + y + z - 8) = 0$$

$$(1 + 2\lambda)x + (-2 + \lambda)y + (1 + \lambda)z - 1 - 8\lambda = 0 \quad (2)$$

For plane the normal is perpendicular to line given parallel to this i.e.

$$A_1A_2 + B_1B_2 + C_1C_2 = 0$$

Where A_1, B_1, C_1 are direction ratios of plane and A_2, B_2, C_2 are of line.

$$(1 + 2\lambda).1 + (-2 + \lambda).2 + (1 + \lambda).1 = 0$$

$$1 + 2\lambda - 4 + 2\lambda + 1 + \lambda = 0$$

$$-2 + 5\lambda = 0$$

$$\lambda = \frac{2}{5}$$

Putting the value of λ in equation (2)

$$\left(1 + 2 \cdot \left(\frac{2}{5}\right)\right)x + \left(-2 + \frac{2}{5}\right)y + \left(1 + \frac{2}{5}\right)z - 1 - 8 \cdot \left(\frac{2}{5}\right) = 0$$

$$\left(\frac{5+4}{5}\right)x + \left(\frac{-10+2}{5}\right)y + \left(\frac{5+2}{5}\right)z + \frac{-5-16}{5} = 0$$

$$9x - 8y + 7z - 21 = 0$$

$$9x - 8y + 7z = 21$$

For the equation of plane $Ax + By + Cz = D$ and point (x_1, y_1, z_1) , a distance of a point from a plane can be calculated as

$$\frac{|Ax_1 + By_1 + Cz_1 - D|}{\sqrt{A^2 + B^2 + C^2}}$$

$$\frac{|9 \cdot 1 - 8 \cdot 1 + 7 \cdot 1 - 21|}{\sqrt{(9)^2 + (-8)^2 + (7)^2}} \Rightarrow \frac{|9 - 8 + 7 - 21|}{\sqrt{81 + 64 + 49}} = \frac{|13|}{\sqrt{194}}$$

So, the required equation of the plane is $9x - 8y + 7z = 21$, and distance of the plane from $(1, 1, 1)$ is

$$d = \frac{13}{\sqrt{194}}$$

Question: 6

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes in Cartesian form

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation 1 we have

$$x + 2y + 3z - 5 + \lambda(3x - 2y - z + 1) = 0$$

$$(1 + 3\lambda)x + (2 - 2\lambda)y + (3 - \lambda)z - 5 + \lambda = 0$$

Now equation of plane in intercept form

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

As given equal intercept means $a=c$

First, we transform equation of a plane in intercept form

$$\frac{\frac{x}{1}}{(1+3\lambda)} + \frac{\frac{y}{1}}{(2-2\lambda)} + \frac{\frac{z}{1}}{(3-\lambda)} = 5-\lambda$$

$$\frac{\frac{x}{5-\lambda}}{(1+3\lambda)} + \frac{\frac{y}{5-\lambda}}{(2-2\lambda)} + \frac{\frac{z}{5-\lambda}}{(3-\lambda)} = 1$$

On comparing with the standard equation of a plane in intercept form

$$a = \frac{5-\lambda}{(1+3\lambda)}, c = \frac{5-\lambda}{(3-\lambda)}$$

Now as $a=b=c$

$$\frac{5-\lambda}{(1+3\lambda)} = \frac{5-\lambda}{(3-\lambda)} \Rightarrow 3-\lambda = 1+3\lambda$$

$$4\lambda = 2 \Rightarrow \lambda = \frac{1}{2}$$

Putting in equation (2), we have

$$\left(1+3\cdot\frac{1}{2}\right)x + \left(2-2\cdot\frac{1}{2}\right)y + \left(3-\frac{1}{2}\right)z - 5 + \frac{1}{2} = 0$$

$$\left(\frac{2+3}{2}\right)x + \left(\frac{4-2}{2}\right)y + \left(\frac{6-1}{2}\right)z + \frac{-10+1}{2} = 0$$

$$5x + 2y + 5z - 9 = 0$$

$$5x + 2y + 5z = 9$$

So, required equation of plane is $5x + 2y + 5z = 9$.

Question: 7

Find the equation

Solution:

Equation of plane through the line of intersection of planes in Cartesian form is

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0 \quad (1)$$

For the standard equation of planes in Cartesian form

$$A_1x + B_1y + C_1z + D_1 \text{ and } A_2x + B_2y + C_2z + D_2$$

So, putting in equation (1), we have

$$3x - 4y + 5z - 10 + \lambda(2x + 2y - 3z - 4) = 0$$

$$(3 + 2\lambda)x + (-4 + 2\lambda)y + (5 - 3\lambda)z - 10 - 4\lambda = 0$$

Given line is parallel to plane then the normal of plane is perpendicular to line,

$$A_1A_2 + B_1B_2 + C_1C_2 = 0$$

Where A_1, B_1, C_1 are direction ratios of plane and A_2, B_2, C_2 are of line.

$$(3 + 2\lambda).6 + (-4 + 2\lambda).3 + (5 - 3\lambda).2 = 0$$

$$18 + 12\lambda - 12 + 6\lambda + 10 - 6\lambda = 0$$

$$16 + 12\lambda = 0$$

$$\lambda = \frac{-16}{12} \Rightarrow \frac{-4}{3}$$

Putting the value of λ in equation (2)

$$\left(3+2\cdot\left(\frac{-4}{3}\right)\right)x + \left(-4+2\cdot\left(\frac{-4}{3}\right)\right)y + \left(5-3\cdot\left(\frac{-4}{3}\right)\right)z - 10 - 4\cdot\left(\frac{-4}{3}\right) = 0$$

$$\left(\frac{9-8}{3}\right)x + \left(\frac{-12-8}{3}\right)y + \left(\frac{15+12}{3}\right)z + \frac{-30+16}{3} = 0$$

$$x - 20y + 27z - 14 = 0$$

So, required equation of plane is $x-20y + 27z-14=0$.

Question: 8

Find the vector e

Solution:

Equation of plane through the line of intersection of two planes in vector form is

$$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 \text{ where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (1)$$

Where the standard equation of planes are

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

Putting values in equation(1)

$$\vec{r}(\hat{i} + 3\hat{j} - \hat{k} + \lambda(\hat{j} + 2\hat{k})) = 0 + \lambda \cdot 0$$

$$\vec{r}(\hat{i} + (3 + \lambda)\hat{j} + (-1 + 2\lambda)\hat{k}) = 0 \quad (2)$$

Now as the plane passes through $(2, 1, -1)$

$$\vec{r} = 2\hat{i} + \hat{j} - \hat{k}$$

Putting in equation (2)

$$(2\hat{i} + \hat{j} - \hat{k})((\hat{i} + (3 + \lambda)\hat{j} + (-1 + 2\lambda)\hat{k})) = 0$$

$$2.1 + 1.(3 + \lambda) + (-1)(-1 + 2\lambda) = 0$$

$$2 + 3 + \lambda + 1 - 2\lambda = 0$$

$$\lambda = 6$$

Putting the value of λ in equation (2)

$$\vec{r}(\hat{i} + (3 + 6)\hat{j} + (-1 + 2(6))\hat{k}) = 0$$

$$\vec{r}(\hat{i} + 9\hat{j} + 11\hat{k}) = 0$$

$$\text{So, required equation of plane is } \vec{r}(\hat{i} + 9\hat{j} + 11\hat{k}) = 0.$$

Question: 9

$$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 \text{ where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (1)$$

Where the standard equation of planes are

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

Putting values in equation(1)

$$\vec{r}(\hat{i} - \hat{j} + 3\hat{k} + \lambda(2\hat{i} + \hat{j} - \hat{k})) = -1 + \lambda \cdot 5$$

$$\vec{r}((1 + 2\lambda)\hat{i} + (-1 + \lambda)\hat{j} + (3 - \lambda)\hat{k}) = -1 + 5\lambda \quad (2)$$

Now as the plane passes through $(1, 1, 1)$

$$\vec{r} = \hat{i} + \hat{j} + \hat{k}$$

Putting in equation (2)

$$(\hat{i} + \hat{j} + \hat{k})(((1 + 2\lambda)\hat{i} + (-1 + \lambda)\hat{j} + (3 - \lambda)\hat{k})) = -1 + 5\lambda$$

$$1.(1 + 2\lambda) + 1.(-1 + \lambda) + 1.(3 - \lambda) = -1 + 5\lambda$$

$$1 + 2\lambda - 1 + \lambda + 3 - \lambda = -1 + 5\lambda$$

$$-3\lambda + 4 = 0$$

$$\lambda = \frac{4}{3}$$

Putting the value of λ in equation (2)

$$\vec{r}\left(\left(1 + 2 \cdot \frac{4}{3}\right)\hat{i} + \left(-1 + \frac{4}{3}\right)\hat{j} + \left(3 - \frac{4}{3}\right)\hat{k}\right) = -1 + 5 \cdot \frac{4}{3}$$

$$\vec{r} \left(\left(\frac{3+8}{3} \right) \hat{i} + \left(\frac{-3+4}{3} \right) \hat{j} + \left(\frac{9-4}{3} \right) \hat{k} \right) = \frac{-3+20}{3}$$

$$\vec{r} (11\hat{i} + \hat{j} + 5\hat{k}) = 17$$

So, required equation of plane is $\vec{r} (11\hat{i} + \hat{j} + 5\hat{k}) = 17$.

Question: 10

Find the vector e

Solution:

Equation of plane through the line of intersection of two planes in vector form is

$$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 \text{ where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (1)$$

Where the standard equation of planes are

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

Putting values in equation (1)

$$\vec{r} (2\hat{i} - 7\hat{j} + 4\hat{k} + \lambda (3\hat{i} - 5\hat{j} + 4\hat{k})) = 3 - \lambda \cdot 11$$

$$\vec{r} ((2+3\lambda)\hat{i} + (-7-5\lambda)\hat{j} + (4+4\lambda)\hat{k}) = 3 - 11\lambda \quad (2)$$

Now as the plane passes through $(-2, 1, 3)$

$$\vec{r} = -2\hat{i} + \hat{j} + 3\hat{k}$$

Putting in equation (2)

$$(-2\hat{i} + \hat{j} + 3\hat{k}) ((2+3\lambda)\hat{i} + (-7-5\lambda)\hat{j} + (4+4\lambda)\hat{k}) = 3 - 11\lambda$$

$$-2 \cdot (2 + 3\lambda) + 1 \cdot (-7 - 5\lambda) + 3 \cdot (4 + 4\lambda) = 3 - 11\lambda$$

$$-4 - 6\lambda - 7 - 5\lambda + 12 + 12\lambda - 3 + 11\lambda = 0$$

$$-14 + 12 + 12\lambda = 0$$

$$\lambda = \frac{1}{6}$$

Putting the value of λ in equation (2)

$$\vec{r} \left(\left(2 + 3 \cdot \frac{1}{6} \right) \hat{i} + \left(-7 - 5 \cdot \frac{1}{6} \right) \hat{j} + \left(4 + 4 \cdot \frac{1}{6} \right) \hat{k} \right) = 3 - 11 \cdot \frac{1}{6}$$

$$\vec{r} \left(\left(\frac{12+3}{6} \right) \hat{i} + \left(\frac{-42-5}{6} \right) \hat{j} + \left(\frac{24+4}{6} \right) \hat{k} \right) = \frac{18-11}{6}$$

$$\vec{r} (15\hat{i} - 47\hat{j} + 28\hat{k}) = 7$$

So, required equation of plane is $\vec{r} (15\hat{i} - 47\hat{j} + 28\hat{k}) = 7$.

Question: 11

Equation of plane through the line of intersection of two planes in vector form is

$$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 \text{ where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (1)$$

Where the standard equation of planes are

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

Putting values in equation (1), we have

$$\vec{r} (2\hat{i} - 3\hat{j} + 4\hat{k} + \lambda (\hat{i} - \hat{j})) = 1 - \lambda \cdot 4$$

$$\vec{r} ((2+\lambda)\hat{i} + (-3-\lambda)\hat{j} + 4\hat{k}) = 1 - 4\lambda \quad (2)$$

Given a plane perpendicular to this plane, So if n_1 and n_2 are normal vectors of planes

$$\vec{n}_1 \cdot \vec{n}_2 = 0$$

$$(2\hat{i} - \hat{j} + \hat{k}) \cdot ((2 + \lambda)\hat{i} + (-3 - \lambda)\hat{j} + 4\hat{k}) = 0$$

$$2.(2 + \lambda) + (-1).(-3 - \lambda) + 1.4 = 0$$

$$4 + 2\lambda + 3 + \lambda + 4 = 0$$

$$11 + 3\lambda = 0$$

$$\lambda = \frac{-11}{3}$$

Putting the value of λ in equation (2)

$$\vec{r} \cdot \left(\left(2 + \frac{-11}{3} \right) \hat{i} + \left(-3 - \frac{-11}{3} \right) \hat{j} + 4\hat{k} \right) = 1 - 4 \cdot \frac{-11}{3}$$

$$\vec{r} \cdot \left(\left(\frac{6 - 11}{3} \right) \hat{i} + \left(\frac{-9 + 11}{3} \right) \hat{j} + 4\hat{k} \right) = \frac{3 + 44}{3}$$

$$\vec{r} \cdot (-5\hat{i} - 2\hat{j} + 12\hat{k}) = 47$$

So required equation of plane is $\vec{r} \cdot (-5\hat{i} - 2\hat{j} + 12\hat{k}) = 47$.

Question: 12

$$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 \text{ where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k} \quad (1)$$

Where the standard equation of planes are

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

Putting values in equation (1)

$$\vec{r} \cdot (\hat{i} - \hat{j} + \lambda(3\hat{i} + 3\hat{j} - 4\hat{k})) = 6 + \lambda \cdot 0$$

$$\vec{r} \cdot ((1 + 3\lambda)\hat{i} + (-1 + 3\lambda)\hat{j} + (-4\lambda)\hat{k}) = 6 \quad (2)$$

For the equation of plane $Ax + By + Cz = D$ and point (x_1, y_1, z_1) , a distance of a point from a plane can be calculated as

$$\left| \frac{Ax_1 + By_1 + Cz_1 - D}{\sqrt{A^2 + B^2 + C^2}} \right|$$

$$\left| \frac{(1 + 3\lambda)0 + (-1 + 3\lambda) \cdot 0 + (-4\lambda) \cdot 0 - 6}{\sqrt{(1 + 3\lambda)^2 + (-1 + 3\lambda)^2 + (-4\lambda)^2}} \right| = 1$$

$$\left| \frac{-6}{\sqrt{1 + 9\lambda^2 + 6\lambda + 1 + 9\lambda^2 - 6\lambda + 16\lambda^2}} \right| = 1$$

$$\sqrt{2 + 34\lambda^2} = -6$$

$$2 + 34\lambda^2 = (-6)^2$$

$$34\lambda^2 = 36 - 2$$

$$34\lambda^2 = 34$$

$$\lambda^2 = 1 \Rightarrow \lambda = 1, -1$$

Putting value of λ in equation (2)

$$\lambda = 1$$

$$\vec{r} \cdot ((1 + 3 \cdot 1)\hat{i} + (-1 + 3 \cdot 1)\hat{j} + (-4 \cdot 1)\hat{k}) = 6$$

$$\vec{r} \cdot (4\hat{i} + 2\hat{j} - 4\hat{k}) = 6 \Rightarrow \vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 3$$

$$\lambda = -1$$

$$\vec{r} \cdot ((1 + 3 \cdot (-1))\hat{i} + (-1 + 3 \cdot (-1))\hat{j} + (-4 \cdot (-1))\hat{k}) = 6$$

$$\vec{r} \cdot (-2\hat{i} - 4\hat{j} + 4\hat{k}) = 6 \Rightarrow \vec{r} \cdot (\hat{i} + 2\hat{j} - 2\hat{k}) = -3$$

For equations in Cartesian form put

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

For $\lambda=1$

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k} - 3) = 0$$

$$x.2 + y.1 + z.(-2) - 3 = 0$$

$$2x + y - 2z - 3 = 0$$

For $\lambda=-1$

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + 2\hat{j} - 2\hat{k} + 3) = 0$$

$$x.1 + y.2 + z.(-2) + 3 = 0$$

$$x + 2y - 2z + 3 = 0$$

So, required equation of plane

$$\text{in vector form are } \vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 3 \text{ for } \lambda = 1$$

$$\vec{r} \cdot (\hat{i} + 2\hat{j} - 2\hat{k}) = -3 \text{ for } \lambda = -1$$

In Cartesian form are $2x + y - 2z - 3 = 0$ & $x + 2y - 2z + 3 = 0$

Exercise : 28F

Question: 1

Find the acute angle

Solution:

To find the angle between two planes, we simply find the angle between the normal vectors of planes. So if n_1 and n_2 are normal vectors and θ is the angle between both then,

$$\cos\theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|}$$

(i) On comparing with the standard equation of planes in vector form

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

$$\vec{n}_1 = \hat{i} + \hat{j} - 2\hat{k} \text{ and } \vec{n}_2 = 2\hat{i} + 2\hat{j} - \hat{k}$$

Then

$$\cos\theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} \Rightarrow \frac{|(\hat{i} + \hat{j} - 2\hat{k}) \cdot (2\hat{i} + 2\hat{j} - \hat{k})|}{|\hat{i} + \hat{j} - 2\hat{k}| |2\hat{i} + 2\hat{j} - \hat{k}|} = \frac{|1.2 + 1.2 + (-2).(-1)|}{(\sqrt{1^2 + 1^2 + (-2)^2}) \cdot (\sqrt{2^2 + 2^2 + (-1)^2})} = \frac{|2 + 2 + 2|}{\sqrt{1+1+4} \sqrt{4+4+1}}$$

$$\Rightarrow \frac{|6|}{\sqrt{6} \cdot \sqrt{9}} = \frac{|\sqrt{6}|}{3}$$

$$\theta = \cos^{-1}\left(\frac{\sqrt{6}}{3}\right)$$

(ii) On comparing with the standard equation of planes in vector form

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

$$\vec{n}_1 = \hat{i} + 2\hat{j} - \hat{k} \text{ and } \vec{n}_2 = 2\hat{i} - \hat{j} - \hat{k}$$

Then

$$\cos\theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} \Rightarrow \frac{|(\hat{i} + 2\hat{j} - \hat{k}) \cdot (2\hat{i} - \hat{j} - \hat{k})|}{|\hat{i} + 2\hat{j} - \hat{k}| |2\hat{i} - \hat{j} - \hat{k}|} = \frac{|1.2 + 2.(-1) + (-1).(-1)|}{(\sqrt{1^2 + 2^2 + (-1)^2}) \cdot (\sqrt{2^2 + (-1)^2 + (-1)^2})} = \frac{|2 - 2 + 1|}{\sqrt{1+4+1} \sqrt{4+1+1}}$$

$$\Rightarrow \frac{|1|}{\sqrt{6} \cdot \sqrt{6}} = \frac{1}{6}$$

$$\theta = \cos^{-1}\left(\frac{1}{6}\right)$$

(iii) On comparing with the standard equation of planes in vector form

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

$$\vec{n}_1 = 2\hat{i} - 3\hat{j} + 4\hat{k} \text{ and } \vec{n}_2 = -\hat{i} + \hat{j}$$

Then

$$\cos\theta = \frac{\left| (2\hat{i} - 3\hat{j} + 4\hat{k}) \cdot (-\hat{i} + \hat{j}) \right|}{\left| 2\hat{i} - 3\hat{j} + 4\hat{k} \right| \left| -\hat{i} + \hat{j} \right|} \Rightarrow \frac{2 \cdot (-1) + (-3) \cdot 1 + 4 \cdot 0}{\left(\sqrt{2^2 + (-3)^2 + 4^2} \right) \cdot \left(\sqrt{(-1)^2 + 1^2} \right)} = \frac{-2 + (-3)}{\left(\sqrt{4 + 9 + 16} \right) \left(\sqrt{1 + 1} \right)}$$

$$\Rightarrow \frac{-5}{\sqrt{29}\sqrt{2}} = \frac{-5}{\sqrt{58}}$$

$$\theta = \cos^{-1}\left(\frac{5}{\sqrt{58}}\right)$$

(iv) On comparing with the standard equation of planes in vector form

$$\vec{r} \cdot \vec{n}_1 = d_1 \text{ and } \vec{r} \cdot \vec{n}_2 = d_2$$

$$\vec{n}_1 = 2\hat{i} - 3\hat{j} + 6\hat{k} \text{ and } \vec{n}_2 = 3\hat{i} + 4\hat{j} - 12\hat{k}$$

Then

$$\cos\theta = \frac{\left| (2\hat{i} - 3\hat{j} + 6\hat{k}) \cdot (3\hat{i} + 4\hat{j} - 12\hat{k}) \right|}{\left| 2\hat{i} - 3\hat{j} + 6\hat{k} \right| \left| 3\hat{i} + 4\hat{j} - 12\hat{k} \right|} \Rightarrow \frac{2 \cdot 3 + (-3) \cdot 4 + 6 \cdot (-12)}{\left(\sqrt{2^2 + (-3)^2 + 6^2} \right) \cdot \left(\sqrt{3^2 + 4^2 + (-12)^2} \right)}$$

$$= \frac{6 + (-12) + (-72)}{\left(\sqrt{4 + 9 + 36} \right) \left(\sqrt{9 + 16 + 144} \right)}$$

$$\Rightarrow \frac{-78}{\sqrt{49}\sqrt{169}} = \frac{-78}{7 \cdot 13}$$

$$\theta = \cos^{-1}\left(\frac{6}{7}\right)$$

Question: 2

To show the right angle between two planes, we simply find the angle between the normal vectors of planes. So if n_1 and n_2 are normal vectors and θ is the angle between both then

$$\cos\theta = \frac{\left| \vec{n}_1 \cdot \vec{n}_2 \right|}{\left| \vec{n}_1 \right| \left| \vec{n}_2 \right|} \text{ for right angle } \theta = 90^\circ$$

$$\cos 90^\circ = 0$$

$$\vec{n}_1 \cdot \vec{n}_2 = 0 \text{ (1)}$$

(i) On comparing with standard equation

$$\vec{n}_1 = 4\hat{i} - 7\hat{j} - 8\hat{k} \text{ and } \vec{n}_2 = 3\hat{i} - 4\hat{j} + 5\hat{k}$$

$$\text{LHS} = \vec{n}_1 \cdot \vec{n}_2 \Rightarrow (4\hat{i} - 7\hat{j} - 8\hat{k}) \cdot (3\hat{i} - 4\hat{j} + 5\hat{k}) = 4 \cdot 3 + (-7) \cdot (-4) + (-8) \cdot 5$$

$$\Rightarrow 12 + 28 - 40 = 40 - 40 \Rightarrow 0 = \text{RHS}$$

Hence proved planes at right angles.

(ii) On comparing with the standard equation of a plane

$$\vec{n}_1 = 2\hat{i} + 6\hat{j} + 6\hat{k} \text{ and } \vec{n}_2 = 3\hat{i} + 4\hat{j} - 5\hat{k}$$

$$\text{LHS} = \vec{n}_1 \cdot \vec{n}_2 \Rightarrow (2\hat{i} + 6\hat{j} + 6\hat{k}) \cdot (3\hat{i} + 4\hat{j} - 5\hat{k}) = 2 \cdot 3 + 6 \cdot 4 + 6 \cdot (-5)$$

$$\Rightarrow 6 + 24 - 30 = 30 - 30 \Rightarrow 0 = \text{RHS}$$

Hence proved planes at right angles.

Question: 3

Find the value of

Solution:

For planes perpendicular $\cos 90^\circ = 0$

$$\vec{n}_1 \cdot \vec{n}_2 = 0 \quad (1)$$

(i) On comparing with the standard equation of a plane

$$\vec{n}_1 = 2\hat{i} - \hat{j} - \lambda\hat{k} \text{ and } \vec{n}_2 = 3\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\vec{n}_1 \cdot \vec{n}_2 = (2\hat{i} - \hat{j} - \lambda\hat{k}) \cdot (3\hat{i} + 2\hat{j} + 2\hat{k}) = 0$$

$$2 \cdot 3 + (-1) \cdot 2 + (-\lambda) \cdot 2 = 0$$

$$6 - 2 - 2\lambda = 0$$

$$2\lambda = 4$$

$$\lambda = 2$$

(ii) On comparing with the standard equation of a plane

$$\vec{n}_1 = \lambda\hat{i} + 2\hat{j} + 3\hat{k} \text{ and } \vec{n}_2 = \hat{i} + 2\hat{j} - 7\hat{k}$$

$$\vec{n}_1 \cdot \vec{n}_2 = (\lambda\hat{i} + 2\hat{j} + 3\hat{k}) \cdot (\hat{i} + 2\hat{j} - 7\hat{k}) = 0$$

$$\lambda \cdot 1 + 2 \cdot 2 + 3 \cdot (-7) = 0 \quad \lambda + 4 - 21 = 0 \quad \lambda = 17$$

Question: 4

$$A_1x + B_1y + C_1z + D_1 = 0 \text{ and } A_2x + B_2y + C_2z + D_2 = 0$$

$$\cos \theta = \left| \frac{A_1A_2 + B_1B_2 + C_1C_2}{\sqrt{(A_1^2 + B_1^2 + C_1^2)} \sqrt{(A_2^2 + B_2^2 + C_2^2)}} \right|$$

(i) On comparing with the standard equation of planes

$$A_1 = 2, B_1 = -1, C_1 = 1 \text{ and } A_2 = 1, B_2 = 1, C_2 = 2$$

$$\cos \theta = \left| \frac{2 \cdot 1 + (-1) \cdot 1 + 1 \cdot 2}{\sqrt{2^2 + (-1)^2 + 1^2} \sqrt{1^2 + 1^2 + 2^2}} \right| \Rightarrow \left| \frac{2 + (-1) + 2}{\sqrt{4 + 1 + 1} \sqrt{1 + 1 + 4}} \right| = \left| \frac{3}{\sqrt{6} \sqrt{6}} \right|$$

$$= \frac{3}{6} \Rightarrow \frac{1}{2}$$

$$\theta = \cos^{-1} \left(\frac{1}{2} \right) \Rightarrow \frac{\pi}{3}$$

(ii) On comparing with the standard equation of planes

$$A_1 = 1, B_1 = 2, C_1 = 2 \text{ and } A_2 = 2, B_2 = -3, C_2 = 6$$

$$\cos \theta = \left| \frac{1 \cdot 2 + 2 \cdot (-3) + 2 \cdot 6}{\sqrt{1^2 + 2^2 + 2^2} \sqrt{2^2 + (-3)^2 + 6^2}} \right| \Rightarrow \left| \frac{2 + (-6) + 12}{\sqrt{1 + 4 + 4} \sqrt{4 + 9 + 36}} \right| = \left| \frac{8}{\sqrt{9} \sqrt{49}} \right|$$

$$= \frac{8}{3 \cdot 7} \Rightarrow \frac{8}{21}$$

$$\theta = \cos^{-1} \left(\frac{8}{21} \right)$$

(iii) On comparing with standard equation of planes

$$A_1 = 1, B_1 = 1, C_1 = -1 \text{ and } A_2 = 1, B_2 = 2, C_2 = 1$$

$$\cos \theta = \left| \frac{1 \cdot 1 + 1 \cdot 2 + (-1) \cdot 1}{\sqrt{1^2 + 1^2 + (-1)^2} \sqrt{1^2 + 2^2 + 1^2}} \right| \Rightarrow \left| \frac{1 + 2 + (-1)}{\sqrt{1 + 1 + 1} \sqrt{1 + 4 + 1}} \right| = \left| \frac{2}{\sqrt{3} \sqrt{6}} \right|$$

$$= \frac{\sqrt{2}}{3}$$

$$\theta = \cos^{-1} \left(\frac{\sqrt{2}}{3} \right)$$

(iv) On comparing with the standard equation of planes

$$A_1 = 1, B_1 = 1, C_1 = -2 \text{ and } A_2 = 2, B_2 = -2, C_2 = 1$$

$$\cos \theta = \frac{1.2 + 1.(-2) + (-2).1}{\sqrt{1^2 + 1^2 + (-2)^2} \sqrt{2^2 + (-2)^2 + 1^2}} \Rightarrow \frac{2 + (-2) + (-2)}{\sqrt{1+1+4} \sqrt{4+4+1}} = \frac{-2}{\sqrt{6}\sqrt{9}}$$

$$= \frac{2}{\sqrt{6}.3}$$

$$\theta = \cos^{-1} \left(\frac{2}{3\sqrt{6}} \right)$$

Question: 5

$$A_1x + B_1y + C_1z + D_1 = 0 \text{ and } A_2x + B_2y + C_2z + D_2 = 0 \quad \cos \theta = \frac{A_1A_2 + B_1B_2 + C_1C_2}{\sqrt{(A_1^2 + B_1^2 + C_1^2)} \sqrt{(A_2^2 + B_2^2 + C_2^2)}}$$

For $\theta = 90^\circ$, $\cos 90^\circ = 0$

$$A_1A_2 + B_1B_2 + C_1C_2 = 0$$

(i) On comparing with the standard equation of a plane

$$A_1 = 3, B_1 = 4, C_1 = -5 \text{ and } A_2 = 2, B_2 = 6, C_2 = 6$$

$$\text{LHS} = A_1A_2 + B_1B_2 + C_1C_2 \Rightarrow 3.2 + 4.6 + (-5).6 = 6 + 24 - 30$$

$$= 0 = \text{RHS}$$

Hence proved that the angle between planes is 90° .

(ii) On comparing with the standard equation of a plane

$$A_1 = 1, B_1 = -2, C_1 = 4 \text{ and } A_2 = 18, B_2 = 17, C_2 = 4$$

$$\text{LHS} = A_1A_2 + B_1B_2 + C_1C_2 \Rightarrow 1.18 + (-2).17 + 4.4 = 18 + (-34) + 16$$

$$= 0 = \text{RHS}$$

Hence proved that angle between planes is 90° .

Question: 6

$$A_1A_2 + B_1B_2 + C_1C_2 = 0$$

Where A_1, B_1, C_1 are direction ratios of plane and A_2, B_2, C_2 are of other plane.

$$2.1 + 2.2 + 4.2 = 2 + 4 + 8 = 14 \neq 0$$

Hence, planes are not perpendicular.

Similarly for the other plane

$$2.5 + 2.6 + 2.7 = 10 + 12 + 14 = 36 \neq 0$$

Hence, planes are not perpendicular.

Question: 7

Show that the planes

Solution:

To show that planes are parallel

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2}$$

On comparing with the standard equation of a plane

$$A_1 = 2, B_1 = -2, C_1 = 4 \text{ and } A_2 = 3, B_2 = -3, C_2 = 6$$

$$\frac{A_1}{A_2} = \frac{2}{3}, \frac{B_1}{B_2} = \frac{-2}{-3} \Rightarrow \frac{2}{3}, \frac{C_1}{C_2} = \frac{4}{6} \Rightarrow \frac{2}{3}$$

So,

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2} = \frac{2}{3}$$

Hence proved that planes are parallel.

Question: 8

Find the value of

Solution:

To find an angle in Cartesian form, for the standard equation of planes

$$A_1x + B_1y + C_1z + D_1 = 0 \text{ and } A_2x + B_2y + C_2z + D_2 = 0$$

$$\cos\theta = \left| \frac{A_1A_2 + B_1B_2 + C_1C_2}{\sqrt{(A_1^2 + B_1^2 + C_1^2)}\sqrt{(A_2^2 + B_2^2 + C_2^2)}} \right|$$

$$\text{For } \theta = 90^\circ, \cos 90^\circ = 0$$

$$A_1A_2 + B_1B_2 + C_1C_2 = 0$$

On comparing with the standard equation of the plane,

$$A_1 = 1, B_1 = -4, C_1 = \lambda \text{ and } A_2 = 2, B_2 = 2, C_2 = 3$$

$$A_1A_2 + B_1B_2 + C_1C_2 \Rightarrow 1.2 + (-4).2 + \lambda.3 = 0$$

$$2 + (-8) + 3\lambda = 0$$

$$-6 + 3\lambda = 0$$

$$\lambda = 2$$

Question: 9

$$A_1x + B_1y + C_1z + D_1 = 0$$

Direction ratios of parallel planes are related to each other as

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2} = k \text{ (constant)}$$

Putting the values from the equation of a given parallel plane,

$$\frac{A_1}{5} = \frac{B_1}{-3} = \frac{C_1}{7} = k$$

$$A_1 = 5k, B_1 = -3k, C_1 = 7k$$

Putting in equation plane

$$5kx - 3ky + 7kz + D_1 = 0$$

As the plane is passing through (0,0,0), it must satisfy the plane,

$$5k.0 - 3k.0 + 7k.0 + D_1 = 0$$

$$D_1 = 0$$

$$5kx - 3ky + 7kz = 0$$

$$5x - 3y + 7z = 0$$

So, required equation of plane is $5x - 3y + 7z = 0$.

Question: 10

Find the equation

Solution:

Let the equation of a plane

$$\vec{r} \cdot (\hat{x}_1\hat{i} + \hat{y}_1\hat{j} + \hat{z}_1\hat{k}) = d \quad (1)$$

Direction ratios of parallel planes are related to each other as

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2} = \lambda \text{ (constant)}$$

Putting the values from the equation of a given parallel plane,

$$\frac{x_1}{1} = \frac{x_1}{1} = \frac{z_1}{1} = \lambda$$

$$x_1 = y_1 = z_1 = \lambda$$

Putting values in equation (1), we have

$$\vec{r} \cdot (\lambda \hat{i} + \lambda \hat{j} + \lambda \hat{k}) = d \quad (2)$$

A plane passes through (a,b,c) then it must satisfy the equation of a plane

$$(\hat{a}\hat{i} + \hat{b}\hat{j} + \hat{c}\hat{k}) \cdot (\lambda \hat{i} + \lambda \hat{j} + \lambda \hat{k}) = d$$

$$\lambda(\hat{a}\hat{i} + \hat{b}\hat{j} + \hat{c}\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k}) = d$$

$$\lambda(a.1 + b.1 + c.1) = d$$

$$\lambda(a + b + c) = d$$

Putting value in equation (2)

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) \cdot \lambda = \lambda(a + b + c)$$

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = a + b + c$$

So, required equation of plane is $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = a + b + c$.

Question: 11

$$A_1x + B_1y + C_1z + D_1 = 0$$

Direction ratios of parallel planes are related to each other as

$$\frac{A_1}{A_2} = \frac{B_1}{B_2} = \frac{C_1}{C_2} = k \text{ (constant)}$$

Putting the values from the equation of a given parallel plane,

$$\frac{A_1}{5} = \frac{B_1}{4} = \frac{C_1}{-11} = k$$

$$A_1 = 5k, B_1 = 4k, C_1 = -11k$$

Putting in the equation of a plane

$$5kx + 4ky - 11kz + D_1 = 0$$

As the plane is passing through (1,-2,7), it must satisfy the plane,

$$5k.1 + 4k.(-2) - 11k.7 + D_1 = 0 \quad (1)$$

$$5k - 8k - 77k + D_1 = 0$$

$$D_1 = 80k$$

Putting value in equation (1), we have

$$5kx + 4ky - 11kz + 80k = 0$$

$$5x + 4y - 11z + 80 = 0$$

So, the required equation of the plane is $5x + 4y - 11z + 80 = 0$.

Question: 12

$$AA_1 + BB_1 + CC_1 = 0$$

Where A, B, C are direction ratios of plane and A_1, B_1, C_1 are of another plane.

$$3.A_1 + 2B_1 - 3C_1 = 0 \quad (1)$$

$$5.A_1 - 4B_1 + C_1 = 0 \quad (2)$$

And plane passes through (-1,-1,2),

$$A(x + 1) + B(y + 1) + C(z - 2) = 0 \quad (3)$$

On solving equation (1) and (2)

$$A = \frac{5B}{9} \text{ and } C = \frac{11B}{9}$$

Putting values in equation (3)

$$\frac{5B}{9} \cdot (x+1) + B(y+1) + \frac{11B}{9} \cdot (z-2) = 0$$

$$B(5x + 5 + 9y + 9 + 11z - 22) = 0$$

$$5x + 9y + 11z - 8 = 0$$

So, required equation of plane is $5x + 9y + 11z = 8$.

Question: 13

Find the equation

Solution:

Applying condition of perpendicularity between planes,

$$AA_1 + BB_1 + CC_1 = 0$$

Where A, B, C are direction ratios of plane and A_1, B_1, C_1 are of other plane.

$$1.A + 2.B - 1.C = 0$$

$$A + 2B - C = 0 \quad (1)$$

$$3.A - 4.B + C = 0$$

$$3A - 4B + C = 0 \quad (2)$$

And plane passes through $(0, 0, 0)$,

$$A(x-0) + B(y-0) + C(z-0) = 0$$

$$Ax + By + Cz = 0 \quad (3)$$

On solving equation (1) and (2)

$$A = \frac{B}{2} \text{ and } C = \frac{5B}{2}$$

Putting values in equation (3)

$$\frac{B}{2} \cdot x + By + \frac{5B}{2} \cdot z = 0$$

$$B(x + 2y + 5z) = 0$$

$$x + 2y + 5z = 0$$

So, required equation of plane is $x + 2y + 5z = 0$.

Question: 14

$$AA_1 + BB_1 + CC_1 = 0$$

Where A, B, C are direction ratios of plane and A_1, B_1, C_1 are of other plane.

$$3.A + 3.B - 2.C = 0$$

$$3A + 3B - 2C = 0 \quad (1)$$

$$1.A + 2.B - 3C = 0$$

$$A + 2B - 3C = 0 \quad (2)$$

And plane contains the point $(1, -1, 2)$,

$$A(x-1) + B(y+1) + C(z-2) = 0 \quad (3)$$

On solving equation (1) and (2)

$$A = \frac{-5B}{7} \text{ and } C = \frac{3B}{7}$$

Putting values in equation (3)

$$\frac{-5B}{7} \cdot (x-1) + B(y+1) + \frac{3B}{7} \cdot (z-2) = 0$$

$$B(-5(x-1) + 7(y+1) + 3(z-2)) = 0$$

$$-5x + 5 + 7y + 7 + 3z - 6 = 0$$

$$-5x + 7y + 3z + 6 = 0$$

$$5x - 7y - 3z - 6 = 0$$

For equation of plane $Ax + By + Cz = D$ and point (x_1, y_1, z_1) , distance of a point from a plane can be calculated as

$$\frac{|Ax_1 + By_1 + Cz_1 - D|}{\sqrt{A^2 + B^2 + C^2}}$$

$$\frac{|5(-2) - 7.5 - 3.5 - 6|}{\sqrt{(5)^2 + (-7)^2 + (-3)^2}} \Rightarrow \frac{|-10 - 35 - 15 - 6|}{\sqrt{25 + 49 + 9}} = \frac{|-66|}{\sqrt{83}} \Rightarrow \frac{66}{\sqrt{83}}$$

Question: 15

$$A(x-1) + B(y-1) + C(z-2) = 0 \quad (1)$$

$$A(x-2) + B(y+2) + C(z-2) = 0 \quad (2)$$

Subtracting (1) from (2),

$$A(x-2-x+1) + B(y+2-y-1) = 0$$

$$A-3B = 0 \quad (3)$$

Now plane is perpendicular to $6x-2y+2z=9$

$$6A-2B+2C=0 \quad (4)$$

Using (3) in (4)

$$18A-2B+2C=0$$

$$16B+2C=0$$

$$C=-8B$$

Putting values in equation (1)

$$3B(x-1) + B(y+2) - 8B(z-2) = 0$$

$$B(3x-3+y+2-8z+16) = 0$$

$$3x+y-8z+15=0$$

Question: 16

$$A(x+1) + B(y-1) + C(z-1) = 0 \quad (1)$$

$$A(x-1) + B(y+1) + C(z-1) = 0 \quad (2)$$

Subtracting (1) from (2),

$$A(x-1-x-1) + B(y+1-y-1) = 0$$

$$-2A+2B=0$$

$$A=B \quad (3)$$

Now plane is perpendicular to $x+2y+2z=5$

$$A+2B+2C=0 \quad (4)$$

Using (3) in (4)

$$B+2B+2C=0$$

$$3B+2C=0$$

$$C = -\frac{3}{2}B$$

Putting values in equation (1)

$$B(x+1) + B(y-1) - \frac{3}{2}B(z-1) = 0$$

$$B(2(x+1) + 2(y-1) - 3(z-1)) = 0$$

$$2x+2y-3z+2-2-3=0$$

$$2x+2y-3z-3=0$$

Question: 17

Find the equation

Solution:

Plane passes through $(3,4,2)$ and $(7,0,6)$,

$$A(x-3) + B(y-4) + C(z-2)=0 \quad (1)$$

$$A(x-7) + B(y-0) + C(z-6)=0 \quad (2)$$

Subtracting (1) from (2),

$$A(x-7-x + 3) + B(y-y + 4) + C(z-6-z + 2)=0$$

$$-4A + 4B-4C=0$$

$$A-B + C=0$$

$$B=A + C \quad (3)$$

Now plane is perpendicular to $2x-5y=15$

$$2A-5B=0 \quad (4)$$

Using (3) in (4)

$$2A-5(A + C)=0$$

$$2A-5A-5C=0$$

$$-3A-5C=0$$

$$C = \frac{-3}{5}A$$

$$B = A + \frac{-3}{5}A \Rightarrow \frac{2}{5}A$$

Putting values in equation (1)

$$A(x-3) + \frac{2}{5}A(y-4) + \frac{-3}{5}A(z-2) = 0$$

$$A(5(x-3) + 2(y-4)-3(z-2))=0$$

$$5x + 2y-3z-15-8 + 6=0$$

$$5x + 2y-3z-17=0$$

So, required equation of plane is $5x + 2y-3z-17=0$.

Question: 18

Plane passes through $(2,1,-1)$ and $(-1,3,4)$,

$$A(x-2) + B(y-1) + C(z+1)=0 \quad (1)$$

$$A(x+1) + B(y-3) + C(z-4)=0 \quad (2)$$

Subtracting (1) from (2),

$$A(x+1-x+2) + B(y-3-y+1) + C(z-4-z-1)=0$$

$$3A-2B-5C=0 \quad (3)$$

Now plane is perpendicular to $x-2y+4z=10$

$$A-2B+4C=0 \quad (4)$$

Using (3) in (4)

$$2A-9C=0$$

$$C = \frac{2}{9}A$$

$$2B = A + 4 \cdot \frac{2}{9}A \Rightarrow \left(\frac{9+8}{9}\right)A = \frac{17}{9}A$$

$$B = \frac{17}{18}A$$

Putting values in equation (1)

$$A(x-2) + \frac{17}{18}A(y-1) + \frac{2}{9}A(z+1) = 0$$

$$A(18(x-2) + 17(y-1) + 4(z+1))=0$$

$$18x + 17y + 4z-36-17 + 4=0$$

$$18x + 17y + 4z-49=0$$

So, the required equation of plane is $18x + 17y + 4z-49=0$

If plane contains $\vec{r} = -\hat{i} + 3\hat{j} + 4\hat{k} + (3\hat{i} - 2\hat{j} - 5\hat{k})$ then $(-1, 3, 4)$ satisfies plane and normal vector of plane is perpendicular to

$$LHS=18(-1) + 17.3 + 4.4-49$$

$$= -18 + 51 + 16 - 49$$

$$= -2 + 2 = 0 = \text{RHS}$$

In vector form normal of plane

$$\vec{n} = 18\hat{i} + 17\hat{j} + 4\hat{k}$$

$$\text{LHS} = 18.3 + 17(-2) + 4(-5) = 54 - 34 - 20 = 0 = \text{RHS}$$

Hence line is contained in plane.

Exercise : 28G

Question: 1

Find the angle be

Solution:

Given - $\vec{r} = (\hat{i} + 2\hat{j} - \hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$ and $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 4$

To find - The angle between the line and the plane

Direction ratios of the line = (1, -1, 1)

Direction ratios of the normal of the plane = (2, -1, 1)

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$= \sin^{-1} \left(\frac{1 \times 2 + (-1) \times (-1) + 1 \times 1}{\sqrt{1^2 + 1^2 + 1^2} \sqrt{2^2 + 1^2 + 1^2}} \right)$$

$$= \sin^{-1} \left(\frac{2 + 1 + 1}{\sqrt{3} \sqrt{6}} \right)$$

$$= \sin^{-1} \left(\frac{4}{3\sqrt{2}} \right)$$

$$= \sin^{-1} \left(\frac{2\sqrt{2}}{3} \right)$$

Question: 2

Find the angle be

Solution:

Given - $\vec{r} = (2\hat{i} - \hat{j} + 3\hat{k}) + \lambda(3\hat{i} - \hat{j} + 2\hat{k})$ and $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 3$

To find - The angle between the line and the plane

Direction ratios of the line = (3, -1, 2)

Direction ratios of the normal of the plane = (1, 1, 1)

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$= \sin^{-1} \left(\frac{3 \times 1 + (-1) \times 1 + 2 \times 1}{\sqrt{3^2 + 1^2 + 2^2} \sqrt{1^2 + 1^2 + 1^2}} \right)$$

$$= \sin^{-1} \left(\frac{3 - 1 + 2}{\sqrt{14} \sqrt{3}} \right)$$

$$= \sin^{-1} \left(\frac{4}{\sqrt{42}} \right)$$

Question: 3

Find the angle be

Solution:

Given - $\vec{r} = (3\hat{i} + \hat{k}) + \lambda(\hat{j} + \hat{k})$ and $\vec{r} \cdot (2\hat{i} - \hat{j} + 2\hat{k}) = 1$

To find - The angle between the line and the plane

Direction ratios of the line = (0, 1, 1)

Direction ratios of the normal of the plane = (2, -1, 2)

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$\begin{aligned} &= \sin^{-1} \left(\frac{0 \times 2 + 1 \times (-1) + 1 \times 2}{\sqrt{0^2 + 1^2 + 1^2} \sqrt{2^2 + 1^2 + 2^2}} \right) \\ &= \sin^{-1} \left(\frac{-1 + 2}{3\sqrt{2}} \right) \\ &= \sin^{-1} \left(\frac{1}{3\sqrt{2}} \right) \end{aligned}$$

Question: 4

Find the angle be

Solution:

Given - $\frac{x-2}{3} = \frac{y+1}{-1} = \frac{z-3}{2}$ and $3x + 4y + z + 5 = 0$

To find - The angle between the line and the plane

Direction ratios of the line = (3, -1, 2)

Direction ratios of the normal of the plane = (3, 4, 1)

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$\begin{aligned} &= \sin^{-1} \left(\frac{3 \times 3 + (-1) \times 4 + 2 \times 1}{\sqrt{3^2 + 1^2 + 2^2} \sqrt{3^2 + 4^2 + 1^2}} \right) \\ &= \sin^{-1} \left(\frac{9 - 4 + 2}{\sqrt{14} \sqrt{26}} \right) \\ &= \sin^{-1} \left(\frac{7}{\sqrt{2} \sqrt{7} \times \sqrt{2} \times \sqrt{13}} \right) \\ &= \sin^{-1} \left(\frac{7}{2\sqrt{91}} \right) \end{aligned}$$

Question: 5

Find the angle be

Solution:

Given - $\frac{x+1}{2} = \frac{y}{3} = \frac{z-3}{6}$ and $10x + 2y - 11z = 3$

To find - The angle between the line and the plane

Direction ratios of the line = (2, 3, 6)

Direction ratios of the normal of the plane = (10, 2, -11)

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$\begin{aligned} &= \sin^{-1} \left(\frac{2 \times 10 + 3 \times 2 + 6 \times (-11)}{\sqrt{2^2 + 3^2 + 6^2} \sqrt{10^2 + 2^2 + 11^2}} \right) \\ &= \sin^{-1} \left(\frac{20 + 6 - 66}{7 \times 15} \right) \\ &= \sin^{-1} \left(\frac{-40}{7 \times 15} \right) \\ &= \sin^{-1} \left(-\frac{8}{21} \right) \end{aligned}$$

Question: 6

Find the angle be

Solution:

Given - $A = (3, -4, -2)$, $B = (12, 2, 0)$ and $3x - y + z = 1$

To find - The angle between the line joining the points A and B and the plane

Tip - If $P = (a, b, c)$ and $Q = (a', b', c')$, then the direction ratios of the line PQ is given by $((a' - a), (b' - b), (c' - c))$

The direction ratios of the line AB can be given by

$$((12 - 3), (2 + 4), (0 + 2))$$

$$= (9, 6, 2)$$

Direction ratios of the normal of the plane = $(3, -1, 1)$

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$= \sin^{-1} \left(\frac{9 \times 3 + 6 \times (-1) + 2 \times 1}{\sqrt{9^2 + 6^2 + 2^2} \sqrt{3^2 + 1^2 + 1^2}} \right)$$

$$= \sin^{-1} \left(\frac{27 - 6 + 2}{11 \times \sqrt{11}} \right)$$

$$= \sin^{-1} \left(\frac{23}{11\sqrt{11}} \right)$$

Question: 7

If the plane $2x -$

Solution:

Given - $y = z = 0$ and $2x - 3y - 6z = 13$

To find - The angle between the line and the plane

Direction ratios of the line = $(1, 0, 0)$

Direction ratios of the normal of the plane = $(2, -3, -6)$

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$= \sin^{-1} \left(\frac{1 \times 2 + 0 \times (-3) + 0 \times (-6)}{\sqrt{1^2 + 0^2 + 0^2} \sqrt{2^2 + 3^2 + 6^2}} \right)$$

$$= \sin^{-1} \left(\frac{2}{7} \right)$$

Question: 8

Show that the lin

Solution:

Given - $\vec{r} = (2\hat{i} + 5\hat{j} + 7\hat{k}) + \lambda(\hat{i} + 3\hat{j} + 4\hat{k})$ and $\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 7$

To prove - The line and the plane are parallel &

To find - The distance between them

Direction ratios of the line = $(1, 3, 4)$

Direction ratios of the normal of the plane = $(1, 1, -1)$

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

The angle between the line and the plane

$$= \sin^{-1} \left(\frac{1 \times 1 + 3 \times 1 + 4 \times (-1)}{\sqrt{1^2 + 3^2 + 4^2} \sqrt{1^2 + 1^2 + 1^2}} \right)$$

$$= \sin^{-1} \left(\frac{1 + 3 - 4}{\sqrt{26}\sqrt{3}} \right)$$

$$= \sin^{-1}(0)$$

$$= 0$$

Hence, **the line and the plane are parallel.**

Now, the equation of the plane may be written as $x + y - z = 7$.

Tip - If $ax + by + c + d = 0$ be a plane and $\vec{r} = (a'\hat{i} + b'\hat{j} + c'\hat{k}) + \lambda(a''\hat{i} + b''\hat{j} + c''\hat{k})$ be a line vector, then the distance between them is given by $\left| \frac{a \times a' + b \times b' + c \times c' + d}{\sqrt{a^2 + b^2 + c^2}} \right|$

The distance between the plane and the line

$$= \left| \frac{1 \times 2 + 1 \times 5 - 1 \times 7 - 7}{\sqrt{1^2 + 1^2 + 1^2}} \right|$$

$$= \left| \frac{2 + 5 - 7 - 7}{\sqrt{3}} \right|$$

$$= \frac{7}{\sqrt{3}} \text{ units}$$

Question: 9

Find the value of

Solution:

Given - $\vec{r} = (\hat{i} + 2\hat{k}) + \lambda(2\hat{i} - m\hat{j} - 3\hat{k})$ and $\vec{r} \cdot (m\hat{i} + 3\hat{j} + \hat{k}) = 4$ and they are parallel

To find - The value of m

Direction ratios of the line = $(2, -m, -3)$

Direction ratios of the normal of the plane = $(m, 3, 1)$

Formula to be used - If (a, b, c) be the direction ratios of a line and (a', b', c') be the direction ratios of the normal to the plane, then, the angle between the two is given by

$$\sin^{-1} \left(\frac{a \times a' + b \times b' + c \times c'}{\sqrt{a^2 + b^2 + c^2} \sqrt{a'^2 + b'^2 + c'^2}} \right)$$

$$\therefore \sin^{-1} \left(\frac{2 \times m + (-m) \times 3 + (-3) \times 1}{\sqrt{2^2 + m^2 + 3^2} \sqrt{m^2 + 3^2 + 1^2}} \right) = 0$$

$$\Rightarrow \sin^{-1} \left(\frac{2m - 3m - 3}{\sqrt{13 + m^2} \sqrt{10 + m^2}} \right) = 0$$

$$\Rightarrow \frac{-m - 3}{\sqrt{13 + m^2} \sqrt{10 + m^2}} = 0$$

$$\Rightarrow m = -3$$

Question: 10

Find the vector e

Solution:

Given - $\vec{r} \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) = 3$

To find - The vector equation of the line passing through the origin and perpendicular to the given plane

Tip - The equation of a plane can be given by $\vec{r} \cdot \hat{n} = d$ where $\hat{n}$ is the normal of the plane

A line parallel to the given plane will be in the direction of the normal and will have the direction ratios same as that of the normal.

Formula to be used - If a line passes through the point (a, b, c) and has the direction ratios as (a', b', c') , then its vector equation is given by $\vec{r} = (a\hat{i} + b\hat{j} + c\hat{k}) + \lambda(a'\hat{i} + b'\hat{j} + c'\hat{k})$ where λ is any scalar constant

The required equation will be $\vec{r} = (0.\hat{i} + 0.\hat{j} + 0.\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$

= $\lambda(\hat{i} + 2\hat{j} + 3\hat{k})$ for some scalar λ

Question: 11

Find the vector e

Solution:

Given - $\vec{r} \cdot (2\hat{i} - 3\hat{j} - \hat{k}) = 0$ and the vector has position vector $(\hat{i} - 2\hat{j} + 5\hat{k})$

To find - The vector equation of the line passing through $(1, -2, 5)$ and perpendicular to the given plane

Tip - The equation of a plane can be given by $\vec{r} \cdot \hat{n} = d$ where $\hat{n}$ is the normal of the plane

A line parallel to the given plane will be in the direction of the normal and will have the direction ratios same as that of the normal.

Formula to be used - If a line passes through the point (a, b, c) and has the direction ratios as (a', b', c') , then its vector equation is given by $\vec{r} = (a\hat{i} + b\hat{j} + c\hat{k}) + \lambda(a'\hat{i} + b'\hat{j} + c'\hat{k})$ where λ is any scalar constant

The required equation will be $\vec{r} = (\hat{i} - 2\hat{j} + 5\hat{k}) + \lambda(2\hat{i} - 3\hat{j} - \hat{k})$ for some scalar λ

Question: 12

Show that the equ

Solution:

Given - The equation of the plane is given by $ax + by + d = 0$

To prove - The plane is parallel to z - axis

Tip - If $ax + by + cz + d$ is the equation of the plane then its angle with the z - axis will be given by $\sin^{-1}\left(\frac{c}{\sqrt{a^2 + b^2 + c^2}}\right)$

Considering the equation, the direction ratios of its normal is given by $(a, b, 0)$

The angle the plane makes with the z - axis $= \sin^{-1}[0/\sqrt{a^2 + b^2}] = 0$

Hence, **the plane is parallel to the z - axis**

To find - Equation of the plane parallel to z - axis and passing through points $A = (2, -3, 1)$ and $B = (-4, 7, 6)$

The given equation $ax + by + d = 0$ passes through $(2, -3, 1)$ & $(-4, 7, 6)$

$$\therefore 2a - 3b + d = 0 \dots\dots(i)$$

$$\therefore -4a + 7b + d = 0 \dots\dots(ii)$$

Solving (i) and (ii),

$$\therefore \frac{a}{\begin{vmatrix} -3 & 1 \\ 7 & 1 \end{vmatrix}} = \frac{b}{\begin{vmatrix} 1 & 2 \\ 1 & -4 \end{vmatrix}} = \alpha [\alpha \rightarrow \text{arbitrary constant}]$$

$$\therefore a = -10\alpha$$

$$\therefore b = -6\alpha$$

Substituting the values of a and b in eqn (i), we get,

$$-2 \times 10\alpha + 3 \times 6\alpha + d = 0 \text{ i.e. } d = -2\alpha$$

Putting the value of a, b and d in the equation $ax + by + d = 0$,

$$(-10\alpha)x + (-6\alpha)y + (-2\alpha) = 0$$

$$\text{i.e. } 5x + 3y + 1 = 0$$

Question: 13

Find the equation

Solution:

Given - A plane passes through points $(1, 2, 3)$ and $(0, -1, 0)$ and is parallel to the line

$$\frac{x-1}{2} = \frac{y+2}{3} = \frac{z}{-3}$$

To find - Equation of the plane

Tip - If a plane passes through points (a', b', c') , then its equation may be given as $a(x - a') + b(y - b') + c(z - c') = 0$

Taking points $(1, 2, 3)$:

$$a(x - 1) + b(y - 2) + c(z - 3) = 0 \dots\dots(i)$$

The plane passes through $(0, -1, 0)$:

$$a(0 - 1) + b(-1 - 2) + c(0 - 3) = 0$$

$$\text{i.e. } a + 3b + 3c = 0 \dots\dots(ii)$$

$$\text{The plane is parallel to the line } \frac{x-1}{2} = \frac{y+2}{3} = \frac{z}{-3}$$

Tip - The normal of the plane will be normal to the given line since both the line and plane are parallel.

Direction ratios of the line is $(2, 3, -3)$

Direction ratios of the normal of the plane is (a, b, c)

$$\text{So, } 2a + 3b - 3c = 0 \dots\dots(iii)$$

Solving equations (ii) and (iii),

$$\therefore \frac{a}{\begin{vmatrix} 3 & 3 \\ 3 & -3 \end{vmatrix}} = -\frac{b}{\begin{vmatrix} 1 & 3 \\ 2 & -3 \end{vmatrix}} = \frac{c}{\begin{vmatrix} 1 & 3 \\ 2 & 3 \end{vmatrix}} = \alpha [\alpha \rightarrow \text{arbitrary constant}]$$

$$\therefore a = -18\alpha$$

$$\therefore b = 9\alpha$$

$$\therefore c = -3\alpha$$

Putting these values in equation (i) we get,

$$-18\alpha(x-1) + 9\alpha(y-2) - 3\alpha(z-3) = 0$$

$$\Rightarrow 18(x-1) - 9(y-2) + 3(z-3) = 0$$

$$\Rightarrow 6(x-1) - 3(y-2) + (z-3) = 0$$

$$\Rightarrow 6x - 3y + z - 3 = 0$$

$$\Rightarrow 6x - 3y + z = 3$$

Question: 14

Find the equation

Solution:

Given - A plane passes through (2, -1, 5), perpendicular to the plane $x + 2y - 3z = 7$ and parallel to the line $\frac{x+5}{3} = \frac{y+1}{-1} = \frac{z-2}{1}$

To find - The equation of the plane

Let the equation of the required plane be $ax + by + cz + d = 0$(a)

The plane passes through (2, -1, 5)

$$\text{So, } 2a - b + 5c + d = 0 \text{.....(i)}$$

The direction ratios of the normal of the plane is given by (a, b, c)

Now, this plane is perpendicular to the plane $x + 2y - 3z = 7$ having direction ratios (1, 2, -3)

$$\text{So, } a + 2b - 3c = 0 \text{.....(ii)}$$

This plane is also parallel to the line having direction ratios (3, -1, 1)

So, the direction of the normal of the required plane is also at right angles to the given line.

$$\text{So, } 3a - b + c = 0 \text{.....(iii)}$$

Solving equations (ii) and (iii),

$$\therefore \frac{a}{\begin{vmatrix} 2 & -3 \\ -1 & 1 \end{vmatrix}} = -\frac{b}{\begin{vmatrix} 1 & -3 \\ 3 & 1 \end{vmatrix}} = \frac{c}{\begin{vmatrix} 1 & 2 \\ 3 & -1 \end{vmatrix}} = \alpha [\alpha \rightarrow \text{arbitrary constant}]$$

$$\therefore a = -\alpha$$

$$\therefore b = -10\alpha$$

$$\therefore c = -7\alpha$$

Putting these values in equation (i) we get,

$$2X(-\alpha) - (-10\alpha) + 5(-7\alpha) + d = 0 \text{ i.e. } d = 27\alpha$$

Substituting all the values of a, b, c and d in equation (a) we get,

$$-\alpha x - 10\alpha y - 7\alpha z + 27\alpha = 0$$

$$\Rightarrow x + 10y + 7z + 27 = 0$$

Question: 15

Find the equation

Solution:

Given - A plane passes through the intersection of $5x - y + z = 10$ and $x + y - z = 4$ and parallel to the line with direction ratios (2, 1, 1)

To find - Equation of the plane

Tip - If $ax + by + cz + d = 0$ and $a'x + b'y + c'z + d' = 0$ be two planes, then the equation of the plane passing through their intersection will be given by

$$(ax + by + cz + d) + \lambda(a'x + b'y + c'z + d') = 0, \text{ where } \lambda \text{ is any scalar constant}$$

So, the equation of the plane maybe written as

$$(5x - y + z - 10) + \lambda(x + y - z - 4) = 0$$

$$= (5 + \lambda)x + (-1 + \lambda)y + (1 - \lambda)z + (-10 - 4\lambda) = 0$$

This is plane parallel to a line with direction ratios (2, 1, 1)

So, the normal of this line with direction ratios ((5 + λ), (-1 + λ), (1 - λ)) will be perpendicular to

the given line.

Hence,

$$2(5 + \lambda) + (-1 + \lambda) + (1 - \lambda) = 0$$

$$\Rightarrow \lambda = -5$$

The equation of the plane will be

$$(5 + (-5))x + (-1 + (-5))y + (1 - (-5))z + (-10 - 4(-5)) = 0$$

$$\Rightarrow -6y + 6z + 10 = 0$$

$$\Rightarrow 3y - 3z = 5$$

To find - Perpendicular distance of point (1, 1, 1) from the plane

Formula to be used - If $ax + by + c + d = 0$ be a plane and (a', b', c') be the point, then the distance between them is given by $\left| \frac{a \times a' + b \times b' + c \times c' + d}{\sqrt{a^2 + b^2 + c^2}} \right|$

The distance between the plane and the line

$$= \left| \frac{0 \times 2 + 3 \times 1 - 3 \times 1 - 5}{\sqrt{0^2 + 3^2 + 3^2}} \right|$$

$$= \left| \frac{3 - 3 - 5}{2\sqrt{3}} \right|$$

$$= \frac{5}{2\sqrt{3}} \text{ units}$$

Exercise : 28H

Question: 1

Find the vector a

Solution:

Given - $\vec{r} = \hat{i} + \hat{j} - \hat{k}$ & $\vec{r'} = 3\hat{i} - \hat{k}$ are two lines to which a plane is parallel and it passes through the origin.

To find - The equation of the plane

Tip - A plane parallel to two vectors will have its normal in a direction perpendicular to both the vectors, which can be evaluated by taking their cross product

$$\therefore \vec{r} \times \vec{r'}$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & -1 \\ 3 & 0 & -1 \end{vmatrix}$$

$$= \hat{i}(-1 - 0) + \hat{j}(-3 + 1) + \hat{k}(0 - 3)$$

$$= -\hat{i} - 2\hat{j} - 3\hat{k}$$

The plane passes through origin (0, 0, 0).

Formula to be used - If a line passes through the point (a, b, c) and has the direction ratios as (a', b', c') , then its vector equation is given by $\vec{r} = (a\hat{i} + b\hat{j} + c\hat{k}) + \lambda(a'\hat{i} + b'\hat{j} + c'\hat{k})$ where λ is any scalar constant

The required plane will be

$$\vec{r} = (0\hat{i} + 0\hat{j} + 0\hat{k}) + \lambda'(-\hat{i} - 2\hat{j} - 3\hat{k})$$

$$\Rightarrow \vec{r} = \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$$

$$\text{The vector equation : } \vec{r} \cdot (\hat{i} + 2\hat{j} + 3\hat{k}) = 0$$

$$\text{The Cartesian equation : } x + 2y + 3z = 0$$

Question: 3

Find the vector a

Solution:

Given - $\vec{r} = (-\hat{j} + 3\hat{k}) + \lambda(2\hat{i} - 5\hat{j} - \hat{k})$ & $\vec{r} = (\hat{i} - 3\hat{j} + \hat{k}) + \mu(-5\hat{i} + 4\hat{j})$. A plane is parallel to both these lines and passes through (3, -1, 2).

To find - The equation of the plane

Tip - A plane parallel to two vectors will have its normal in a direction perpendicular to both the vectors, which can be evaluated by taking their cross product

$$\vec{R} = 2\hat{i} - 5\hat{j} - \hat{k} \text{ \& } \vec{R'} = -5\hat{i} + 4\hat{j}, \text{ where the two vectors represent the directions}$$

$$\therefore \vec{R} \times \vec{R'}$$

$$\begin{aligned}
&= \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -5 & -1 \\ -5 & 4 & 0 \end{bmatrix} \\
&= \hat{i}(0 + 4) + \hat{j}(5 - 0) + \hat{k}(8 - 25) \\
&= 4\hat{i} + 5\hat{j} - 17\hat{k}
\end{aligned}$$

The equation of the plane maybe represented as $4x + 5y - 17z + d = 0$

Now, this plane passes through the point $(3, -1, 2)$

Hence,

$$4 \times 3 + 5 \times (-1) - 17 \times 2 + d = 0$$

$$= d = 27$$

The Cartesian equation of the plane : $4x + 5y - 17z + 27 = 0$

The vector equation : $\vec{r} \cdot (4\hat{i} + 5\hat{j} - 17\hat{k}) + 27 = 0$

Question: 3

Find the vector e

Solution:

Given - The lines have direction ratios of $(1, -1, -2)$ and $(-1, 0, 2)$. The plane parallel to these lines passes through $(1, 2, 3)$

To find - The vector equation of the plane

Tip - A plane parallel to two vectors will have its normal in a direction perpendicular to both the vectors, which can be evaluated by taking their cross product

$$\vec{R} = \hat{i} - \hat{j} - 2\hat{k} \text{ \& } \vec{R'} = -\hat{i} + 2\hat{k}, \text{ where the two vectors represent the directions}$$

$$\therefore \vec{R} \times \vec{R'}$$

$$\begin{aligned}
&= \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & -2 \\ -1 & 0 & 2 \end{bmatrix} \\
&= \hat{i}(-2 - 0) + \hat{j}(2 - 2) + \hat{k}(0 - 1) \\
&= -2\hat{i} - \hat{k}
\end{aligned}$$

The equation of the plane maybe represented as $-2x - z + d = 0$

Now, this plane passes through the point $(1, 2, 3)$

Hence,

$$(-2) \times 1 - 3 + d = 0$$

$$= d = 5$$

The Cartesian equation of the plane : $-2x - z + 5 = 0$ i.e. $2x + z = 5$

The vector equation : $\vec{r} \cdot (2\hat{i} + \hat{k}) = 5$

Question: 4

Find the Cartesian

Solution:

Given - $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z+1}{6}$ & $\frac{x-1}{1} = \frac{y+3}{1} = \frac{z}{-1}$. A plane is parallel to both these lines and passes through $(1, 2, -4)$.

To find - The equation of the plane

Tip - A plane parallel to two vectors will have its normal in a direction perpendicular to both the vectors, which can be evaluated by taking their cross product

The direction ratios of the given lines are $(2, 3, 6)$ and $(1, 1, -1)$

$$\therefore \vec{R} = 2\hat{i} + 3\hat{j} + 6\hat{k} \text{ \& } \vec{R'} = \hat{i} + \hat{j} - \hat{k}$$

$$\therefore \vec{R} \times \vec{R'}$$

$$\begin{aligned}
&= \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 6 \\ 1 & 1 & -1 \end{bmatrix} \\
&= \hat{i}(-3 - 6) + \hat{j}(6 + 2) + \hat{k}(2 - 3) \\
&= -9\hat{i} + 8\hat{j} - \hat{k}
\end{aligned}$$

The equation of the plane maybe represented as $-9x + 8y - z + d = 0$

Now, this plane passes through the point $(1, 2, -4)$

Hence,

$$(-9) \times 1 + 8 \times 2 - (-4) + d = 0$$

$$\Rightarrow d = -11$$

The Cartesian equation of the plane : $-9x + 8y - z - 11 = 0$ i.e. **$9x - 8y + z + 11 = 0$**

The vector equation : **$\vec{r} \cdot (9\hat{i} - 8\hat{j} + \hat{k}) + 11 = 0$**

Question: 5

Find the vector e

Solution:

Given - $\vec{r} = \hat{i} + 2\hat{j} + 3\hat{k}$ & $\vec{r'} = \hat{i} - \hat{j} + \hat{k}$ are two lines to which a plane is parallel and it passes through the point $3\hat{i} + 4\hat{j} + 2\hat{k}$

To find - The equation of the plane

Tip - A plane parallel to two vectors will have its normal in a direction perpendicular to both the vectors, which can be evaluated by taking their cross product

$$\therefore \vec{r} \times \vec{r'}$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 1 & -1 & 1 \end{vmatrix}$$

$$= \hat{i}(2 + 3) + \hat{j}(3 - 1) + \hat{k}(-1 - 2)$$

$$= 5\hat{i} + 2\hat{j} - 3\hat{k}$$

The equation of the plane maybe represented as $5x + 2y - 3z + d = 0$

Now, this plane passes through the point (3, 4, 2)

Hence,

$$5 \times 3 + 2 \times 4 - 3 \times 2 + d = 0$$

$$\Rightarrow d = -17$$

The Cartesian equation of the plane : $5x + 2y - 3z - 17 = 0$ i.e. $5x + 2y - 3z = 17$

The vector equation : **$\vec{r} \cdot (5\hat{i} + 2\hat{j} - 3\hat{k}) = 17$**

Exercise : 28I

Question: 1

Show that the lin

Solution:

Given : Equations of lines -

$$\vec{r}_1 = (2\hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$$

$$\vec{r}_2 = (2\hat{i} + 6\hat{j} + 3\hat{k}) + \mu(2\hat{i} + 3\hat{j} + 4\hat{k})$$

To Prove : $\vec{r}_1$ & $\vec{r}_2$ are coplanar.

To Find : Equation of plane.

Formulae :

1) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (\vec{a}_1 \times \vec{b}_1) + (\vec{a}_2 \times \vec{b}_2) + (\vec{a}_3 \times \vec{b}_3)$$

3) Coplanarity of two lines :

If two lines $\vec{r}_1 = \vec{a} + \lambda \vec{b}$ & $\vec{r}_2 = \vec{c} + \mu \vec{d}$ are coplanar then

$$\vec{a} \cdot (\vec{b} \times \vec{d}) = \vec{c} \cdot (\vec{b} \times \vec{d})$$

4) Equation of plane :

If two lines $\vec{r}_1 = \vec{a}_1 + \lambda \vec{b}_1$ & $\vec{r}_2 = \vec{a}_2 + \lambda \vec{b}_2$ are coplanar then equation of the plane containing them is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Answer :

Given equations of lines are

$$\vec{r}_1 = (2\hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$$

$$\vec{r}_2 = (2\hat{i} + 6\hat{j} + 3\hat{k}) + \mu(2\hat{i} + 3\hat{j} + 4\hat{k})$$

$$\text{Let, } \vec{r}_1 = \vec{a}_1 + \lambda \vec{b}_1 \text{ \& } \vec{r}_2 = \vec{a}_2 + \lambda \vec{b}_2$$

Where,

$$\vec{a}_1 = 2\hat{j} - 3\hat{k}$$

$$\vec{b}_1 = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{a}_2 = 2\hat{i} + 6\hat{j} + 3\hat{k}$$

$$\vec{b}_2 = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix}$$

$$= \hat{i}(8 - 9) - \hat{j}(4 - 6) + \hat{k}(3 - 4)$$

$$\therefore (\vec{b}_1 \times \vec{b}_2) = -\hat{i} + 2\hat{j} - \hat{k}$$

Therefore,

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = (0 \times (-1)) + (2 \times 2) + ((-3) \times (-1))$$

$$= 0 + 4 + 3$$

$$= 7$$

$$\therefore \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = 7 \dots\dots\dots \text{eq(1)}$$

And

$$\vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2) = (2 \times (-1)) + (6 \times 2) + (3 \times (-1))$$

$$= -2 + 12 - 3$$

$$= 7$$

$$\therefore \vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2) = 7 \dots\dots\dots \text{eq(2)}$$

From eq(1) and eq(2)

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Hence lines $\vec{r}_1$ & $\vec{r}_2$ are coplanar.

Equation of plane containing lines $\vec{r}_1$ & $\vec{r}_2$ is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = -\hat{i} + 2\hat{j} - \hat{k}$$

From eq(1)

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = 7$$

Therefore, equation of required plane is

$$\vec{r} \cdot (-\hat{i} + 2\hat{j} - \hat{k}) = 7$$

$$\therefore \vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) = -7$$

$$\therefore \vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) + 7 = 0$$

$$\vec{r} \cdot (\hat{i} - 2\hat{j} + \hat{k}) + 7 = 0$$

Question: 2

Find the vector a

Solution:

Given : Equations of lines -

$$\vec{r}_1 = (\hat{i} + 2\hat{j} - 4\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k})$$

$$\vec{r}_2 = (9\hat{i} + 5\hat{j} - \hat{k}) + \mu(-2\hat{i} + 3\hat{j} + 8\hat{k})$$

To Find : Equation of plane.

Formulae :

1) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

If two lines $\vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$ & $\vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$ are coplanar then equation of the plane containing them is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Where,

Given equations of lines are

$$\vec{r}_1 = (\hat{i} + 2\hat{j} - 4\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k})$$

$$\vec{r}_2 = (9\hat{i} + 5\hat{j} - \hat{k}) + \mu(-2\hat{i} + 3\hat{j} + 8\hat{k})$$

$$\text{Let, } \vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1 \text{ \& } \vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$$

Where,

$$\vec{a}_1 = \hat{i} + 2\hat{j} - 4\hat{k}$$

$$\vec{b}_1 = 2\hat{i} + 3\hat{j} + 6\hat{k}$$

$$\vec{a}_2 = 9\hat{i} + 5\hat{j} - \hat{k}$$

$$\vec{b}_2 = -2\hat{i} + 3\hat{j} + 8\hat{k}$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 6 \\ -2 & 3 & 8 \end{vmatrix}$$

$$= \hat{i}(24 - 18) - \hat{j}(16 + 12) + \hat{k}(6 + 6)$$

$$\therefore (\vec{b}_1 \times \vec{b}_2) = 6\hat{i} - 28\hat{j} + 12\hat{k}$$

Therefore,

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = (1 \times 6) + (2 \times (-28)) + ((-4) \times 12)$$

$$= 6 - 56 - 48$$

$$= -98$$

$$\therefore \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = -98 \dots\dots\dots eq(1)$$

Equation of plane containing lines $\vec{r}_1$ & $\vec{r}_2$ is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = 6\hat{i} - 28\hat{j} + 12\hat{k}$$

From eq(1)

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = -98$$

Therefore, equation of required plane is

$$\vec{r} \cdot (6\hat{i} - 28\hat{j} + 12\hat{k}) = -98$$

$$\therefore \vec{r} \cdot (6\hat{i} - 28\hat{j} + 12\hat{k}) + 98 = 0$$

This vector equation of plane.

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore \vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = (x \times 6) + (y \times (-28)) + (z \times 12)$$

$$= 6x - 28y + 12z$$

Therefore, equation of plane is

$$6x - 28y + 12z = -98$$

$$6x - 28y + 12z + 98 = 0$$

This Cartesian equation of plane.

Question: 3

Find the vector a

Solution:

Given : Equations of lines -

$$\vec{r}_1 = (2\hat{i} + \hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 5\hat{k})$$

$$\vec{r}_2 = (3\hat{i} + 3\hat{j} + 2\hat{k}) + \mu(3\hat{i} - 2\hat{j} + 5\hat{k})$$

To Prove : $\vec{r}_1$ & $\vec{r}_2$ are coplanar.

To Find : Equation of plane.

Formulae :

1) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Coplanarity of two lines :

If two lines $\vec{r}_1 = \vec{a} + \lambda\vec{b}$ & $\vec{r}_2 = \vec{c} + \mu\vec{d}$ are coplanar then

$$\vec{a} \cdot (\vec{b} \times \vec{d}) = \vec{c} \cdot (\vec{b} \times \vec{d})$$

4) Equation of plane :

If two lines $\vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1$ & $\vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$ are coplanar then equation of the plane containing them is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Where,

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Answer :

Given equations of lines are

$$\vec{r}_1 = (2\hat{i} + \hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 5\hat{k})$$

$$\vec{r}_2 = (3\hat{i} + 3\hat{j} + 2\hat{k}) + \mu(3\hat{i} - 2\hat{j} + 5\hat{k})$$

$$\text{Let, } \vec{r}_1 = \vec{a}_1 + \lambda\vec{b}_1 \text{ \& } \vec{r}_2 = \vec{a}_2 + \lambda\vec{b}_2$$

Where,

$$\vec{a}_1 = 2\hat{i} + \hat{j} - 3\hat{k}$$

$$\vec{b}_1 = \hat{i} + 2\hat{j} + 5\hat{k}$$

$$\vec{a}_2 = 3\hat{i} + 3\hat{j} + 2\hat{k}$$

$$\vec{b}_2 = 3\hat{i} - 2\hat{j} + 5\hat{k}$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 5 \\ 3 & -2 & 5 \end{vmatrix}$$

$$= \hat{i}(10 + 10) - \hat{j}(5 - 15) + \hat{k}(-2 - 6)$$

$$\therefore (\vec{b}_1 \times \vec{b}_2) = 20\hat{i} + 10\hat{j} - 8\hat{k}$$

Therefore,

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = (2 \times 20) + (1 \times 10) + ((-3) \times (-8))$$

$$= 40 + 10 + 24$$

$$= 74$$

$$\therefore \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = 74 \text{ eq(1)}$$

And

$$\vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2) = (3 \times 20) + (3 \times 10) + (2 \times (-8))$$

$$= 60 + 30 - 16$$

$$= 74$$

$$\therefore \vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2) = 74 \text{ eq(2)}$$

From eq(1) and eq(2)

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_2 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Hence lines $\vec{r}_1$ & $\vec{r}_2$ are coplanar.

Equation of plane containing lines $\vec{r}_1$ & $\vec{r}_2$ is

$$\vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$$

Now,

$$\vec{b}_1 \times \vec{b}_2 = 20\hat{i} + 10\hat{j} - 8\hat{k}$$

From eq(1)

$$\vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2) = 74$$

Therefore, equation of required plane is

$$\vec{r} \cdot (20\hat{i} + 10\hat{j} - 8\hat{k}) = 74$$

$$\therefore \vec{r} \cdot (10\hat{i} + 5\hat{j} - 4\hat{k}) = 37$$

$$\therefore \vec{r} \cdot (10\hat{i} + 5\hat{j} - 4\hat{k}) - 37 = 0$$

This vector equation of plane.

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore \vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = (x \times 20) + (y \times 10) + (z \times (-8))$$

$$= 20x + 10y - 8z$$

Therefore, equation of plane is

$$20x + 10y - 8z = 74$$

$$20x + 10y - 8z - 74 = 0$$

$$10x + 5y - 4z - 37 = 0$$

This Cartesian equation of plane.

Question: 4

Prove that the li

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3}$$

$$\text{Line 2 : } \frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$$

To Prove : Line 1 & line 2 are coplanar.

To Find : Equation of plane.

Formulae :

1) Coplanarity of two lines :

If two lines are given by,

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}, \text{ then these lines are coplanar, if}$$

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

2) Equation of plane :

The equation of plane containing two coplanar lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$

& $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3}$$

$$\text{Line 2 : } \frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$$

Here, $x_1 = 0$, $y_1 = 2$, $z_1 = -3$, $a_1 = 1$, $b_1 = 2$, $c_1 = 3$

$x_2 = 2$, $y_2 = 6$, $z_2 = 3$, $a_2 = 2$, $b_2 = 3$, $c_2 = 4$

Now,

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = \begin{vmatrix} 2 - 0 & 6 - 2 & 3 + 3 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix}$$

$$= 2(8 - 9) - 4(4 - 6) + 6(3 - 4)$$

$$= 2(-1) - 4(-2) + 6(-1)$$

$$= -2 + 8 - 6$$

$$= 0$$

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Hence, given two lines are coplanar.

Equation of plane passing through line1 and line 2 is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x - 0 & y - 2 & z + 3 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix} = 0$$

$$\therefore (x - 0) \times (8 - 9) - (y - 2) \times (4 - 6) + (z + 3) \times (3 - 4) = 0$$

$$\therefore -1(x) - (y-2)(-2) + (z+3)(-1) = 0$$

$$-x + 2y - 4 - z - 3 = 0$$

$$-x + 2y - z - 7 = 0$$

$$x - 2y + z + 7 = 0$$

Therefore, equation of plane is

$$x - 2y + z + 7 = 0$$

Question: 5

Prove that the li

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x-2}{1} = \frac{y-4}{4} = \frac{z-6}{7}$$

$$\text{Line 2 : } \frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$$

To Prove : Line 1 & line 2 are coplanar.

To Find : Equation of plane.

Formulae :

1) Coplanarity of two lines :

If two lines are given by,

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}, \text{ then these lines are coplanar, if}$$

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

2) Equation of plane :

The equation of plane containing two coplanar lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$

& $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x-2}{1} = \frac{y-4}{4} = \frac{z-6}{7}$$

$$\text{Line 2 : } \frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$$

Here, $x_1 = 2$, $y_1 = 4$, $z_1 = 6$, $a_1 = 1$, $b_1 = 4$, $c_1 = 7$

$x_2 = -1$, $y_2 = -3$, $z_2 = -5$, $a_2 = 3$, $b_2 = 5$, $c_2 = 7$

Now,

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = \begin{vmatrix} -1 - 2 & -3 - 4 & -5 - 6 \\ 1 & 4 & 7 \\ 3 & 5 & 7 \end{vmatrix}$$

$$= \begin{vmatrix} -3 & -7 & -11 \\ 1 & 4 & 7 \\ 3 & 5 & 7 \end{vmatrix}$$

$$= -3(28 - 35) - (-7)(7 - 21) - 11(5 - 12)$$

$$= -3(-7) + 7(-14) - 11(-7)$$

$$= 21 - 98 + 77$$

$$= 0$$

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Hence, given two lines are coplanar.

Equation of plane passing through line 1 and line 2 is given by,

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x-2 & y-4 & z-6 \\ 1 & 4 & 7 \\ 3 & 5 & 7 \end{vmatrix} = 0$$

$$\therefore (x-2) \times (28-35) - (y-4) \times (7-21) + (z-6) \times (5-12) = 0$$

$$\therefore -7(x-2) - (y-4)(-14) + (z-6)(-7) = 0$$

$$-7x + 14 + 14y - 56 - 7z + 42 = 0$$

$$-7x + 14y - 7z = 0$$

$$x - 2y + z = 0$$

Therefore, equation of plane is

$$x - 2y + z = 0$$

Question: 6

Show that the line

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x-5}{-4} = \frac{y-7}{4} = \frac{z+3}{-5} \text{ or } \frac{x-5}{4} = \frac{y-7}{4} = \frac{z+3}{-5}$$

$$\text{Line 2 : } \frac{x-8}{7} = \frac{y-4}{2} = \frac{z-5}{3} \text{ or } \frac{x-8}{7} = \frac{y-4}{1} = \frac{z-5}{3}$$

To Prove : Line 1 & line 2 are coplanar.

To Find : Equation of plane.

Formulae :

1) Coplanarity of two lines :

If two lines are given by,

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}, \text{ then these lines are coplanar, if}$$

$$\begin{vmatrix} x_2-x_1 & y_2-y_1 & z_2-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

2) Equation of plane :

The equation of plane containing two coplanar lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$

& $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ is given by,

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x-5}{4} = \frac{y-7}{4} = \frac{z+3}{-5}$$

$$\text{Line 2 : } \frac{x-8}{7} = \frac{y-4}{1} = \frac{z-5}{3}$$

Here, $x_1 = 5$, $y_1 = 7$, $z_1 = -3$, $a_1 = 4$, $b_1 = 4$, $c_1 = -5$

$x_2 = 8$, $y_2 = 4$, $z_2 = 5$, $a_2 = 7$, $b_2 = 1$, $c_2 = 3$

Now,

$$\begin{vmatrix} x_2-x_1 & y_2-y_1 & z_2-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = \begin{vmatrix} 8-5 & 4-7 & 5+3 \\ 4 & 4 & -5 \\ 7 & 1 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 3 & -3 & 8 \\ 4 & 4 & -5 \\ 7 & 1 & 3 \end{vmatrix}$$

$$= 3(12+5) - (-3)(12+35) + 8(4-28)$$

$$= 3(17) + 3(47) + 8(-24)$$

$$= 51 + 141 - 192$$

$$= 0$$

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Hence, given two lines are coplanar.

Equation of plane passing through line1 and line 2 is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x - 5 & y - 7 & z + 3 \\ 4 & 4 & -5 \\ 7 & 1 & 3 \end{vmatrix} = 0$$

$$\therefore (x - 5) \times (12 + 5) - (y - 7) \times (12 + 35) + (z + 3) \times (4 - 28) = 0$$

$$\therefore 17(x - 5) - 47(y - 7) + (z + 3)(-24) = 0$$

$$17x - 85 - 47y + 329 - 24z - 72 = 0$$

$$17x - 47y - 24z + 172 = 0$$

Therefore, equation of plane is

$$17x - 47y - 24z + 172 = 0$$

Question: 7

Show that the lin

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$$

$$\text{Line 2 : } \frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2}$$

To Prove : Line 1 & line 2 are coplanar.

To Find : Equation of plane.

Formulae :

1) Coplanarity of two lines :

If two lines are given by,

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}, \text{ then these lines are coplanar, if}$$

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

2) Equation of plane :

The equation of plane containing two coplanar lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$

& $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x+1}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$$

$$\text{Line 2 : } \frac{x}{1} = \frac{y-7}{-3} = \frac{z+7}{2}$$

Here, $x_1 = -1$, $y_1 = 3$, $z_1 = -2$, $a_1 = -3$, $b_1 = 2$, $c_1 = 1$

$x_2 = 0$, $y_2 = 7$, $z_2 = -7$, $a_2 = 1$, $b_2 = -3$, $c_2 = 2$

Now,

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = \begin{vmatrix} 0 + 1 & 7 - 3 & -7 + 2 \\ -3 & 2 & 1 \\ 1 & -3 & 2 \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 4 & -5 \\ -3 & 2 & 1 \\ 1 & -3 & 2 \end{vmatrix}$$

$$= 1(4 + 3) - 4(-6 - 1) - 5(9 - 2)$$

$$= 1(7) - 4(-7) - 5(7)$$

$$= 7 + 28 - 35$$

$$= 0$$

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Hence, given two lines are coplanar.

Equation of plane passing through line1 and line 2 is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x + 1 & y - 3 & z + 2 \\ -3 & 2 & 1 \\ 1 & -3 & 2 \end{vmatrix} = 0$$

$$\therefore (x + 1) \times (4 + 3) - (y - 3) \times (-6 - 1) + (z + 2) \times (9 - 2) = 0$$

$$\therefore 7(x + 1) - (y - 3)(-7) + (z + 2)(7) = 0$$

$$7x + 7 + 7y - 21 + 7z + 14 = 0$$

$$7x + 7y + 7z = 0$$

$$x + y + z = 0$$

Therefore, equation of plane is

Question: 8

Show that the line

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x-1}{2} = \frac{y-3}{-1} = \frac{z}{-1}$$

$$\text{Line 2 : } \frac{x-4}{3} = \frac{y-1}{-2} = \frac{z-1}{-1}$$

To Prove : Line 1 & line 2 are coplanar.

To Find : Equation of plane.

Formulae :

1) Coplanarity of two lines :

If two lines are given by,

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}, \text{ then these lines are coplanar, if}$$

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

2) Equation of plane :

The equation of plane containing two coplanar lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$

& $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x-1}{2} = \frac{y-3}{-1} = \frac{z}{-1}$$

$$\text{Line 2 : } \frac{x-4}{3} = \frac{y-1}{-2} = \frac{z-1}{-1}$$

Here, $x_1 = 1$, $y_1 = 3$, $z_1 = 0$, $a_1 = 2$, $b_1 = -1$, $c_1 = -1$

$x_2 = 4$, $y_2 = 1$, $z_2 = 1$, $a_2 = 3$, $b_2 = -2$, $c_2 = -1$

Now,

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = \begin{vmatrix} 4 - 1 & 1 - 3 & 1 - 0 \\ 2 & -1 & -1 \\ 3 & -2 & -1 \end{vmatrix}$$

$$= \begin{vmatrix} 3 & -2 & 1 \\ 2 & -1 & -1 \\ 3 & -2 & -1 \end{vmatrix}$$

$$= 3(1-2) - (-2)(-2+3) + 1(-4+3)$$

$$= 3(-1) + 2(1) + 1(-1)$$

$$= -2$$

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} \neq 0$$

Hence, given two lines are not coplanar.

Question: 9

Find the equation

Solution:

Given : Equations of lines -

$$\text{Line 1 : } \frac{x-3}{1} = \frac{y+2}{-4} = \frac{z}{5}$$

$$\text{Line 2 : } \frac{x-4}{1} = \frac{y-3}{-4} = \frac{z-2}{5}$$

To Find : Equation of plane.

Formulae :

Equation of plane :

The equation of plane containing two parallel lines $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$

& $\frac{x-x_2}{a} = \frac{y-y_2}{b} = \frac{z-z_2}{c}$ is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a & b & c \end{vmatrix} = 0$$

Answer :

Given lines -

$$\text{Line 1 : } \frac{x-3}{1} = \frac{y+2}{-4} = \frac{z}{5}$$

$$\text{Line 2 : } \frac{x-4}{1} = \frac{y-3}{-4} = \frac{z-2}{5}$$

Here, $x_1 = 3$, $y_1 = -2$, $z_1 = 0$, $a = 1$, $b = -4$, $c = 5$

$x_2 = 4$, $y_2 = 3$, $z_2 = 2$

Therefore, equation of plane containing line 1 & line 2 is given by,

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a & b & c \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x - 3 & y + 2 & z - 0 \\ 4 - 3 & 3 + 2 & 2 - 0 \\ 1 & -4 & 5 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} x - 3 & y + 2 & z \\ 1 & 5 & 2 \\ 1 & -4 & 5 \end{vmatrix} = 0$$

$$\therefore (x-3) \times (25+8) - (y+2) \times (5-2) + (z) \times (-4-5) = 0$$

$$\therefore 33(x-3) - (y+2)(3) + (z)(-9) = 0$$

$$33x - 99 - 3y - 6 - 9z = 0$$

$$33x - 3y - 9z - 105 = 0$$

$$11x - y - 3z = 35$$

Therefore, equation of plane is

Exercise : 28J

Question: 1

Find the directio

Solution:

Given :

Equation of plane : $x + 2y - 3z = 5$

To Find : direction ratios of normal

Answer :

Given equation of plane : $x + 2y - 3z = 5$

It can be written as

$$(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} + 2\hat{j} - 3\hat{k}) = 5$$

Comparing with $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$

Therefore, normal vector is $\vec{n} = \hat{i} + 2\hat{j} - 3\hat{k}$

Hence, direction ratios of normal are (1, 2, -3).

Question: 2

Find the directio

Solution:

Given :

Equation of plane : $2x + 3y - z = 4$

To Find : Direction cosines of the normal i.e. $l, m \text{ \& } n$

Formula :

1) Direction cosines :

If $a, b \text{ \& } c$ are direction ratios of the vector then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

Answer :

For the given equation of plane

$$2x + 3y - z = 4$$

Direction ratios of normal vector are (2, 3, -1)

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{2^2 + 3^2 + (-1)^2}$$

$$= \sqrt{4 + 9 + 1}$$

$$= \sqrt{14}$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{2}{\sqrt{14}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{3}{\sqrt{14}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{-1}{\sqrt{14}}$$

$$(l, m, n) = \left(\frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}, \frac{-1}{\sqrt{14}} \right)$$

Question: 3

Find the directio

Solution:

Given :

Equation of plane : $y = 3$

To Find : Direction cosines of the normal i.e. $l, m \text{ \& } n$

Formula :

1) Direction cosines :

If $a, b \text{ \& } c$ are direction ratios of the vector then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

Answer :

For the given equation of plane

$$y = 3$$

Direction ratios of normal vector are $(0, 1, 0)$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{0^2 + 1^2 + 0^2}$$

$$= \sqrt{0 + 1 + 0}$$

$$= \sqrt{1}$$

$$= 1$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{1} = 0$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{1}{1} = 1$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{1} = 0$$

$$(l, m, n) = (0, 1, 0)$$

Question: 4

Find the directio

Solution:

Given :

$$\text{Equation of plane : } 3x + 4 = 0$$

To Find : Direction cosines of the normal i.e. $l, m \text{ \& } n$

Formula :

1) Direction cosines :

If a, b & c are direction ratios of the vector then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

Answer :

For the given equation of plane

$$-3x = 4$$

Direction ratios of normal vector are $(-3, 0, 0)$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{(-3)^2 + 0^2 + 0^2}$$

$$= \sqrt{9 + 0 + 0}$$

$$= \sqrt{9}$$

$$= 3$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{-3}{3} = -1$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{3} = 0$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{0}{3} = 0$$

$$(l, m, n) = (-1, 0, 0)$$

Question: 5

Write the equatio

Solution:

Given :

Point : (4, -2, 3)

To Find : equation of plane

Formula :

1) Equation of plane :

Equation of plane passing through point A with position vector $\vec{a}$ and perpendicular to vector $\vec{n}$ is given by,

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

Answer :

Position vector for given point A $\equiv$ (4, -2, 3) is

$$\vec{a} = 4\hat{i} - 2\hat{j} + 3\hat{k}$$

As required plane is parallel to XY plane, therefore Z-axis is perpendicular to the plane.

$$\therefore \vec{n} = \hat{k}$$

Therefore, equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{k}) = (4\hat{i} - 2\hat{j} + 3\hat{k}) \cdot (\hat{k})$$

$$\therefore (x \times 0) + (y \times 0) + (z \times 1) = (4 \times 0) + (-2 \times 0) + (3 \times 1)$$

$$\therefore z = 3$$

This is required equation of plane.

Question: 6

Write the equatio

Solution:

Given :

Point : (-3, 2, 0)

To Find : equation of plane

Formula :

1) Equation of plane :

Equation of plane passing through point A with position vector $\vec{a}$ and perpendicular to vector $\vec{n}$ is given by,

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

Answer :

Position vector for given point A $\equiv$ (-3, 2, 0) is

$$\vec{a} = -3\hat{i} + 2\hat{j} + 0\hat{k}$$

As required plane is parallel to YZ plane, therefore X-axis is perpendicular to the plane.

$$\therefore \vec{n} = \hat{i}$$

Therefore, equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i}) = (-3\hat{i} + 2\hat{j} + 0\hat{k}) \cdot (\hat{i})$$

$$\therefore (x \times 1) + (y \times 0) + (z \times 0) = (-3 \times 1) + (2 \times 0) + (0 \times 0)$$

$$\therefore x = -3$$

This is required equation of plane.

Question: 7

Write the general

Solution:

Let, normal vector of plane be

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

Equation of plane is given by,

$$\vec{r} \cdot \vec{n} = d$$

$$\therefore (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (a\hat{i} + b\hat{j} + c\hat{k}) = d$$

$$\therefore ax + by + cz = d$$

As the required plane is parallel to the given plane, hence normal vector of plane is perpendicular to x-axis.

$$\therefore \vec{n} \cdot \hat{i} = 0$$

$$\therefore (a\hat{i} + b\hat{j} + c\hat{k}) \cdot \hat{i} = 0$$

$$a = 0$$

Therefore, equation of plane is

$$by + cz = d$$

Question: 8

Write the interce

Solution:

Given :

$$\text{Equation of plane : } 2x + y - z = 5$$

To Find : Intercept made by the plane with the X-axis.

Formula :

$$\text{If } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

is the equation of plane in intercept form then intercept made by it with co-ordinate axes are

$$\text{X-intercept} = a$$

$$\text{Y-intercept} = b$$

$$\text{Z-intercept} = c$$

Answer :

Given equation of plane:

$$2x + y - z = 5$$

Dividing above equation throughout by 5

$$\therefore \frac{2x}{5} + \frac{y}{5} + \frac{-z}{5} = 1$$

$$\therefore \frac{x}{5/2} + \frac{y}{5} + \frac{z}{-5} = 1$$

Comparing above equation with

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

We get,

$$a = 5/2$$

Therefore, intercepts made by plane with X-axis are

$$\text{X-intercept} = 5/2$$

Question: 9

Write the interce

Solution:

Given :

$$\text{Equation of plane : } 4x - 3y + 2z = 12$$

To Find :

1) Equation of plane in intercept form

2) Intercepts made by the plane with the co-ordinate axes.

Formula :

$$\text{If } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

is the equation of plane in intercept form then intercept made by it with co-ordinate axes are

$$\text{X-intercept} = a$$

Y-intercept = b

Z-intercept = c

Answer :

Given equation of plane:

$$4x - 3y + 2z = 12$$

Dividing above equation throughout by 12

$$\therefore \frac{4x}{12} + \frac{-3y}{12} + \frac{2z}{12} = 1$$

$$\therefore \frac{x}{3} + \frac{y}{-4} + \frac{z}{6} = 1$$

This is the equation of plane in intercept form.

Comparing above equation with

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

We get,

$$a = 3$$

$$b = -4$$

$$c = 6$$

Therefore, intercepts made by plane with co-ordinate axes are

X-intercept = 3

Y-intercept = -4

Z-intercept = 6

Question: 10

Reduce the equati

Solution:

Given :

$$\text{Equation of plane : } 2x - 3y + 5z + 4 = 0$$

To Find :

1) Equation of plane in intercept form

2) Intercepts made by the plane with the co-ordinate axes.

Formula :

$$\text{If } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

is the equation of plane in intercept form then intercept made by it with co-ordinate axes are

X-intercept = a

Y-intercept = b

Z-intercept = c

Answer :

Given equation of plane:

$$2x - 3y + 5z = -4$$

Dividing above equation throughout by -4

$$\therefore \frac{2x}{-4} + \frac{-3y}{-4} + \frac{5z}{-4} = 1$$

$$\therefore \frac{x}{-2} + \frac{y}{4/3} + \frac{z}{-4/5} = 1$$

This is the equation of plane in intercept form.

Comparing above equation with

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

We get,

$$a = -2$$

$$b = \frac{4}{3}$$

$$c = \frac{-4}{5}$$

Therefore, intercepts made by plane with co-ordinate axes are

$$X\text{-intercept} = -2$$

$$Y\text{-intercept} = \frac{4}{3}$$

$$Z\text{-intercept} = -\frac{4}{5}$$

Question: 11

Find the equation

Solution:

Given : Plane is passing through points

$$A \equiv (a, 0, 0)$$

$$B \equiv (0, b, 0)$$

$$C \equiv (0, 0, c)$$

To Find : Equation of plane

Formulae :

Equation of plane making intercepts (a, b, c) on X, Y & Z axes respectively is given by,

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Answer : As plane is passing through points $A \equiv (a, 0, 0)$,

$$B \equiv (0, b, 0) \text{ \& } C \equiv (0, 0, c)$$

Therefore, intercepts made by it on X, Y & Z axes respectively are

a, b & c .

hence, equation of plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Question: 12

Write the value o

Solution:

Given : equations of perpendicular planes-

$$2x - 5y + kz = 4$$

$$x + 2y - z = 6$$

To Find : k

Formulae :

Normal vector to the plane :

If equation of the plane is $ax + by + cz = d$ then,

Vector normal to the plane is given by,

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

Answer :

For given planes -

$$2x - 5y + kz = 4$$

$$x + 2y - z = 6$$

normal vectors are

$$\vec{n}_1 = 2\hat{i} - 5\hat{j} + k\hat{k}$$

$$\vec{n}_2 = \hat{i} + 2\hat{j} - \hat{k}$$

As given vectors are perpendicular, hence their normal vectors are also perpendicular to each other.

$$\therefore \vec{n}_1 \cdot \vec{n}_2 = 0$$

$$\therefore (2\hat{i} - 5\hat{j} + k\hat{k}) \cdot (\hat{i} + 2\hat{j} - \hat{k}) = 0$$

$$(2 \times 1) + (-5 \times 2) + (k \times (-1)) = 0$$

$$2 - 10 - k = 0$$

$$-8 - k = 0$$

$$k = -8$$

Question: 13

Find the angle between

Solution:

Given : equations of planes-

$$2x + y - 2z = 5$$

$$3x - 6y - 2z = 7$$

To Find : angle between two planes

Formulae :

1) Normal vector to the plane :

If equation of the plane is $ax + by + cz = d$ then,

Vector normal to the plane is given by,

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

2) Angle between two planes :

The angle θ between the planes $\vec{r} \cdot \vec{n}_1 = p_1$ and $\vec{r} \cdot \vec{n}_2 = p_2$ is given by

$$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| \cdot |\vec{n}_2|}$$

Answer :

For given planes

$$2x + y - 2z = 5$$

$$3x - 6y - 2z = 7$$

Normal vectors are

$$\vec{n}_1 = 2\hat{i} + \hat{j} - 2\hat{k} \text{ and}$$

$$\vec{n}_2 = 3\hat{i} - 6\hat{j} - 2\hat{k}$$

$$\therefore |\vec{n}_1| = \sqrt{2^2 + 1^2 + (-2)^2} = \sqrt{4 + 1 + 4} = \sqrt{9} = 3$$

$$\therefore |\vec{n}_2| = \sqrt{3^2 + (-6)^2 + (-2)^2} = \sqrt{9 + 36 + 4} = \sqrt{49} = 7$$

Therefore, angle between two planes is

$$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| \cdot |\vec{n}_2|}$$

$$\therefore \cos \theta = \frac{(2\hat{i} + \hat{j} - 2\hat{k}) \cdot (3\hat{i} - 6\hat{j} - 2\hat{k})}{3 \times 7}$$

$$\therefore \cos \theta = \frac{(2 \times 3) + (1 \times (-6)) + ((-2) \times (-2))}{21}$$

$$\therefore \cos \theta = \frac{6 - 6 + 4}{21}$$

$$\therefore \cos \theta = \frac{4}{21}$$

$$\therefore \theta = \cos^{-1}\left(\frac{4}{21}\right)$$

Question: 14

Find the angle between

Solution:

Given : equations of planes-

$$\vec{r} \cdot (\hat{i} + \hat{j}) = 1$$

$$\vec{r} \cdot (\hat{j} + \hat{k}) = 3$$

To Find : angle between two planes

Formulae :

Angle between two planes :

The angle θ between the planes $\vec{r} \cdot \vec{n}_1 = p_1$ and $\vec{r} \cdot \vec{n}_2 = p_2$ is given by

$$\cos \theta = \frac{\overline{n_1} \cdot \overline{n_2}}{|\overline{n_1}| \cdot |\overline{n_2}|}$$

Answer :

For given planes

$$\vec{r} \cdot (\hat{i} + \hat{j}) = 1$$

$$\vec{r} \cdot (\hat{j} + \hat{k}) = 3$$

Normal vectors are

$$\overline{n_1} = \hat{i} + \hat{j} \text{ and}$$

$$\overline{n_2} = \hat{j} + \hat{k}$$

$$\therefore |\overline{n_1}| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{1 + 1 + 0} = \sqrt{2}$$

$$\therefore |\overline{n_2}| = \sqrt{0^2 + 1^2 + 1^2} = \sqrt{0 + 1 + 1} = \sqrt{2}$$

Therefore, angle between two planes is

$$\cos \theta = \frac{\overline{n_1} \cdot \overline{n_2}}{|\overline{n_1}| \cdot |\overline{n_2}|}$$

$$\therefore \cos \theta = \frac{(\hat{i} + \hat{j}) \cdot (\hat{j} + \hat{k})}{\sqrt{2} \times \sqrt{2}}$$

$$\therefore \cos \theta = \frac{(1 \times 0) + (1 \times 1) + (0 \times 1)}{2}$$

$$\therefore \cos \theta = \frac{0 + 1 + 0}{2}$$

$$\therefore \cos \theta = \frac{1}{2}$$

$$\therefore \theta = \cos^{-1}\left(\frac{1}{2}\right)$$

$$\therefore \theta = \frac{\pi}{3}$$

Question: 15

Find the angle be

Solution:

Given : equations of planes-

$$\vec{r} \cdot (3\hat{i} - 4\hat{j} + 5\hat{k}) = 0$$

$$\vec{r} \cdot (2\hat{i} - \hat{j} - 2\hat{k}) = 7$$

To Find : angle between two planes

Formulae :

Angle between two planes :

The angle θ between the planes $\vec{r} \cdot \overline{n_1} = p_1$ and $\vec{r} \cdot \overline{n_2} = p_2$ is given by

$$\cos \theta = \frac{\overline{n_1} \cdot \overline{n_2}}{|\overline{n_1}| \cdot |\overline{n_2}|}$$

Answer :

For given planes

$$\vec{r} \cdot (3\hat{i} - 4\hat{j} + 5\hat{k}) = 0$$

$$\vec{r} \cdot (2\hat{i} - \hat{j} - 2\hat{k}) = 7$$

Normal vectors are

$$\overline{n_1} = 3\hat{i} - 4\hat{j} + 5\hat{k} \text{ and}$$

$$\overline{n_2} = 2\hat{i} - \hat{j} - 2\hat{k}$$

$$\therefore |\overline{n_1}| = \sqrt{3^2 + (-4)^2 + 5^2} = \sqrt{9 + 16 + 25} = \sqrt{50} = 5\sqrt{2}$$

$$\therefore |\overline{n_2}| = \sqrt{2^2 + (-1)^2 + (-2)^2} = \sqrt{4 + 1 + 4} = \sqrt{9} = 3$$

Therefore, angle between two planes is

$$\cos \theta = \frac{\overline{n_1} \cdot \overline{n_2}}{|\overline{n_1}| \cdot |\overline{n_2}|}$$

$$\therefore \cos \theta = \frac{(3\hat{i} - 4\hat{j} + 5\hat{k}) \cdot (2\hat{i} - \hat{j} - 2\hat{k})}{5\sqrt{2} \times 3}$$

$$\therefore \cos \theta = \frac{(3 \times 2) + ((-4) \times (-1)) + (5 \times (-2))}{15\sqrt{2}}$$

$$\therefore \cos \theta = \frac{6 + 4 - 10}{2}$$

$$\therefore \cos \theta = 0$$

$$\therefore \theta = \cos^{-1}(0)$$

$$\therefore \theta = \frac{\pi}{2}$$

Question: 16

Find the angle between

Solution:

Given :

$$\text{Equation of line : } \frac{x+1}{2} = \frac{y}{3} = \frac{z-3}{6}$$

$$\text{Equation of plane : } 10x + 2y - 11z = 3$$

To Find : angle between line and plane

Formulae :

1) Parallel vector to the line :

$$\text{If equation of the line is } \frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ then,}$$

Vector parallel to the line is given by,

$$\vec{b} = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$$

2) Normal vector to the plane :

If equation of the plane is $ax + by + cz = d$ then,

Vector normal to the plane is given by,

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

3) Angle between a line and a plane :

If θ is an angle between the line $\vec{r} = \vec{a} + \lambda\vec{b}$ and the plane $\vec{r} \cdot \vec{n} = p$, then

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

Where, $\vec{b}$ is vector parallel to the line and

$\vec{n}$ is the vector normal to the plane.

Answer :

For given equation of line,

$$\frac{x+1}{2} = \frac{y}{3} = \frac{z-3}{6}$$

Parallel vector to the line is

$$\vec{b} = 2\hat{i} + 3\hat{j} + 6\hat{k}$$

$$\therefore |\vec{b}| = \sqrt{2^2 + 3^2 + 6^2} = \sqrt{4 + 9 + 36} = \sqrt{49} = 7$$

For given equation of plane,

$$10x + 2y - 11z = 3$$

normal vector to the plane is

$$\vec{n} = 10\hat{i} + 2\hat{j} - 11\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{10^2 + 2^2 + (-11)^2} = \sqrt{100 + 4 + 121} = \sqrt{225} = 15$$

Therefore, angle between given line and plane is

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

$$\therefore \sin \theta = \frac{(2\hat{i} + 3\hat{j} + 6\hat{k}) \cdot (10\hat{i} + 2\hat{j} - 11\hat{k})}{7 \times 15}$$

$$\therefore \sin \theta = \frac{(2 \times 10) + (3 \times 2) + (6 \times (-11))}{105}$$

$$\therefore \sin \theta = \frac{20 + 6 - 66}{105}$$

$$\therefore \sin \theta = \frac{-40}{105}$$

$$\therefore \sin \theta = \frac{-8}{21}$$

$$\therefore \theta = \sin^{-1}\left(\frac{-8}{21}\right)$$

Question: 17

Find the angle be

Solution:

Given :

$$\text{Equation of line : } \vec{r} = (\hat{i} + \hat{j} - 2\hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$$

$$\text{Equation of plane : } \vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 4$$

To Find : angle between line and plane

Formulae :

1) Angle between a line and a plane :

If θ is a angle between the line $\vec{r} = \vec{a} + \lambda\vec{b}$ and the plane $\vec{r} \cdot \vec{n} = p$, then

$$\sin \theta = \frac{\vec{b} \cdot \vec{n}}{|\vec{b}| \cdot |\vec{n}|}$$

Where, $\vec{b}$ is vector parallel to the line and

$\vec{n}$ is the vector normal to the plane.

Answer :

For given equation of line,

$$\vec{r} = (\hat{i} + \hat{j} - 2\hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$$

Parallel vector to the line is

$$\vec{b} = \hat{i} - \hat{j} + \hat{k}$$

$$\therefore |\vec{b}| = \sqrt{1^2 + (-1)^2 + 1^2} = \sqrt{1 + 1 + 1} = \sqrt{3}$$

For given equation of plane,

$$\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 4$$

normal vector to the plane is

$$\vec{n} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\therefore |\vec{n}| = \sqrt{2^2 + (-1)^2 + 1^2} = \sqrt{4 + 1 + 1} = \sqrt{6}$$

Therefore, angle between given line and plane is

$$\sin \theta = \frac{\vec{b} \cdot \vec{n}}{|\vec{b}| \cdot |\vec{n}|}$$

$$\therefore \sin \theta = \frac{(\hat{i} - \hat{j} + \hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k})}{\sqrt{3} \times \sqrt{6}}$$

$$\therefore \sin \theta = \frac{(1 \times 2) + ((-1) \times (-1)) + (1 \times 1)}{\sqrt{18}}$$

$$\therefore \sin \theta = \frac{2 + 1 + 1}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{4}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{2 \times 2}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{2\sqrt{2}}{3}$$

$$\therefore \theta = \sin^{-1}\left(\frac{2\sqrt{2}}{3}\right)$$

Question: 18

Find the value of

Solution:

Given :

$$\text{Equation of line : } \frac{x-2}{6} = \frac{y-1}{\lambda} = \frac{z+5}{4}$$

$$\text{Equation of plane : } 3x - y - 2z = 7$$

To Find : λ

Formulae :

1) Parallel vector to the line :

If equation of the line is $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ then,

Vector parallel to the line is given by,

$$\vec{b} = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$$

2) Normal vector to the plane :

If equation of the plane is $ax + by + cz = d$ then,

Vector normal to the plane is given by,

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

3) Cross Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Answer :

For given equation of line,

$$\frac{x-2}{6} = \frac{y-1}{\lambda} = \frac{z+5}{4}$$

Parallel vector to the line is

$$\vec{b} = 6\hat{i} + \lambda\hat{j} + 4\hat{k}$$

For given equation of plane,

$$3x - y - 2z = 7$$

normal vector to the plane is

$$\vec{n} = 3\hat{i} - \hat{j} - 2\hat{k}$$

As given line and plane are perpendicular to each other.

$$\therefore \vec{b} \times \vec{n} = 0$$

$$\therefore \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 6 & \lambda & 4 \\ 3 & -1 & -2 \end{vmatrix} = 0$$

$$\therefore \hat{i}(-2\lambda + 4) - \hat{j}(-12 - 12) + \hat{k}(-6 - 3\lambda) = 0\hat{i} + 0\hat{j} + 0\hat{k}$$

Comparing coefficients of $\hat{k}$ on both sides

$$\therefore -6 - 3\lambda = 0$$

$$3\lambda = -6$$

$$\lambda = -2$$

Question: 19

Write the equatio

Solution:

Given :

$$A \equiv (a, b, c)$$

Equation of plane parallel to required plane

$$\therefore \vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$$

To Find : Equation of plane

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

If a plane is passing through point A, then equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{a}$ = position vector of A

$\vec{n}$ = vector perpendicular to the plane

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Answer :

For point A $\equiv (a, b, c)$, position vector is

$$\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$$

As plane $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$ is parallel to the required plane, the vector normal to required plane is

$$\vec{n} = \hat{i} + \hat{j} + \hat{k}$$

$$\text{Now, } \vec{a} \cdot \vec{n} = (a \times 1) + (b \times 1) + (c \times 1)$$

$$= a + b + c$$

Equation of the plane passing through point A and perpendicular to vector $\vec{n}$ is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore \vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = a + b + c$$

Question: 20

Find the length o

Solution:

Given :

$$\text{Equation of plane : } 2x - 3y + 6z + 21 = 0$$

To Find :

Length of perpendicular drawn from origin to the plane = d

Formulae :

1) Distance of the plane from the origin :

Distance of the plane from the origin is given by,

$$d = \frac{p}{|\vec{n}|}$$

Answer :

For the given equation of plane

$$2x - 3y + 6z = -21$$

Direction ratios of normal vector are (2, -3, 6)

Therefore, equation of normal vector is

$$\vec{n} = 2\hat{i} - 3\hat{j} + 6\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{2^2 + (-3)^2 + 6^2}$$

$$= \sqrt{4 + 9 + 36}$$

$$= \sqrt{49}$$

$$= 7$$

From given equation of plane,

$$p = -21$$

Now, distance of the plane from the origin is

$$d = \frac{p}{|\vec{n}|}$$

$$\therefore d = \frac{-21}{7}$$

$$d = 3 \text{ units}$$

Question: 21

Find the direction

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (6\hat{i} - 3\hat{j} - 2\hat{k}) + 1 = 0$$

To Find :

Direction cosines of the normal i.e. $l, m \text{ \& } n$

Formulae :

1) Direction cosines :

If $a, b \text{ \& } c$ are direction ratios of the vector then its direction cosines are given by

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

Answer :

For the given equation of plane

$$\vec{r} \cdot (6\hat{i} - 3\hat{j} - 2\hat{k}) + 1 = 0$$

Equation of normal vector is

$$\vec{n} = 6\hat{i} - 3\hat{j} - 2\hat{k}$$

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{6^2 + (-3)^2 + (-2)^2}$$

$$= \sqrt{36 + 9 + 4}$$

$$= \sqrt{49}$$

$$= 7$$

Therefore, direction cosines are

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}} = \frac{6}{7}$$

$$m = \frac{b}{\sqrt{a^2 + b^2 + c^2}} = \frac{-3}{7}$$

$$n = \frac{c}{\sqrt{a^2 + b^2 + c^2}} = \frac{-2}{7}$$

$$(l, m, n) = \left(\frac{6}{7}, \frac{-3}{7}, \frac{-2}{7}\right)$$

Question: 22

Show that the line

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (5\hat{i} + 4\hat{j} - 4\hat{k}) = 7$$

Equation of line :

$$\vec{r} = (4\hat{i} - 7\hat{k}) + \lambda(4\hat{i} - 2\hat{j} + 3\hat{k})$$

To Prove : Given line is parallel to the given plane.

Answer :

Comparing given plane i.e.

$$\vec{r} \cdot (5\hat{i} + 4\hat{j} - 4\hat{k}) = 7$$

with $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$, we get,

$$\vec{n} = 5\hat{i} + 4\hat{j} - 4\hat{k}$$

This is the vector perpendicular to the given plane.

Now, comparing given equation of line i.e.

$$\vec{r} = (4\hat{i} - 7\hat{k}) + \lambda(4\hat{i} - 2\hat{j} + 3\hat{k})$$

with $\vec{r} = \vec{a} + \lambda\vec{b}$, we get,

$$\vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$$

Now,

$$\vec{n} \cdot \vec{b} = (5\hat{i} + 4\hat{j} - 4\hat{k}) \cdot (4\hat{i} - 2\hat{j} + 3\hat{k})$$

$$= (5 \times 4) + (4 \times (-2)) + ((-4) \times 3)$$

$$= 20 - 8 - 12$$

$$= 0$$

$$\therefore \vec{n} \cdot \vec{b} = 0$$

Therefore, vector normal to the plane is perpendicular to the vector parallel to the line.

Hence, the given line is parallel to the given plane.

Question: 23

Find the length o

Solution:

Given :

$$\text{Equation of plane : } \vec{r} \cdot (2\hat{i} - 3\hat{j} + 6\hat{k}) + 14 = 0$$

To Find : Length of perpendicular = d

Formulae :

1) Unit Vector :

Let $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ be any vector

Then unit vector of $\vec{a}$ is

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Where, } |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

2) Length of perpendicular :

The length of the perpendicular from the origin to the plane

$\vec{r} \cdot \vec{n} = p$ is given by,

$$d = \frac{p}{|\vec{n}|}$$

Answer :

Given equation of the plane is

$$\vec{r} \cdot (2\hat{i} - 3\hat{j} + 6\hat{k}) + 14 = 0$$

$$\therefore \vec{r} \cdot (2\hat{i} - 3\hat{j} + 6\hat{k}) = -14$$

$$\therefore \vec{r} \cdot (-2\hat{i} + 3\hat{j} - 6\hat{k}) = 14$$

Comparing above equation with

$$\vec{r} \cdot \vec{n} = p$$

We get,

$$\vec{n} = -2\hat{i} + 3\hat{j} - 6\hat{k} \text{ \& } p = 14$$

Therefore,

$$|\vec{n}| = \sqrt{(-2)^2 + 3^2 + (-6)^2}$$

$$= \sqrt{4 + 9 + 36}$$

$$= \sqrt{49}$$

$$= 7$$

The length of the perpendicular from the origin to the given plane is

$$d = \frac{p}{|\vec{n}|}$$

$$\therefore d = \frac{14}{7}$$

$$\therefore d = 2 \text{ units}$$

Question: 24

Find the value of

Solution:

Given :

$$\text{Equation of line : } \frac{x-1}{2} = \frac{y-1}{3} = \frac{z-1}{\lambda}$$

$$\text{Equation of plane : } \vec{r} \cdot (2\hat{i} + 3\hat{j} + 4\hat{k}) = 4$$

To Find : λ

Formulae :

1) Parallel vector to the line :

$$\text{If equation of the line is } \frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ then,}$$

Vector parallel to the line is given by,

$$\vec{b} = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$$

2) Angle between a line and a plane :

If θ is a angle between the line $\vec{r} = \vec{a} + \lambda\vec{b}$ and the plane $\vec{r} \cdot \vec{n} = p$, then

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

Where, $\vec{b}$ is vector parallel to the line and

$\vec{n}$ is the vector normal to the plane.

Answer :

For given equation of line,

$$\frac{x-1}{2} = \frac{y-1}{3} = \frac{z-1}{\lambda}$$

Parallel vector to the line is

$$\vec{b} = 2\hat{i} + 3\hat{j} + \lambda\hat{k}$$

For given equation of plane,

$$\vec{r} \cdot (2\hat{i} + 3\hat{j} + 4\hat{k}) = 4$$

normal vector to the plane is

$$\vec{n} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

Therefore, angle between given line and plane is

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

As given line is parallel too the given plane, angle between them is 0.

$$\therefore \theta = 0$$

$$\therefore \sin \theta = 0$$

$$\therefore \vec{b} \cdot \vec{n} = 0$$

$$\therefore (2\hat{i} + 3\hat{j} + \lambda\hat{k}) \cdot (2\hat{i} + 3\hat{j} + 4\hat{k}) = 0$$

$$\therefore (2 \times 2) + (3 \times 3) + (\lambda \times 4) = 0$$

$$4 + 9 + 4\lambda = 0$$

$$13 + 4\lambda = 0$$

$$4\lambda = -13$$

$$\therefore \lambda = -\frac{13}{4}$$

$$\lambda = -\frac{13}{4}$$

Question: 25

Write the angle θ

Solution:

Given :

$$\text{Equation of line : } \frac{x-1}{2} = \frac{y-2}{1} = \frac{z+3}{-2}$$

$$\text{Equation of plane : } x + y + 4 = 0$$

To Find : angle between line and plane

Formulae :

1) Parallel vector to the line :

If equation of the line is $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ then,

Vector parallel to the line is given by,

$$\vec{b} = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$$

2) Normal vector to the plane :

If equation of the plane is $ax + by + cz = d$ then,

Vector normal to the plane is given by,

$$\vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

3) Angle between a line and a plane :

If θ is a angle between the line $\vec{r} = \vec{a} + \lambda\vec{b}$ and the plane $\vec{r} \cdot \vec{n} = p$, then

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

Where, $\vec{b}$ is vector parallel to the line and

$\vec{n}$ is the vector normal to the plane.

Answer :

For given equation of line,

$$\frac{x-1}{2} = \frac{y-2}{1} = \frac{z+3}{-2}$$

Parallel vector to the line is

$$\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$$

$$\therefore |\vec{b}| = \sqrt{2^2 + 1^2 + (-2)^2} = \sqrt{4 + 1 + 4} = \sqrt{9} = 3$$

For given equation of plane,

$$x + y + 4 = 0$$

normal vector to the plane is

$$\vec{n} = \hat{i} + \hat{j} + 0\hat{k}$$

$$\therefore |\vec{n}| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{1 + 1 + 0} = \sqrt{2}$$

Therefore, angle between given line and plane is

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

$$\therefore \sin \theta = \frac{(2\hat{i} + \hat{j} - 2\hat{k}) \cdot (\hat{i} + \hat{j} + 0\hat{k})}{3 \times \sqrt{2}}$$

$$\therefore \sin \theta = \frac{(2 \times 1) + (1 \times 1) + ((-2) \times 0)}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{2 + 1 - 0}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{3}{3\sqrt{2}}$$

$$\therefore \sin \theta = \frac{1}{\sqrt{2}}$$

$$\therefore \theta = \sin^{-1}\left(\frac{1}{\sqrt{2}}\right)$$

$$\theta = \frac{\pi}{4}$$

Question: 26

Write the equatio

Solution:

Given :

$$A \equiv (2, -1, 1)$$

Plane parallel to the required plane : $3x + 2y - z = 7$

To Find : Equation of plane

Formulae :

1) Position vectors :

If A is a point having co-ordinates (a_1, a_2, a_3) , then its position vector is given by,

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

2) Dot Product :

If $\vec{a}$ & $\vec{b}$ are two vectors

$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

then,

$$\vec{a} \cdot \vec{b} = (a_1 \times b_1) + (a_2 \times b_2) + (a_3 \times b_3)$$

3) Equation of plane :

If a plane is passing through point A, then equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

Where, $\vec{a}$ = position vector of A

$\vec{n}$ = vector perpendicular to the plane

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

Answer :

For point A $\equiv (2, -1, 1)$, position vector is

$$\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$$

As required plane is parallel to $3x + 2y - z = 7$.

Therefore, normal vector of given plane is also perpendicular to required plane

$$\vec{n} = 3\hat{i} + 2\hat{j} - \hat{k}$$

$$\text{Now, } \vec{a} \cdot \vec{n} = (2 \times 3) + ((-1) \times 2) + (1 \times (-1))$$

$$= 6 - 2 - 1$$

$$= 3$$

Equation of the plane passing through point A and perpendicular to vector $\vec{n}$ is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\therefore \vec{r} \cdot (3\hat{i} + 2\hat{j} - \hat{k}) = 3$$

$$\text{As } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\therefore \vec{r} \cdot (3\hat{i} + 2\hat{j} - \hat{k}) = (x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + 2\hat{j} - \hat{k})$$

$$= 3x + 2y - z$$

Therefore, equation of the plane is

$$3x + 2y - z = 3$$

$$3x + 2y - z - 3 = 0$$

Exercise : OBJECTIVE QUESTIONS

Question: 1

Mark against the

Solution:

Given: Equation of plane is $\vec{r} \cdot (6\hat{i} - 3\hat{j} + 2\hat{k}) + 1 = 0$

Formula Used: Equation of a plane is $\hat{u} \cdot \vec{r} = p$ where $\hat{u}$ is the unit vector normal to the plane, $\vec{r}$ represents a point on the plane and p is the distance of the plane from the origin.

Explanation:

The equation of the given plane is $\vec{r} \cdot (6\hat{i} - 3\hat{j} + 2\hat{k}) = -1 \dots (1)$

Now, $|6\hat{i} - 3\hat{j} + 2\hat{k}| = \sqrt{36 + 9 + 4}$

$= 7$

$\therefore \frac{6}{7}\hat{i} - \frac{3}{7}\hat{j} + \frac{2}{7}\hat{k}$ is a unit vector.

(1) can be rewritten as

$$\vec{r} \cdot \left(\frac{6}{7}\hat{i} - \frac{3}{7}\hat{j} + \frac{2}{7}\hat{k} \right) = -\frac{1}{7}$$

$$\Rightarrow \vec{r} \cdot \left(-\frac{6}{7}\hat{i} + \frac{3}{7}\hat{j} - \frac{2}{7}\hat{k} \right) = \frac{1}{7}$$

which is of the form $\hat{u} \cdot \vec{r} = p$

Perpendicular vector from the origin to the plane is

$$\hat{u} = -\frac{6}{7}\hat{i} + \frac{3}{7}\hat{j} - \frac{2}{7}\hat{k}$$

So, direction cosines of the vector perpendicular from the origin to the plane is $\left(-\frac{6}{7}, \frac{3}{7}, -\frac{2}{7} \right)$

Question: 2

Mark against the

Solution:

Given: Equation of plane is $5y + 4 = 0$

Formula Used: Equation of a plane is $lx + my + nz = p$ where (l, m, n) are the direction cosines of the normal to the plane and (x, y, z) is a point on the plane and p is the distance of plane from origin.

Explanation:

Given equation is $5y = -4$

Dividing by -5,

$$-y = \frac{4}{5}$$

which is of the form $lx + my + nz = p$ where $l = 0, m = -1, n = 0$

Therefore, direction cosines of the normal to the plane is $(0, -1, 0)$

Question: 3

Mark against the

Solution:

Given: Equation of plane is $\vec{r} \cdot (3\hat{i} - 4\hat{j} - 12\hat{k}) + 39 = 0$

Formula Used: Equation of a plane is $\hat{u} \cdot \vec{r} = p$ where $\hat{u}$ is the unit vector normal to the plane, $\vec{r}$ represents a point on the plane and p is the distance of the plane from the origin.

Explanation:

Given equation is $\vec{r} \cdot (3\hat{i} - 4\hat{j} - 12\hat{k}) = -39 \dots (1)$

Now, $|3\hat{i} - 4\hat{j} - 12\hat{k}| = \sqrt{9 + 16 + 144} = \sqrt{169}$

$= 13$

Dividing (1) by 13 and multiplying by -1,

$$\vec{r} \cdot \left(-\frac{3}{13}\hat{i} + \frac{4}{13}\hat{j} + \frac{12}{13}\hat{k} \right) = 3$$

which is of the form $\hat{u} \cdot \vec{r} = p$

Therefore, length of perpendicular from origin to plane is 3 units.

Question: 4

Mark against the

Solution:

Given: A(2, -3, 7) is a point on the plane making equal intercepts on the axes.

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x-axis, y-axis and z-axis respectively.

Explanation:

Let the equation of the plane be

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1 \dots (1)$$

Here $a = b = c = p$ (let's say)

Since (2, -3, 7) is a point on the plane,

(1) becomes

$$\frac{2 - 3 + 7}{p} = 1$$

$$p = 6$$

Therefore equation of the plane is

$$x + y + z = 6$$

Question: 5

Mark against the

Solution:

Given: Plane makes intercepts 3, -4 and 6 with the coordinate axes.

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x-axis, y-axis and z-axis respectively.

Normal Form of a plane $= lx + my + nz = p$ where (l, m, n) is the direction cosines and p is the distance of perpendicular to the plane from the origin.

Explanation:

Equation of the given plane is

$$\frac{x}{3} + \frac{y}{-4} + \frac{z}{6} = 1$$

$$\text{i.e., } 4x - 3y + 2z = 12 \dots (1)$$

which is of the form $ax + by + cz = d$

Direction ratios are (4, -3, 12)

$$\text{So, } \sqrt{4^2 + (-3)^2 + 2^2} = \sqrt{16 + 9 + 4}$$

$$= \sqrt{29}$$

Dividing (1) by 13,

$$\frac{4}{\sqrt{29}}x - \frac{3}{\sqrt{29}}y + \frac{2}{\sqrt{29}}z = \frac{12}{\sqrt{29}}$$

which is in the normal form

Therefore length of perpendicular from the origin is $\frac{12}{\sqrt{29}}$ units

Question: 6

Mark against the

Solution:

Given:

$$1. \text{ Equation of line is } \frac{x+1}{3} = \frac{y-2}{4} = \frac{z+6}{5}$$

$$2. \text{ Equation of plane is } 2x - 3y + kz = 0$$

Formula Used: If two direction ratios are perpendicular, then

$$a_1a_2 + b_1b_2 + c_1c_2 = 0$$

Explanation:

Direction ratios of given line is (3, 4, 5)

Direction ratios of given plane is $(2, -3, k)$

Since the given line is parallel to the plane, the normal to the plane is perpendicular to the line.

So direction ratio of line is perpendicular to direction ratios of plane.

$$= 3 \times 2 + 4 \times -3 + 5 \times k = 0$$

$$= 6 - 12 + 5k = 0$$

$$\Rightarrow k = \frac{6}{5}$$

$$\text{Therefore, } k = \frac{6}{5}$$

Question: 7

Mark against the

Solution:

Given: $P(1, 2, -3)$ is a point on the plane. OP is perpendicular to the plane.

Explanation:

Let equation of plane be $ax + by + cz = d \dots (1)$

Substituting point P ,

$$= a + 2b - 3c = d \dots (2)$$

$$\overrightarrow{OP} = \hat{i} + 2\hat{j} - 3\hat{k}$$

Since OP is perpendicular to the plane, direction ratio of the normal is $(1, 2, -3)$

Substituting in (2)

$$1 + 4 + 9 = d$$

$$d = 14$$

Substituting the direction ratios and value of 'd' in (1), we get

$$x + 2y - 3z = 14$$

Therefore equation of plane is $x + 2y - 3z = 14$

Question: 8

Mark against the

Solution:

Given: Equation of plane is $2x - 4y + z = 7$

Line $\frac{(x-4)}{1} = \frac{(y-2)}{1} = \frac{(z-k)}{2}$ lies on the given plane.

Formula Used: Equation of a line is

$$\frac{(x - x_1)}{b_1} = \frac{(y - y_1)}{b_2} = \frac{(z - z_1)}{b_3} = \lambda$$

Where (x_1, y_1, z_1) is a point on the line and b_1, b_2, b_3 : direction ratios of line.

Explanation:

$$\text{Let } \frac{x-4}{1} = \frac{y-2}{1} = \frac{z-k}{2} = \lambda$$

So the given line passes through the point $(4, 2, k)$

Since the line lies on the given plane, $(4, 2, k)$ is a point on the plane.

Therefore, substituting the point on the equation for the plane,

$$= 8 - 8 + k = 7$$

$$= k = 7$$

Question: 9

Mark against the

Solution:

Given: The plane $2x + 3y + 4z = 12$ meets coordinate axes at A, B and C .

To find: Centroid of ΔABC

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x -axis, y -axis and z -axis respectively.

$$\text{Centroid of a triangle} = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$$

Explanation:

Equation of given plane is $2x + 3y + 4z = 12$

Dividing by 12,

$$\frac{x}{6} + \frac{y}{4} + \frac{z}{3} = 1$$

Therefore the intercepts on x, y and z-axis are 6, 6 and 3 respectively.

So, the vertices of ΔABC are (6, 0, 0), (0, 4, 0) and (0, 0, 3)

$$\text{Centroid} = \left(\frac{6+0+0}{3}, \frac{0+4+0}{3}, \frac{0+0+3}{3} \right)$$

$$= (2, 4/3, 1)$$

Therefore, the centroid of ΔABC is (2, 4/3, 1)

Question: 10

Mark against the

Solution:

Given: Centroid of ΔABC is (1, 2, 4)

To find: Equation of plane.

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x-axis, y-axis and z-axis respectively.

$$\text{Centroid of a triangle} = \left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}, \frac{z_1+z_2+z_3}{3} \right)$$

Explanation:

Let the equation of plane be

$$\frac{x}{A} + \frac{y}{B} + \frac{z}{C} = 1 \dots (1)$$

Therefore, $A = 3a$, $B = 3b$, $C = 3c$ where (a, b, c) is the centroid of the triangle with vertices (A, 0, 0), (0, B, 0) and (0, 0, C)

Substituting in (1),

$$\Rightarrow \frac{x}{3a} + \frac{y}{3b} + \frac{z}{3c} = 1$$

Here $a = 1$, $b = 2$ and $c = 4$

$$\Rightarrow \frac{x}{3} + \frac{y}{6} + \frac{z}{12} = 1$$

Multiplying by 12,

$$4x + 2y + z = 12$$

Therefore equation of required plane is $4x + 2y + z = 12$

Question: 11

Given: Plane passes through the point A(1, 0, -1).

Plane is perpendicular to the line

$$\frac{x+1}{2} = \frac{y+3}{4} = \frac{z+7}{-3}$$

To find: Equation of the plane.

Formula Used: Equation of a plane is $ax + by + cz = d$ where (a, b, c) are the direction ratios of the normal to the plane.

Explanation:

Let the equation of the plane be

$$ax + by + cz = d \dots (1)$$

Substituting point A,

$$a - z = d$$

Since the given line is perpendicular to the plane, it is the normal.

Direction ratios of line is 2, 4, -3

$$\text{Therefore, } 2 + 3 = d$$

$$d = 5$$

So the direction ratios of perpendicular to plane is 2, 4, -3 and $d = 5$

Substituting in (1),

$$2x + 4y - 3z = 5$$

Therefore, equation of plane is $2x + 4y - 3z = 5$

Question: 12

Mark against the

Solution:

Given: Line $\frac{x-1}{2} = \frac{y-2}{4} = \frac{z-3}{-3}$ meets plane $2x + 3y - z = 14$

To find: Point of intersection of line and plane.

Explanation:

Let the equation of the line be

$$\frac{x-1}{2} = \frac{y-2}{4} = \frac{z-3}{-3} = \lambda$$

Therefore, any point on the line is $(2\lambda + 1, 4\lambda + 2, -3\lambda + 3)$

Since this point also lies on the plane,

$$2(2\lambda + 1) + 3(4\lambda + 2) - (-3\lambda + 3) = 14$$

$$4\lambda + 2 + 12\lambda + 6 + 3\lambda - 3 = 14$$

$$19\lambda + 5 = 14$$

$$\lambda = \frac{19}{19} = 1$$

Therefore the required point is $(3, 5, 7)$.

Question: 13

Mark against the

Solution:

Given: Plane passes through $A(2, 2, 1)$ and $B(9, 3, 6)$. Plane is perpendicular to $2x + 6y + 6z = 1$

To find: Equation of the plane

Formula Used: Equation of a plane is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

where $a:b:c$ is the direction ratios of the normal to the plane.

(x_1, y_1, z_1) is a point on the plane.

Explanation:

Let the equation of plane be $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$

Since $(2, 2, 1)$ is a point in the plane,

$$a(x - 2) + b(y - 2) + c(z - 1) = 0 \dots (1)$$

Since $B(9, 3, 6)$ is another point on the plane,

$$a(9 - 2) + b(3 - 2) + c(6 - 1) = 0$$

$$7a + b + 5c = 0 \dots (2)$$

Since this plane is perpendicular to the plane $2x + 6y + 6z = 1$, the direction ratios of the normal to the plane will also be perpendicular.

$$\text{So, } 2a + 6b + 6c = 0 \Rightarrow a + 3b + 3c = 0 \dots (3)$$

Solving (1) and (2),

$$\frac{a}{\begin{vmatrix} 1 & 5 \\ 3 & 3 \end{vmatrix}} = \frac{-b}{\begin{vmatrix} 7 & 5 \\ 1 & 3 \end{vmatrix}} = \frac{c}{\begin{vmatrix} 7 & 1 \\ 1 & 3 \end{vmatrix}}$$

$$\frac{a}{-12} = \frac{b}{-16} = \frac{c}{20}$$

$$\frac{a}{3} = \frac{b}{4} = \frac{c}{-5}$$

$$a : b : c = 3 : 4 : -5$$

Substituting in (1),

$$3x - 6 + 4y - 8 - 5z + 5 = 0$$

$$3x + 4y - 5z - 9 = 0$$

Therefore the equation of the plane is $3x + 4y - 5z - 9 = 0$

Question: 14

Mark against the

Solution:

Given: Plane passes through the intersection of planes $3x - y + 2z - 4 = 0$ and $x + y + z - 2 = 0$.
Point $A(2, 2, 1)$ lies on the plane.

To find: Equation of the plane.

Formula Used: Equation of plane passing through the intersection of 2 planes P_1 and P_2 is given by $P_1 + \lambda P_2 = 0$

Explanation:

Equation of plane is

$$3x - y + 2z - 4 + \lambda (x + y + z - 2) = 0 \dots (1)$$

Since $A(2, 2, 1)$ lies on the plane,

$$6 - 2 + 2 - 4 + \lambda (2 + 2 + 1 - 2) = 0$$

$$2 + 3\lambda = 0$$

$$\lambda = \frac{-2}{3}$$

Substituting in (1) and multiplying by 3,

$$9x - 3y + 6z - 12 - 2(x + y + z - 2) = 0$$

$$9x - 3y + 6z - 12 - 2x - 2y - 2z + 4 = 0$$

$$7x - 5y + 4z - 8 = 0$$

Therefore the equation of the plane is $7x - 5y + 4z - 8 = 0$

Question: 15

Mark against the

Solution:

Given: Plane passes through $A(0, -1, 0)$, $B(2, 1, -1)$ and $C(1, 1, 1)$

To find: Equation of the plane

Formula Used: Equation of a plane is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

where $a:b:c$ is the direction ratios of the normal to the plane.

(x_1, y_1, z_1) is a point on the plane.

Explanation:

Let the equation of plane be $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$

Substituting point A,

$$ax + b(y + 1) + cz = 0 \dots (1)$$

Substituting points B and C,

$$2a + 2b - c = 0 \text{ and } a + 2b + c = 0$$

Solving,

$$\frac{a}{\begin{vmatrix} 2 & -1 \\ 2 & 1 \end{vmatrix}} = \frac{-b}{\begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix}} = \frac{c}{\begin{vmatrix} 2 & 2 \\ 1 & 2 \end{vmatrix}}$$

$$\frac{a}{4} = \frac{b}{-3} = \frac{c}{2}$$

Therefore, $a : b : c = 4 : -3 : 2$

Substituting in (1),

$$4x - 3(y + 1) + 2z = 0$$

$$4x - 3y + 2z - 3 = 0$$

Therefore equation of plane is $4x - 3y + 2z - 3 = 0$

Question: 16

Mark against the

Solution:

Given: Plane $2x - y + z = 0$ is parallel to the line

$$\frac{2x-1}{2} = \frac{2-y}{2} = \frac{z+1}{a},$$

To find: value of a

Formula Used: If two lines with direction ratios $a_1:a_2:a_3$ and $b_1:b_2:b_3$ are perpendicular, then

$$a_1b_1 + a_2b_2 + a_3b_3 = 0$$

Explanation:

Since the plane is parallel to the line, the normal to the plane will be perpendicular to the line.

Equation of the line can be rewritten as

$$\frac{x - \frac{1}{2}}{1} = \frac{y - 2}{-2} = \frac{z - (-1)}{a}$$

Direction ratio of the normal to the plane is $2 : -1 : 1$

Direction ratio of line is $1 : -2 : a$

Therefore,

$$2 + 2 + a = 0$$

$$a = -4$$

Therefore, $a = -4$

Question: 17

Mark against the

Solution:

$$\text{Given: Equation of line is } \frac{x+1}{1} = \frac{y}{2} = \frac{z-1}{1}$$

Equation of plane is $x - y + z = 0$

To find: Angle between a line and the normal to a plane.

Formula Used: If θ is the angle between two lines with direction ratios $b_1:b_2:b_3$ and $c_1:c_2:c_3$, then

$$\cos \theta = \frac{b_1c_1 + b_2c_2 + b_3c_3}{\sqrt{b_1^2 + b_2^2 + b_3^2} \times \sqrt{c_1^2 + c_2^2 + c_3^2}}$$

Explanation:

Direction ratios of given line is $1 : 2 : 1$

Direction ratios of the normal to the plane is $1 : -1 : 1$

Therefore,

$$\cos \theta = \frac{1 - 2 + 1}{\sqrt{1 + 4 + 1} \times \sqrt{1 + 1 + 1}}$$

$$\cos \theta = 0$$

$$\theta = 90^\circ$$

Therefore angle between them is 90°

Question: 18

Mark against the

Solution:

$$\text{Given: Line } \frac{x-1}{3} = \frac{y+2}{4} = \frac{z-3}{-2} \text{ meets plane } 2x - y + 3z - 1 = 0$$

To find: Point of intersection of line and plane.

Explanation:

Let the equation of the line be

$$\frac{x-1}{3} = \frac{y+2}{4} = \frac{z-3}{-2} = \lambda$$

Therefore, any point on the line is $(3\lambda + 1, 4\lambda - 2, -2\lambda + 3)$

Since this point also lies on the plane,

$$2(3\lambda + 1) - (4\lambda - 2) + 3(-2\lambda + 3) = 1$$

$$6\lambda + 2 - 4\lambda + 2 - 6\lambda + 9 = 1$$

$$-4\lambda = -12$$

$$\lambda = 3$$

Therefore required point is (10, 10, -3)

Question: 19

Mark against the

Solution:

Given: Plane passes through the points $A(a, 0, 0)$, $B(0, b, 0)$ and $C(0, 0, c)$

To find: Equation of plane.

Explanation:

The given points lie on the co-ordinate axes.

Therefore, the plane makes intercepts of a , b and c on the x , y and z -axis respectively.

Equation of the plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Question: 20

Mark against the

Solution:

Given: Equation of two planes are $2x - y + 2z = 3$ and $6x - 2y + 3z = 5$

To find: $\cos \theta$ where θ : angle between the planes

Formula Used: Angle between two planes $a_1x + b_1y + c_1z = 0$ and $a_2x + b_2y + c_2z = 0$ is

$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \times \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

where θ : angle between the planes,

Explanation:

Here $a_1 = 2$, $b_1 = -1$, $c_1 = 2$

$a_2 = 6$, $b_2 = -2$, $c_2 = 3$

$$\Rightarrow \cos \theta = \frac{12 + 2 + 6}{\sqrt{4 + 1 + 4} \times \sqrt{36 + 4 + 9}}$$

$$\Rightarrow \cos \theta = \frac{20}{3 \times 7}$$

$$\Rightarrow \cos \theta = \frac{20}{21}$$

Therefore, $\cos \theta = \frac{20}{21}$

Question: 21

Mark against the

Solution:

Given: Equation of two planes are $2x - y + z = 6$ and $x + y + 2z = 7$

To find: Angle between the two planes

Formula Used: Angle between two planes $a_1x + b_1y + c_1z = 0$ and $a_2x + b_2y + c_2z = 0$ is

$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \times \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

where θ : angle between the planes,

Explanation:

Here $a_1 = 2$, $b_1 = -1$, $c_1 = 1$

$a_2 = 1$, $b_2 = 1$, $c_2 = 2$

$$\Rightarrow \cos \theta = \frac{2 - 1 + 2}{\sqrt{4 + 1 + 1} \times \sqrt{1 + 1 + 4}}$$

$$\Rightarrow \cos \theta = \frac{3}{\sqrt{6} \times \sqrt{6}}$$

$$\Rightarrow \cos \theta = \frac{3}{6}$$

$$\Rightarrow \cos \theta = \frac{1}{2}$$

$$\Rightarrow \theta = \frac{\pi}{3}$$

Therefore angle between the planes is $\frac{\pi}{3}$

Question: 22

Mark against the

Solution:

Given: Equation of two planes are $\vec{r} \cdot (3\hat{i} - 6\hat{j} + 2\hat{k}) = 4$ and $\vec{r} \cdot (2\hat{i} - \hat{j} + 2\hat{k}) = 3$

To find: Angle between the two planes

Formula Used: Angle between two planes $a_1x + b_1y + c_1z = 0$ and $a_2x + b_2y + c_2z = 0$ is

$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \times \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

where θ : angle between the planes,

Explanation:

Since $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, the given equation of planes can be rewritten as:

$$3x - 6y + 2z = 4 \text{ and } 2x - y + 2z = 3$$

$$\text{Here } a_1 = 3, b_1 = -6, c_1 = 2$$

$$a_2 = 2, b_2 = -1, c_2 = 2$$

$$\Rightarrow \cos \theta = \frac{6 + 6 + 4}{\sqrt{9 + 36 + 4} \times \sqrt{4 + 1 + 4}}$$

$$\Rightarrow \cos \theta = \frac{16}{7 \times 3}$$

$$\Rightarrow \cos \theta = \frac{16}{21}$$

$$\Rightarrow \theta = \cos^{-1} \frac{16}{21}$$

Therefore angle between the planes is $\cos^{-1} \frac{16}{21}$

Question: 23

Mark against the

Solution:

Given: Plane passes through the points $A(2, 3, 1)$ and $B(4, -5, 3)$ and is parallel to x-axis

To find: Equation of plane

Formula Used: Equation of a plane parallel to x-axis is

$$b(y - y_1) + c(z - z_1) = 0$$

Explanation:

Let the equation of the plane be

$$b(y - y_1) + c(z - z_1) = 0$$

Since $A(2, 3, 1)$ lies on the plane,

$$b(y - 3) + c(z - 1) = 0 \dots (1)$$

Since $B(4, -5, 3)$ lies on the plane,

$$b(-5 - 3) + c(3 - 1) = 0$$

$$-8b + 2c = 0 \text{ or } -4b + c = 0$$

$$b : c = 1 : 4$$

Substituting in (1),

$$y - 3 + 4z - 4 = 0$$

$$y + 4z = 7$$

The equation of the plane is $y + 4z = 7$

Question: 24

Mark against the

Solution:

Given: Variable plane moves so that the sum of the reciprocals of its intercepts on the coordinate

axes is $(1/2)$

Formula Used: Equation of a plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Explanation:

Let the intercepts made by the plane on the co-ordinate axes be a , b and c .

$$\Rightarrow \frac{1}{a} + \frac{1}{b} + \frac{1}{c} = \frac{1}{2}$$

Let the equation of the plane be

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

On solving for each of the given options,

$$(0, 0, 0) \Rightarrow LHS \neq RHS$$

$$(1, 1, 1) \Rightarrow LHS \neq RHS$$

$$\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) \Rightarrow LHS = \frac{1}{2a} + \frac{1}{2b} + \frac{1}{2c} = \frac{1}{2} \times \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) = \frac{1}{4} \neq RHS$$

$$(2, 2, 2) \Rightarrow LHS = \frac{2}{a} + \frac{2}{b} + \frac{2}{c} = 2 \times \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) = 1 = RHS$$

Therefore, plane passes through the point $(2, 2, 2)$

Question: 25

Mark against the

Solution:

Given: Plane is perpendicular to $(2\hat{i} - 3\hat{j} + \hat{k})$ and is at a distance of 5 units from origin.

To find: Equation of plane

Formula Used: Equation of a plane is $lx + my + nz = p$ where p is the distance from the origin and l , m and n are the direction cosines of the normal to the plane

Explanation:

Direction ratio of normal to the plane is $2:-3:1$

$$|2\hat{i} - 3\hat{j} + \hat{k}| = \sqrt{4+9+1} = \sqrt{14}$$

Therefore, direction cosines of the normal to the plane is

$$l = \frac{2}{\sqrt{14}}, m = \frac{-3}{\sqrt{14}}, n = \frac{1}{\sqrt{14}}$$

Since the equation of a plane is $lx + my + nz = p$ where p is the distance from the origin,

$$2x - 3y + z = 5\sqrt{14}$$

Therefore, equation of the plane is $2x - 3y + z = 5\sqrt{14}$

Question: 26

Mark against the

Solution:

Given: Point $A(2, 3, 4)$ lies on a plane which is parallel to $5x - 6y + 7z = 3$

To find: Equation of the plane

Formula Used: Equation of a plane is

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

where $a:b:c$ is the direction ratios of the normal to the plane

(x_1, y_1, z_1) is a point on the plane.

Explanation:

Since the plane (say P_1) is parallel to the plane $5x - 6y + 7z = 3$ (say P_2), the direction ratios of the normal to P_1 is same as the direction ratios of the normal to P_2 .

i.e., direction ratios of P_1 is $5 : -6 : 7$

Let the equation of the required plane be

$$a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$$

Here $a = 5$, $b = -6$ and $c = 7$

Since $(2, 3, 4)$ lies on the plane,

$$5(x - 2) - 6(y - 3) + 7(z - 4) = 0$$

$$5x - 6y + 7z - 10 + 18 - 28 = 0$$

$$5x - 6y + 7z = 20$$

The equation of the plane is $5x - 6y + 7z = 20$

Question: 27

Mark against the

Solution:

Given: Perpendicular dropped from A(7, 14, 5) on to the plane $2x + 4y - z = 2$

To find: co-ordinates of the foot of perpendicular

Formula Used: Equation of a line is

$$\frac{x - x_1}{b_1} = \frac{y - y_2}{b_2} = \frac{z - z_3}{b_3} = \lambda$$

Where $b_1:b_2:b_3$ is the direction ratio and (x_1, x_2, x_3) is a point on the line.

Explanation:

Let the foot of the perpendicular be (a, b, c)

Since this point lies on the plane,

$$2a + 4b - c = 2 \dots (1)$$

Direction ratio of the normal to the plane is $2 : 4 : -1$

Direction ratio perpendicular = direction ratio of normal to the plane

So, equation of the perpendicular is

$$\frac{x - x_1}{2} = \frac{y - y_1}{4} = \frac{z - z_1}{-1} = \lambda$$

Since (a, b, c) is a point on the perpendicular,

$$\frac{x - a}{2} = \frac{y - b}{4} = \frac{z - c}{-1} = \lambda$$

$(7, 14, 5)$ is a point on the perpendicular.

$$\frac{7 - a}{2} = \frac{14 - b}{4} = \frac{5 - c}{-1} = \lambda$$

$$\text{So, } a = 7 - 2\lambda, b = 14 - 4\lambda, c = 5 + \lambda$$

Substituting in (1),

$$14 - 4\lambda + 56 - 16\lambda - 5 - \lambda = 2$$

$$21\lambda = 70 - 7 = 63$$

$$\lambda = 3$$

Therefore, foot of the perpendicular is $(1, 2, 8)$

Question: 28

Mark against the

Solution:

Given: Centroid of triangle is (α, β, γ)

To find: Equation of plane.

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x-axis, y-axis and z-axis respectively.

$$\text{Centroid of a triangle} = \left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3}, \frac{z_1 + z_2 + z_3}{3} \right)$$

Explanation:

Let the equation of plane be

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1 \dots (1)$$

Therefore, $A = 3\alpha, B = 3\beta, C = 3\gamma$ where (a, b, c) is the centroid of the triangle with vertices $(A, 0, 0), (0, B, 0)$ and $(0, 0, C)$

Substituting in (1),

$$\Rightarrow \frac{x}{3a} + \frac{y}{3b} + \frac{z}{3c} = 1$$

Here $a = \alpha, b = \beta$ and $c = \gamma$

$$\Rightarrow \frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 3$$

Therefore equation of required plane is $\frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 3$

Question: 29

Mark against the

Solution:

Given: Equation of plane is $\vec{r} \cdot (2\hat{i} - 3\hat{j} + 4\hat{k}) = 12$

To find: Intercepts made by the plane.

Formula Used: Equation of plane is $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$ where (x, y, z) is a point on the plane and a, b, c are intercepts on x-axis, y-axis and z-axis respectively.

Explanation:

The equation of the plane can be written as

$$2x - 3y + 4z = 12$$

Dividing by 12,

$$\frac{x}{6} + \frac{y}{-4} + \frac{z}{3} = 1 \text{ which is of the form } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Therefore the intercepts made by the plane are 6, -4, 3

Question: 30

Mark against the

Solution:

$$\text{Given: Equation of line is } \frac{x-2}{1} = \frac{y+3}{-2} = \frac{z+4}{-3}$$

Equation of the plane is $2x - 3y + z = 5$

To find: angle between line and plane

Formula Used: If θ is the angle between a line with direction ratio $b_1:b_2:b_3$ and a plane with direction ratio of normal $n_1:n_2:n_3$, then

$$\sin \theta = \frac{n_1 b_1 + n_2 b_2 + n_3 b_3}{\sqrt{n_1^2 + n_2^2 + n_3^2} \times \sqrt{b_1^2 + b_2^2 + b_3^2}}$$

Explanation:

Here direction ratio of the line is 1 : -2 : -3

Direction ratio of normal to the plane is 2 : -3 : 1

Therefore,

$$\sin \theta = \frac{2 + 6 - 3}{\sqrt{1 + 4 + 9} \times \sqrt{4 + 9 + 1}}$$

$$\sin \theta = \frac{5}{\sqrt{14} \times \sqrt{14}}$$

$$\theta = \sin^{-1} \frac{5}{14}$$

Therefore, angle between the line and plane is $\sin^{-1} \frac{5}{14}$

Question: 31

Mark against the

Solution:

Given: Equation of line is $\vec{r} \cdot (\hat{i} + \hat{j} - 3\hat{k}) + \lambda(2\hat{i} + 2\hat{j} + \hat{k})$

Equation of plane is $\vec{r} \cdot (6\hat{i} - 3\hat{j} + 2\hat{k}) = 5$

To find: angle between line and plane

Formula Used: If θ is the angle between a line with direction ratio $b_1:b_2:b_3$ and a plane with direction ratio of normal $n_1:n_2:n_3$, then

$$\sin \theta = \frac{n_1 b_1 + n_2 b_2 + n_3 b_3}{\sqrt{n_1^2 + n_2^2 + n_3^2} \times \sqrt{b_1^2 + b_2^2 + b_3^2}}$$

Explanation:

Here direction ratio of the line is 2 : 2 : 1

Direction ratio of normal to the plane is 6 : -3 : 2

Therefore,

$$\sin \theta = \frac{12 - 6 + 2}{\sqrt{4+4+1} \times \sqrt{36+9+4}}$$

$$\sin \theta = \frac{8}{3 \times 7}$$

$$\theta = \sin^{-1} \frac{8}{21}$$

Therefore, angle between the line and plane is $\sin^{-1} \frac{8}{21}$

Question: 32

Mark against the

Solution:

Given: Point is at $(\hat{i} + 2\hat{j} + 5\hat{k})$ and equation of plane is $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) + 17 = 0$

To find: distance of point from plane

Formula Used: Perpendicular distance from (x_1, y_1, z_1) to the plane $ax + by + cz + d = 0$ is

$$\frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Explanation:

The point is at $(1, 2, 5)$ and equation of plane is $x + y + z + 17 = 0$

$$\text{Distance} = \frac{1+2+5+17}{\sqrt{1+1+1}}$$

$$= \frac{25}{\sqrt{3}}$$

Therefore, distance = $\frac{25}{\sqrt{3}}$ units

Question: 33

Mark against the

Solution:

Given: The equations of the parallel planes are $2x - 3y + 6z = 5$ and $6x - 9y + 18z + 20 = 0$

To find: distance between the planes

Formula Used: Distance between two parallel planes $ax + by + cz + d_1 = 0$ and $ax_1 + by_1 + cz_1 + d_1 = 0$ is

$$\left| \frac{d_2 - d_1}{\sqrt{a^2 + b^2 + c^2}} \right|$$

Explanation:

The equations of the parallel planes are:

$$2x - 3y + 6z - 5 = 0$$

$$2x - 3y + 6z + \frac{20}{3} = 0$$

Therefore distance between them is

$$= \left| \frac{\frac{20}{3} + 5}{\sqrt{4 + 9 + 36}} \right|$$

$$= \left| \frac{35}{3 \times \sqrt{49}} \right|$$

$$= \frac{5}{3}$$

Therefore distance between the planes is $\frac{5}{3}$ units

Question: 34

Mark against the

Solution:

Given: The equations of the planes are $x + 2y - 2z + 1 = 0$ and $2x + 4y - 4z + 5 = 0$

To find: distance between the planes

Formula Used: Distance between two parallel planes $ax + by + cz + d_1 = 0$ and $ax_1 + by_1 + cz_1 + d_1 = 0$ is

$$\left| \frac{d_2 - d_1}{\sqrt{a^2 + b^2 + c^2}} \right|$$

Explanation:

The equations of the planes are:

$$x + 2y - 2z + 1 = 0 \text{ and } 2x + 4y - 4z - 4z + 5 = 0$$

Multiplying the equation of first plane by 2,

$$2x + 4y - 4z + 2 = 0$$

Therefore distance between them is

$$= \left| \frac{\frac{20}{3} + 5}{\sqrt{4 + 9 + 36}} \right|$$

$$= \left| \frac{35}{3 \times \sqrt{49}} \right|$$

$$= \frac{5}{3}$$

Therefore distance between the planes is $\frac{5}{3}$ units

Question: 35


Mark against the

Solution:

Given: Equation of plane is $2x - y + z + 3 = 0$. P is at $(1, 3, 4)$

To find: image of P

Explanation:



From the figure, P' is the image of P and B is the midpoint of PP'

If B is (a, b, c) , then

$$a = \frac{1 + x_1}{2}$$

$$b = \frac{3 + y_1}{2}$$

$$c = \frac{4 + z_1}{2}$$

B lies on the plane.

$$\text{So, } 2a - b + c + 3 = 0 \dots (1)$$

Also, equation of line PB is

$$\frac{x-1}{2} = \frac{y-3}{-1} = \frac{z-4}{1} = \lambda$$

So any point on the line PB will be of the form

$$x = 2\lambda + 1, y = -\lambda + 3, z = \lambda + 4$$

Since (a, b, c) will also be of this form we can substitute these values in (1)

$$= 2(2\lambda + 1) - (-\lambda + 3) + (\lambda + 4) + 3 = 0$$

$$= 4\lambda + 2 + \lambda - 3 + \lambda + 4 + 3 = 0$$

$$= 6\lambda = -6$$

$$= \lambda = -1$$

So $(a, b, c) = (-1, 4, 3)$

Substituting these values in the equations of a, b and c ,

$$-1 = \frac{1 + x_1}{2}$$

$$4 = \frac{3 + y_1}{2}$$

$$3 = \frac{4 + z_1}{2}$$

Therefore,

$$x_1 = -3, y_1 = 5, z_1 = 2$$

Therefore, the image is $(-3, 5, 2)$