Sample Question Paper - 10 Mathematics-Standard (041) Class- X, Session: 2021-22 TERM II

Time Allowed: 2 hours

General Instructions:

- 1. The question paper consists of 14 questions divided into 3 sections A, B, C.
- 2. All questions are compulsory.
- 3. Section A comprises of 6 questions of 2 marks each. Internal choice has been provided in two questions.
- 4. Section B comprises of 4questions of 3 marks each. Internal choice has been provided in one question.
- 5. Section C comprises of 4 questions of 4 marks each. An internal choice has been provided in one question. It contains two case study-based questions.

Section A

1. Find the 12th term from the end of the arithmetic progression 1,4,7,10,..., 88.[2]

OR

Determine the A.P. whose fourth term is 18 and the difference of the ninth term from the fifteenth term is 30.

- 2. Solve the given quadratic equation by factorization $y^2 3 = 0$ [2]
- 3. In the adjoining figure, O is the centre of the circle, AB is a chord and AT is tangent at A. If (2) $\angle AOB = 100^{\circ}$, then find $\angle BAT$.



- 4. If the heights of two right circular cones are in the ratio 1 : 2 and the perimeters of their bases [2] are in the ratio 3 : 4, what is the ratio of their volumes?
- 5. i. Find the mode of the following data: 25, 16, 19, 48, 19, 20, 34, 15, 19, 20, 21, 24, 19,16, 22, 16, [2]
 18, 20, 16, 19.
 - ii. If one of the 19's is changed to 16 in the above data, find the new mode.
- 6. In a rectangular part of dimensions, 50 m \times 40 m a rectangular pond is constructed so that [2] the area of grass strip of uniform breadth surrounding the pond would be 1184 m². Find the length and breadth of the pond.



Maximum Marks: 40

Solve the quadratic equation by factorization:

 $a^2b^2x^2 + b^2x - a^2x - 1 = 0$

Section **B**

7. Find the mean marks from the following data:

| Marks | Number of students |
|-----------|--------------------|
| Below 10 | 5 |
| Below 20 | 9 |
| Below 30 | 17 |
| Below 40 | 29 |
| Below 50 | 45 |
| Below 60 | 60 |
| Below 70 | 70 |
| Below 80 | 78 |
| Below 90 | 83 |
| Below 100 | 85 |

- 8. Draw a circle of radius 3 cm. Take two points P and Q on one of its extended diameter each at [3] a distance of 7 cm from its centre. Draw tangents to the circle from these two points P and Q.
- 9. Compute the median for the following cumulative frequency distribution:

| Less | Less than |
|---------|---------|---------|---------|---------|---------|---------|---------|-----------|
| than 20 | than 30 | than 40 | than 50 | than 60 | than 70 | than 80 | than 90 | 100 |
| 0 | 4 | 16 | 30 | 46 | 66 | 82 | 92 | 100 |

10. The lower window of a house is at a height of 2 m above the ground and its upper window is 4 [3] m vertically above the lower window. At certain instant the angles of elevation of a balloon from these windows are observed to be 60° and 30° respectively. Find the height of the balloon above the ground.

OR

A man on the deck of a ship, 12 m above water level, observes that the angle of elevation of the top of a cliff is 60° and the angle of depression of the base of the cliff is 30°. Find the distance of the cliff from the ship and the height of the cliff. [Use $\sqrt{3}$ = 1.732]

Section C

- 11. A toy is in the form of a cone mounted on a hemisphere of radius 3.5 cm. The total height of [4] the toy is 15.5 cm; find the total surface area and volume of the toy.
- 12. In the given figure, PT is tangent to the circle at T. If PA = 4 cm and AB = 5 cm, find PT.



[3]

[3]

[4]

In figure, PQ is a chord of length 16 cm, of a circle of radius 10 cm. The tangents at P and Q intersect at a point T. Find the length of TP.



13. Marine Insight uses extensive utilization of wind energy to move a vessel in the seawater. [4] They allow the towing kite to gain a height of anything between 100 metres - 300 metres. The sailing kite is made in such a way that it can be raised to its proper elevation and then brought back with the help of a 'telescopic mast' that enables the kite to be raised properly and effectively.

Based on the following figure, answer the question that follows:



- i. In the given figure, if sin θ = cos(3 θ 30°), where θ and 3 θ 30° are acute angles, then find the value of θ .
- ii. What should be the length of the rope of the kite sail in order to pull the ship at the angle θ (calculated above) and be at a vertical height of 150m?
- Saving money is a good habit and it should be inculcated in children from the beginning. Mrs. [4]
 Pushpa brought a piggy bank for her child Akshar. He puts one five-rupee coin of his savings in the piggy bank on the first day. He increases his savings by one five-rupee coin daily.



- i. If the piggy bank can hold 190 coins of five rupees in all, find the number of days he can contribute to put the five-rupee coins into it
- ii. Find the total money he saved.

Solution

MATHEMATICS STANDARD 041

Class 10 - Mathematics

Section A

```
1. Let the total number of terms be n.
```

So, First term (a) = 1 Last term $(a_n) = 88$ Common difference, *d*= 4-1=3 Now, as we know, $a_n = a + (n-1)d$ So, for the last term, 88=1+(n-1)3 88=1+3n-3 88=-2+3n 88+2=3n Further simplifying, 90=3n $n=\frac{90}{3}$ n=30 So, the 12^{th} term from the end means the 19^{th} term from the beginning.So, for the 19^{th} term (n = 19) a_{19} =1+(19-1)3 =1+(18)3 =1+54 =55 Therefore, the 12th term from the end of the given A.P. is 55. OR Let the given AP be a,a+d,a+2d,a+3d,..... 4th term = 18 (given) a+3d=18(1) Difference between 15^{th} term and 9^{th} term =30 (given) (a + 14d) - (a + 8d) = 30a + 14d - a -8d = 30 6d = 30 $d = \frac{30}{6}$ d = 5 put d=5 in eq. (1),we get, a + 3 × 5=18 a + 15 = 18 a = 18 - 15 a = 3 A.P. is a, a + d, a + 2d,.... i.e 3, 3 + 5, 3 + 10 i.e 3, 8, 13. 2. $y^2 - 3 = 0$ $\implies y^2 - (\sqrt{3})^2 = 0$ $\implies (y+\sqrt{3})(y-\sqrt{3})=0$ (Using Identity a² – b²= (a + b)(a – b)) Either $y + \sqrt{3} = 0$ or $y - \sqrt{3} = 0$ $\therefore y = -\sqrt{3}, \sqrt{3}$ Thus, $y = -\sqrt{3}$ and $\sqrt{3}$ are the roots of the given equation.

3. From the given figure AO and BO are the radius of the circle. So, AO = BO (both are radii) Hence, in \triangle AOB: $\angle OBA = \angle OAB$ (angles opposite to equal sides are equal).....(1) Also given , $\angle AOB = 100^{\circ}$ (2) In \triangle AOB, We have $igtriangle AOB + igtriangle OBA + igtriangle OAB = 180^\circ$ (sum of angles in a triangle is 180°) \Rightarrow 100° + $\angle OAB$ + $\angle OAB$ = 180° [from(1) & (2)] $\Rightarrow 2 \angle OAB = 80^{\circ}$ $\Rightarrow \angle OAB = 40^{\circ}$ We know that the radius and tangent at their point of contact are perpendicular to each other

 $\therefore \angle OAT = 90^{\circ}$ $\Rightarrow \angle OAB + \angle BAT = 90^{\circ}$ $\Rightarrow \angle BAT = 90^{\circ} - 40^{\circ} = 50^{\circ}$

4. Ratio in the heights of two cones = 1:2 and ratio in the perimeter of their bases = 3:4Let r_1 , r_2 be the radii of two cones and h_1 and h_2 be their heights

$$\therefore \frac{h_1}{h_2} = \frac{1}{2}$$

and $\frac{2\pi r_1}{2\pi r_2} = \frac{3}{4}$
$$\Rightarrow \frac{r_1}{r_2} = \frac{3}{4}$$

Now $\frac{\text{Volume of first cone}}{\text{Volume of second cone}} = \frac{\frac{1}{3}\pi r_1^2 h_1}{\frac{1}{3}\pi r_2^2 h_2}$
$$= \frac{r_1^2 h_1}{r_2^2 h_2} = \left(\frac{r_1}{r_2}\right)^2 \times \frac{h_1}{h_2}$$
$$= \left(\frac{3}{4}\right)^2 \times \frac{1}{2} = \frac{9}{16} \times \frac{1}{2} = \frac{9}{32}$$
$$\therefore \text{ Ratio in their volumes = 9 : 32}$$

5. The given data is 25, 16, 19, 48, 19, 20, 34, 15, 19, 20, 21, 24, 19, 16, 22, 16, 18, 20, 16, 19.

i. mode is 19

- ii. If 19 is changed to 16, then frequency of 16 is 5 and frequency of 19 becomes 4 i.e., 16 has maximum frequency. Therefore, 16 is the new mode.
- 6. Let width of grass strip be x mts.

: Length of pond = (50 - 2x) mt and Breadth of pond = (40 - 2x) mt And area of park - area of pond = area of grass strip $\Rightarrow (50 \times 40) - (50 - 2x)(40 - 2x) = 1184$ $2000 - 2000 + 180x - 4x^2 = 1184$ \Rightarrow $x^2 - 45x + 296 = 0$ \Rightarrow $x^2 - 37x - 8x + 296 = 0$ \Rightarrow \Rightarrow x(x - 37) - 8(x - 37) = 0 or, (x - 37) = 0 or (x - 8) = 0 \Rightarrow x = 8, 37 37 is rejected Length of pond = $50 - 2 \times 8 = 50 - 16 = 34$ m and breadth of pond 40 - 2(8) = 40 - 16 = 24 m. (

Given,
$$a^2b^2x^2 + b^2x - a^2x - 1 = 0$$

or, $b^2x(a^2x+1) - 1(a^2x+1) = 0$
or, $(b^2x-1)(a^2x+1) = 0$
 $(b^2x-1) = 0$ or $(a^2x+1) = 0$
 $b^2x = 1$ or $a^2x = -1$
 $\therefore \quad x = \frac{1}{b^2}$, or, $x = -\frac{1}{a^2}$
Hence, roots $= \frac{1}{b^2}, -\frac{1}{a^2}$

7. Calculation of Mean

Section **B**

| iculation of Mean. | | |
|--------------------|--|--|
| | | |

| C.I. | fi | xi | $u_i=rac{x_i-a}{h}$ | f _i u _i |
|----------|-------------------|----|----------------------|-------------------------------|
| 0 - 10 | 5 | 5 | -4 | -20 |
| 10 - 20 | 4 | 15 | -3 | -12 |
| 20 - 30 | 8 | 25 | -2 | -16 |
| 30 - 40 | 12 | 35 | -1 | -12 |
| 40 - 50 | 16 | 45 | 0 | 0 |
| 50 - 60 | 15 | 55 | 1 | 15 |
| 60 - 70 | 10 | 65 | 2 | 20 |
| 70 - 80 | 8 | 75 | 3 | 24 |
| 80 - 90 | 5 | 85 | 4 | 20 |
| 90 - 100 | 2 | 95 | 5 | 10 |
| Total | Σf_i = 85 | | | $\Sigma f_i u_i$ = 29 |

a = 45, h = 10

$$ar{x} = a + rac{\Sigma f_i u_i}{\Sigma f} imes h$$

$$=45+\frac{29}{25}\times 10$$

$$=45+3.417=48.41$$

Therefore, the mean marks are 48.41.

8. Required: To draw a circle of radius 3 cm and take two points P and Q on one of its extended its centre and then draw tangent o the circle from these two point P and Q.

Steps of construction:

i. Bisect PO. Let M be the mid-point of PO.

ii. Taking M as centre and MO as radius, draw a circle. Let it intersect the Given circle at the point A and B. iii. Join PA and PB

Then, PA and PB are the required two tangents.



iv. Bisect QO. Let N be the mid-point of QO.

v. Taking N as centre and NO as radius, draw a circle. Let it intersect the given circle at the point C and D.

vi. Join QC and QD

Then QC and QD

Justification: Join OA and OB

Then \angle PAO is an Angle in the semicircle and, therefore,

∠ PAO = 90°

 \Rightarrow PA \perp OA

Since OA is a radius of given circle, PA has to be tangent to the circle.

Similarly, PB is also a tangent to the circle.

Again, Join OC and OD.

Then \angle QCO is an angle on the semicircle and, therefore \angle QOC = 90°

Since OC is a radius of the given circle, QC has to be a tangent to the circle.

Similarly, QD is also a tangent to the circle.

9. We are given the cumulative frequency distribution. So, we first construct a frequency table from the given cumulative frequency distribution and then we will make necessary computations to compute median

| Class intervals | Frequency (f) | Cumulative frequency (c.f.) |
|-----------------|---------------|-----------------------------------|
| 20-30 | 4 | 4 |
| 30-40 | 12 | 16 |
| 40-50 | 14 | 30 |
| 50-60 | 16 | 46 |
| 60-70 | 20 | 66 |
| 70-80 | 16 | 82 |
| 80-90 | 10 | 92 |
| 90-100 | 8 | 100 |
| | | \mathbf{N} = Σf_i = 100 |

Here, N = Σf_i = 100 $\therefore \frac{N}{2}$ = 50

We observe that the cumulative frequency just greater than $\frac{N}{2}$ = 50 is 66 and the corresponding class is 60-70. So, 60-70 is the median class.



From the figure,

let the height of balloon is = x+4+2 = x+6 $\tan 30^\circ = \frac{x}{y}$ $\Rightarrow \frac{1}{\sqrt{3}} = \frac{x}{y}$ $\Rightarrow y = \sqrt{3}x$ (i) And, $\tan 60^\circ = \frac{x+4}{y}$ $\Rightarrow \tan 60^\circ = \frac{x+4}{y}$ $\Rightarrow \sqrt{3} = \frac{x+4}{y}$ $\Rightarrow \sqrt{3} = \frac{x+4}{\sqrt{3}x}$ $\Rightarrow 3x = x + 4$ $\Rightarrow 2x = 4 \Rightarrow x = 2$ Thus, height of balloon $= x + 4 + 2 \Rightarrow 2 + 4 + 2 = 8m$ A is the position of the man, OA = 12 m, BC is cliff.



Section C

11. The Radius of the toy (r) = 3.5 cm Total height of the toy = 15.5 cm ∴ Height of the conical part is = 15.5 - 3.5 = 12 cm Slant height of the conical part (l) $= \sqrt{r^2 + h^2} = \sqrt{(3.5)^2 + (12)^2}$ $= \sqrt{12.25 + 144} = \sqrt{156.25} = 12.5 cm$



i. Now total surface area of the toy = curved surface area of conical part + curved surface area of hemipherical part

$$= \pi r l + 2\pi r^{2} = \pi r (l + 2r)$$

= $\frac{22}{7} \times 3.5(12.5 + 2 \times 3.5) \text{ cm}^{2}$
= 11 (12.5 + 7) = 11 × 19.5 cm²
= 214.5 cm²

ii. Volume of the toy
$$=$$
 $\frac{1}{3}\pi r^2h + \frac{2}{3}\pi r^3$

$$= \frac{1}{3}\pi r^{2}(h+2r)$$

$$= \frac{1}{3} \times \frac{22}{7}(3.5)^{2}(12+2\times 3.5)\text{cm}^{3}$$

$$= \frac{1}{3} \times \frac{22}{7} \times 12.25(12+7)\text{cm}^{3}$$

$$= \frac{22}{3} \times 1.75 \times 19\text{cm}^{3}$$

$$= \frac{731.5}{3} = 243.83 \text{ cm}^{3}$$

12. Given, PT is tangent to the circle at T. PA = 4 cm and AB = 5 cm.



Construction: Draw OC \perp AB and join OP, OT and OA. Proof: In right \triangle OCP $\mathrm{OP}^2 = \mathrm{PC}^2 + \mathrm{OC}^2$ $\mathrm{OP}^2 = [\mathrm{AP} + \mathrm{AC}]^2 + \mathrm{OC}^2$ $\mathbf{OP}^2 = \left[4 + \frac{1}{2}\mathbf{AB}\right]^2 + \mathbf{OC}^2 \text{ [OC } \perp \text{ AB, AC = BC]}$ $\Rightarrow OP^{2} = \left(4 + \frac{5}{2}\right)^{2} + OC^{2}$ $\Rightarrow OP^{2} = \left(\frac{13}{2}\right)^{2} + OC^{2} \quad ..(i)$ In right \triangle OCA, $OA^2 = OC^2 + AC^2$ $OA^2 - AC^2 = OC^2$ $OA^{2} - \left(\frac{5}{2}\right)^{2} = OC^{2}$...,(ii) ∴ eq (i) becomes. $OP^{2} = \left(\frac{13}{2}\right)^{2} + OA^{2} - \left(\frac{5}{2}\right)^{2}$ $OP^{2} = \frac{169}{4} - \frac{25}{4} + OA^{2}$ $OP^{2} = \frac{144}{4} + OA^{2}$ $\Rightarrow OP^{2} = 36 + OA^{2} \dots (iii)$ Also, $OP^2 = OT^2 + PT^2$...(iv) from (iv) and (iii), $\mathrm{PT}^2 + \mathrm{OT}^2 = 36 + \mathrm{OA}^2$ $\Rightarrow \mathrm{PT}^2 = 36$ [\therefore OT = OA = radii] PT = 6cm.

OR



Given, PQ is a chord of length 16 cm, of a circle of radius 10 cm. The tangents at P and Q intersect at a point T.

Construction: Join OP and OQ. In triangles OTP and OTQ, OT is common OP = OQ(radii) TP = TQ $\therefore \Delta OPT \cong \Delta OQT$ $\therefore \angle POT = \angle QOT$ Consider, triangles OPR and OQR; OP = OQ (radii); OP is common $\angle POR = \angle QOR$ [from(i)] $\therefore \Delta OPR \cong \Delta OQR$ (SAS cong. rule) $\therefore PR = RQ = \frac{1}{2} \times 16 = 8$ cm...(ii)

 $\angle \text{ORP} = \angle \text{ORQ} = 90^{\circ}$ (iii) In right angled triangle TRP, $TR^2 = TP^2 - (8)^2 = TP^2 - 64$ [from (iii)] ,..(iv)] Also $OT^2 = TP^2 + (10)^2$ $(TR+6)^2 = TP^2 + 100$ $[: OR = \sqrt{100 - 64} = 6]$ $TR^2 + 12TR + 36 = TP^2 + 100$ $TP^2 - 64 + 12TR + 36 = TP^2 + 100$ $12\text{TR} = 128 \Rightarrow \text{TR} = \frac{32}{3} \text{cm}$ From (iv) $\left(\frac{32}{3}\right)^2 = TP^2 - 64$ $\Rightarrow TP^{2} = \frac{1024}{9} + 64 = \frac{1024 + 576}{9} = \frac{1600}{9}$ $\Rightarrow TP = \frac{40}{3} \text{ cm}$ 13. i. $\sin \theta = \cos(3\theta - 30^{\circ})$ $\cos(90^\circ - \theta) = \cos(3\theta - 30^\circ)$ $\Rightarrow 90^{\circ} - heta = 3 heta - 30^{\circ} \Rightarrow heta = 30^{\circ}$ ii. $rac{AB}{AC} = \sin 30^\circ$: Length of rope, AC = $\frac{AB}{sin30^{\circ}} = \frac{150}{\frac{1}{2}} = 150 \times 2 = 300 \text{ m}$ 14. Child's savings day wise are, 5, 10, 15, 20, 25, to *n* days $1 \operatorname{coins}$, $2 \operatorname{coins}$, $3 \operatorname{coins}$, $4 \operatorname{coins}$, $5 \operatorname{coins}$, $5 \operatorname{coins}$ We can have at most 190 coins i.e., 1 + 2 + 3 + 4 + 5 + to n term = 190 $\Rightarrow rac{n}{2}[2 imes 1+(n-1)1]=190$ \Rightarrow n(n + 1) = 380 \Rightarrow n² + n - 380 = 0 \Rightarrow (n + 20)(n-19) = 0 \Rightarrow (n + 20)(n - 19) = 0 \Rightarrow n = -20 or n = 19 \Rightarrow n = -20 or n = 19 But number of coins cannot be negative \therefore n = 19 (rejecting n = -20) So, number of days = 19 Total money she saved = 5 + 10 + 15 + 20 + ... = 5 + 10 + 15 + 20 + ... upto 19 terms $=\frac{19}{2}[2 \times 5 + (19 - 1)5]$ $=\frac{19}{2}[100]=\frac{1900}{2}=950$ So, number of days = 19 and total money she shaved = Rs. 950