

WBBSE Class 10 Maths Madhyamik Question Paper 2018

2018

MATHEMATICS

Time - Three Hours Fifteen Minutes

(First fifteen minutes for reading the question paper only)

Full Marks - 90

(For Regular and Sightless Regular Candidates)

Full Marks - 100

(For External and sightless External Candidates)

Special credits will be given for answers which are brief and to the point.

Marks will be deducted for spelling mistakes, untidiness and bad handwriting.

General Instructions :

The answers of the Question Nos 1, 2, 3, 4 are to be written at the beginning of the answer-script mentioning the question numbers in the serial order. Necessary calculation and drawing must be given on the right hand side by drawing margins on the first pages on the Answer - script. Tables and Calculators of any type are not allowed. Approximate value of $\pi = \frac{22}{7}$, if necessary. Graph paper will be supplied if required. Arithmetic problems may be solved by algebraic method.

1. Choose the correct option in each case from the following questions :

[1 x 6 = 6]

(i) Interest on Rs. a at the simple interest 10% per annum for b months is :

(a) Rs. $\frac{ab}{100}$ (b) Rs. $\frac{ab}{120}$ (c) Rs. $\frac{ab}{1200}$ (d) Rs. $\frac{ab}{10}$

(ii) If $x \propto y$ then

(a) $x^2 \propto y^2$ (b) $x^3 \propto y^2$ (c) $x \propto y^2$ (d) $x^2 \propto y^2$

(iii) If $\angle A = 100^\circ$ of a cyclic quadrilateral ABCD, then the value of $\angle C$ is :

- (a) 50° (b) 20° (c) 80° (d) 180°

(iv) The sexagesimal value of $\frac{7\pi}{12}$ is :

- (a) 115° (b) 150° (c) 135° (d) 105°

(v) If the side of a cube is a unit and the diagonal of the cube is d unit then the relation between a and d will be.

- (a) $\sqrt{2}a = d$ (b) $\sqrt{3}a = d$ (c) $a = \sqrt{3}d$ (d) $a = \sqrt{2}d$

(vi) If the mean of the numbers 6, 7, x, 8, y, 16 is 9 then :

- (a) $x + y = 21$ (b) $x + y = 17$ (c) $x - y = 21$ (d) $x - y = 9$

2. Fill up the blanks (any five) :

[1 x 5 = 5]

(i) If the simple interest of a principal for n years at r% p.a. be Rs $\frac{pnr}{25}$, then

the principal will be Rs _____ .

(ii) The equation $(a - 2)x^2 + 3x + 5 = 0$ will not be a quadratic equation for a
= _____.

(iii) if ABCD IS a cyclic parallelogram then $\angle A$ is _____.

(iv) If $\tan 35^\circ \tan 55^\circ = \sin \theta$, then the lowest positive value of θ will be
_____.

(v) The shape of a pencil with one end sharpened is the combination of a
cylinder and a _____

(vi) The measures of central tendency are Mean, Median and _____.

3. Write True or False (any five):

[1 x 5 = 5]

(i) At same rate of interest the simple interest for 2 years is more than the
compound interest on the same principal.

(ii) x^3y , x^2y^2 and xy^3 are in continued proportion.

(iii) The angle in the segment of a circle which is less than a semi-circle is

an obtuse angle.

(iv) Simplest value of $\sec^2 27^\circ - \cot^2 63^\circ$ is 1.

(v) If the radius of a sphere is twice that of 1st sphere then the volume of the sphere will be twice that of the 1st sphere.

(vi)

Score	1	2	3	4	5
No. Of students	3	6	4	7	5

The mode of the distribution is 3.

4. Answer the following questions (any ten) :

[2 x 10 = 20]

(i) The rate of simple interest per annum reduces from 4% to $3\frac{3}{4}\%$ and for this, a person's annual income decreases by Rs. 60. Determine the principal of that person.

(ii) A and B start a business with Rs. 15,000 and Rs. 45,000 respectively. After 6 months B received Rs 3,030 as profit. What is A's profit?

(iii) If $2x + \frac{1}{x} = 2$ the find the value of $\frac{x}{2x^2 + x + 1}$.

(iv) If the roots of a quadratic equation be 2 and -3, then write the equation.

(v) The line parallel to BC of $\triangle ABC$ meets AB and AC at P and Q respectively. If AP = 4 cm, QC = 9 cm and PB = AQ, then find the length of PB.

(vi) The radius of a circle with centre O is 5 cm. P is a point at a distance 13 cm from O. PQ and PR are two tangents to this circle. Find the area of the quadrilateral PQOR .

(vii) The two chords AB and CD of a circle are at equal distance from the centre O. If $\angle AOB = 60^\circ$ and CD = 6 cm, then calculate the length of the radius of the circle.



- (viii) If $\tan \theta + \cot \theta = 2$, then find the value of $\tan^7 \theta + \cot^7 \theta$.
- (ix) If the ratio of length of shadow of a tower and height of the tower is $\sqrt{3} : 1$, find the angle of elevation of the Sun.
- (x) The volumes of two right circular cylinders are same. The ratio of their height is $1 : 2$. Find the ratio of their radii.
- (xi) The volume of a solid hemisphere is 144π cu. cm, then find the diameter of the sphere.
- (xii) The mean of a frequency distribution is 8.1, if $\sum f_i x_i = 132 + 5K$ and $\sum f_i = 20$ then what is the value of K?

5. Answer any one question :

[5 x 1 = 5]

- (a) Aminur has taken a loan of Rs. 64, 000 from a bank. If the rate of interest be 2.5 paise per rupee per annum, calculate the compound interest payable after 2 years.
- (b) A, B and C start a business with the capital of Rs. 6,000, Rs 8,000 and Rs. 9, 000 respectively. After few months A invests Rs 3, 000 more in the business. At the end of the year they gained Rs 30,000 and C got Rs. 10,800 as share of profit. When did A invest Rs. 3,000 more?

6. Answer any one question :

[3 x 1 = 3]

- (a) solve : $\left(\frac{x+4}{x-4}\right)^2 - 5\left(\frac{x+4}{x-4}\right) + 6 = 0$, ($x \neq 4$).
- (b) The digit in the unit's place of a two digit number is 6 more than that at the ten's place. The product of the digits is 12 less than the number. Find the possible values of the digit in the unit place.

7. Answer any one question :

[3 x 1 = 3]

- (a) Find the simplest value of : $\sqrt{7}(\sqrt{5}-\sqrt{2})-\sqrt{5}(\sqrt{7}-\sqrt{2})+\frac{2\sqrt{2}}{\sqrt{5}+\sqrt{7}}$

(b) If $x \propto y$, and $y \propto z$ then prove that: $(x^2 + y^2 + z^2) \propto (xy + yz + zx)$

8. Answer any one question :

[3 x 1 = 3]

(a) If $\frac{a+b-c}{a+b} = \frac{b+c-a}{b+c} = \frac{c+a-b}{c+a}$ and $a+b+c \neq 0$ then prove that $a+b=c$

(b) If $x:a = y:b = z:c$ then show that: $(a^2 + b^2 + c^2)(x^2 + y^2 + z^2) = (ax + by + cz)^2$

9. Answer any one question :

[5 x 1 = 5]

(a) Prove that, if a perpendicular is drawn on the hypotenuse from the right angular point of a right angled triangle, two triangles so formed on the two sides of the perpendicular are each similar to the original triangle and also similar to each other.

(b) Prove that the tangent and the radius through the point of contact of a circle are perpendicular to each other.

10. Answer any one question :

[3 x 1 = 3]

(a) In $\triangle ABC$, AD is perpendicular on BC and $AD^2 = BD \cdot DC$, prove that $\angle BAC$ is a right angle.

(b) A straight line intersects one of the two concentric circles at the points A and B and other at the points C and D. Prove that $AC = BD$.

11. Answer any one question :

[5 x 1 = 5]

(a) Construct two circles of radii 4 cm and 2 cm and the distance between their centres is 7 cm. Construct a direct common tangent of the circles. (only traces of construction are required).

(b) Construct a triangle whose two sides are 9 cm and 7 cm and the angle between them is 60° . Construct the incircle of the triangle. (only traces of construction are required).

12. Answer any two questions :

[3 x 2 = 6]

(a) An arc of length 220 cm of a circle makes an angle 60° at the centre.

Find the radius of the circle.

(b) If $\cos^2 \theta - \sin^2 \theta = \frac{1}{2}$, then find the value of $\tan^2 \theta$

(c) Find the value of : $\frac{\sec 17^\circ}{\operatorname{cosec} 73^\circ} + \frac{\tan 68^\circ}{\cot 22^\circ} + \cos^2 44^\circ + \cos^2 46^\circ$

13. Answer any one question :

[5 x 1 = 5]

(a) The length of the shadow of a post becomes 3 meters smaller when the angle of elevation of the Sun increases from 45° to 60° . Find the height of the post.

(b) A man standing on a railway bridge $5\sqrt{3}$ meters high, observes the engine of a train at an angle of depression 30° . But after 2 seconds, he observes the engine at an angle of depression 45° on the other side of the bridge. Find the speed of the train.

14. Answer any TWO questions :

[4 x 2 = 8]

(a) Each side of a cube is decreased by 50%. Calculate the ratio of the volumes of original and changed cube.

(b) The total surface area of a right circular cylindrical pot without lid be 200 sq.cm. If the radius of the base be 7 cm find the quantity of water in litres contained in the pot. (1 litre = 1 cu. dm)

(c) A tank of length 21 dcm, breadth 11 dcm and 6 dcm deep is half filled with water. If 100 solid iron balls of diameter 21 cm are completely immersed in the tank, then how much dcm of water level is raised?

15. Answer any two questions :

[4 x 2 = 8]

(a) Find the mode from the following frequency distribution table of ages

of examinees of an entrance examination :

Age (in year)	16 - 18	18-20	20 - 22	22 - 24	24 -26
No. Of examinees	45	75	38	22	20

(b) Find the median of given data :

Class interval	1 - 5	6 - 10	11 - 15	16 - 20	21 - 25	26 - 30	31 - 35
Frequency	2	3	6	7	5	4	3

(c) From the frequency distribution table given below, draw less than ogive :

Marks obtained	50 - 60	60 - 70	70 - 80	80 - 90	90 - 100
Frequency	4	8	12	6	10

WBBSE Class 10th Maths Question Paper With Solutions 2018

Question 1: Choose the correct option in each case from the following questions: **[1 x 6 = 6]**

(i) Interest on Rs. a at the simple interest 10% per annum for b months is:

- (a) Rs. $ab / 100$ (b) Rs. $ab / 120$ (c) Rs. $ab / 1200$ (d) Rs. $ab / 10$

Answer: (b)

$$R = 10\%$$

$$T = b \text{ months} = b / 12 \text{ years}$$

$$SI = PTR / 100$$

$$= a * b * (10) / 100 * 12$$

$$= ab / 120$$

(ii) If $x \propto y$ then

- (a) $x^2 \propto y^2$ (b) $x^3 \propto y^2$ (c) $x \propto y^2$ (d) $x^2 \propto y^2$

Answer: (a)

$$x \propto y$$

$$x = ky$$


$$x^2 = k^2 y^2$$

$$x^2 \propto y^2$$

(iii) If $\angle A = 100^\circ$ of a cyclic quadrilateral ABCD, then the value of $\angle C$ is:

- (a) 50° (b) 20° (c) 80° (d) 180°

Answer: (c)



$$\angle A = 100^\circ$$

$$\angle A + \angle C = 180^\circ$$

$$\angle C = 180^\circ - 100$$

$$\angle C = 80^\circ$$

(iv) The sexagesimal value of $7\pi / 12$ is:

(a) 115°

(b) 150°

(c) 135°

(d) 105°

Answer: (d)

$$7\pi / 12$$

$$= (7 * 180) / 12$$

$$= 105^\circ$$

(v) If the side of a cube is a unit and the diagonal of the cube is d unit then the relation between a and d will be.

(a) $\sqrt{2}a = d$

(b) $\sqrt{3}a = d$

(c) $a = \sqrt{3}d$

(d) a

$= \sqrt{2}d$

Answer: (b)

(vi) If the mean of the numbers 6, 7, x, 8, y, 16 is 9 then:

(a) $x + y = 21$ (b) $x + y = 17$ (c) $x - y = 21$ (d) $x - y = 9$

Answer: (b)

6, 7, x, 8, y, 16 is 9

Mean = 9

$$9 = [6 + 7 + x + 8 + y + 16] / 6$$

$$54 = 37 + x + y$$

$$x + y = 17$$

Question 2: Fill up the blanks (any five):

[1 x 5 = 5]

(i) If the simple interest of a principal for n years at r% p.a. be Rs. $\frac{pnr}{25}$, then the principal will be Rs _____. **[4P]**

(ii) The equation $(a - 2)x^2 + 3x + 5 = 0$ will not be a quadratic equation for a = _____. **[a = 2]**

(iii) if ABCD is a cyclic parallelogram then A is _____. **[90°]**

(iv) If $\tan 35^\circ \tan 55^\circ = \sin \theta$, then the lowest positive value of θ will be _____. **[90°]**

$$\tan 35^\circ \tan 55^\circ = \sin \theta$$

$$\tan 35^\circ \tan (90^\circ - 35^\circ) = \sin \theta$$

$$\tan 35^\circ * \cot 35^\circ = \sin \theta$$

$$1 = \sin \theta$$

$$\theta = 90^\circ$$

(v) The shape of a pencil with one end sharpened is the combination of a cylinder and a _____. **[cone]**

(vi) The measures of central tendency are Mean, Median and _____. **[Mode]**

Question 3: Write True or False (any five):

[1 x 5 = 5]

(i) At the same rate of interest, the simple interest for 2 years is more than the compound interest on the same principal. **[False]**

(ii) x^3y , x^2y^2 and xy^3 are in continued proportion. **[True]**

(iii) The angle in the segment of a circle which is less than a semicircle is an obtuse angle. **[True]**

(iv) Simplest value of $\sec^2 27^\circ - \cot^2 63^\circ$ is 1. **[True]**

$$\sec^2 27^\circ - \cot^2 63^\circ$$

$$= \sec^2 27^\circ - \cot^2 [90 - 27]$$

$$= \sec^2 27^\circ - \tan^2 27^\circ$$

$$= 1$$

(v) If the radius of a sphere is twice that of the 1st sphere then the volume of the sphere will be twice that of the 1st sphere. **[False]**

(vi)

Score	1	2	3	4	5
Number of students	3	6	4	7	5

The mode of the distribution is 3. **[False]**

Question 4: Answer any one question:

[3 x 1 = 3]

[i] The rate of simple interest per annum reduces from 4% to 3 (3 / 4) % and for this, a person's annual income decreases by Rs. 60. Determine the principal of that person.

Solution:

$$SI (1) = PTR / 100$$

$$= P * 4 * 1 / 100$$

$$= P / 25$$

$$\text{Total income} = P + (P / 25) = 26P / 25$$

$$\begin{aligned}
SI(2) &= PTR / 100 \\
&= P * (15 / 4) * 1 / 100 \\
&= 3P / 80 \\
\text{Total income} &= P + (3P / 80) = 83P / 80 \\
[26P / 25] - 60 &= 83P / 80 \\
26P - 1500 / 25 &= 83P / 80 \\
415P &= 416P - 24000 \\
P &= 24000
\end{aligned}$$

[ii] A and B start a business with Rs. 15,000 and Rs. 45,000, respectively. After 6 months B received Rs 3,030 as profit. What is A's profit?

Solution:

Amount invested by A = Rs. 15000

Amount invested by B = Rs. 45000

The ratio of their profits after 6 months would be

A:B

15000:45000

15:45

1:3

Profit earned by B = Rs. 3030

According to the question, it becomes,

$$3x = 3010$$

$$x = 3010 / 3$$

$$x = 1010$$

So, A's profit after 6 months would be Rs. 1010.

[iii] If $2x + [1 / x] = 2$, then find the value of $x / [2x^2 + x + 1]$.

Solution:

$$2x + [1 / x] = 2$$

$$2x^2 + 1 = 2x$$

$$\text{LHS} = x / [2x^2 + x + 1]$$

$$= x / 2x^2 + 1 + x$$

$$= x / 2x + x$$

$$= x / 3x$$

$$= 1 / 3$$

[iv] If the roots of a quadratic equation are 2 and -3, then write the equation.

Solution:

$$a = 2, b = -3$$


$$x^2 - (a + b)x + ab = 0$$

$$x^2 - (2 + (-3))x + (2 * -3) = 0$$

$$x^2 + x - 6 = 0$$

[v] The line parallel to BC of $\triangle ABC$ meets AB and AC at P and Q respectively. If $AP = 4$ cm, $QC = 9$ cm and $PB = AQ$, then find the length of PB.

Solution:



Since $PQ \parallel BC$

By basic proportionality theorem,

$$AP / PB = AQ / QC$$

$$4 / x = x / 9$$


$$36 = x^2$$

$$x = 6\text{cm}$$

Length of PB is 6cm.

[vi] The radius of a circle with centre O is 5 cm. P is a point at a distance 13 cm from O. PQ and PR are two tangents to this circle. Find the area of the quadrilateral PQOR.

Solution:



PQ & PR are 2 tangents and OQ & OR are 2 radii at contact point Q & R.

$\angle PQO = 90^\circ$ [a tangent to a circle is perpendicular to the radius through the point of contact]

By Pythagoras theorem

$$PQ^2 = OP^2 - OQ^2$$

$$PQ^2 = 13^2 - 5^2$$

$$= 169 - 25$$

$$= 144$$

$$PQ = \sqrt{144} = 12$$

$$PQ = 12\text{cm}$$

$PQ = PR = 12\text{cm}$ [The lengths of two tangents drawn from an external point to a circle are equal]

In $\triangle OPQ$ & $\triangle OPR$

$OQ = OR$ (5cm) given

$OP = OP$ (Common)

$PQ = PR$ (12cm)

Hence, $\triangle OPQ$ and $\triangle OPR$ are congruent. (by SSS congruence)

Area of $\triangle OPQ$ = Area $\triangle OPR$

Area of quadrilateral QORP = $2 \times$ (area of $\triangle OPR$)


Area of quadrilateral QORP = $2 \times \frac{1}{2} \times \text{base} \times \text{altitude}$

Area of quadrilateral QORP = $OR \times PR$

$$\begin{aligned}\text{Area of quadrilateral QORP} &= 12 \times 5 \\ &= 60 \text{ cm}^2\end{aligned}$$

[vii] The two chords AB and CD of a circle are at equal distance from the centre O. If $\angle AOB = 60^\circ$ and $CD = 6 \text{ cm}$, then calculate the length of the radius of the circle.

Solution:



In $\triangle AOB$ and $\triangle COD$,

$$AB = CD$$

$$OA = OC = OB = OD$$

All the angles and sides should be equal.

$$AB = 6 \text{ cm}$$

$$\angle COD = 60^\circ$$

$$AE = AB / 2$$

$$AE = 6 / 2 = 3 \text{ cm}$$

In $\triangle AOE$ and $\triangle BOE$,

$$OA = OB$$

$$OE = OE$$

$$AE = BE$$

By SSS congruence,

$$\triangle AOE \cong \triangle BOE$$

$$\angle AOE = \angle BOE$$

$$\angle AOE = 30^\circ$$

$\sin \theta = \text{perpendicular} / \text{hypotenuse}$

$$\sin 30^\circ = AE / OA$$

$$[1 / 2] = AE / OA$$

$$OA = 3 * 2$$

$$OA = 6\text{cm}$$

[viii] If $\tan \theta + \cot \theta = 2$, then find the value of $\tan^7 \theta + \cot^7 \theta$.

Solution:

$$\tan \theta + \cot \theta = 2$$

$$\Rightarrow \tan \theta + 1 / \tan \theta = 2$$

$$\Rightarrow \tan^2 \theta + 1 = 2 \tan \theta$$

$$\Rightarrow \tan^2 \theta - 2 \tan \theta + 1 = 0$$

$$\Rightarrow (\tan \theta - 1)^2 = 0$$

$$\Rightarrow \tan \theta = 1$$

$$\cot \theta = 1 / \tan \theta = 1$$

$$\tan^7 \theta + \cot^7 \theta$$

$$= (\tan \theta)^7 + (\cot \theta)^7$$

$$= (1)^7 + (1)^7$$

$$= 1 + 1$$

$$= 2$$

[ix] If the ratio of the length of the shadow of a tower and height of the tower is $\sqrt{3}:1$, find the angle of elevation of the sun.

Solution:

The ratio of the height of a tower and the length of its shadow is given by $\sqrt{3}:1$.

$$\tan \theta = \text{Height of tower} / \text{Length of shadow}$$

$$\tan \theta = \sqrt{3} / 1$$

$$\tan \theta = \tan 60^\circ$$

$$\theta = 60^\circ$$

Hence, the angle of the elevation of the sun is 60° .

[x] The volumes of two right circular cylinders are the same. The ratio of their height is 1:2. Find the ratio of their radii.

Solution:

The volume of a right circular cylinder with radius r and height h is $V = \pi r^2 h$.

It is given that the ratio of the heights of two circular cylinders is 1:2 that is $h_1 / h_2 = 1 / 2$

$$V_1 = V_2$$

$$\Rightarrow \pi r_1^2 h_1 = \pi r_2^2 h_2$$

$$\Rightarrow r_1^2 / r_2^2 = h_2 / h_1$$

$$\Rightarrow r_1^2 / r_2^2 = 1 / [h_1 / h_2]$$

$$\Rightarrow r_1^2 / r_2^2 = 1 / (1 / 2)$$

$$\Rightarrow (r_1 / r_2)^2 = 2$$

$$\Rightarrow r_1 / r_2 = \sqrt{2}$$

Hence, the ratio of their radius is $\sqrt{2}:1$.

[xi] The volume of a solid hemisphere is 144π cubic cm, then find the diameter of the sphere.

Solution:

The volume of the hemisphere = $2\pi r^3 / 3$

$$144\pi = 2 * (22 / 7) * r^3 / 3$$

$$144 * 3 = 2 * r^3$$

$$216 = r^3$$

$$r = 6\text{cm}$$

$$d = 2 * r = 2 * 6 = 12\text{cm}$$

[xii] The mean of a frequency distribution is 8.1 if $\sum f_i x_i = 132 + 5K$ and $\sum f_i = 20$ then what is the value of K ?

Solution:

$$\text{Mean} = \sum f_i x_i / \sum f_i$$

$$\text{Mean} = 8.1$$

$$\sum f_i x_i = 132 + 5K$$

$$\sum f_i = 20$$

$$8.1 = (132 + 5k) / 20$$

$$8.1 \times 20 = 132 + 5k$$

$$162 = 132 + 5k$$

$$162 - 132 = 5k$$

$$30 = 5k$$

$$k = 6$$

Question 5: Answer any one question :

[5 x 1 = 5]

(a) Aminur has taken a loan of Rs. 64, 000 from a bank. If the rate of interest is 2.5 paise per rupee per annum, calculate the compound interest payable after 2 years.

(b) A, B and C start a business with the capital of Rs. 6,000, Rs 8,000 and Rs. 9, 000, respectively. After a few months, A invests Rs 3, 000 more in the business. At the end of the year, they gained Rs 30,000 and C got Rs. 10,800 as a share of profit. When did A invest Rs. 3,000 more?

Solution:

$$[a] P = \text{Rs. } 64000$$

$$r = 2.5 \text{ paise per rupee per annum (given)}$$

$$= 0.025 \text{ rupee per rupee per annum}$$

$$= 0.025 \times 100 \text{ rupee per hundred rupee per annum}$$

$$= 0.025 \times 100 \text{ per cent per annum}$$

$$= 2.5 \text{ percent per annum}$$

$$t = 2 \text{ years}$$

$$C.I. = 64000 [(1 + 2.5 / 100)^2]$$

$$= 64000 [(1.025)^2]$$

$$= 64000 \times 1.050625$$

$$= 67240$$

$$= \text{Rs. } 67240$$

$$CI = 67240 - 64000 = \text{Rs. } 3240$$

[b] A invests Rs 3, 000 more in the business.

$$A = 6000 + 3000 = 9000$$

$$= 6000 * x + 9000 (12 - x)$$

$$= 6000x + 108000 - 9000x$$

$$= 108000 - 3000x$$

$$= 3000 (36 - x)$$

B invested Rs. 8000

$$= (8000 * 12)$$

$$= \text{Rs. } 96000$$

C invested Rs. 9000

$$= (9000 * 12)$$

$$= \text{Rs. } 108000$$

Ratio of A, B and C together

$$= 3000 (36 - x) : 96000 : 108000$$

$$= (36 - x) : 32 : 36$$

Gain = Rs. 30000

$$C = 30000 * [36 / (36 - x) + 32 + 36]$$

$$= 30000 * [36 / 104 - x]$$

$$30000 * [36 / 104 - x] = 10800$$

$$36 / 104 - x = 10800 / 30000$$

$$936 - 9x = 900$$

$$-9x = -36$$

$$x = 4$$

Question 6: Answer any one question:

[3 x 1 = 3]

(a) Solve: $\{[x + 4] / [x - 4]\}^2 - 5 [x + 4 / x - 4] + 6 = 0$, ($x \neq 4$)

(b) The digit in the unit's place of a two-digit number is 6 more than that at the ten's place. The product of the digits is 12 less than the number. Find the possible values of the digit in the unit place.

Solution:

$$[a] \{[x + 4] / [x - 4]\}^2 - 5 [x + 4 / x - 4] + 6 = 0,$$

$$\text{Take } [x + 4 / x - 4] = a$$

$$a^2 - 5a + 6 = 0$$

$$a^2 - 3a - 2a + 6 = 0$$

$$a (a - 3) - 2 (a - 3) = 0$$

$$(a - 3)(a - 2) = 0$$

$$a = 3, 2$$

$$a = 3$$

$$x + 4 / x - 4 = 3$$

$$x + 4 = 3(x - 4)$$

$$x + 4 = 3x - 12$$

$$12 + 4 = 3x - x$$

$$16 = 2x$$

$$x = 8$$

$$a = 2$$

$$x + 4 / x - 4 = 2$$

$$x + 4 = 2(x - 4)$$

$$x + 4 = 2x - 8$$

$$8 + 4 = 2x - x$$

$$12 = x$$

$$[b] \ x(x + 6) = 10x + x + 6 - 12$$

$$x^2 + 6x = 11x - 6$$

$$x^2 - 5x + 6 = 0$$

$$(x - 3)(x - 2) = 0$$

$$x = 3, 2$$

$$3 + 6 = 9$$

$$2 + 6 = 8$$

Question 7: Answer any one question:

[3 x 1 = 3]

(a) Find the simplest value of $\sqrt{7}(\sqrt{5} - \sqrt{2}) - \sqrt{5}(\sqrt{7} - \sqrt{2}) + 2\sqrt{2} / \sqrt{5} + \sqrt{7}$.

(b) If $x \propto y$ and $y \propto z$, then prove that: $(x^2 + y^2 + z^2) \propto (xy + yz + xz)$

Solution:

$$[a] \ \sqrt{7}(\sqrt{5} - \sqrt{2}) - \sqrt{5}(\sqrt{7} - \sqrt{2}) + 2\sqrt{2} / \sqrt{5} + \sqrt{7}$$

$$= \sqrt{35} - \sqrt{14} - \sqrt{35} + \sqrt{10} + [2\sqrt{2}(\sqrt{7} - \sqrt{5}) / (\sqrt{5} + \sqrt{7})(\sqrt{7} - \sqrt{5})]$$

$$\begin{aligned}
&= \sqrt{35} - \sqrt{14} - \sqrt{35} + \sqrt{10} + 2\sqrt{2} (\sqrt{7} - \sqrt{5}) / ((\sqrt{7})^2 - (\sqrt{5})^2) \\
&= \sqrt{35} - \sqrt{14} - \sqrt{35} + \sqrt{10} + 2\sqrt{2} (\sqrt{7} - \sqrt{5}) / 2 \\
&= \sqrt{35} - \sqrt{14} - \sqrt{35} + \sqrt{10} + \sqrt{14} - \sqrt{10} \\
&= 0
\end{aligned}$$

$$[b] \mathbf{x} \propto \mathbf{y}$$

$$x = k_1 y$$

$$\mathbf{y} \propto \mathbf{z}$$

$$y = k_2 z$$

$$\text{Hence, } x = k_1 k_2 z$$

$$(x^2 + y^2 + z^2) \propto (\mathbf{xy} + \mathbf{yz} + \mathbf{xz})$$

$$= (k_1 k_2 z)^2 + (k_2 z)^2 + z^2 / [k_1 y * k_2 z + k_2 z * z + k_1 y * z]$$

$$= z^2 (k_1^2 k_2^2 + k_2^2 + 1) / [k_1 k_2^2 z^2 + k_2 z^2 + k_1 k_2 z^2]$$

$$= z^2 (k_1^2 k_2^2 + k_2^2 + 1) / z^2 [k_1 k_2^2 + k_2 + k_1 k_2]$$

$$= (k_1^2 k_2^2 + k_2^2 + 1) / [k_1 k_2^2 + k_2 + k_1 k_2]$$

$$\text{So, } (x^2 + y^2 + z^2) \propto (\mathbf{xy} + \mathbf{yz} + \mathbf{xz})$$

Question 8: Answer any one question:

[3 x 1 = 3]

[a] If $[a + b - c] / [a + b] = [b + c - a] / [b + c] = [c + a - b] / [c + a]$ and $a + b + c \neq 0$ then prove that $a = b = c$.

[b] If $x:a, y:b, z:c$ that show that $(a^2 + b^2 + c^2)(x^2 + y^2 + z^2) = (ax + by + cz)^2$.

Solution:

$$[a] [a + b - c] / [a + b] = [b + c - a] / [b + c] = [c + a - b] / [c + a]$$

$$[a + b] / [a + b] - [c] / [a + b] = [b + c] / [b + c] - [a] / [b + c] = [c + a] / [c + a] - [b] / [c + a]$$

$$1 - [c] / [a + b] = 1 - [a] / [b + c] = 1 - [b] / [c + a]$$

$$[c] / [a + b] = [a] / [b + c] = [b] / [c + a]$$

$$a + b / c = b + c / a = c + a / b$$

$$a + b + c / c = b + c + a / a = c + a + b / b$$

$$1 / c = 1 / a = 1 / b$$

$$c = a = b \text{ or } a = b = c$$

[b] $x:a, y:b, z:c$

$x/a = y/b = z/c = k$ [Say]

Let $x = ka, y = kb, z = kc$

$$\text{LHS} = (a^2 + b^2 + c^2)(x^2 + y^2 + z^2)$$

$$= (a^2 + b^2 + c^2)(k^2a^2 + k^2b^2 + k^2c^2)$$

$$= k^2(a^2 + b^2 + c^2)(a^2 + b^2 + c^2)$$

$$= k^2(a^2 + b^2 + c^2)^2$$

$$\text{RHS} = (ax + by + cz)^2$$

$$= (a * ka + b * kb + c * kc)^2$$

$$= (k^2a^2 + k^2b^2 + k^2c^2)^2$$

$$= k^2(a^2 + b^2 + c^2)^2$$

Question 9: Answer any one question:


[5 x 1 = 5]

(a) Prove that, if a perpendicular is drawn on the hypotenuse from the right angular point of a right-angled triangle, two triangles so formed on the two sides of the perpendicular are each similar to the original triangle and also similar to each other.

(b) Prove that the tangent and the radius through the point of contact of a circle are perpendicular to each other.

Solution:

[a]



Given a right angle triangle, right-angled at A.

AD is the perpendicular drawn to the hypotenuse BC from vertex A.

To Prove:

(i) $\triangle BDA \sim \triangle BAC$

(ii) $\triangle ADC \sim \triangle BAC$

(iii) $\triangle BDA \sim \triangle ADC$

Proof:

In $\triangle BDA$ and $\triangle BAC$:

$$\angle ADB = \angle A = 90^\circ$$

$$\angle B = \angle B \quad [\text{common}]$$

Therefore, by using AA similar condition,

$\triangle BDA \sim \triangle BAC \quad \dots(i)$

Now, in $\triangle ADC$ and $\triangle BAC$,

$$\angle ADC = \angle A = 90^\circ$$


$$\angle C = \angle C \quad [\text{common}]$$

Therefore, by using AA similar condition,

$\triangle ADC \sim \triangle BAC \quad \dots(ii)$

Comparing (i) and (ii), $\triangle BDA \sim \triangle ADC$.

[b]



Given: A circle C (0, r) and a tangent l at point A.

To prove: $OA \perp l$

Construction: Take a point B, other than A, on the tangent l. Join OB. Suppose OB meets the circle in C.

Proof: We know that, among all line segments joining the point O to a point on l, the perpendicular is shortest to l.

$$OA = OC \quad (\text{Radius of the same circle})$$

$$\text{Now, } OB = OC + BC.$$

$$\therefore OB > OC$$

$$\Rightarrow OB > OA$$

$$\Rightarrow OA < OB$$

B is an arbitrary point on the tangent l. Thus, OA is shorter than any other line segment joining O to any point on l.

Here, $OA \perp l$.

Question 10: Answer any one question:

[3 x 1 = 3]

(a) In triangle ABC, AD is perpendicular on BC and $AD^2 = BD \cdot DC$, prove that $\angle BAC$ is a right angle.

(b) A straight line intersects one of the two concentric circles at the points A and B and another at the points C and D. Prove that $AC = BD$.

Solution:

[a] Given: In triangle ABC, AD is perpendicular to BC and $AD^2 = BD \cdot DC$

To prove: $\angle BAC = 90^\circ$

Proof: In right triangles $\triangle ADB$ and $\triangle ADC$, Pythagoras theorem should be applied,

$$AB^2 = AD^2 + BD^2 \text{ ----- (1)}$$

$$AC^2 = AD^2 + DC^2 \text{ ----- (2)}$$


$$\begin{aligned} AB^2 + AC^2 &= 2AD^2 + BD^2 + DC^2 \\ &= 2BD \cdot CD + BD^2 + CD^2 \quad [\because \text{given } AD^2 = BD \cdot CD] \\ &= (BD + CD)^2 = BC^2 \end{aligned}$$

Thus in triangle ABC, $AB^2 + AC^2 = BC^2$

Hence triangle ABC is a right triangle right angled at A.

$$\angle BAC = 90^\circ$$

[b]



Given: O is the centre and a straight line intersects one of the two concentric circles at the points A and B and other at the points C and D.

To prove $AC = BD$

Construction: OM is drawn perpendicular to AB

Proof:

$CM = DM$ [perpendicular drawn from the centre of the circle to the chord bisects the chord]

$AM - CM = BM - DM$

$AC = BD$

Question 11: Answer any one question:

[5 x 1 = 5]

(a) Construct two circles of radii 4 cm and 2 cm and the distance between their centres is 7 cm. Construct a direct common tangent of the circles. (only traces of construction are required).

(b) Construct a triangle whose two sides are 9 cm and 7 cm and the angle between them is 60° . Construct the incircle of the triangle. (only traces of construction are required).

Solution:

[a]

Solution:

$$[a] \text{ Arc length} = \theta / 360 * 2\pi r$$

Given that an arc of length 220cm of a circle makes an angle 60° at the centre,

$$220 = \{60\} / \{360\} * 2 * \{22\} / \{7\} r$$

$$220 = (1 / 6) * (44 / 7) * r$$

$$220 / 1.048 = r$$

$$210 = r$$

Hence the radius of the circle is 210 cm.

$$[b] \cos^2 \theta - \sin^2 \theta = 1 / 2$$

$$[\cos^2 \theta - \sin^2 \theta + 1] / [\cos^2 \theta - \sin^2 \theta - \sin^2 \theta - \cos^2 \theta] = [1 + 2] / [1 - 2]$$

$$[\cos^2 \theta - \sin^2 \theta + \sin^2 \theta + \cos^2 \theta] / [\cos^2 \theta - \sin^2 \theta - \sin^2 \theta - \cos^2 \theta] = 3 / -1$$

$$2\cos^2 \theta / -2 \sin^2 \theta = 3 / -1$$

$$\sin^2 \theta / \cos^2 \theta = 1 / 3$$

$$\tan^2 \theta = 1 / 3$$

$$[c] \sec 17^\circ / \operatorname{cosec} 73^\circ + \tan 68^\circ / \cot 22^\circ + \cos^2 44^\circ + \cos^2 46^\circ$$

$$= [\sec 17^\circ / \operatorname{cosec} (90^\circ - 73^\circ)] + [(\tan 90^\circ - 22^\circ) / \cot 22^\circ] + \cos^2 (90^\circ - 44^\circ) + \cos^2 46^\circ$$

$$= [\sec 17^\circ / \sec 17^\circ] + [\cot 22^\circ / \cot 22^\circ] + [\sin^2 46^\circ + \cos^2 46^\circ]$$

$$= 1 + 1 + 1$$

$$= 3$$

Question 13: Answer any one question:


[5 x 1 = 5]

(a) The length of the shadow of a post becomes 3 meters smaller when the angle of elevation of the Sun increases from 45° to 60° . Find the height of the post.

(b) A man standing on a railway bridge $5\sqrt{3}$ meters high, observes the engine of a train at an angle of depression 30° . But after 2 seconds, he observes the engine at an angle of depression 45° on the other side of the bridge. Find the speed of the train.

Solution:

[a]



$$\angle BCA = 45^\circ$$

$$\angle BDA = 60^\circ$$

$$CD = 3\text{m}$$

To find BD

In $\triangle ABD$,

$$AB / BD = \tan 60^\circ$$

$$AB / x = 1 / \sqrt{3}$$

$$AB = x * \sqrt{3}\text{m}$$

In $\triangle ABC$,

$$AB / BC = \tan 45^\circ$$

$$x\sqrt{3} / x + 3 = 1$$

$$x\sqrt{3} = x + 3$$

$$x\sqrt{3} - x = 3$$

$$x(\sqrt{3} - 1) = 3$$

$$x = 3 / (\sqrt{3} - 1)$$

By rationalising the denominator,

$$x = 3\sqrt{3} + 3 / 2$$

$$AB = \sqrt{3}(3\sqrt{3} + 3 / 2)$$

$$AB = 9 + 3\sqrt{3} / 2$$

$$= 9 + 5.193 / 2$$

$$= 7.098\text{m}$$

[b] Height(h) of the bridge = $AB = 5\sqrt{3}$ m

The angle of depression from one side = $30^\circ = \angle ACB$

The angle of depression from other side = $45^\circ = \angle ADB$

Required time (t) = 2 seconds

Speed of the train = ?

Now, from the $\triangle ADB$,

$$\tan 45^\circ = AB / BD$$

$$1 = 5\sqrt{3} / BD$$

$$BD = 5\sqrt{3} \text{ m}$$

Now, from the $\triangle ACB$,

$$\tan 30^\circ = AB / BC$$

$$1 / \sqrt{3} = 5\sqrt{3} / BC$$

$$BC = 15 \text{ m}$$

$$CD = BC + BD$$

$$CD = 15 + 5\sqrt{3} \text{ m}$$

Therefore, the distance covered by the train in 2 seconds is
 $= (15 + 5\sqrt{3}) \text{ m}$

$$\text{Speed of the train} = (15 + 5\sqrt{3}) / 2 = 11.83 \text{ m / s}$$

Question 14: Answer any TWO questions:

[4 x 2 = 8]

(a) Each side of a cube is decreased by 50%. Calculate the ratio of the volumes of the original and changed cube.

(b) The total surface area of a right circular cylindrical pot without a lid be 200 sq.cm. If the radius of the base is 7 cm find the quantity of water in litres contained in the pot. (1 litre = 1 cubic dm)

(c) A tank of length 21 dcm, breadth 11 dcm and 6 dcm deep is half-filled with water. If 100 solid iron balls of diameter 21 cm are completely immersed in the tank, then how much dcm of water level is raised?

Solution:

[a] Let length of the cube be x unit

$$V = (\text{Side})^3$$

$$V = (x)^3 \text{ unit}^3$$

Now, when the length of cube is reduced by 50%

$$\text{New length} = x - x * 50 / 100$$

$$= x - [x / 2]$$

$$= (2x - x) / 2$$

$$= x / 2 \text{ unit}$$

$$\text{New volume} = (\text{side})^3$$

$$= (x / 2)^3 \text{ unit}^3$$

$$\text{Ratio} = \text{Original cube volume} : \text{New cube volume}$$

$$= x^3 / (x^3 / 8)$$

$$= 8 : 1$$

$$[b] 2\pi rh = \pi r^2$$

$$\pi r(2h + r) = 2002$$

$$(22 / 7) * 7(2h + 7) = 2002$$

$$2h + 7 = 91$$

$$2h = 84$$

$$h = 42$$

$$V = \pi r^2 h$$

$$= (22 / 7) * 7^2 * (42)$$

$$= 22 * 7 * 42$$

$$= 6.468$$

$$= 6.468 / 1000$$

$$= 6.468 \text{ dcm}$$

$$[c] \text{ Length} = 21 \text{ dcm}$$

$$\text{Breadth} = 11 \text{ dcm}$$

$$6 \text{ dcm deep}$$

$$\text{Water level raised} = x \text{ dcm}$$

$$\text{Volume of the tank} = (21 * 11 * x) \text{ cm}^3$$

$$d = 21 / 2 \text{ cm}$$

$$= 21 / 20 \text{ d cm}$$

100 iron balls immersed in the tank.

$$= 100 * (4 / 3) * (22 / 7) * (21 / 20)^3 \text{ dcm}^3$$

$$(21 * 11 * x) = 100 * (4 / 3) * (22 / 7) * (21 / 20)^3$$

$$231x = 485.1$$

$$x = 231 / 485.1$$

$$x = 2.1 \text{ dcm}$$

Question 15: Answer any two questions:

[4 x 2 = 8]

(a) Find the mode from the following frequency distribution table of ages of examinees of an entrance examination:

Age (in years)	16 - 18	18 - 20	20 - 22	22 - 24	24 - 26
Number of examinees	45	75	38	22	20

(b) Find the median of the given data:

Class Interval	1 - 5	6 - 10	11 - 15	16 - 20	21 - 25	26 - 30	31 - 35
Frequency	2	3	6	7	5	4	3

(c) From the frequency distribution table given below, draw less than ogive:

Marks obtained	50 - 60	60 - 70	70 - 80	80 - 90	90 - 100
Frequency	4	8	12	6	10

Solution:

[a] The most frequently occurring frequency = 75

The modal class = 18 - 20

$$\text{Mode} = Z = L_1 + (F_1 - F_0) / (2F_1 - F_0 - F_2) * i$$

$$= 18 + [75 - 45] / [2 * 75 - 45 - 38] * 2$$

$$= 18 + [30 / 67] * 2$$

$$= 18.9$$

[b]

Class Interval	1 - 5	6 - 10	11 - 15	16 - 20	21 - 25	26 - 30	31 - 35
Class Interval	0.5 - 5.5	5.5 - 10.5	10.5 - 15.5	15.5 - 20.5	20.5 - 25.5	25.5 - 30.5	30.5 - 35.5
Frequency	2	3	6	7	5	4	3
CF	2	5	11	18	23	27	30

$$n = 30$$

$$\text{Median} = n / 2 = 30 / 2 = 15$$

$$\text{Median class} = 15.5 - 20.5$$

$$\text{Median} = m = l + [(n / 2 - CF) / f] * h$$

$$= 15.5 + [(30 / 2) - 11] / 7 * 5$$

$$= 15.5 + 2.86$$

$$= 18.36$$

[c]

Marks obtained	50 - 60	60 - 70	70 - 80	80 - 90	90 - 100
Frequency	4	8	12	6	10
CF	4	12	24	30	40

