

# Differential Equations

## Multiple Choice Questions

Choose and write the correct option in the following questions.

- The degree of the differential equation  $x^2 \frac{d^2 y}{dx^2} = \left(x \frac{dy}{dx} - y\right)^3$  is [CBSE 2020 (65/3/1)]  
(a) 1 (b) 2 (c) 3 (d) 6
- The degree of the differential equation  $\frac{d^2 y}{dx^2} + 3\left(\frac{dy}{dx}\right)^2 = x^2 \log\left(\frac{d^2 y}{dx^2}\right)$  is [NCERT Exemplar]  
(a) 1 (b) 2 (c) 3 (d) Not defined
- The order and degree of differential equation  $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^2 = \frac{d^2 y}{dx^2}$  respectively, are  
(a) 1, 2 (b) 2, 2 (c) 2, 1 (d) 4, 2
- The solution of the differential equation  $2x \cdot \frac{dy}{dx} - y = 3$  represents a family of  
(a) straight lines (b) circles (c) parabolas (d) ellipses
- The integrating factor of the differential equation  $\frac{dy}{dx}(x \log x) + y = 2 \log x$  is [NCERT Exemplar]  
(a)  $e^x$  (b)  $\log x$  (c)  $\log(\log x)$  (d)  $x$
- A solution of the differential equation  $\left(\frac{dy}{dx}\right)^2 - x \frac{dy}{dx} + y = 0$  is [NCERT Exemplar]  
(a)  $y = 2$  (b)  $y = 2x$  (c)  $y = 2x - 4$  (d)  $y = 2x^2 - 4$
- Which of the following is not a homogeneous function of  $x$  and  $y$ ?  
(a)  $x^2 + 2xy$  (b)  $2x - y$  (c)  $\cos^2\left(\frac{y}{x}\right) + \frac{y}{x}$  (d)  $\sin x - \cos y$
- The solution of the differential equation  $\frac{dx}{x} + \frac{dy}{y} = 0$  is [CBSE 2023 (65/3/2)]  
(a)  $\frac{1}{x} + \frac{1}{y} = C$  (b)  $\log x - \log y = C$  (c)  $xy = C$  (d)  $x + y = C$
- The solution of the differential equation  $x \frac{dy}{dx} + 2y = x^2$  is  
(a)  $y = \frac{x^2 + C}{4x^2}$  (b)  $y = \frac{x^2}{4} + C$  (c)  $y = \frac{x^4 + C}{x^2}$  (d)  $y = \frac{x^4 + C}{4x^2}$
- Degree of the differential equation  $\sin x + \cos\left(\frac{dy}{dx}\right) = y^2$  is [CBSE 2023 (65/2/1)]  
(a) 2 (b) 1 (c) not defined (d) 0
- The degree of the differential equation  $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2} = \frac{d^2 y}{dx^2}$  is  
(a) 4 (b)  $\frac{3}{2}$  (c) Not defined (d) 2

12. The order and degree of a differential equation  $\frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^{\frac{1}{4}} + x^{\frac{1}{5}} = 0$ , respectively, are  
 (a) 2 and not defined (b) 2 and 2  
 (c) 2 and 3 (d) 3 and 3
13. The solution of the differential equation  $\frac{dy}{dx} + \frac{2y}{x} = 0$  with  $y(1) = 1$  is given by  
 (a)  $y = \frac{1}{x^2}$  (b)  $x = \frac{1}{y^2}$  (c)  $x = \frac{1}{y}$  (d)  $y = \frac{1}{x}$
14. The general solution of  $\frac{dy}{dx} = e^{x-y} + x^2e^{-y}$  is  
 (a)  $y = e^{x-y} - x^2e^{-y} + C$  (b)  $e^y - e^x = \frac{x^3}{3} + C$   
 (c)  $e^x + e^y = \frac{x^3}{3} + C$  (d)  $e^x - e^y = \frac{x^3}{3} + C$
15. The integrating factor of the differential equation  $(1 - y^2) \frac{dx}{dy} + yx = ay$ ,  $(-1 < y < 1)$  is  
 [CBSE 2023 (65/2/1)]  
 (a)  $\frac{1}{y^2 - 1}$  (b)  $\frac{1}{\sqrt{y^2 - 1}}$  (c)  $\frac{1}{1 - y^2}$  (d)  $\frac{1}{\sqrt{1 - y^2}}$
16. Solution of  $\frac{dy}{dx} - y = 1$ ,  $y(0) = 1$  is given by  
 (a)  $xy = -e^x$  (b)  $xy = -e^{-x}$  (c)  $xy = -1$  (d)  $y = 2e^x - 1$
17. The number of solution of  $\frac{dy}{dx} = \frac{y+1}{x-1}$  when  $y(1) = 2$  is  
 (a) none (b) one (c) two (d) infinite
18. Integrating factor of the differential equation  $(1 - x^2) \frac{dy}{dx} - xy = 1$  is  
 (a)  $-x$  (b)  $\frac{x}{1+x^2}$  (c)  $\sqrt{1-x^2}$  (d)  $\frac{1}{2} \log(1-x^2)$
19. What is the product of the order and degree of the differential equation  $\frac{d^2y}{dx^2} \sin y + \left(\frac{dy}{dx}\right)^3 = \sqrt{y}$ ?  
 [CBSE 2023 (65/3/2)]  
 (a) 3 (b) 2 (c) 6 (d) not defined
20. The sum of the order and the degree of the differential equation  $\frac{d^2y}{dx^2} + \left[\frac{dy}{dx}\right]^3 = \sin y$  is:  
 [CBSE 2023 (65/1/1)]  
 (a) 5 (b) 2 (c) 3 (d) 4
21. The general solution of the differential equation  $x dy - (1 + x^2) dx = dx$  is: [CBSE 2023 (65/1/1)]  
 (a)  $y = 2x + \frac{x^3}{3} + C$  (b)  $y = 2 \log x + \frac{x^3}{3} + C$   
 (c)  $y = \frac{x^2}{2} + C$  (d)  $y = 2 \log x + \frac{x^2}{2} + C$

## Answers

- |         |         |         |         |         |         |
|---------|---------|---------|---------|---------|---------|
| 1. (a)  | 2. (d)  | 3. (c)  | 4. (c)  | 5. (b)  | 6. (c)  |
| 7. (d)  | 8. (c)  | 9. (d)  | 10. (c) | 11. (d) | 12. (a) |
| 13. (a) | 14. (b) | 15. (d) | 16. (d) | 17. (b) | 18. (c) |
| 19. (b) | 20. (c) | 21. (d) |         |         |         |

## Solutions of Selected Multiple Choice Questions

2. The given differential equation is not a polynomial equation in terms of its derivatives, so its degree is not defined.

∴ Option (d) is correct.

8. Given differential equation be

$$\frac{dx}{x} + \frac{dy}{y} = 0 \Rightarrow \frac{dx}{x} = -\frac{dy}{y}$$

$$\Rightarrow \int \frac{dx}{x} = -\int \frac{dy}{y} \quad (\text{on integrating both sides})$$

$$\Rightarrow \log x = -\log y + \log C$$

$$\Rightarrow \log x + \log y = \log C \Rightarrow \log xy = \log C$$

$$\Rightarrow xy = C$$

∴ Option (c) is correct.

9. Given differential equation is

$$x \frac{dy}{dx} + 2y = x^2 \Rightarrow \frac{dy}{dx} + \frac{2}{x}y = x$$

It is of the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \frac{2}{x}$ ,  $Q = x$ .

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \frac{2}{x} dx} = e^{2 \log x} = e^{\log x^2} = x^2$$

$$\therefore \text{Solution is } y \times x^2 = \int x \times x^2 dx + C_1 = \int x^3 dx + C_1$$

$$\Rightarrow yx^2 = \frac{x^4}{4} + C_1$$

$$\Rightarrow yx^2 = \frac{x^4 + 4C_1}{4} = \frac{x^4 + C}{4} \text{ where } C = 4C_1$$

$$\Rightarrow y = \frac{x^4 + C}{4x^2}$$

∴ Option (d) is correct.

10. Given differential equation be

$$\sin x + \cos\left(\frac{dy}{dx}\right) = y^2, \text{ which is not a polynomial in } \frac{dy}{dx}.$$

Thus, degree is not defined.

∴ Option (c) is correct.

14. Given,  $\frac{dy}{dx} = e^x - y + x^2 e^{-y}$

$$\Rightarrow \frac{dy}{dx} = e^{-y}(e^x + x^2) \Rightarrow \frac{dy}{e^{-y}} = (e^x + x^2) dx$$

$$\Rightarrow e^y dy = (e^x + x^2) dx = e^x dx + x^2 dx$$

Integrating, we get

$$\int e^y dy = \int e^x dx + \int x^2 dx$$

$$\Rightarrow e^y = e^x + \frac{x^3}{3} + C \quad \Rightarrow e^y - e^x = \frac{x^3}{3} + C$$

$\therefore$  Option (b) is correct.

15. Given differential equation be

$$(1-y^2) \frac{dx}{dy} + yx = ay, \quad -1 < y < 1$$

$$\Rightarrow \frac{dx}{dy} + \frac{y}{1-y^2} \cdot x = \frac{ay}{1-y^2}$$

It is a linear differential equation of the form

$$\frac{dx}{dy} + P \cdot x = Q, \text{ where } P, Q \text{ be the function of } y \text{ or constant.}$$

$$\therefore P = \frac{y}{1-y^2} \text{ and } Q = \frac{ay}{1-y^2}$$

$$\therefore IF = e^{\int P dy} = e^{\int \frac{y}{1-y^2} dy} = e^{-\frac{1}{2} \int \frac{2y}{1-y^2} dy}$$

$$= e^{-\frac{1}{2} \log|1-y^2|} = e^{\log|1-y^2|^{-\frac{1}{2}}}$$

$$= (1-y^2)^{-\frac{1}{2}} = \frac{1}{\sqrt{1-y^2}}$$

$\therefore$  Option (d) is correct.

16. Given that,  $\frac{dy}{dx} - y = 1$

$$\Rightarrow \frac{dy}{dx} = 1 + y \quad \Rightarrow \frac{dy}{1+y} = dx$$

On integrating both sides, we get  $\log(1+y) = x + C$

When  $x = 0$  and  $y = 1$ , then

$$\log 2 = 0 + C \Rightarrow C = \log 2$$

The required solution is  $\log(1+y) = x + \log 2$

$$\Rightarrow \log\left(\frac{1+y}{2}\right) = x \quad \Rightarrow \frac{1+y}{2} = e^x$$

$$\Rightarrow 1+y = 2e^x \quad \Rightarrow y = 2e^x - 1$$

$\therefore$  Option (d) is correct.

18. Given that,  $(1-x^2) \frac{dy}{dx} - xy = 1$

$$\Rightarrow \frac{dy}{dx} - \frac{x}{1-x^2} y = \frac{1}{1-x^2}, \text{ which is a linear differential equation.}$$

$$\therefore IF = e^{\int \frac{-x}{1-x^2} dx}, \text{ Let } 1-x^2 = t \Rightarrow -2x dx = dt \Rightarrow -x dx = \frac{dt}{2}$$

$$\Rightarrow e^{\frac{1}{2} \int \frac{dt}{t}} = e^{\frac{1}{2} \log t} = e^{\frac{1}{2} \log(1-x^2)} = \sqrt{1-x^2}$$

$\therefore$  Option (c) is correct.

19. Given differential equation be

$$\frac{d^2 y}{dx^2} \sin y + \left(\frac{dy}{dx}\right)^3 \cos y = \sqrt{y}$$

Order = 2, degree = 1

⇒ Product of order and degree =  $2 \times 1 = 2$

∴ Option (b) is correct.

20. Given differential equation be

$$\frac{d^2 y}{dx^2} + \left(\frac{dy}{dx}\right)^3 = \sin y$$

Its order = 2 and degree = 1

⇒ Sum of the order and the degree =  $2 + 1 = 3$

∴ Option (c) is correct.

21. Given differential equation,

$$x \, dy - (1 + x^2) dx = dx \quad \Rightarrow \quad x \, dy = (1 + x^2 + 1) dx = (2 + x^2) dx$$

$$\Rightarrow \quad dy = \frac{2 + x^2}{x} dx$$

On integrating both sides, we have

$$\Rightarrow \quad \int dy = \int \frac{2 + x^2}{x} dx = 2 \int \frac{1}{x} dx + \int x \, dx \quad \Rightarrow \quad y = 2 \log x + \frac{x^2}{2} + C$$

∴ Option (d) is correct.

## Assertion-Reason Questions

The following questions consist of two statements—Assertion(A) and Reason(R). Answer these questions selecting the appropriate option given below:

- (a) Both A and R are true and R is the correct explanation for A.
- (b) Both A and R are true but R is not the correct explanation for A.
- (c) A is true but R is false.
- (d) A is false but R is true.

1. **Assertion (A) :** The degree of the differential equation  $\frac{d^2 y}{dx^2} = 1 + \sqrt{\frac{dy}{dx}}$  is 2.

**Reason (R) :** The degree of a differential equation is the degree of the highest order derivative occurring in the equation, when differential co-efficients are made free from radicals, fractions and it is written as a polynomial in differential coefficient.

2. **Assertion (A) :** Solution of the differential equation  $(1 + x^2) \frac{dy}{dx} + y = \tan^{-1} x$  is

$$y e^{\tan^{-1} x} = (\tan^{-1} x - 1) e^{\tan^{-1} x} + C$$

**Reason (R) :** The differential equation of the form  $\frac{dy}{dx} + Py = Q$ , where P, Q be the functions of x or constant, is a linear type differential equation.

3. **Assertion (A)** : The integrating factor of differential equation  $\frac{dx}{dy} + (\tan y) \cdot x = \sec^2 y$  is  $\sec y$ .

**Reason (R)** : Linear differential equation of the form  $\frac{dx}{dy} + Px = Q$ , where  $P, Q = f(y)$  or constant has integrating factor,  $IF = e^{\int P dy}$

4. **Assertion (A)** : General solution of differential equation  $\frac{dy}{dx} = \frac{y}{x}$  is  $y = Cx$ .

**Reason (R)** : The differential equation  $\frac{d^2y}{dx^2} + y = 0$  has order 2.

5. **Assertion (A)** : Solution of the differential equation  $e^{dy/dx} = x^2$  is  $y = 2(x \log x - x) + C$ .

**Reason (R)** : The integrating factor of the differential equation  $(1 + x^2) \frac{dy}{dx} + y = \tan^{-1} x$  is  $e^{\tan^{-1} x}$ .

## Answers

1. (a)

2. (b)

3. (a)

4. (b)

5. (b)

## Solutions of Assertion-Reason Questions

1. We have,  $\frac{d^2y}{dx^2} = 1 + \sqrt{\frac{dy}{dx}}$

$$\Rightarrow \left( \frac{d^2y}{dx^2} - 1 \right)^2 = \left( \sqrt{\frac{dy}{dx}} \right)^2 \quad [\text{Squaring both sides}]$$

$$\Rightarrow \left( \frac{d^2y}{dx^2} \right)^2 - 2 \frac{d^2y}{dx^2} + 1 = \frac{dy}{dx} \quad \Rightarrow \quad \left( \frac{d^2y}{dx^2} \right)^2 - 2 \frac{d^2y}{dx^2} - \frac{dy}{dx} + 1 = 0$$

$\therefore$  Degree = 2

Clearly, both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion (A).

$\therefore$  Option (a) is correct.

2.  $(1 + x^2) \frac{dy}{dx} + y = \tan^{-1} x \Rightarrow \frac{dy}{dx} + \frac{1}{1 + x^2} \cdot y = \frac{\tan^{-1} x}{1 + x^2}$

$$\therefore \quad IF = e^{\int \frac{1}{1+x^2} dx} = e^{\tan^{-1} x}$$

$$\text{Solution will be } y \times e^{\tan^{-1} x} = \int \frac{\tan^{-1} x}{1 + x^2} \times e^{\tan^{-1} x} dx \quad \dots (i)$$

$$\text{Let } e^{\tan^{-1} x} = t \Rightarrow \frac{e^{\tan^{-1} x}}{1 + x^2} dx = dt \text{ and } \log(e^{\tan^{-1} x}) = \log t \Rightarrow \tan^{-1} x = \log t$$

$$\text{From equation (i), } yx e^{\tan^{-1} x} = \int \log t \cdot dt = t \log t - t + C = t (\log t - 1) + C$$

$$y e^{\tan^{-1} x} = e^{\tan^{-1} x} (\log(\tan^{-1} x) - 1) + C$$

Clearly, both Assertion (A) and Reason (R) are true but Reason (R) is not the correct explanation of Assertion (A).

$\therefore$  Option (b) is correct.



$$3. \frac{dx}{dy} + (\tan y) \cdot x = \sec^2 y$$

$$\text{Here, IF} = e^{\int \tan y \, dy} = e^{\log \sec y} = \sec y$$

Clearly, both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion (A).

∴ Option (a) is correct.

4. We have

$$\frac{dy}{dx} = \frac{y}{x} \Rightarrow \frac{dy}{y} = \frac{dx}{x}$$

Integrating, we get

$$\log y = \log x + \log C$$

$$\Rightarrow \log y = \log (Cx) \Rightarrow y = Cx$$

So statement A is true.

Also statement R is true but R is not correct explanation of A.

∴ Option (b) is correct.

5. We have,  $e^{dy/dx} = x^2$

Taking logarithm both sides, we get

$$\log(e^{dy/dx}) = \log x^2 \Rightarrow \frac{dy}{dx} = \log x^2$$

$$\Rightarrow \frac{dy}{dx} = 2 \log x \Rightarrow dy = 2 \log x \, dx$$

Integrating, we get

$$\begin{aligned} y &= 2 \left[ \int \log x \times 1 \, dx \right] \\ &= 2 \left[ \log x \int dx - \int \left\{ \frac{d}{dx} (\log x) \int dx \right\} dx \right] = 2 \left[ \log x \times x - \int \frac{1}{x} \times x \, dx \right] \end{aligned}$$

$$\Rightarrow y = 2[x \log x - x] + C$$

$$\Rightarrow y = 2[x \log x - x] + C$$

$$\Rightarrow y = 2x(\log x - 1) + C$$

So statement A is correct.

For statement R,

$$(1+x^2) \frac{dy}{dx} + y = \tan^{-1} x$$

$$\Rightarrow \frac{dy}{dx} + \frac{1}{1+x^2} y = \frac{1}{1+x^2} \tan^{-1} x$$

It is of the form

$$\frac{dy}{dx} + Py = Q$$

$$\text{Where, } P = \frac{1}{1+x^2}, Q = \frac{1}{1+x^2} \tan^{-1} x$$

$$\therefore \text{IF} = e^{\int P \, dx} = e^{\int \frac{1}{1+x^2} \, dx} = e^{\tan^{-1} x}$$

So statement R is also correct, but R is not correct explanation of statement A.

∴ Option (b) is correct.

$$\Rightarrow x \frac{dv}{dx} = \frac{v^2 - 1}{2v} - v = \frac{v^2 - 1 - 2v^2}{2v} = -\frac{(1 + v^2)}{2v}$$

$$\Rightarrow \frac{2v}{1 + v^2} dv = -\frac{dx}{x}$$

Integrating, we get

$$\int \frac{2v}{1 + v^2} dv = -\int \frac{dx}{x}$$

$$\Rightarrow \log|1 + v^2| = -\log|x| + \log C$$

$$\Rightarrow \log|1 + v^2| + \log|x| = \log C$$

$$\Rightarrow \log|(1 + v^2)x| = \log C$$

$$\Rightarrow (1 + v^2)x = C$$

$$\Rightarrow \left\{ 1 + \left( \frac{y}{x} \right)^2 \right\} x = C \Rightarrow \left( \frac{x^2 + y^2}{x^2} \right) x = C$$

$$\Rightarrow x^2 + y^2 = Cx$$

2. Read the following passage and answer the following questions.

If an equation is of the form

$$\frac{dy}{dx} + Py = Q$$

Where  $P, Q$  are functions of  $x$  then such equation is known as linear differential equation. Its solution is given by

$$y \times \text{IF} = \int Q \times \text{IF} dx + C$$

Where  $\text{IF} = e^{\int P dx}$

Now suppose we have equation.  $\frac{dy}{dx} + \frac{y}{x} = x^2$

- (i) Write the value of  $P$ .
- (ii) Write the value of  $Q$ .
- (iii) (a) Find the general solution of given differential equation.

OR

- (iii) (b) If the value of  $Q$  replace by  $\sin x$  find the solution.

Sol. Given differential equation is  $\frac{dy}{dx} + \frac{y}{x} = x^2$

It is of the form  $\frac{dy}{dx} + Py = Q$

$$(i) \text{ Here } P = \frac{1}{x}$$

$$(ii) \text{ Here } Q = x^2$$

$$(iii) (a) \text{ IF} = e^{\int P dx} = e^{\int \frac{1}{x} dx} = e^{\log x} = x$$

Solution is given by

$$y \times x = \int x^2 \times x dx + C \Rightarrow yx = \int x^3 dx + C$$

$$\Rightarrow yx = \frac{x^4}{4} + C \Rightarrow y = \frac{x^3}{4} + \frac{C}{x}$$



OR

(iii) (b) If  $Q = \sin x$

From (iii) above IF =  $x$

Solution is given by

$$y \times x = \int x \sin x \, dx + C$$

$$yx = x \int \sin x \, dx - \int \left\{ \frac{d}{dx}(x) \int \sin x \, dx \right\} dx + C$$

$$yx = -x \cos x - \int (-\cos x) dx + C$$

$$yx = -x \cos x + \int \cos x \, dx + C$$

$$\Rightarrow yx = -x \cos x + \sin x + C$$

$$\Rightarrow y = -\cos x + \frac{\sin x}{x} + \frac{C}{x}$$

3. Read the following passage and answer the following questions.

Polio drops are delivered to 50K children in a district. The rate at which polio drops are given is directly proportional to the number of children who have not been administered the drops. By the end of 2<sup>nd</sup> week half the children have been given the polio drops. How many will have been given the drops by the end of 3<sup>rd</sup> week can be estimated using the solution to the differential equation  $\frac{dy}{dx} = k(50 - y)$  where  $x$  denotes the number of weeks and  $y$  the number of children who have been given the drops.

(i) (a) Find the solution of the differential equation  $\frac{dy}{dx} = k(50 - y)$ .

(b) Find the value of  $C$  in the particular solution given that  $y(0) = 0$  and  $k = 0.049$ .

(ii) Find the solution that may be used to find the number of children who have been given the polio drops.

Sol. (i) (a) We have,

$$\frac{dy}{dx} = k(50 - y)$$

$$\Rightarrow \int \frac{dy}{50 - y} = \int k dx \quad \Rightarrow \quad -\log |50 - y| = kx + C$$

(b) Given  $y(0) = 0$  and  $k = 0.049$

$$\therefore -\log |50 - y| = kx + C$$

$$\Rightarrow -\log |50 - 0| = 0.049 \times 0 + C$$

$$\Rightarrow -\log 50 = C \quad \Rightarrow \quad C = \log \frac{1}{50}$$

(ii) We have,

$$-\log |50 - y| = kx + \log \frac{1}{50} \quad [\text{from (i) (a), (b)}]$$

$$\Rightarrow -kx = \log |50 - y| + \log \frac{1}{50} \quad \Rightarrow -kx = \log \frac{50 - y}{50}$$

$$\Rightarrow e^{-kx} = \frac{50 - y}{50} = 1 - \frac{y}{50} \quad \Rightarrow \frac{y}{50} = 1 - e^{-kx} \Rightarrow y = 50(1 - e^{-kx})$$

This is the required solution to find the number of children who have been given the polio drops.

## CONCEPTUAL QUESTIONS

1. How many arbitrary constants are there in the particular solution of the differential equation

$$\frac{dy}{dx} = -4xy^2; y(0) = 1?$$

[CBSE Sample Paper 2021]

Sol. 0

2. For what value of  $n$  is the following a homogeneous differential equation?

$$\frac{dy}{dx} = \frac{x^3 - y^n}{x^2y + xy^2}$$

[CBSE Sample Paper 2021]

Sol. 3

3. Find the general solution of the differential equation  $e^{y-x} \frac{dy}{dx} = 1$ . [CBSE 2020 (65/2/1)]

Sol. Given differential equation is  $e^y dy = e^x dx$  ½

Integrating to get  $e^y = e^x + C$  ½

[CBSE Marking Scheme 2020 (65/2/1)]

4. Find the integrating factor of the differential equation

$$x \frac{dy}{dx} = 2x^2 + y$$

[CBSE 2020 (65/2/2)]

Sol. Integrating factor is  $e^{\int \frac{-1}{x} dx}$  or  $\begin{cases} \text{writing given equation as} \\ \frac{dy}{dx} - \frac{y}{x} = 2x \end{cases}$  ½

$= \frac{1}{x}$  ½

[CBSE Marking Scheme 2020 (65/2/2)]

5. Find the order and degree of differential equation:

$$\frac{d^4 y}{dx^4} + \sin\left(\frac{d^3 y}{dx^3}\right) = 0$$

[NCERT Exemplar]

Sol. Order is 4 but degree is not defined because given differential equation cannot be written in the form of polynomial in differential co-efficient.

6. Find the general solution of the differential equation  $e^{y-x} \frac{dy}{dx} = 1$ .

Sol.  $e^{y-x} \frac{dy}{dx} = 1 \Rightarrow \frac{e^y dy}{e^x dx} = 1$

$$\Rightarrow e^y dy = e^x dx$$

On integrating we have

$$\int e^y dx = \int e^x dx$$

$$\Rightarrow e^y = e^x + C \Rightarrow y = \log(e^x + C)$$

7. Write the sum of the order and degree of the following differential equation:

$$\frac{d}{dx} \left\{ \left( \frac{dy}{dx} \right)^3 \right\} = 0$$

[CBSE Allahabad 2015]

Sol. Given differential equation is

$$\frac{d}{dx} \left\{ \left( \frac{dy}{dx} \right)^3 \right\} = 0 \Rightarrow 3 \left( \frac{dy}{dx} \right)^2 \cdot \frac{d^2 y}{dx^2} = 0$$

i.e., order = 2, degree = 1

∴ Required sum = 2 + 1 = 3.

8. Solve the differential equation  $(y + 3x^2) \frac{dx}{dy} = x$ .

**Sol.**  $(y + 3x^2)dx = xdy$

$$\Rightarrow ydx + 3x^2dx = xdy$$

$$\Rightarrow 3x^2dx = xdy - ydx$$

$$\Rightarrow 3dx = \frac{xdy - ydx}{x^2} = d\left(\frac{y}{x}\right)$$

Integrating, we get

$$\Rightarrow 3x = \frac{y}{x} + C \quad \Rightarrow \quad 3x^2 = y + Cx$$

$$\Rightarrow y - 3x^2 + Cx = 0.$$

9. Solve the following differential equation:

$$\frac{dy}{dx} + y = \cos x - \sin x$$

**Sol.** Given differential equation,

$$\frac{dy}{dx} + y = \cos x - \sin x$$

It is a linear differential equation of the type

$$\frac{dy}{dx} + Py = Q, \text{ where } P, Q \text{ be the function of } x \text{ or constants.}$$

$$\therefore P = 1, Q = \cos x - \sin x$$

Now, integrating factor,  $IF = e^{\int P dx} = e^{\int 1 dx} = e^x$

$$\therefore \text{Solution be } y \times IF = \int Q \times IF dx$$

$$\Rightarrow y e^x = \int e^x (\cos x - \sin x) dx$$

$$\Rightarrow y e^x = e^x \cos x + C$$

$$\Rightarrow y = \cos x + C e^{-x}$$

## Very Short Answer Questions

1. Find the general solution of  $y^2 dx + (x^2 - xy + y^2) dy = 0$ .

[NCERT Exemplar]

**Sol.** Given, differential equation is  $y^2 dx + (x^2 - xy + y^2) dy = 0$ .

$$\Rightarrow y^2 dx = -(x^2 - xy + y^2) dy$$

$$\Rightarrow y^2 \frac{dx}{dy} = -(x^2 - xy + y^2)$$

$$\Rightarrow \frac{dx}{dy} = -\left(\frac{x^2}{y^2} - \frac{x}{y} + 1\right) \quad \dots(i)$$

Which is a homogeneous differential equation.

Put  $\frac{x}{y} = v$  or  $x = vy$

$$\Rightarrow \frac{dx}{dy} = v + y \frac{dv}{dy}$$

On substituting these values in equation (i), we get

$$v + y \frac{dv}{dy} = -[v^2 - v + 1]$$

$$\Rightarrow y \frac{dv}{dy} = -v^2 + v - 1 - v$$

$$\Rightarrow y \frac{dv}{dy} = -v^2 - 1 \Rightarrow \frac{dv}{v^2 + 1} = -\frac{dy}{y}$$

On integrating both sides, we get

$$\tan^{-1}(v) = -\log y + C$$

$$\Rightarrow \tan^{-1}\left(\frac{x}{y}\right) + \log y = C \quad \left[ \because v = \frac{x}{y} \right]$$

2. Solve the differential equation  $(y + 3x^2) \frac{dx}{dy} = x$ .

[CBSE 2019 (65/5/2)]

Sol.  $(y + 3x^2)dx = xdy \Rightarrow ydx + 3x^2dx = xdy$

$$\Rightarrow 3x^2dx = xdy - ydx$$

$$\Rightarrow 3dx = \frac{xdy - ydx}{x^2} = d\left(\frac{y}{x}\right)$$

Integrating, we get

$$\Rightarrow 3x = \frac{y}{x} + C \quad \Rightarrow 3x^2 = y + Cx$$

$$\Rightarrow y - 3x^2 + Cx = 0.$$

3. Write the integrating factor of the following differential equation:

$$(1 + y^2) + (2xy - \cot y) \frac{dy}{dx} = 0$$

[CBSE Allahabad 2015]

Sol.  $(1 + y^2) + (2xy - \cot y) \frac{dy}{dx} = 0$

$$\Rightarrow (2xy - \cot y) \frac{dy}{dx} = -(1 + y^2)$$

$$\Rightarrow \frac{dy}{dx} = -\frac{1 + y^2}{2xy - \cot y}$$

$$\Rightarrow \frac{dx}{dy} = -\frac{(2xy - \cot y)}{1 + y^2}$$

$$\Rightarrow \frac{dx}{dy} + \frac{2y}{1 + y^2} \cdot x = \frac{\cot y}{1 + y^2}$$

It is in the form  $\frac{dx}{dy} + Px = Q$ , where  $P$  and  $Q$  are function of  $y$ .

$$\Rightarrow \text{IF} = e^{\int P dy} = e^{\int \frac{2y}{1+y^2} dy} = e^{\log |1+y^2|} = 1 + y^2$$

Solve the following differential equations Q(4 – 6).

4.  $\frac{dy}{dx} = y \tan x$ ;  $y = 1$  when  $x = 0$ .

**Sol.** The given equation is  $\frac{dy}{dx} = y \tan x$

$$\Rightarrow \frac{1}{y} dy = \tan x \, dx$$

$$\Rightarrow \int \frac{1}{y} dy = \int \tan x \, dx$$

$$\Rightarrow \log |y| = -\log |\cos x| + \log C$$

$$\Rightarrow \log |y| + \log |\cos x| = \log C$$

$$\Rightarrow \log |y \cos x| = \log C \Rightarrow y \cos x = C$$

Putting  $y = 1$  and  $x = 0$

$$\text{We have, } 1 \cdot \cos(0) = C \Rightarrow C = 1$$

$$\therefore y \cos x = 1 \Rightarrow y = \frac{1}{\cos x} \Rightarrow y = \sec x$$

5.  $\cos\left(\frac{dy}{dx}\right) = a$  ( $-1 \leq a \leq 1$ );  $y = 1$  when  $x = 0$ .

**Sol.** The given equation is  $\cos\left(\frac{dy}{dx}\right) = a$ .

$$\Rightarrow \frac{dy}{dx} = \cos^{-1} a \Rightarrow dy = \cos^{-1} a \, dx$$

$$\Rightarrow \int dy = \int \cos^{-1} a \, dx$$

$$\Rightarrow \int dy = \cos^{-1} a \int dx \Rightarrow y = (\cos^{-1} a) x + C$$

Putting  $y = 1$  and  $x = 0$

We have,

$$1 = \cos^{-1}(a) \times 0 + C \Rightarrow C = 1$$

$$\therefore y = (\cos^{-1} a) x + 1 \Rightarrow \cos^{-1} a = \frac{y-1}{x}$$

$$\Rightarrow a = \cos\left(\frac{y-1}{x}\right).$$

6.  $\frac{dy}{dx} + \frac{1+y^2}{2y} = 0$ .

**Sol.** We have,

$$\frac{dy}{dx} + \frac{1+y^2}{2y} = 0 \Rightarrow \frac{dy}{dx} = -\frac{1+y^2}{2y} \Rightarrow \frac{2y}{1+y^2} dy = -dx$$

$$\Rightarrow \int \frac{2y}{1+y^2} dy = -\int dx \text{ putting } 1+y^2 = t \Rightarrow 2y \, dy = dt$$

$$\Rightarrow \int \frac{1}{t} dt = -\int dx + C$$

$$\Rightarrow \log t = -x + C$$

$$\Rightarrow \log |1+y^2| + x = C.$$

7. Solve the differential equation  $\cos x \frac{dy}{dx} = \cos 3x - \cos 2x$ .

Sol.  $\cos x \frac{dy}{dx} = \cos 3x - \cos 2x$

$$\Rightarrow \cos x \frac{dy}{dx} = (4 \cos^3 x - 3 \cos x) - (2 \cos^2 x - 1)$$

$$\Rightarrow \frac{dy}{dx} = 4 \cos^2 x - 3 - 2 \cos x + \frac{1}{\cos x}$$

$$\Rightarrow dy = \left( \frac{4(1 + \cos 2x)}{2} - 3 - 2 \cos x + \sec x \right) dx$$

$$\Rightarrow \int dy = \int (2 + 2 \cos 2x - 3 - 2 \cos x + \sec x) dx = \int (2 \cos 2x - 1 - 2 \cos x - \sec x) dx$$

$$\Rightarrow y = \frac{2 \sin 2x}{2} - x - 2 \sin x - \log |\sec x + \tan x| + C$$

$$\Rightarrow y = \sin 2x - x - 2 \sin x - \log |\sec x + \tan x| + C$$

8. Find the general solution of the differential equation

$$\log \left( \frac{dy}{dx} \right) = ax + by$$

[CBSE 2021-22 (Term-2)]

Sol.

$$\log \left( \frac{dy}{dx} \right) = ax + by$$

$$\Rightarrow \frac{dy}{dx} = e^{ax+by}$$

$$\Rightarrow \frac{dy}{dx} = e^{ax} \cdot e^{by} \quad [e^{a+b} = e^a \cdot e^b]$$

$$\Rightarrow \frac{dy}{e^{by}} = e^{ax} dx$$

$$\Rightarrow e^{-by} dy = e^{ax} dx$$

on integrating both sides.

$$\int e^{-by} dy = \int e^{ax} dx$$

$$\Rightarrow \frac{-e^{-by}}{b} = \frac{e^{ax}}{a} + C$$

$$\frac{e^{ax}}{a} + \frac{e^{-by}}{b} = C' \quad [C' = -C]$$

where  $C$  &  $C'$  are constants.

$$\text{Answer} \quad \frac{e^{ax}}{a} + \frac{e^{-by}}{b} = C'$$

[Topper's Answer 2022]



9. Find the sum of the order and degree of the differential equation:

$$\left(x + \frac{dy}{dx}\right)^2 = \left(\frac{dy}{dx}\right)^2 + 1$$

[CBSE 2021-22 (Term-2) (65/1/1)]

**Sol.** Given differential equation can be written as

$$x^2 + \left(\frac{dy}{dx}\right)^2 + 2x \frac{dy}{dx} = \left(\frac{dy}{dx}\right)^2 + 1$$

$$\text{i.e., } x^2 + 2x \frac{dy}{dx} = 1; \text{ Order} = 1, \text{ degree} = 1$$

$\frac{1}{2} + \frac{1}{2}$

$$\text{Sum of order and degree} = 1 + 1 = 2$$

1

[CBSE Marking Scheme 2022 (65/1/1)]

## Short Answer Questions

1. Find the general solution of the differential equation  $x \frac{dy}{dx} = y(\log y - \log x + 1)$ .

[CBSE 2021-22 (Term-2) (65/3/2)]

**Sol.**

$$\begin{aligned}
 & x \frac{dy}{dx} = y(\log y - \log x + 1) \\
 \Rightarrow & \frac{dy}{dx} = \frac{y}{x} \left( \log \left( \frac{y}{x} \right) + 1 \right) \quad \left[ \log a - \log b = \log \left( \frac{a}{b} \right) \right] \\
 & \text{on putting } x = \lambda x, \quad y = \lambda y \\
 & \text{Let } f(\lambda x, \lambda y) = \frac{\lambda y}{\lambda x} \left( \log \left( \frac{\lambda y}{\lambda x} \right) + 1 \right) \\
 & f(\lambda x, \lambda y) = \frac{y}{x} \left( \log \left( \frac{y}{x} \right) + 1 \right) \\
 & = f(x, y) \\
 & \text{Thus, this equation is homogenous equation.} \\
 & \text{Let } \frac{y}{x} = t \quad \text{or } y = tx \\
 & \frac{dy}{dx} \text{ on differentiating with respect to } x \\
 & \frac{dy}{dx} = t + x \frac{dt}{dx} \\
 & \frac{dy}{dx} = \frac{y}{x} \left( \log \left( \frac{y}{x} \right) + 1 \right)
 \end{aligned}$$

$$t + x \frac{dt}{dx} = t (\log t + 1) \quad \left[ \frac{y}{x} = t \right]$$

$$t + x \frac{dt}{dx} = t \log t + t$$

$$x \frac{dt}{dx} = t \log t$$

$$\frac{dt}{t \log t} = \frac{dx}{x}$$

on integrating both sides

$$\int \frac{dt}{t \log t} = \int \frac{dx}{x}$$

$$\int \frac{dt}{t \log t} = \log x + c$$

$$\text{let } \log t = u$$

on differentiating,

$$\frac{1}{t} dt = du$$

$$\int \frac{du}{u} = \ln x + c \quad [\log_e x = \ln x]$$

$$\ln u = \ln x + c \quad [c \text{ is integration constant}]$$

$$\ln(\ln t) = \ln x + c \quad [u = \log t = \ln t]$$

$$\ln(\ln(y/x)) = \ln x + c$$

$$\ln(\ln(y/x)) - \ln x = c$$

$$\ln\left(\frac{\ln(y/x)}{x}\right) = c$$

$$[\log a - \log b = \log(a/b)]$$

$$\text{Answer: } \ln\left(\frac{\ln(y/x)}{x}\right) = c$$

$$[\text{where } \ln x = \log_e x]$$

[Topper's Answer 2022]

2. Find the particular solution of the differential equation  $x \frac{dy}{dx} + x \cos^2\left(\frac{y}{x}\right) = y$ ; given that when  $x = 1$ ,  $y = \frac{\pi}{4}$ .  
[CBSE 2021-22 (65/1/1) (Term-2)]

**Sol.** Given differential equation can be written as

$$\frac{dy}{dx} = \frac{y}{x} - \cos^2 \frac{y}{x} \quad \dots(1) \quad 1$$

$$\text{Let } y = vx \quad \therefore \frac{dy}{dx} = v + x \frac{dv}{dx} \quad \frac{1}{2}$$

$$\text{Equation (1), Becomes } v + x \frac{dv}{dx} = v - \cos^2 v$$

$$\sec^2 v \, dv = -\frac{dx}{x}$$

Integrating both sides we get 1/2

$$\tan v = -\log |x| + c$$

$$\tan \frac{y}{x} = -\log |x| + c$$

$$x = 1, y = \frac{\pi}{4} \Rightarrow c = 1 \quad 1$$

$$\therefore \text{Particular solution is } \tan \frac{y}{x} = -\log |x| + 1 \quad \frac{1}{2}$$

[CBSE Marking Scheme 2022]

3. Find the particular solution of the differential equation

$$x \frac{dy}{dx} = y - x \tan\left(\frac{y}{x}\right), \text{ given that } y = \frac{x}{4} \text{ at } x = 1.$$

[CBSE 2020 (65/1/1)]

**Sol.**

$$\begin{aligned} x \frac{dy}{dx} &= y - x \tan\left(\frac{y}{x}\right) \\ \Rightarrow \frac{dy}{dx} &= \frac{y}{x} - \tan\left(\frac{y}{x}\right) = f\left(\frac{y}{x}\right) \quad \dots(i) \end{aligned}$$

$\therefore$  It is a homogeneous function.

$$\text{let } \frac{y}{x} = v \Rightarrow y = vx$$

Differentiating with respect to  $x$

$$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$\therefore$  equation (i) can be written as

$$v + x \frac{dv}{dx} = v - \tan v$$

$$\Rightarrow x \frac{dv}{dx} = -\tan v$$

$$\Rightarrow \int -\cot v \, dv = \int \frac{dx}{x}$$

$$\Rightarrow -\log \sin v = \log x + \log c, \quad -\log c \text{ is integration constant}$$

$$\Rightarrow \log \sin v + \log x = \log c$$

$$\Rightarrow x \sin v = c$$

$$\Rightarrow x \sin\left(\frac{y}{x}\right) = c$$

At  $x=1, y=\pi/4$

$$\sin\left(\frac{\pi}{4}\right) = c = \frac{1}{\sqrt{2}}$$

$$\Rightarrow x \sin\left(\frac{y}{x}\right) = \frac{1}{\sqrt{2}} \quad \underline{\text{Answer}}$$

[Topper's Answer 2020]

4. Solve the differential equation  $(1+x^2) \frac{dy}{dx} + 2xy - 4x^2 = 0$ , subject to the initial condition  $y(0) = 0$ . [CBSE 2019 (65/1/1)]

Sol.

$$(1+x^2) \frac{dy}{dx} + 2xy = 4x^2$$

$$\frac{dy}{dx} + \frac{2x}{1+x^2} y = \frac{4x^2}{1+x^2}$$

It is linear DE of form  $\frac{dy}{dx} + Py = Q$

$$P = \frac{2x}{1+x^2} \quad Q = \frac{4x^2}{1+x^2}$$

$$\text{I.F.} = e^{\int P dx} = e^{\int \frac{2x}{1+x^2} dx} = 1+x^2$$

Soln of DE:

$$y(1+x^2) = \int \frac{4x^2}{1+x^2} dx + C$$

$$= \int 4x^2 dx + C$$

$$y(1+x^2) = \frac{4x^3}{3} + C$$

$x=0, y=0$

$$\therefore C=0$$

$$\Rightarrow 3y(1+x^2) = 4x^3$$

[Topper's Answer 2019]

5. Find the general solution of the differential equation:  $\frac{d}{dx}(xy^2) = 2y(1+x^2)$  [CBSE 2023 (65/3/2)]

Sol. Given differential equation be  $\frac{d(xy^2)}{dx} = 2y(1+x^2)$

$$\Rightarrow x \frac{dy^2}{dx} + y^2 \frac{dx}{dx} = 2y(1+x^2)$$

$$\Rightarrow 2xy \frac{dy}{dx} + y^2 = 2y(1+x^2)$$

$$\Rightarrow 2x \frac{dy}{dx} + y = 2(1+x^2)$$

$$\Rightarrow \frac{dy}{dx} + \frac{1}{2x} \cdot y = \frac{2(1+x^2)}{2x}$$

$$\Rightarrow \frac{dy}{dx} + \frac{1}{2x} y = \frac{1+x^2}{x}$$

It is a linear differential equation of the type

$$\frac{dy}{dx} + Py = Q, \text{ where } P, Q \text{ be the function of a constant}$$

$$\therefore P = \frac{1}{2x}, Q = \frac{1+x^2}{x}$$

$$\therefore IF = e^{\int P dx} = e^{\int \frac{1}{2x} dx} = e^{\frac{1}{2} \log x} = e^{\log \sqrt{x}} = \sqrt{x}$$

Its solution be

$$y \times IF = \int Q \times IF dx$$

$$\Rightarrow y \times \sqrt{x} = \int \frac{1+x^2}{x} \times \sqrt{x} dx = \int \frac{1+x^2}{\sqrt{x}} dx = \int (x^{-\frac{1}{2}} + x^{\frac{3}{2}}) dx$$

$$\Rightarrow y\sqrt{x} = 2x^{\frac{1}{2}} + \frac{2}{5}x^{\frac{5}{2}} + C$$

$$\Rightarrow y = 2 + \frac{2}{5}x^2 + C$$

6. Solve the following differential equation :  $xe^{\frac{y}{x}} - y + x \frac{dy}{dx} = 0$

[CBSE 2023 (65/3/2)]

Sol. Given differential equation be

$$xe^{\frac{y}{x}} - y + x \frac{dy}{dx} = 0$$

$$\Rightarrow x \frac{dy}{dx} = y - xe^{\frac{y}{x}} \Rightarrow \frac{dy}{dx} = \frac{y}{x} - e^{\frac{y}{x}}$$

It is a homogeneous differential equation.

Put  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$ , we have

$$v + x \frac{dv}{dx} = v - e^v \Rightarrow x \frac{dv}{dx} = -e^v$$

$$\Rightarrow \int \frac{dv}{e^v} = -\int \frac{dx}{x} \text{ (on integrating)}$$

$$\Rightarrow \int e^{-v} dv = -\int \frac{dx}{x}$$

$$\Rightarrow -e^{-v} = -\log x + C$$

$$\Rightarrow \log x - e^{-\frac{y}{x}} = C$$

7. Solve the differential equation given by  $x dy - y dx - \sqrt{x^2 + y^2} dx = 0$ .

[CBSE 2023 (65/1/1)]

Sol. Given differential equation be

$$x dy - y dx - \sqrt{x^2 + y^2} dx = 0$$

$$\Rightarrow x dy = (y + \sqrt{x^2 + y^2}) dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x}$$



It is a homogeneous differential equation.

Put  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$ , we get

$$v + x \frac{dv}{dx} = \frac{vx + \sqrt{x^2 + v^2 x^2}}{x} = v + \sqrt{1 + v^2}$$
$$\Rightarrow x \frac{dv}{dx} = \sqrt{1 + v^2} \Rightarrow \frac{dv}{\sqrt{1 + v^2}} = \frac{dx}{x}$$

On integrating both sides, we have

$$\int \frac{dv}{\sqrt{1 + v^2}} = \int \frac{dx}{x} \Rightarrow \log |v + \sqrt{1 + v^2}| = \log x + \log C$$
$$\Rightarrow \log \left| \frac{v + \sqrt{1 + v^2}}{x} \right| = \log C$$
$$\Rightarrow \frac{\frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}}}{x} = C \Rightarrow y + \sqrt{x^2 + y^2} = Cx^2$$

8. Find the particular solution of the differential equation  $\frac{dy}{dx} + \sec^2 x \cdot y = \tan x \cdot \sec^2 x$ , given that  $y(0) = 0$ . [CBSE 2023 (65/1/1)]

**Sol.** Given differential equation be

$$\frac{dy}{dx} + \sec^2 x \cdot y = \tan x \cdot \sec^2 x, \text{ given } y(0) = 0$$

It is a linear differential equation of type

$$\frac{dy}{dx} + Py = Q, \text{ where } P, Q \text{ be the function of } x \text{ or constant}$$

$$\therefore P = \sec^2 x, Q = \tan x \sec^2 x$$

$$\therefore IF = e^{\int P dx} = e^{\int \sec^2 x dx} = e^{\tan x}$$

Its solution be

$$y \times IF = \int Q \times IF dx$$
$$\Rightarrow y \times e^{\tan x} = \int \tan x \cdot \sec^2 x \cdot e^{\tan x} dx$$

$$\text{Let } \tan x = t \Rightarrow \sec^2 x dx = dt$$

$$= \int t \cdot e^t dt = t \int e^t dt - \int \left( \frac{dt}{dt} \cdot \int e^t dt \right) dt$$
$$= t \cdot e^t - \int e^t dt = te^t - e^t + C = e^t (t - 1) + C$$

$$\Rightarrow y e^{\tan x} = e^{\tan x} (\tan x - 1) + C$$

$$\Rightarrow y = \tan x - 1 + C e^{-\tan x}$$

$$\text{Given, } y(0) = 0 \Rightarrow 0 = \tan 0 - 1 + C e^{\tan 0}$$

$$\Rightarrow 0 = C - 1 \Rightarrow C = 1$$

Particular solution is given by

$$y = \tan x - 1 + e^{-\tan x}$$

9. Find the general solution of the differential equation:

$$(xy - x^2) dy = y^2 dx.$$

[CBSE 2023 (65/2/1)]

**Sol.** Given differential equation be

$$(xy - x^2) dy = y^2 dx$$



$$\Rightarrow \frac{dy}{dx} = \frac{y^2}{xy - x^2}$$

$$\Rightarrow \frac{dx}{dy} = \frac{xy - x^2}{y^2}$$

It is a homogeneous differential equation.

$$\text{Put } x = vy \Rightarrow \frac{dx}{dy} = v + y \frac{dv}{dy}$$

$$\Rightarrow v + y \frac{dv}{dy} = \frac{vy^2 - v^2y^2}{y^2} = \frac{v - v^2}{1} = v - v^2$$

$$\Rightarrow v + y \frac{dv}{dy} = v - v^2$$

$$\Rightarrow y \frac{dv}{dy} = -v^2 \quad \Rightarrow \frac{dv}{v^2} = -\frac{dy}{y}$$

On integrating both sides, we have

$$\int \frac{dv}{v^2} = - \int \frac{dy}{y} \quad \Rightarrow \int v^{-2} dv = -\log y + C$$

$$\Rightarrow \frac{v^{-1}}{-1} = -\log y + C \quad \Rightarrow \log y - \frac{1}{v} = C$$

$$\Rightarrow \log y - \frac{y}{x} = C$$

**10. Find the general solution of the differential equation:**

$$(x^2 + 1) \frac{dy}{dx} + 2xy = \sqrt{x^2 + 4}$$

[CBSE 2023 (65/2/1)]

**Sol.** Given differential equation be

$$(x^2 + 1) \frac{dy}{dx} + 2xy = \sqrt{x^2 + 4}$$

$$\Rightarrow \frac{dy}{dx} + \frac{2x}{x^2 + 1} \cdot y = \frac{\sqrt{x^2 + 4}}{x^2 + 1}$$

It is a linear differential equation of the type

$$\frac{dy}{dx} + Py = Q$$

$$\text{Hence, } P = \frac{2x}{x^2 + 1}, Q = \frac{\sqrt{x^2 + 4}}{x^2 + 1}$$

$$\therefore IF = e^{\int P dx} = e^{\int \frac{2x}{x^2 + 1} dx} = e^{\log(x^2 + 1)} = (x^2 + 1)$$

Thus, its solution be

$$y \times IF = \int Q \times IF dx$$

$$\Rightarrow y \times (x^2 + 1) = \int \frac{\sqrt{x^2 + 4}}{x^2 + 1} \times (x^2 + 1) dx = \int \sqrt{x^2 + 4} dx$$

$$\begin{aligned} \Rightarrow y(x^2 + 1) &= \frac{x}{2} \sqrt{x^2 + 4} + \frac{4}{2} \log |x + \sqrt{x^2 + 4}| + C \\ &= \frac{x}{2} \sqrt{x^2 + 4} + 2 \log |x + \sqrt{x^2 + 4}| + C \end{aligned}$$

11. Find the particular solution of the differential equation  $\frac{dy}{dx} + 2y \tan x = \sin x$ , given that  $y = 0$  when  $x = \frac{\pi}{3}$ . [CBSE 2018]

Sol.

$$\frac{dy}{dx} + 2y \tan x = \sin x$$

on comparing the above equation with the standard linear equation  $\frac{dy}{dx} + Py = Q$

we get,  $P = 2 \tan x$ ,  $Q = \sin x$

Therefore, I.F. =  $e^{\int P dx}$

$$\begin{aligned} \text{I.F.} &= e^{\int 2 \tan x dx} \\ &= e^{2(\log \sec x)} \\ &= e^{\log (\sec x)^2} = \sec^2 x \end{aligned}$$

Now,

$$y \cdot \text{I.F.} = \int Q \times \text{I.F.} dx$$

$$y \cdot \sec^2 x = \int \sin x \sec^2 x dx$$

$$y \cdot \sec^2 x = \int \sin x \times \frac{1}{\cos^2 x} dx$$

$$y \cdot \sec^2 x = \int \tan x \cdot \sec x dx$$

Put  $\sec x = t$   
 $(\sec x \tan x) dx = dt$

$$y \cdot \sec^2 x = \int t dt$$

$$y \cdot \sec^2 x = \frac{t^2}{2} + C$$

$$y \cdot \sec^2 x = \sec x + C$$

Now when  $y = 0$ ,  $x = \pi/3$

$$0 = \sec \frac{\pi}{3} + C, \quad 0 = 2 + C$$

$$C = -2$$

Therefore,

$$y \cdot \sec^2 x = \sec x - 2$$

or  $y = \frac{1}{\sec x} - \frac{2}{\sec^2 x}$

or  $y = \sec^{-1} x - 2(\sec^2 x)^{-1}$  Ans.

[Topper's Answer 2018]

12. Solve the following differential equation:  $(1 + e^{y/x})dy + e^{y/x}\left(1 - \frac{y}{x}\right)dx = 0$ , ( $x \neq 0$ ). [CBSE 2020 (65/2/1)]

Sol. Given differential equation

$$(1 + e^{y/x})dy + e^{y/x}\left(1 - \frac{y}{x}\right)dx = 0$$

$$\Rightarrow (1 + e^{y/x}) dy = \left( \frac{y}{x} - 1 \right) e^{y/x} dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{\left( \frac{y}{x} - 1 \right) e^{y/x}}{(1 + e^{y/x})}$$

It is a homogeneous differential equation.

Put  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$

We have,

$$v + x \frac{dv}{dx} = \frac{(v-1)e^v}{1+e^v} = \frac{ve^v - e^v}{1+e^v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{ve^v - e^v}{1+e^v} - v = \frac{ve^v - e^v - v - ve^v}{1+e^v}$$

$$\Rightarrow x \frac{dv}{dx} = -\frac{(v+e^v)}{1+e^v}$$

$$\Rightarrow \frac{1+e^v}{v+e^v} dv = -\frac{dx}{x}$$

On integrating both sides, we have

$$\int \frac{1+e^v}{v+e^v} dv = -\int \frac{dx}{x}$$

$$\Rightarrow \log |v+e^v| = -\log |x| + \log |C|$$

$$\Rightarrow \log |v+e^v| + \log |x| = \log |C|$$

$$\Rightarrow \log |x(v+e^v)| = \log |C|$$

$$\Rightarrow x(v+e^v) = C \Rightarrow x\left(\frac{y}{x} + e^{y/x}\right) = C$$

$$\Rightarrow y + x e^{y/x} = C$$

13. Solve the differential equation  $(1+x^2)\frac{dy}{dx} + y = e^{\tan^{-1}x}$ .

[CBSE (AI) 2014]

**Sol.** Given differential equation is

$$(1+x^2)\frac{dy}{dx} + y = e^{\tan^{-1}x}$$

$$\Rightarrow \frac{dy}{dx} + \frac{1}{1+x^2}y = \frac{e^{\tan^{-1}x}}{1+x^2} \quad \dots(i)$$

Equation (i) is of the form

$$\frac{dy}{dx} + Py = Q, \text{ where } P = \frac{1}{1+x^2}, Q = \frac{e^{\tan^{-1}x}}{1+x^2}$$

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \frac{1}{1+x^2} dx} = e^{\tan^{-1}x}$$

Therefore, general solution of required differential equation is

$$y \cdot e^{\tan^{-1}x} = \int e^{\tan^{-1}x} \cdot \frac{e^{\tan^{-1}x}}{1+x^2} dx + C$$

$$\Rightarrow y \cdot e^{\tan^{-1}x} = \int \frac{e^{2\tan^{-1}x}}{1+x^2} dx + C \quad \dots(ii)$$

$$\text{Let } \tan^{-1} x = z \quad \Rightarrow \quad \frac{1}{1+x^2} dx = dz$$

(ii) becomes

$$y \cdot e^{\tan^{-1} x} = \int e^{2z} dz + C \quad \Rightarrow \quad y \cdot e^{\tan^{-1} x} = \frac{e^{2z}}{2} + C$$

$$\Rightarrow \quad y \cdot e^{\tan^{-1} x} = \frac{e^{2 \tan^{-1} x}}{2} + C \quad [\text{Putting } z = \tan^{-1} x]$$

$$\Rightarrow \quad y = \frac{e^{\tan^{-1} x}}{2} + C \cdot e^{-\tan^{-1} x} \quad [\text{Dividing both sides by } e^{\tan^{-1} x}]$$

It is the required solution.

14. Find the particular solution of the differential equation  $e^x \sqrt{1-y^2} dx + \frac{y}{x} dy = 0$  given that  $y = 1$  when  $x = 0$ . [CBSE Delhi 2014]

**Sol.** We have,  $e^x \sqrt{1-y^2} dx + \frac{y}{x} dy = 0$

$$\Rightarrow \quad e^x \sqrt{1-y^2} dx = -\frac{y}{x} dy \quad \Rightarrow \quad x e^x dx = -\frac{y}{\sqrt{1-y^2}} dy$$

$$\Rightarrow \quad \int x e^x dx = -\int \frac{y}{\sqrt{1-y^2}} dy$$

$$\Rightarrow \quad x e^x - \int e^x dx = \frac{1}{2} \int \frac{dt}{\sqrt{t}}, \text{ where } t = 1-y^2 \Rightarrow \frac{dt}{2} = -y dy \quad (\text{Using ILATE on LHS})$$

$$\Rightarrow \quad x e^x - e^x = \frac{1}{2} \left( \frac{t^{1/2}}{1/2} \right) + C \quad \Rightarrow \quad x e^x - e^x = \sqrt{t} + C$$

$$\Rightarrow \quad x e^x - e^x = \sqrt{1-y^2} + C, \text{ is the general solution.}$$

Putting  $y = 1$  and  $x = 0$ , we get

$$0e^0 - e^0 = \sqrt{1-1^2} + C \quad \Rightarrow \quad C = -1$$

Therefore, required particular solution is  $x e^x - e^x = \sqrt{1-y^2} - 1$ .

15. Solve the differential equation:

[CBSE (AI) 2013; Delhi 2015]

$$(\tan^{-1} y - x) dx = (1 + y^2) dy$$

**Sol.** The given differential equation can be written as

$$\frac{dx}{dy} + \frac{x}{1+y^2} = \frac{\tan^{-1} y}{1+y^2} \quad \dots (i)$$

$$\text{Now, (i) is of the form } \frac{dx}{dy} + Px = Q, \text{ where } P = \frac{1}{1+y^2} \text{ and } Q = \frac{\tan^{-1} y}{1+y^2}$$

$$\text{Therefore, IF} = e^{\int \frac{1}{1+y^2} dy} = e^{\tan^{-1} y}$$

Thus, the solution of the given differential equation is

$$x e^{\tan^{-1} y} = \int \left( \frac{\tan^{-1} y}{1+y^2} \right) e^{\tan^{-1} y} dy + C \quad \dots (ii)$$

$$\text{Let } I = \int \left( \frac{\tan^{-1} y}{1+y^2} \right) e^{\tan^{-1} y} dy$$

Substituting  $\tan^{-1} y = t$  so that  $\left( \frac{1}{1+y^2} \right) dy = dt$ , we get

$$I = \int t e^t dt = t e^t - \int 1 \cdot e^t dt = t e^t - e^t \equiv e^t (t - 1)$$

$$\text{or } I = e^{\tan^{-1}y}(\tan^{-1}y - 1)$$

Substituting the value of  $I$  in equation (ii), we get

$$x \cdot e^{\tan^{-1}y} = e^{\tan^{-1}y}(\tan^{-1}y - 1) + C$$

or  $x = (\tan^{-1}y - 1) + C e^{-\tan^{-1}y}$  is the required solution.

16. Solve the differential equation  $(x^2 - 1) \frac{dy}{dx} + 2xy = \frac{2}{x^2 - 1}$ , where  $x \in (-\infty, -1) \cup (1, \infty)$ .

[CBSE Delhi 2014; (AI) 2010; (F) 2009, 2011]

**Sol.** The given differential equation is  $(x^2 - 1) \frac{dy}{dx} + 2xy = \frac{2}{x^2 - 1}$ .

$$\Rightarrow \frac{dy}{dx} + \frac{2x}{x^2 - 1}y = \frac{2}{(x^2 - 1)^2} \quad \dots(i)$$

This is a linear differential equation of the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \frac{2x}{x^2 - 1}$  and  $Q = \frac{2}{(x^2 - 1)^2}$

$$\therefore \text{IF} = e^{\int P dx} = e^{\int 2x/(x^2-1) dx} = e^{\log|x^2-1|} = x^2 - 1$$

Multiplying both sides of (i) by IF  $= x^2 - 1$ , we get  $(x^2 - 1) \frac{dy}{dx} + 2xy = \frac{2}{x^2 - 1}$

$$\Rightarrow d(y(x^2 - 1)) = \frac{2}{x^2 - 1}$$

Integrating both sides, we get

$$y(x^2 - 1) = \int \frac{2}{x^2 - 1} dx + C \quad [\text{Using: } y(\text{IF}) = \int Q \cdot (\text{IF}) dx + C]$$

$$\Rightarrow y(x^2 - 1) = \frac{2}{2} \log \left| \frac{x-1}{x+1} \right| + C \Rightarrow y(x^2 - 1) = \log \left| \frac{x-1}{x+1} \right| + C$$

This is the required solution.

17. Find the particular solution of the differential equation  $\frac{dy}{dx} = 1 + x + y + xy$  given that  $y = 0$  when  $x = 1$ . [CBSE (AI) 2014]

**Sol.** Given differential equation is  $\frac{dy}{dx} = 1 + x + y + xy$ .

$$\Rightarrow \frac{dy}{dx} = (1 + x) + y(1 + x) \Rightarrow \frac{dy}{dx} = (1 + x)(1 + y) \Rightarrow \frac{dy}{1 + y} = (1 + x) dx$$

Integrating both sides, we get  $\log |1 + y| = \int (1 + x) dx$

$$\Rightarrow \log |1 + y| = x + \frac{x^2}{2} + C \text{ is the general solution.}$$

Putting  $x = 1, y = 0$ , we get

$$\log 1 = 1 + \frac{1}{2} + C \Rightarrow 0 = \frac{3}{2} + C \Rightarrow C = -\frac{3}{2}$$

$$\text{Hence, particular solution is } \log |1 + y| = x + \frac{x^2}{2} - \frac{3}{2}.$$

18. Solve the differential equation  $x \log x \frac{dy}{dx} + y = \frac{2}{x} \log x$ .

[CBSE (F) 2014]

**Sol.** Given differential equation is  $x \log x \frac{dy}{dx} + y = \frac{2}{x} \log x$

$$\Rightarrow \frac{dy}{dx} + \left( \frac{1}{x \cdot \log x} \right) y = \frac{2}{x^2} \quad (\text{Divide each term by } x \log x)$$

It is in the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \frac{1}{x \cdot \log x}$  and  $Q = \frac{2}{x^2}$ .

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \frac{dx}{x \log x}}$$

$$\text{Put } \log x = z \Rightarrow \frac{dx}{x} = dz$$

$$\text{IF } e^{\int \frac{1}{z} dz} = e^{\log z} = z = \log x$$

$\therefore$  General solution is

$$y \cdot \log x = \int \log x \cdot \frac{2}{x^2} dx + C \quad \Rightarrow \quad y \log x = 2 \int \frac{\log x}{x^2} dx + C$$

$$\text{Let } \log x = z \Rightarrow \frac{1}{x} dx = dz \quad \text{also } \log x = z \Rightarrow x = e^z$$

$$\therefore y \log x = 2 \int \frac{z}{e^z} dz + C \quad \Rightarrow \quad y \log x = 2 \int z \cdot e^{-z} dz + C$$

$$\Rightarrow y \log x = 2 \left[ z \cdot \frac{e^{-z}}{-1} - \int \frac{e^{-z}}{-1} dz \right] + C \Rightarrow y \log x = 2 \left[ -ze^{-z} + \int e^{-z} dz \right] + C$$

$$\Rightarrow y \log x = -2ze^{-z} - 2e^{-z} + C \quad \Rightarrow \quad y \log x = -2 \log x \cdot e^{-\log x} - 2e^{-\log x} + C$$

$$\Rightarrow y \log x = -2 \log x \cdot \frac{1}{x} - \frac{2}{x} + C \quad \left[ \because e^{-\log x} = e^{\log \frac{1}{x}} = \frac{1}{x} \right]$$

$$\Rightarrow y \log x = -\frac{2}{x} (1 + \log x) + C$$

19. Show that the differential equation  $x \frac{dy}{dx} \sin\left(\frac{y}{x}\right) + x - y \sin\left(\frac{y}{x}\right) = 0$  is homogeneous. Find the particular solution of this differential equation, given that  $x = 1$  when  $y = \frac{\pi}{2}$ . [CBSE Delhi 2013]

**Sol.** Given differential equation is  $x \frac{dy}{dx} \sin\left(\frac{y}{x}\right) + x - y \sin\left(\frac{y}{x}\right) = 0$ .

Dividing both sides by  $x \sin\left(\frac{y}{x}\right)$ , we get

$$\frac{dy}{dx} + \operatorname{cosec} \frac{y}{x} \cdot \frac{y}{x} - \frac{y}{x} = 0 \quad \Rightarrow \quad \frac{dy}{dx} = \frac{y}{x} - \operatorname{cosec} \frac{y}{x} \quad \dots (i)$$

$$\text{Let } F(x, y) = \frac{y}{x} - \operatorname{cosec} \frac{y}{x}$$

$$\therefore F(\lambda x, \lambda y) = \frac{\lambda y}{\lambda x} - \operatorname{cosec} \frac{\lambda y}{\lambda x} = \lambda^0 \left[ \frac{y}{x} - \operatorname{cosec} \frac{y}{x} \right] = \lambda^0 F(x, y)$$

Hence, differential equation (i) is homogeneous.

$$\text{Let } y = vx \quad \Rightarrow \quad \frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$$

Now, equation (i) becomes

$$v + x \cdot \frac{dv}{dx} = \frac{vx}{x} - \operatorname{cosec} \frac{vx}{x}$$

$$v + x \cdot \frac{dv}{dx} = v - \operatorname{cosec} v \quad \Rightarrow \quad x \cdot \frac{dv}{dx} = -\operatorname{cosec} v$$



$$\Rightarrow -\sin v dv = \frac{dx}{x} \quad \Rightarrow \quad -\int \sin v dv = \int \frac{dx}{x}$$

$$\Rightarrow \cos v = \log |x| + C \quad \Rightarrow \quad \cos \frac{y}{x} = \log |x| + C \quad \dots(ii)$$

Putting  $y = \frac{\pi}{2}$ ,  $x = 1$  in (ii), we get

$$\therefore \cos \frac{\pi}{2} = \log 1 + C \quad \Rightarrow \quad 0 = 0 + C \quad \Rightarrow \quad C = 0$$

Hence, particular solution is

$$\cos \frac{y}{x} = \log |x| + 0 \quad \text{i.e.,} \quad \cos \frac{y}{x} = \log |x|$$

**20. Solve the differential equation:**

$$\sqrt{1+x^2+y^2+x^2y^2} + xy \frac{dy}{dx} = 0$$

[CBSE (AI) 2010; (F) 2015]

**Sol.** Given  $\sqrt{1+x^2+y^2+x^2y^2} + xy \frac{dy}{dx} = 0$

By simplifying the equation, we get

$$xy \frac{dy}{dx} = -\sqrt{1+x^2+y^2+x^2y^2} = -\sqrt{1+x^2+y^2(1+x^2)}$$

$$\Rightarrow xy \frac{dy}{dx} = -\sqrt{(1+x^2)(1+y^2)} = -\sqrt{(1+x^2)} \sqrt{(1+y^2)}$$

$$\Rightarrow \frac{y}{\sqrt{(1+y^2)}} dy = -\frac{\sqrt{(1+x^2)}}{x} dx$$

Integrating both sides, we get

$$\int \frac{y}{\sqrt{(1+y^2)}} dy = -\int \frac{\sqrt{(1+x^2)}}{x} dx = -\int \frac{\sqrt{1+x^2}}{x^2} \times x dx \quad \dots(i)$$

$$\text{Let } 1+y^2 = t \Rightarrow 2y dy = dt \quad \text{and} \quad 1+x^2 = m^2 \Rightarrow 2x dx = 2m dm \Rightarrow x dx = m dm$$

$$\therefore (i) \Rightarrow \frac{1}{2} \int \frac{1}{\sqrt{t}} dt = -\int \frac{m}{m^2-1} \cdot m dm$$

$$\Rightarrow \frac{1}{2} \frac{t^{1/2}}{1/2} + \int \frac{m^2}{m^2-1} dm = 0 \quad \Rightarrow \quad \sqrt{t} + \int \frac{m^2+1-1}{m^2-1} dm = 0$$

$$\Rightarrow \sqrt{t} + \int \left(1 + \frac{1}{m^2-1}\right) dm = 0 \quad \Rightarrow \quad \sqrt{t} + m + \frac{1}{2} \log \left| \frac{m-1}{m+1} \right| = 0$$

Now, substituting the value of  $t$  and  $m$ , we get

$$\sqrt{1+y^2} + \sqrt{1+x^2} + \frac{1}{2} \log \left| \frac{\sqrt{1+x^2}-1}{\sqrt{1+x^2}+1} \right| + C = 0$$

**21.  $(x^2 + y^2)dy = xy dx$ . If  $y(1) = 1$  and  $y(x_0)e$ , then find the value of  $x_0$ .** [CBSE Bhubneshwar 2015]

**Sol.** Given differential equation is  $(x^2 + y^2)dy = xy dx$

It is also written as

$$\frac{dy}{dx} = \frac{xy}{x^2 + y^2} \quad \dots(i)$$

Now, to solve let  $y = vx$ .

[ $\because$  (i) is a homogeneous equation]

Differentiating  $y = vx$  with respect to  $x$ , we get

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

Putting  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$  in (i), we get

$$v + x \frac{dv}{dx} = \frac{x \cdot vx}{x^2 + (vx)^2}$$

$$v + x \frac{dv}{dx} = \frac{vx^2}{x^2 + v^2 x^2} \Rightarrow v + x \frac{dv}{dx} = \frac{vx^2}{x^2(1+v^2)} = \frac{v}{1+v^2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v}{(1+v^2)} - v \Rightarrow x \frac{dv}{dx} = \frac{v - v - v^3}{(1+v^2)}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{-v^3}{(1+v^2)} \Rightarrow \frac{(1+v^2)dv}{v^3} = -\frac{dx}{x}$$

Integrating both sides, we get

$$\int \frac{(1+v^2)dv}{v^3} = -\int \frac{dx}{x}$$

$$\Rightarrow \int \frac{dv}{v^3} + \int \frac{dv}{v} = -\log |x| + C \Rightarrow -\frac{1}{2v^2} + \log |v| = -\log |x| + C$$

$$\Rightarrow -\frac{x^2}{2y^2} + \log \left| \frac{y}{x} \right| = -\log |x| + C \Rightarrow -\frac{x^2}{2y^2} + \log |y| - \log |x| = -\log |x| + C$$

$$\Rightarrow -\frac{x^2}{2y^2} + \log |y| = C \quad \dots (ii)$$

Given,  $x = 1, y = 1$

$$\Rightarrow -\frac{1}{2 \times 1} + \log |1| = C \Rightarrow -\frac{1}{2} = C \quad [\because \log 1 = 0]$$

Now (ii) becomes

$$-\frac{x^2}{2y^2} + \log |y| = -\frac{1}{2} \Rightarrow \log |y| = \frac{x^2}{2y^2} - \frac{1}{2} \Rightarrow \log |y| = \frac{x^2 - y^2}{2y^2} \quad \dots (iii)$$

Putting  $x = x_0$  and  $y = e$  in (iii), we get

$$\begin{aligned} \log |e| &= \frac{x_0^2 - e^2}{2e^2} \Rightarrow 1 = \frac{x_0^2 - e^2}{2e^2} \Rightarrow x_0^2 - e^2 = 2e^2 \\ \Rightarrow x_0^2 &= 3e^2 \Rightarrow x_0 = \sqrt{3}e \end{aligned}$$

22. Show that the differential equation  $(x - y) \frac{dy}{dx} = x + 2y$  is homogeneous and solve it.

[CBSE (AI) 2010, 2017; (F) 2013; Ajmer 2015]

**Sol.** Given,  $(x - y) \frac{dy}{dx} = x + 2y$

By simplifying the above equation, we get

$$\frac{dy}{dx} = \frac{x + 2y}{x - y} \quad \dots (i)$$

$$\text{Let } F(x, y) = \frac{x+2y}{x-y}$$

$$\text{then } F(\lambda x, \lambda y) = \frac{\lambda x + 2\lambda y}{\lambda x - \lambda y} = \frac{\lambda(x+2y)}{\lambda(x-y)} = \lambda^0 F(x, y)$$

$F(x, y)$  is homogeneous function and hence given differential equation is homogeneous.

$$\text{Now, let } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

Substituting these values in equation (i), we get

$$\begin{aligned} v + x \frac{dv}{dx} &= \frac{x+2vx}{x-vx} \\ \Rightarrow x \frac{dv}{dx} &= \frac{1+2v}{1-v} - v = \frac{1+2v-v+v^2}{1-v} = \frac{1+v+v^2}{1-v} \\ \Rightarrow \frac{1-v}{1+v+v^2} dv &= \frac{dx}{x} \end{aligned}$$

By integrating both sides, we get

$$\int \frac{1-v}{1+v+v^2} dv = \int \frac{dx}{x} \quad \dots(ii)$$

$$\text{LHS} = \int \frac{1-v}{v^2+v+1} dv$$

$$\text{Let } 1-v = A(2v+1) + B = 2Av + (A+B)$$

Comparing coefficients of both sides, we get

$$2A = -1, \quad A+B = 1 \quad \text{or} \quad A = -\frac{1}{2}, \quad B = \frac{3}{2}$$

$$\begin{aligned} \therefore \int \frac{1-v}{v^2+v+1} dv &= \int \frac{-\frac{1}{2}(2v+1) + \frac{3}{2}}{v^2+v+1} dv \\ &= -\frac{1}{2} \int \frac{2v+1}{v^2+v+1} dv + \frac{3}{2} \int \frac{dv}{v^2+v+1} \\ &= -\frac{1}{2} \int \frac{2v+1}{v^2+v+1} dv + \frac{3}{2} \int \frac{dv}{\left(v+\frac{1}{2}\right)^2 + \frac{3}{4}} \\ &= -\frac{1}{2} \log |v^2+v+1| + \frac{3}{2} \times \frac{2}{\sqrt{3}} \tan^{-1} \left( \frac{v+\frac{1}{2}}{\frac{\sqrt{3}}{2}} \right) \end{aligned}$$

Now, substituting it in equation (ii), we get

$$\begin{aligned} -\frac{1}{2} \log |v^2+v+1| + \sqrt{3} \tan^{-1} \left( \frac{2v+1}{\sqrt{3}} \right) &= \log x + C \\ \Rightarrow -\frac{1}{2} \log \left| \frac{y^2}{x^2} + \frac{y}{x} + 1 \right| + \sqrt{3} \tan^{-1} \left( \frac{\frac{2y}{x} + 1}{\sqrt{3}} \right) &= \log x + C \\ \Rightarrow -\frac{1}{2} \log |x^2 + xy + y^2| + \frac{1}{2} \log x^2 + \sqrt{3} \tan^{-1} \left( \frac{2y+x}{\sqrt{3}x} \right) &= \log x + C \\ \Rightarrow -\frac{1}{2} \log |x^2 + xy + y^2| + \sqrt{3} \tan^{-1} \left( \frac{2y+x}{\sqrt{3}x} \right) &= C \end{aligned}$$

23. Solve  $\frac{dy}{dx} = \cos(x+y) + \sin(x+y)$ .

[NCERT Exemplar]

Sol. Given,  $\frac{dy}{dx} = \cos(x+y) + \sin(x+y)$

Put  $x+y=z \Rightarrow 1 + \frac{dy}{dx} = \frac{dz}{dx}$

On substituting these values in equation (i), we get

$$\left(\frac{dz}{dx} - 1\right) = \cos z + \sin z \Rightarrow \frac{dz}{dx} = (\cos z + \sin z + 1) \Rightarrow \frac{dz}{\cos z + \sin z + 1} = dx$$

On integrating both sides, we get

$$\int \frac{dz}{\cos z + \sin z + 1} = \int 1 dx$$

$$\Rightarrow \int \frac{dz}{\frac{1 - \tan^2 z/2}{1 + \tan^2 z/2} + \frac{2 \tan z/2}{1 + \tan^2 z/2} + 1} = \int dx$$

$$\Rightarrow \int \frac{dz}{\frac{1 - \tan^2 z/2 + 2 \tan z/2 + 1 + \tan^2 z/2}{(1 + \tan^2 z/2)}} = \int dx$$

$$\Rightarrow \int \frac{(1 + \tan^2 z/2) dz}{2 + 2 \tan z/2} = \int dx \Rightarrow \int \frac{\sec^2 z/2 dz}{2(1 + \tan z/2)} = \int dx$$

Put  $1 + \tan z/2 = t \Rightarrow \left(\frac{1}{2} \sec^2 z/2\right) dz = dt$

$$\Rightarrow \int \frac{dt}{t} = \int dx \Rightarrow \log |t| = x + C$$

$$\Rightarrow \log |1 + \tan z/2| = x + C \Rightarrow \log \left| 1 + \tan \frac{(x+y)}{2} \right| = x + C$$

24. Find the particular solution of the differential equation:

$(1-y^2)(1+\log x) dx + 2xy dy = 0$  given that  $y = 0$  when  $x = 1$

[CBSE Delhi 2016]

Sol. We have  $(1-y^2)(1+\log x) dx + 2xy dy = 0$

$$\Rightarrow 2xy dy = -(1-y^2)(1+\log x) dx \Rightarrow \frac{2y dy}{1-y^2} = -\frac{(1+\log x) dx}{x}$$

Integrating both sides, we get

$$\Rightarrow \int \frac{2y}{1-y^2} dy = -\int \frac{(1+\log x)}{x} dx \Rightarrow -\log |1-y^2| = -\int \frac{(1+\log x)}{x} dx$$

$$\Rightarrow -\log |1-y^2| = -\int x dz \left[ \text{Let } 1+\log x = z \Rightarrow \frac{1}{x} dx = dz \right]$$

$$\Rightarrow \log |1-y^2| = \frac{z^2}{2} + C \Rightarrow \log |1-y^2| = \frac{(1+\log x)^2}{2} + C$$

Putting  $x = 1$  and  $y = 0$ , we get

$$\Rightarrow \log 1 = \frac{(1+\log 1)^2}{2} + C \Rightarrow 0 = \frac{1}{2} + C \Rightarrow C = -\frac{1}{2}$$

Hence, particular solution is  $\log |1-y^2| = \frac{(1+\log x)^2}{2} - \frac{1}{2}$ .

25. Find the general solution of the following differential equation:

$$(1 + y^2) + (x - e^{\tan^{-1}y}) \frac{dy}{dx} = 0$$

[CBSE Delhi 2016]

Sol. We have  $(1 + y^2) + (x - e^{\tan^{-1}y}) \frac{dy}{dx} = 0$

$$\Rightarrow (x - e^{\tan^{-1}y}) \frac{dy}{dx} = -(1 + y^2)$$

$$\Rightarrow \frac{dy}{dx} = -\left(\frac{1 + y^2}{x - e^{\tan^{-1}y}}\right) \Rightarrow \frac{dx}{dy} = -\left(\frac{x - e^{\tan^{-1}y}}{1 + y^2}\right)$$

$$\Rightarrow \frac{dx}{dy} = -\frac{x}{1 + y^2} + \frac{e^{\tan^{-1}y}}{1 + y^2} \Rightarrow \frac{dx}{dy} + \frac{1}{1 + y^2}x = \frac{e^{\tan^{-1}y}}{1 + y^2}$$

It is in the form  $\frac{dx}{dy} + Px = Q$ , where  $P = \frac{1}{1 + y^2}$  and  $Q = \frac{e^{\tan^{-1}y}}{1 + y^2}$ .

$$\therefore \text{IF} = e^{\int P \cdot dy} = e^{\int \frac{1}{1 + y^2} dy} = e^{\tan^{-1}y}$$

Therefore, general solution is  $x \cdot e^{\tan^{-1}y} = \int \frac{e^{\tan^{-1}y}}{1 + y^2} \cdot e^{\tan^{-1}y} dy + C$ .

$$\Rightarrow x \cdot e^{\tan^{-1}y} = \int e^z \cdot e^z dz + C \quad \left[ \text{Let } \tan^{-1}y = z \Rightarrow \frac{1}{1 + y^2} dy = dz \right]$$

$$\Rightarrow x \cdot e^{\tan^{-1}y} = \int e^{2z} dz + C$$

$$\Rightarrow x \cdot e^{\tan^{-1}y} = \frac{e^{2z}}{2} + C \Rightarrow x \cdot e^{\tan^{-1}y} = \frac{e^{2 \tan^{-1}y}}{2} + C$$

$$\Rightarrow x = \frac{1}{2} e^{\tan^{-1}y} + C \cdot e^{-\tan^{-1}y}$$

26. Find the particular solution of differential equation:  $\frac{dy}{dx} = -\frac{x + y \cos x}{1 + \sin x}$  given that  $y = 1$  when  $x = 0$ . [CBSE (North) 2016]

Sol. We have

$$\frac{dy}{dx} = -\frac{x + y \cos x}{1 + \sin x}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{x}{1 + \sin x} - \frac{y \cos x}{1 + \sin x} \Rightarrow \frac{dy}{dx} + \frac{\cos x}{1 + \sin x} y = -\frac{x}{1 + \sin x}$$

It is in the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \frac{\cos x}{1 + \sin x}$ ,  $Q = -\frac{x}{1 + \sin x}$ .

$$\text{Now IF} = e^{\int \frac{\cos x}{1 + \sin x} dx} = e^{\log |1 + \sin x|} = 1 + \sin x$$

Therefore, general solution is

$$y(1 + \sin x) = \int -\frac{x}{1 + \sin x} (1 + \sin x) dx + C = -\int x dx + C$$

$$\Rightarrow y(1 + \sin x) = -\frac{x^2}{2} + C$$

$$1(1 + \sin 0) = 0 + C$$

$$[\text{Given } y = 1 \text{ and } x = 0]$$

$$\Rightarrow C = 1$$

Hence, particular solution is

$$y(1 + \sin x) = -\frac{x^2}{2} + 1$$

$$\Rightarrow y = \frac{2 - x^2}{2(1 + \sin x)}$$

## Long Answer Questions

1. Solve the following differential equation:

$$3e^x \tan y \, dx + (2 - e^x) \sec^2 y \, dy = 0, \text{ given that when } x = 0, y = \frac{\pi}{4} \quad [\text{CBSE (F) 2012}]$$

Sol. Given,  $3e^x \tan y \, dx + (2 - e^x) \sec^2 y \, dy = 0$

$$\Rightarrow (2 - e^x) \sec^2 y \, dy = -3e^x \tan y \, dx$$

$$\Rightarrow \frac{\sec^2 y}{\tan y} dy = \frac{-3e^x}{2 - e^x} dx$$

$$\Rightarrow \int \frac{\sec^2 y}{\tan y} dy = 3 \int \frac{-e^x dx}{2 - e^x}$$

$$\Rightarrow \log |\tan y| = 3 \log |2 - e^x| + \log C$$

$$\Rightarrow \log |\tan y| = \log |C \cdot (2 - e^x)^3|$$

$$\Rightarrow \tan y = C (2 - e^x)^3$$

$$\text{Putting } x = 0, y = \frac{\pi}{4}, \text{ we get}$$

$$\Rightarrow \tan \frac{\pi}{4} = C(2 - e^0)^3 \quad \Rightarrow \quad 1 = C(2 - 1)^3 \quad \Rightarrow \quad 1 = C$$

$$\text{Therefore, particular solution is } \tan y = (2 - e^x)^3.$$

2. Solve:  $x \, dy - y \, dx = \sqrt{x^2 + y^2} \, dx$

[CBSE (AI) 2011]

Sol. The given differential equation can be written as

$$\frac{dy}{dx} = \frac{\sqrt{x^2 + y^2} + y}{x}, x \neq 0$$

Clearly, it is a homogeneous differential equation.

Putting  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$  in it, we get

$$v + x \frac{dv}{dx} = \frac{\sqrt{x^2 + v^2 x^2} + vx}{x} \quad \Rightarrow \quad v + x \frac{dv}{dx} = \sqrt{1 + v^2} + v$$

$$\Rightarrow x \frac{dv}{dx} = \sqrt{1 + v^2} \quad \Rightarrow \quad \frac{dv}{\sqrt{1 + v^2}} = \frac{dx}{x}$$

Integrating both sides, we get

$$\int \frac{1}{\sqrt{1 + v^2}} dv = \int \frac{1}{x} dx \quad \Rightarrow \quad \log |v + \sqrt{1 + v^2}| = \log |x| + \log C$$



$$\Rightarrow |v + \sqrt{1+v^2}| = |Cx| \Rightarrow \left| \frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}} \right| = |Cx| \quad [\because v = y/x]$$

$$\Rightarrow \{y + \sqrt{x^2 + y^2}\}^2 = C^2 x^2 \quad [\text{Squaring both sides}]$$

Hence,  $\{y + \sqrt{x^2 + y^2}\}^2 = C^2 x^2$  gives the required solution.

3. Show that the differential equation  $(xe^{y/x} + y) dx = x dy$  is homogeneous. Find the particular solution of this differential equation, given that  $x = 1$  when  $y = 1$ . [CBSE Delhi 2013]

Sol. Given differential equation is  $(x.e^{y/x} + y)dx = x dy \Rightarrow \frac{dy}{dx} = \frac{x.e^{y/x} + y}{x} \dots(i)$

Let  $F(x, y) = \frac{x.e^{y/x} + y}{x}$

$$\Rightarrow F(\lambda x, \lambda y) = \frac{\lambda x.e^{\frac{\lambda y}{\lambda x}} + \lambda y}{\lambda x} = \lambda^0 \frac{x.e^{y/x} + y}{x} = \lambda^0 F(x, y)$$

Hence, given differential equation (i) is homogeneous.

Let  $y = vx \Rightarrow \frac{dy}{dx} = v + x \cdot \frac{dv}{dx}$

Now, given differential equation (i) would become

$$v + x \cdot \frac{dv}{dx} = \frac{x.e^{v} + vx}{x} \Rightarrow v + x \cdot \frac{dv}{dx} = e^v + v \Rightarrow x \cdot \frac{dv}{dx} = e^v$$

$$\Rightarrow \frac{dv}{e^v} = \frac{dx}{x} \Rightarrow \int e^{-v} dv = \int \frac{dv}{v} \Rightarrow \frac{e^{-v}}{-1} = \log x + C$$

$$\Rightarrow -e^{-v} = \log x + C \Rightarrow -\frac{1}{e^{y/x}} = \log x + C \Rightarrow e^{y/x} \cdot \log x + C e^{y/x} + 1 = 0$$

Putting  $x = 1, y = 1$ , we get

$$\therefore e \log 1 + C e + 1 = 0 \Rightarrow C = -\frac{1}{e}$$

$\therefore$  The required particular solution is

$$e^{y/x} \cdot \log x - \frac{1}{e} e^{y/x} + 1 = 0 \quad \text{or} \quad e^{y/x} \log x - e^{y/x-1} + 1 = 0$$

4. Show that the differential equation  $\left[x \sin^2\left(\frac{y}{x}\right) - y\right]dx + x dy = 0$  is homogeneous. Find the particular solution of this differential equation, given that  $y = \frac{\pi}{4}$  when  $x = 1$ . [CBSE (AI) 2013]

Sol. Given differential equation is  $\left[x \sin^2\left(\frac{y}{x}\right) - y\right]dx + x dy = 0 \Rightarrow \frac{dy}{dx} = \frac{y - x \sin^2\left(\frac{y}{x}\right)}{x} \dots(i)$

Let  $F(x, y) = \frac{y - x \sin^2\left(\frac{y}{x}\right)}{x}$

Then  $F(\lambda x, \lambda y) = \frac{\lambda y - \lambda x \sin^2\left(\frac{\lambda y}{\lambda x}\right)}{\lambda x} = \lambda^0 \frac{y - x \sin^2\left(\frac{y}{x}\right)}{x} = \lambda^0 F(x, y)$

Hence, differential equation (i) is homogeneous.

Now, let  $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

Putting these value in (i), we get

$$v + x \frac{dv}{dx} = \frac{vx - x \sin^2 \frac{vx}{x}}{x} \Rightarrow v + x \frac{dv}{dx} = \frac{x\{v - \sin^2 v\}}{x}$$

$$\Rightarrow v + x \frac{dv}{dx} = v - \sin^2 v \Rightarrow x \frac{dv}{dx} = -\sin^2 v \Rightarrow \frac{dv}{\sin^2 v} = -\frac{dx}{x}$$

Integrating both sides, we get

$$\Rightarrow \int \operatorname{cosec}^2 v dv = -\int \frac{1}{x} dx \Rightarrow -\cot v = -\log x + C \Rightarrow \log x - \cot\left(\frac{y}{x}\right) = C \quad \dots(ii)$$

Putting  $y = \frac{\pi}{4}$  and  $x = 1$  in (ii), we get

$$\log 1 - \cot \frac{\pi}{4} = C \Rightarrow 0 - 1 = C \Rightarrow C = -1$$

Hence, particular solution is

$$\log x - \cot\left(\frac{y}{x}\right) = -1 \Rightarrow \log x - \cot\left(\frac{y}{x}\right) + 1 = 0$$

5. Find the particular solution of the differential equation  $\frac{dy}{dx} = \frac{x(2 \log x + 1)}{\sin y + y \cos y}$  given that  $y = \frac{\pi}{2}$  when  $x = 1$ . [CBSE Delhi 2014] [HOTS]

Sol. Given differential equation is  $\frac{dy}{dx} = \frac{x(2 \log x + 1)}{\sin y + y \cos y}$

$$\Rightarrow (\sin y + y \cos y) dy = x(2 \log x + 1) dx$$

$$\Rightarrow \int \sin y dy + \int y \cos y dy = 2 \int x \log x dx + \int x dx$$

$$\Rightarrow \int \sin y dy + [y \sin y - \int \sin y dy] = 2 \left[ \log x \frac{x^2}{2} - \int \frac{1}{x} \cdot \frac{x^2}{2} dx \right] + \int x dx$$

$$\Rightarrow \int \sin y dy + y \sin y - \int \sin y dy = x^2 \log x - \int x dx + \int x dx + C$$

$$\Rightarrow y \sin y = x^2 \log x + C, \text{ is general solution.} \quad \dots(i)$$

For particular solution, we put  $y = \frac{\pi}{2}$  when  $x = 1$

$$(i) \text{ becomes } \frac{\pi}{2} \sin \frac{\pi}{2} = 1 \cdot \log 1 + C \Rightarrow \frac{\pi}{2} = C \quad [\because \log 1 = 0]$$

Putting the value of  $C$  in (i), we get the required particular solution

$$y \sin y = x^2 \log x + \frac{\pi}{2}$$

6. If a curve  $y = f(x)$ , passing through the point  $(1, 2)$  is the solution of the differential equation  $2x^2 dy = (2xy + y^2) dx$ .

Find the value of  $f(1/2)$ .

Sol. We have

$$2x^2 dy = (2xy + y^2) dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{2xy + y^2}{2x^2} \quad \dots(i)$$

$$f(x, y) = \frac{2xy + y^2}{2x^2}$$

$$\Rightarrow f(\lambda x, \lambda y) = \frac{2\lambda^2 xy + \lambda^2 y^2}{2\lambda^2 x^2} = \frac{2xy + y^2}{2x^2} = \lambda^0 f(x, y)$$

$\Rightarrow$  Given differential equation is a homogeneous differential equation.

Put  $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$

Using these values in (i), we get

$$v + x \frac{dv}{dx} = \frac{2x vx + v^2 x^2}{2x^2} = \frac{2x^2 v + v^2 x^2}{2x^2}$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{2v + v^2}{2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{2v + v^2}{2} - v = \frac{2v + v^2 - 2v}{2} = \frac{v^2}{2}$$

$$\Rightarrow \frac{dv}{v^2} = \frac{1}{2} \frac{dx}{x} \Rightarrow 2 \frac{dv}{v^2} = \frac{dx}{x}$$

Integrating, we get

$$\Rightarrow \frac{-2}{v} = \log|x| + C$$

$$\Rightarrow \frac{-2x}{y} = \log|x| + C \quad \dots(ii)$$

Since (ii) passes through (1, 2).

$$\therefore \frac{-2}{2} = \log|1| + C \Rightarrow C = -1$$

From (ii)

$$\frac{-2x}{y} = \log x - 1$$

$$\Rightarrow \frac{2x}{y} = 1 - \log x \Rightarrow y = \frac{2x}{1 - \log x}$$

$$\therefore y\left(\frac{1}{2}\right) = \frac{2 \times \frac{1}{2}}{1 - \log \frac{1}{2}} = \frac{1}{1 + \log 2}$$

$$\text{i.e., } y\left(\frac{1}{2}\right) = f\left(\frac{1}{2}\right) = \frac{1}{1 + \log 2}.$$

7. If  $\cos x \frac{dy}{dx} - y \sin x = 6x$ ,  $0 < x < \frac{\pi}{2}$  and  $y\left(\frac{\pi}{3}\right) = 0$ . Find the value of  $y\left(\frac{\pi}{6}\right)$ .

Sol. We have

$$\cos x \frac{dy}{dx} - y \sin x = 6x$$

$$\Rightarrow \frac{dy}{dx} - \frac{y \sin x}{\cos x} = \frac{6x}{\cos x} \Rightarrow \frac{dy}{dx} + \left(\frac{-\sin x}{\cos x}\right)y = \frac{6x}{\cos x}$$

It is of the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \frac{-\sin x}{\cos x}$ ,  $Q = \frac{6x}{\cos x}$ .

$$\therefore \text{IF} = e^{\int P dx} = e^{\int \frac{-\sin x}{\cos x} dx} = e^{\log|\cos x|} = \cos x$$

$$\therefore \text{Solution is } y \cos x = \int \frac{6x}{\cos x} \cos x \, dx + C$$

$$\Rightarrow y \cos x = \int 6x \, dx + C = \frac{6x^2}{2} + C = 3x^2 + C$$

$$\therefore y\left(\frac{\pi}{3}\right) = 0$$

$$\Rightarrow 0 = 3 \times \frac{\pi^2}{9} + C = \frac{\pi^2}{3} + C \Rightarrow C = -\frac{\pi^2}{3}$$

$$\therefore y \cos x = 3x^2 - \frac{\pi^2}{3}$$

$$\Rightarrow y = \left(3x^2 - \frac{\pi^2}{3}\right) \sec x$$

$$\begin{aligned} \therefore y\left(\frac{\pi}{6}\right) &= \left(\frac{3\pi^2}{36} - \frac{\pi^2}{3}\right) \sec \frac{\pi}{6} \\ &= \left(\frac{\pi^2}{12} - \frac{\pi^2}{3}\right) \times \frac{2}{\sqrt{3}} = \frac{-3\pi^2}{12} \times \frac{2}{\sqrt{3}} = \frac{-\pi^2}{2\sqrt{3}} \end{aligned}$$

8. Solve:  $x \frac{dy}{dx} + y - x + xy \cot x = 0 (x \neq 0)$

[NCERT][CBSE (East) 2016]

Sol. The given differential equation  $x \frac{dy}{dx} + y - x + xy \cot x = 0 (x \neq 0)$

$$\Rightarrow \frac{dy}{dx} + \left(\cot x + \frac{1}{x}\right)y = 1 \quad \text{(Dividing both sides by } x) \quad \dots(i)$$

This is a linear differential equation of the form  $\frac{dy}{dx} + Py = Q$ , where  $P = \cot x + \frac{1}{x}$  and  $Q = 1$ .

$$\begin{aligned} \text{So, IF} &= e^{\int \left(\cot x + \frac{1}{x}\right) dx} = e^{\log |\sin x| + \log |x|} \\ &= e^{\log |x \sin x|} = x \sin x \end{aligned}$$

Multiplying both sides by IF in equation (i), we get

$$x \sin x \frac{dy}{dx} + x \sin x \left(\cot x + \frac{1}{x}\right)y = x \sin x$$

$$\Rightarrow x \sin x \frac{dy}{dx} + (x \cos x + \sin x)y = x \sin x \quad \Rightarrow \quad \frac{d}{dx}(yx \sin x) = x \sin x \quad \text{[By product rule]}$$

On integrating both sides, we get

$$\begin{aligned} \Rightarrow d(yx \sin x) &= x \sin x \, dx \\ yx \sin x &= \int x \sin x \, dx + C \quad \dots(ii) \end{aligned}$$

Let  $I = \int x \sin x \, dx = x \times (-\cos x) - \int 1 \cdot (-\cos x) \, dx$  (Using by parts)

$$I = -x \cos x + \sin x$$

Putting the value of  $I$  in (ii), we get

$$yx \sin x = -x \cos x + \sin x + C$$

$$\Rightarrow yx \sin x = \sin x - x \cos x + C$$

Hence,  $y = \frac{1}{x} - \cot x + \frac{C}{x \sin x}$  is the required solution.

9. Find the particular solution of the differential equation

$$(1 + e^{2x})dy + (1 + y^2) e^x dx = 0 \text{ given that } y = 1 \text{ when } x = 0.$$

[NCERT] [CBSE (F) 2011]

Sol. We have,  $(1 + e^{2x}) dy + (1 + y^2) e^x dx = 0$  and given that  $y = 1$ , when  $x = 0$

$$\therefore \frac{dy}{dx} = \frac{-(1 + y^2)e^x}{1 + e^{2x}} \Rightarrow \frac{dy}{-(1 + y^2)} = \frac{e^x dx}{1 + e^{2x}}$$

Integrating both sides, we get

$$-\int \frac{dy}{1 + y^2} = \int \frac{e^x dx}{1 + e^{2x}} \Rightarrow -\tan^{-1} y = \int \frac{e^x dx}{1 + (e^x)^2}$$

$$\Rightarrow -\tan^{-1} y = \int \frac{dt}{1 + t^2} \quad [\text{Putting } e^x = t \Rightarrow e^x dx = dt]$$

$$\Rightarrow -\tan^{-1} y = \tan^{-1}(t) + C$$

$$\Rightarrow -\tan^{-1} y = \tan^{-1}(e^x) + C \quad \dots(i)$$

Put  $x = 0, y = 1$  in (i), we get

$$-\tan^{-1} 1 = \tan^{-1}(e^0) + C \Rightarrow -\frac{\pi}{4} = \frac{\pi}{4} + C \Rightarrow C = -\frac{\pi}{2}$$

Putting the value of C in (i), we get

$$-\tan^{-1} y = \tan^{-1}(e^x) - \frac{\pi}{2} \Rightarrow \frac{\pi}{2} = \tan^{-1}(e^x) + \tan^{-1} y$$

Hence,  $\tan^{-1}(e^x) + \tan^{-1} y = \frac{\pi}{2}$  is the required solution.

## Questions for Practice

### ■ Objective Type Questions

1. Choose and write the correct option in each of the following questions.

(i) The degree of the differential equation is  $\left(1 + \frac{dy}{dx}\right)^3 = \left(\frac{d^2y}{dx^2}\right)^2$  [NCERT Exemplar]

(a) 1

(b) 2

(c) 3

(d) 4

(ii) The order and degree of the differential equation  $\frac{d^4y}{dx^4} = y + \left(\frac{dy}{dx}\right)^4$  are respectively

(a) 4, 1

(b) 4, 2

(c) 2, 2

(d) 2, 4

(iii) The integrating factor of the differential equation  $x \frac{dy}{dx} - y = 2x^2$  is

(a)  $e^{-x}$

(b)  $e^{-y}$

(c)  $\frac{1}{x}$

(d)  $x$

(iv) Solution of the differential equation  $x \frac{dy}{dx} + y = x e^x$  is

(a)  $xy = e^x(1 - x) + C$

(b)  $xy = e^x(x + 1) + C$

(c)  $xy = e^y(y - 1) + C$

(d)  $xy = e^x(x - 1) + C$

(v) The general solution of the differential equation  $e^x dy + (y e^x + 2x) dx = 0$  is

(a)  $x e^y + x^2 = C$

(b)  $x e^y + y^2 = C$

(c)  $y e^x + x^2 = C$

(d)  $y e^y + x^2 = C$

### ■ Conceptual Questions

2. What is the degree of the following differential equation:

$$5x\left(\frac{dy}{dx}\right)^2 - \frac{d^2y}{dx^2} - 6y = \log x ? \quad [\text{CBSE Delhi 2010}]$$

3. Write the degree of the following differential equation:

$$x^3\left(\frac{d^2y}{dx^2}\right)^2 + x\left(\frac{dy}{dx}\right)^4 = 0 \quad [\text{CBSE Delhi 2013}]$$

4. Write the sum of the order and degree of the following differential equation:

$$\left(\frac{d^2y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^3 + x^4 = 0 \quad [\text{CBSE (F) 2015}]$$

5. Find the product of the order and degree of  $x\left(\frac{d^2y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^2 + y^2 = 0$ . [CBSE Chennai 2015]

6. Write the integrating factor of  $\frac{dy}{dx}(x \log x) + y = 2 \log x$ . [CBSE Punchkula 2015]

7. Solve:  $e^{dy/dx} = x^2$

8. State whether  $y = e^{-x}(x + a)$  is the solution of differential equation:

$$\frac{dy}{dx} + y = e^{-x}$$

9. Solve:  $\frac{dy}{dx} - \frac{y(x+1)}{x} = 0$

### ■ Very Short Answer Questions

10. Write the general solution of the differential equation  $\frac{dy}{dx} = \frac{y}{x}$ .

11. Write the integrating factor of  $\frac{dy}{dx} + y = \frac{1+y}{x}$ .

12. Given that  $\frac{dy}{dx} = e^{-2y}$  and  $y = 0$  when  $x = 5$ . Find the value of  $x$  when  $y = 3$ .

13. Find the general solution of the differential equation  $\frac{dy}{dx} = 2^{y-x}$ .

### ■ Short Answer Questions

14. Solve the following differential equation:

$$\cos^2 x \frac{dy}{dx} + y = \tan x \quad [\text{CBSE Delhi 2008, 2011; (AI) 2009}]$$

15. Solve the differential equation:

$$(x^2 + 1) \frac{dy}{dx} + 2xy = \sqrt{x^2 + 4} \quad [\text{CBSE (AI) 2010}]$$

16. Solve the differential equation:

$$(x^2 + 3xy + y^2)dx - x^2dy = 0 \text{ given that } y = 0, \text{ when } x = 1 \quad [\text{CBSE (East) 2016}]$$

17. Solve the differential equation:  $(x+1)\frac{dy}{dx} - y = e^{3x}(x+1)^3$  [CBSE (South) 2016]

18. Find the particular solution of this differential equation  $x^2 \frac{dy}{dx} - xy = 1 + \cos\left(\frac{y}{x}\right), x \neq 0$ . Find the particular solution of this differential equation, given that when  $x = 1, y = \frac{\pi}{2}$ . [CBSE (F) 2013]



19. Find the particular solution of the differential equation  $\frac{dy}{dx} = \frac{xy}{x^2 + y^2}$  given that  $y = 1$ , when  $x = 0$ .  
[CBSE Delhi 2015]

20. Solve the differential equation  $\frac{dy}{dx} + y \cot x = 2 \cos x$ , given that  $y = 0$ , when  $x = \frac{\pi}{2}$ .  
[CBSE (F) 2014]

21. Solve the differential equation  $(x^2 - yx^2) dy + (y^2 + x^2y^2) dx = 0$ , given that  $y = 1$ , when  $x = 1$ .  
[CBSE (F) 2014]

22. Solve the differential equation:

$$\frac{dy}{dx} = \frac{x+y}{x-y} \quad \text{[CBSE 2019 (65/3/1)]}$$

23. Solve the differential equation:

$$(1 + x^2) dy + 2xy dx = \cot x dx \quad \text{[CBSE 2019 (65/3/1)]}$$

24. Find the general solution of the differential equation  $x^2y dx - (x^3 + y^3) dy = 0$ . [CBSE 2020 (65/3/1)]

25. Find the particular solution of the differential equation  $\log \left( \frac{dy}{dx} \right) = 3x + 4y$ , given that  $y = 0$  when  $x = 0$ .  
[CBSE (AI) 2014]

#### ■ Long Answer Questions

26. Solve the following differential equation, given that  $y = 0$ , when  $x = \frac{\pi}{4}$ :

$$\sin 2x \frac{dy}{dx} - y = \tan x \quad \text{[CBSE Chennai 2015]}$$

27. Solve the following differential equation:

$$\left[ y - x \cos \left( \frac{y}{x} \right) \right] dy + \left[ y \cos \left( \frac{y}{x} \right) - 2x \sin \left( \frac{y}{x} \right) \right] dx = 0 \quad \text{[CBSE (F) 2015]}$$

28. Find the particular solution of the differential equation  $(1 + x^2) \frac{dy}{dx} = (e^{m \tan^{-1} x} - y)$  given that  $y = 1$ , when  $x = 0$ .  
[CBSE Panchkula 2015]

29. Find the particular solution of the following differential equation:

$$xy \frac{dy}{dx} = (x+2)(y+2); y = -1 \text{ when } x = 1 \quad \text{[CBSE Delhi 2012]}$$

30. Find the particular solution of the differential equation

$$\frac{dx}{dy} + x \cot y = 2y + y^2 \cot y, (y \neq 0) \text{ given that } x = 0 \text{ when } y = \frac{\pi}{2}. \quad \text{[CBSE (AI) 2013]}$$

31. Find the particular solution of the differential equation  $x(1 + y^2)dx - y(1 + x^2)dy = 0$  given that  $y = 1$  when  $x = 0$ .  
[CBSE (AI) 2014]

32. Find the particular solution of the differential equation satisfying the given conditions

$$x^2 dy + (xy + y^2) dx = 0; y = 1 \text{ when } x = 1. \quad \text{[CBSE Delhi 2010]}$$

33.  $(x^2 + y^2) dy = xy dx$ . If  $y(1) = 1$  and  $y(x_0) = e$ , then find the value of  $x_0$ . [CBSE Bhubaneswar 2015]

34. Find the particular solution of the differential equation  $(y - \sin x)dx + (\tan x)dy = 0$  satisfying the condition that  $y = 0$  when  $x = 0$ .  
[CBSE Gurwahati 2015]

35. If  $\frac{y dx - x dy}{y} = 0$ ,  $x, y > 0$  and  $y(1) = x^2$  then find the value of  $y(5)$

## Answers

1. (i) (b)      (ii) (a)      (iii) (c)      (iv) (d)      (v) (c)
2. 1      3. 2      4. 4      5. 4      6.  $\log x$
7.  $y = 2(x \log x - x) + C$       8. Yes      9.  $y = x e^{x+C}$       10.  $y = Cx$       11.  $\frac{e^x}{x}$
12.  $\frac{e^6 + 9}{2}$       13.  $2^{-x} - 2^{-y} = C$       14.  $y = \tan x - 1 + C e^{-\tan x}$
15.  $(x^2 + 1)y = \frac{x}{2}\sqrt{x^2 + 4} + 2\log|x + \sqrt{x^2 + 4}|$       16.  $y = \frac{x \log|x|}{1 - \log|x|}$
17.  $\frac{y}{x+1} = (x+1)\frac{e^{3x}}{3} - \frac{e^{3x}}{9} + C$       18.  $\tan\left(\frac{y}{2x}\right) = -\frac{1}{2x^2} + \frac{3}{2}$       19.  $-\frac{x^2}{2y^2} + \log|y| = 0$
20.  $2y \sin x = -(1 + \cos 2x)$       21.  $\log|y| + \frac{1}{y} = -\frac{1}{x} + x + 1$
22.  $\tan^{-1}\left(\frac{y}{x}\right) = \frac{1}{2}\log(x^2 + y^2) + C$       23.  $y = \frac{1}{1+x^2}\log|\sin x| + \frac{C}{1+x^2}$       24.  $\log|y| = \frac{x^3}{3y^3} + C$
25.  $4e^{3x} + 3e^{-4y} = 7$       26.  $y = \tan x - \sqrt{\tan x}$       27.  $y^2 - 2x^2 \cos\left(\frac{y}{x}\right) = C$
28.  $y e^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{m+1} + \frac{m}{m+1}$       29.  $x + 2 \log|x| - 2$       30.  $x \sin y = y^2 \sin y - \frac{\pi^2}{4}$
31.  $y^2 = 2x^2 + 1$       32.  $3x^2y = y + 2x$       33.  $x_0 = \sqrt{3}e$
34.  $y = \frac{1}{2}\sin x$       35. 25

