

11

Applications of Definite Integrals

Definite integrals have a wide range of applications. In this chapter, we shall use definite integrals in computing the *areas of bounded regions*.

11.1 AREAS OF BOUNDED REGIONS

If the function f is continuous and non-negative in the closed interval $[a, b]$, then the area of the region below the curve $y = f(x)$, above the x -axis and between the ordinates $x = a$ and $x = b$ or briefly the area of the region bounded by the curve $y = f(x)$, the x -axis and the

ordinates $x = a$, $x = b$ is given by $\int_a^b f(x) dx$ or $\int_a^b y dx$.

Proof. Let AB be the curve $y = f(x)$ between $AC(x = a)$ and $BD(x = b)$, then the required area is the area of the shaded region $ACDB$.

Let $P(x, y)$ be a point on the curve $y = f(x)$ and $Q(x + \delta x, y + \delta y)$ be a neighbouring point on the curve, then $MP = y$, $NQ = y + \delta y$ and $MN = \delta x$. Let A be the area of the region $ACMP$ and $A + \delta A$ be the area of the region $ACNQ$, then $\delta A = \text{area of region } PMNQ$.

Area of rectangle $PMNR = y\delta x$ and area of rectangle $SMNQ = (y + \delta y)\delta x$.

From fig. 11.1, area of rectangle $PMNR \leq \text{area of region } PMNQ \leq \text{area of rectangle } SMNQ$

$$\Rightarrow y\delta x \leq \delta A \leq (y + \delta y)\delta x$$

$$\Rightarrow y \leq \frac{\delta A}{\delta x} \leq y + \delta y$$

...(i)

When $P \rightarrow Q$, $\delta x \rightarrow 0$, $\delta y \rightarrow 0$ and $\frac{\delta A}{\delta x} \rightarrow \frac{dA}{dx}$.

From (i), $\lim_{\delta x \rightarrow 0} y \leq \lim_{\delta x \rightarrow 0} \frac{\delta A}{\delta x} \leq \lim_{\delta y \rightarrow 0} (y + \delta y)$

$$\Rightarrow y \leq \frac{dA}{dx} \leq y \Rightarrow y = \frac{dA}{dx}.$$

Integrating both sides w.r.t. x between the limits a to b , we get

$$\begin{aligned} \int_a^b y dx &= \int_a^b \frac{dA}{dx} dx = [A]_a^b \\ &= (\text{value of area } A \text{ when } x = b) - (\text{value of area } A \text{ when } x = a) \\ &= \text{area } ACDB - 0 = \text{area } ACDB. \end{aligned}$$




Fig. 11.1.

If a function f is continuous and non-positive in the closed interval $[a, b]$, then the curve $y = f(x)$ lies below the x -axis and the definite integral $\int_a^b f(x) dx$ is negative. Since the area of a region is always non-negative, the area of the region bounded by the curve $y = f(x)$, the x -axis and the ordinates $x = a$, $x = b$ is given by $\left| \int_a^b f(x) dx \right|$ or $\left| \int_a^b y dx \right|$.




Fig. 11.2.

Hence, if the curve $y = f(x)$ is continuous and does not cross the x -axis, then the area of the region bounded by the curve $y = f(x)$, the x -axis and the ordinates $x = a$ and $x = b$ is given by $\left| \int_a^b f(x) dx \right|$ or

$$\left| \int_a^b y dx \right|.$$

Similarly, if the curve $x = g(y)$ is continuous and does not cross the y -axis, then the area of the region bounded by the curve $x = g(y)$, the y -axis and the abscissae $y = c$, $y = d$ is given by

$$\left| \int_c^d g(y) dy \right| \text{ or } \left| \int_c^d x dy \right|.$$




Fig. 11.3.

Remark. It may be noted that when sign of $f(x)$ is not known, then $\int_a^b f(x) dx$ may not represent the area enclosed between the curve $y = f(x)$, the x -axis and the ordinates $x = a$ and $x = b$, whereas $\int_a^b |f(x)| dx$ equals the area enclosed between the graph of the curve $y = f(x)$, the x -axis and the ordinates $x = a$ and $x = b$.

For example, let us consider the integrals $\int_{-1}^1 x dx$ and $\int_{-1}^1 |x| dx$.

$$\begin{aligned} \text{First integral} &= \int_{-1}^1 x dx = \left[\frac{x^2}{2} \right]_{-1}^1 = \frac{1}{2} (1^2 - (-1)^2) = 0, \text{ whereas second integral} \\ &= \int_{-1}^1 |x| dx = \int_{-1}^0 (-x) dx + \int_0^1 x dx \end{aligned}$$

(Common sense suggests this division as $|x| = -x$ in $[-1, 0]$ and $|x| = x$ in $[0, 1]$).

$$= \left[-\frac{x^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_0^1 = -\frac{1}{2} (0 - 1) + \frac{1}{2} (1 - 0) = 1.$$




Fig. 11.4.

Clearly, the area enclosed between $y = x$, the x -axis and the ordinates $x = -1$ and $x = 1$ is not zero.

It follows that if the graph of a function f is continuous in $[a, b]$ and crosses the x -axis at finitely many points in $[a, b]$, then the area enclosed between the graph of the

curve $y = f(x)$, the x -axis and the ordinates $x = a$, $x = b$ is given by $\int_a^b |f(x)| dx$ or $\int_a^b |y| dx$.

11.1.1 Area bounded between curves

If $f(x)$, $g(x)$ are both continuous in $[a, b]$ and $0 \leq g(x) \leq f(x)$ for all $x \in [a, b]$, then the area of the region between the graphs of $y = f(x)$, $y = g(x)$ and the ordinates $x = a$, $x = b$ is given by

$$\begin{aligned} \int_a^b f(x) dx - \int_a^b g(x) dx \\ = \int_a^b (f(x) - g(x)) dx. \end{aligned}$$

Similarly, the area of the region between the graphs of $x = f(y)$, $x = g(y)$ and the abscissae $y = c$, $y = d$ is

$$\text{given by } \int_c^d (f(y) - g(y)) dy.$$




Fig. 11.5.




Fig. 11.6.

Remarks

1. If $f(x)$, $g(x)$ are both continuous in $[a, b]$ and $g(x) \leq f(x)$ for all $x \in [a, b]$, then the above formula also holds when one or both of the curves $y = f(x)$ and $y = g(x)$ lie partially or completely below the x -axis.
2. If the graphs of the curves $y = f(x)$ and $y = g(x)$ cross each other at finitely many points, then the area enclosed between the graphs of the two curves and the

ordinates $x = a$ and $x = b$ is given by $\int_a^b |f(x) - g(x)| dx$.

3. Similarly, the area of the region between the graphs of $x = f(y)$, $x = g(y)$ and the

abscissae $y = c$, $y = d$ is given by $\int_c^d |f(y) - g(y)| dy$.

ILLUSTRATIVE EXAMPLES

Example 1. Find the area of the region bounded by $y^2 = 4x$, $x = 1$, $x = 4$ and the x -axis in the first quadrant.

Solution. The given curve is $y^2 = 4x$ which represents a right hand parabola with vertex at $(0, 0)$. The area bounded by $y^2 = 4x$, $x = 1$, $x = 4$ and the x -axis is shown shaded in the figure.

$$\begin{aligned}
 \text{Required area} &= \int_1^4 y \, dx = \int_1^4 2\sqrt{x} \, dx \\
 (\because y^2 = 4x \Rightarrow y &= 2\sqrt{x} \text{ in the first quadrant}) \\
 &= 2 \cdot \left[\frac{x^{3/2}}{\frac{3}{2}} \right]_1^4 = \frac{4}{3} [4^{3/2} - 1^{3/2}] \text{ sq. units} \\
 &= \frac{4}{3} [8 - 1] \text{ sq. units} = \frac{28}{3} \text{ sq. units.}
 \end{aligned}$$




Fig. 11.7.

Example 2. Draw a rough sketch of the curve $x^2 + y = 9$ and find the area enclosed by the curve, the x -axis and the lines $x + 1 = 0$ and $x - 2 = 0$. (I.S.C. 2009)

Solution. The given curve is $x^2 + y = 9$

It can be written as $x^2 = 9 - y$

$$\Rightarrow (x - 0)^2 = -(y - 9)$$

which represents a downward parabola with vertex at $(0, 9)$.

The parabola meets the x -axis i.e. $y = 0$ at $x^2 = 9$ i.e. at $x = -3, 3$.

A rough sketch of the curve is shown in fig. 11.8.

The given lines are $x + 1 = 0$ and $x - 2 = 0$ i.e. $x = -1$ and $x = 2$.

The area enclosed by the curve, the x -axis and the given lines is shown shaded in fig. 11.8.




Fig. 11.8.

$$\begin{aligned}
 \therefore \text{Required area} &= \int_{-1}^2 y \, dx = \int_{-1}^2 (9 - x^2) \, dx \quad \text{(using (i))} \\
 &= \left[9x - \frac{x^3}{3} \right]_{-1}^2 = \left(\left(18 - \frac{8}{3} \right) - \left(-9 + \frac{1}{3} \right) \right) \text{ sq. units} \\
 &= \left(27 - \frac{8}{3} - \frac{1}{3} \right) \text{ sq. units} = 24 \text{ sq. units.}
 \end{aligned}$$

Example 3. Determine the area enclosed between the curve $y = 4x - x^2$ and the x -axis.

Solution. Given curve is $y = 4x - x^2$.

It can be written as $x^2 - 4x = -y \Rightarrow (x - 2)^2 = -(y - 4)$

which represents a downward parabola with vertex at $(2, 4)$.

The parabola meets x -axis i.e. $y = 0$ at $4x - x^2 = 0$ i.e. at $x = 0, x = 4$.

$\therefore$ The area enclosed between the curve and the x -axis

$$\begin{aligned}
 &= \int_0^4 y \, dx = \int_0^4 (4x - x^2) \, dx = \left[4 \cdot \frac{x^2}{2} - \frac{x^3}{3} \right]_0^4 \\
 &= \left(32 - \frac{64}{3} \right) - (0 - 0) = \frac{32}{3} \text{ sq. units.}
 \end{aligned}$$




Fig. 11.9.

Alternatively. Since the parabola is symmetrical about the line $x = 2$,

$$\begin{aligned}\text{required area} &= 2 \int_0^2 (4x - x^2) dx = 2 \left[4 \cdot \frac{x^2}{2} - \frac{x^3}{3} \right]_0^2 \\ &= 2 \left[\left(8 - \frac{8}{3} \right) - (0 - 0) \right] \text{ sq. units} = 2 \cdot \frac{16}{3} \text{ sq. units} = \frac{32}{3} \text{ sq. units.}\end{aligned}$$

Remark. In case of symmetrical closed area, find the area of the smallest part and multiply the result by the number of symmetrical parts.

Example 4. Draw a rough sketch of the curve $y^2 + 1 = x$, $x \leq 2$. Find the area enclosed by the curve and the line $x = 2$. (I.S.C. 2008)

Solution. Given curve is $y^2 + 1 = x$.

It can be written as $y^2 = x - 1$, which represents a right hand parabola with vertex at A(1, 0).

The parabola meets the line $x = 2$ at when $y^2 = 1$ i.e. $y = 1, -1$.

A rough sketch of the curve $y^2 + 1 = x$, $x \leq 2$ is shown in fig. 11.10. The area bounded by the curve $y^2 = x - 1$ and the line $x = 2$ is shown shaded in the figure. Since the given area is symmetrical about x -axis,




Fig. 11.10.

required area = 2(area of the region bounded by the curve $y^2 = x - 1$, the x -axis and the line $x = 2$)

$$\begin{aligned}&= 2 \int_1^2 y dx = 2 \int_1^2 \sqrt{x-1} dx \quad (\because y^2 = x - 1 \Rightarrow y = \sqrt{x-1} \text{ in the first quadrant}) \\ &= 2 \cdot \left[\frac{(x-1)^{3/2}}{\frac{3}{2}} \right]_1^2 = \frac{4}{3} [1^{3/2} - 0] \text{ sq. units} = \frac{4}{3} \text{ sq. units.}\end{aligned}$$

Example 5. Draw a rough sketch of the curve $y = x^2 - 5x + 6$ and find the area bounded by the curve and the x -axis. (I.S.C. 2010)

Solution. The given curve is $y = x^2 - 5x + 6$.

It can be written as $x^2 - 5x + \frac{25}{4} = y + \frac{1}{4}$

$\Rightarrow \left(x - \frac{5}{2} \right)^2 = y + \frac{1}{4}$, which represents an

upward parabola with vertex at $\left(\frac{5}{2}, -\frac{1}{4} \right)$.

A rough sketch of the curve is shown in fig. 11.11.

The parabola meets the x -axis i.e. $y = 0$ at

$$x^2 - 5x + 6 = 0 \text{ i.e. at } (x - 2)(x - 3) = 0$$

i.e. at $x = 2$, $x = 3$.

As the required portion of the curve lies below x -axis, y is negative.




Fig. 11.11.

$$\begin{aligned}\therefore \text{ Required area} &= \left| \int_2^3 y dx \right| = \left| \int_2^3 (x^2 - 5x + 6) dx \right| \\ &= \left| \left[\frac{x^3}{3} - 5 \cdot \frac{x^2}{2} + 6x \right]_2^3 \right| = \left| \left(9 - \frac{45}{2} + 18 \right) - \left(\frac{8}{3} - 10 + 12 \right) \right| \text{ sq. units} \\ &= \left| \frac{9}{2} - \frac{14}{3} \right| \text{ sq. units} = \left| -\frac{1}{6} \right| \text{ sq. units} = \frac{1}{6} \text{ sq. units.}\end{aligned}$$

Example 6. Find the area of the region bounded by the curve $y^2 = 4x$, y -axis and the line $y = 3$.

Solution. The given curve is $y^2 = 4x$ which represents a right hand parabola with vertex $(0, 0)$. The area bounded by the curve $y^2 = 4x$, y -axis and the line $y = 3$ is shown shaded in fig. 11.12.

$$\begin{aligned}\text{Required area} &= \int_0^3 x \, dy = \int_0^3 \frac{y^2}{4} \, dy \\ &\quad (\because y^2 = 4x \Rightarrow x = \frac{y^2}{4}) \\ &= \frac{1}{4} \cdot \left[\frac{y^3}{3} \right]_0^3 = \frac{1}{12} [27 - 0] \\ &= \frac{9}{4} \text{ sq. units.}\end{aligned}$$




Fig. 11.12.

Example 7. Find the area of the region bounded by the curve $y = x^2$ and the line $y = 4$.

Solution. The given curve is $y = x^2$ which represents an upward parabola with vertex at $(0, 0)$. The area bounded by the curve and the line $y = 4$ is shown shaded in fig. 11.13.

Since the area is symmetrical about y -axis,

required area = 2 (area of the region bounded by $y = x^2$, the y -axis and the line $y = 4$)

$$\begin{aligned}&= 2 \int_0^4 x \, dy = 2 \int_0^4 \sqrt{y} \, dx \\ &\quad (\because x^2 = y \Rightarrow x = \sqrt{y} \text{ in the first quadrant}) \\ &= 2 \cdot \left[\frac{x^{3/2}}{\frac{3}{2}} \right]_0^4 = \frac{4}{3} [4^{3/2} - 0] = \frac{4}{3} [8 - 0] = \frac{32}{3} \text{ sq. units.}\end{aligned}$$




Fig. 11.13.

Example 8. Sketch and shade the area of the region lying in the first quadrant and bounded by $y = 9x^2$, $x = 0$, $y = 1$ and $y = 4$. Find the area of the shaded region. (I.S.C. 2004)

Solution. The given curve is $y = 9x^2$. It can be written

as $x^2 = \frac{y}{9}$ which represents an upward parabola with vertex at $(0, 0)$. The area lying in the first quadrant and bounded by $y = 9x^2$, $x = 0$, $y = 1$ and $y = 4$ is shown shaded in fig. 11.14.

$$\begin{aligned}\text{The required area} &= \int_1^4 x \, dy = \int_1^4 \sqrt{\frac{y}{9}} \, dy \\ &\quad (\because x^2 = \frac{y}{9} \Rightarrow x = \sqrt{\frac{y}{9}} \text{ in the first quadrant.}) \\ &= \frac{1}{3} \left[\frac{y^{3/2}}{\frac{3}{2}} \right]_1^4 = \frac{2}{9} [4^{3/2} - 1^{3/2}] \\ &= \frac{2}{9} (8 - 1) = \frac{14}{9} \text{ sq. units.}\end{aligned}$$




Fig. 11.14.

Example 9. Find the area bounded by the curve $x = 8 + 2y - y^2$, the y -axis and the lines $y = -1$, $y = 3$.

Solution. The given curve is $x = 8 + 2y - y^2$.

It can be written as

$$y^2 - 2y = -x + 8$$

$\Rightarrow (y - 1)^2 = -(x - 9)$ which represents a left hand parabola with vertex at $(9, 1)$.

Required area

$$\begin{aligned} &= \int_{-1}^3 x \, dy = \int_{-1}^3 (8 + 2y - y^2) \, dy \\ &= \left[8y + 2 \cdot \frac{y^2}{2} - \frac{y^3}{3} \right]_{-1}^3 \\ &= (24 + 9 - 9) - \left(-8 + 1 + \frac{1}{3} \right) = \frac{92}{3} \text{ sq. units.} \end{aligned}$$




Fig. 11.15.

Example 10. Draw a rough sketch of the graph of the function $y = 2\sqrt{1-x^2}$, $x \in [0, 1]$ and evaluate the area enclosed between the curve and the axes.

Solution. The given curve is $y = 2\sqrt{1-x^2}$

$\Rightarrow \frac{y^2}{4} = 1 - x^2 \Rightarrow \frac{x^2}{1} + \frac{y^2}{4} = 1$, which represents an ellipse of the second standard form. Hence, the given equation $y = 2\sqrt{1-x^2}$ represents the portion of the ellipse lying in the first quadrant. Its rough sketch is shown in fig. 11.16.

The required area = the area of the shaded region

$$\begin{aligned} &= \int_0^1 y \, dx = \int_0^1 2\sqrt{1-x^2} \, dx \\ &= 2 \left[\frac{x\sqrt{1-x^2}}{2} + \frac{1}{2} \sin^{-1} x \right]_0^1 = \left[x\sqrt{1-x^2} + \sin^{-1} x \right]_0^1 \\ &= (0 + \sin^{-1} 1) - (0 + \sin^{-1} 0) = \frac{\pi}{2} \text{ sq. units.} \end{aligned}$$




Fig. 11.16.

Example 11. Find the area bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the ordinates $x = 0$ and $x = ae$ where $b^2 = a^2(1 - e^2)$ and $0 < e < 1$.

Solution. The given ellipse is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\Rightarrow \frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$\Rightarrow y = \pm \frac{b}{a} \sqrt{a^2 - x^2}$$

The required area is shown shaded in fig. 11.17.

Since the area is symmetrical about the x -axis,

required area = 2 (area of the region bounded by the given ellipse, x -axis and the lines $x = 0$ and $x = ae$)

$$= 2 \int_0^{ae} y \, dx = 2 \int_0^{ae} \frac{b}{a} \sqrt{a^2 - x^2} \, dx$$




Fig. 11.17.

($\because y \geq 0$ in the first quadrant)

$$\begin{aligned}
 &= 2 \frac{b}{a} \left[\frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]_0^{ae} \\
 &= \frac{b}{a} \left[\left(ae \sqrt{a^2 - a^2 e^2} + a^2 \sin^{-1} e \right) - \left(0 + \frac{a^2}{2} \sin^{-1} 0 \right) \right] \\
 &= ab \left(e \sqrt{1 - e^2} + \sin^{-1} e \right).
 \end{aligned}$$

Example 12. The area between $x = y^2$ and $x = 4$ is divided into two equal parts by the line $x = a$, find the value of a .

Solution. The given curve is $y^2 = x$ which represents a right hand parabola with vertex $(0, 0)$.

The area bounded by the parabola and the line $x = 4$ is shown shaded in the fig. 11.18.




Fig. 11.18.

This area $= 2 \int_0^4 y \, dx = 2 \int_0^4 \sqrt{x} \, dx$

$$= 2 \cdot \left[\frac{x^{3/2}}{\frac{3}{2}} \right]_0^4 = \frac{4}{3} (4^{3/2} - 0) = \frac{4}{3} (8 - 0) = \frac{32}{3}.$$

Since the line $x = a$ divides this area into two equal parts, therefore,

$$\begin{aligned}
 2 \int_0^a \sqrt{x} \, dx &= \frac{1}{2} \cdot \frac{32}{3} \Rightarrow \int_0^a \sqrt{x} \, dx = \frac{8}{3} \\
 \Rightarrow \left[\frac{x^{3/2}}{\frac{3}{2}} \right]_0^a &= \frac{8}{3} \Rightarrow \frac{2}{3} (a^{3/2} - 0) = \frac{8}{3} \\
 \Rightarrow a^{3/2} &= 4 \Rightarrow a = 4^{2/3} \\
 \Rightarrow a &= \sqrt[3]{16}.
 \end{aligned}$$

Example 13. Find the area of the smaller region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the straight line $\frac{x}{a} + \frac{y}{b} = 1$.

Solution. The given ellipse is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow \frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$

$$\Rightarrow y = \frac{b}{a} \sqrt{a^2 - x^2} \quad (\because \text{In first quadrant, } y \geq 0)$$

The given line is $\frac{x}{a} + \frac{y}{b} = 1$

$$\Rightarrow \frac{y}{b} = 1 - \frac{x}{a} = \frac{a-x}{a}$$

$$\Rightarrow y = \frac{b}{a} (a-x)$$

The area of the smaller region bounded by the given ellipse and the given line is shown shaded in the figure.




Fig. 11.19.

$$\begin{aligned}
 \text{Required area} &= \int_0^a \left(\frac{b}{a} \sqrt{a^2 - x^2} - \frac{b}{a} (a-x) \right) dx && \text{(Article 11.1.1)} \\
 &= \frac{b}{a} \left[\frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} - ax + \frac{x^2}{2} \right]_0^a
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{b}{a} \left[\left(0 + \frac{a^2}{2} \sin^{-1} 1 - a^2 + \frac{a^2}{2} \right) - \left(0 + \frac{a^2}{2} \sin^{-1} 0 - 0 + 0 \right) \right] \\
 &= \frac{b}{a} \left[\left(\frac{a^2}{2} \cdot \frac{\pi}{2} - \frac{a^2}{2} \right) - 0 \right] = \frac{1}{4} (\pi - 2) ab \text{ sq. units.}
 \end{aligned}$$

Example 14. Find the area of the region included between the curve $4y = 3x^2$ and the line $2y = 3x + 12$.

Solution. The given curve is $4y = 3x^2$... (i)

It can be written as $y = \frac{3}{4}x^2$, which represents an upward parabola with vertex at $(0, 0)$.

The given line is $3x - 2y + 12 = 0$

$$\Rightarrow y = \frac{3x + 12}{2} \quad \dots (ii)$$

Solving (i) and (ii), we get

$$\frac{3x + 12}{2} = \frac{3}{4}x^2$$

$$\Rightarrow 6x + 24 = 3x^2$$

$$\Rightarrow x^2 - 2x - 8 = 0 \Rightarrow (x + 2)(x - 4) = 0$$

$$\Rightarrow x = -2, x = 4.$$

$\therefore$ The points of intersection are $P(-2, 3)$ and $Q(4, 12)$.

$\therefore$ Required area = area of the shaded region

$$= \int_{-2}^4 \left(\frac{3x + 12}{2} - \frac{3}{4}x^2 \right) dx$$

[Article 11.1.1]

$$= \left[\frac{3}{2} \cdot \frac{x^2}{2} + 6x - \frac{3}{4} \cdot \frac{x^3}{3} \right]_{-2}^4 = \frac{1}{4} [3x^2 + 24x - x^3]_{-2}^4$$

$$= \frac{1}{4} [(48 + 96 - 64) - (12 - 48 + 8)]$$

$$= \frac{1}{4} \cdot 108 = 27 \text{ sq. units.}$$




Fig. 11.20.

Example 15. Find the area enclosed by the parabola $y^2 = x$ and the line $y + x = 2$.

Solution. The given parabola is $y^2 = x$... (i)

It represents a right hand parabola with vertex at $(0, 0)$.

The given line is $y + x = 2$

$$\text{i.e. } x = 2 - y \quad \dots (ii)$$

Solving (i) and (ii), we get

$$y^2 = 2 - y \Rightarrow y^2 + y - 2 = 0$$

$$\Rightarrow (y - 1)(y + 2) = 0 \Rightarrow y = 1, -2$$

When $y = 1$, $x = 1$, when $y = -2$, $x = 4$

The points of intersection are $P(1, 1)$ and $Q(4, -2)$.

The required area = area of the shaded region

$$= \int_{-2}^1 ((2 - y) - y^2) dy = \left[2y - \frac{y^2}{2} - \frac{y^3}{3} \right]_{-2}^1$$




Fig. 11.21.

$$\begin{aligned}
 &= \left(2 - \frac{1}{2} - \frac{1}{3}\right) - \left(-4 - 2 + \frac{8}{3}\right) \\
 &= 2 - \frac{1}{2} - \frac{1}{3} + 6 - \frac{8}{3} = 4\frac{1}{2} \text{ sq. units.}
 \end{aligned}$$

Example 16. Find the area bounded by the curve $y = 2x - x^2$ and the line $y = x$. (I.S.C. 2013)

Solution. The given curve is $y = 2x - x^2$... (i)

It can be written as $y = -(x^2 - 2x + 1) + 1$

i.e. $(y - 1) = -(x - 1)^2$, which represents a downward parabola with vertex at (1, 1).

The given line is $y = x$

Solving (i) and (ii), we get

$$x = 2x - x^2 \Rightarrow x^2 - x = 0$$

$$\Rightarrow x = 0, 1.$$

$\therefore$ The points of intersection are O (0, 0) and P (1, 1).

$\therefore$ Required area = area of the shaded region

$$\begin{aligned}
 &= \int_0^1 ((2x - x^2) - x) dx = \int_0^1 (x - x^2) dx \\
 &= \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \left(\frac{1}{2} - \frac{1}{3} \right) - (0 - 0) = \frac{1}{6} \text{ sq. units.}
 \end{aligned}$$




Fig. 11.22.

Example 17. Find the area enclosed by the curve $y = -x^2$ and the line $x + y + 2 = 0$.

Solution. The given curve is $y = -x^2$... (i)

It represents a downward parabola with vertex O(0, 0).

The given line is $x + y + 2 = 0$

$$\Rightarrow y = -(x + 2) \quad \dots (ii)$$

Solving (i) and (ii), we get

$$-x^2 = -(x + 2) \Rightarrow x^2 - x - 2 = 0$$

$$\Rightarrow (x + 1)(x - 2) = 0 \Rightarrow x = -1, 2.$$

When $x = -1$, $y = -1$ and when $x = 2$, $y = -4$.

$\therefore$ The points of intersection are P(-1, -1) and Q(2, -4).

The required area is shown shaded in fig. 11.23. We note that the required area lies below the x-axis, therefore,




Fig. 11.23.

$$\begin{aligned}
 \text{required area} &= \left| \int_{-1}^2 (-(x + 2) - (-x^2)) dx \right| \\
 &= \left| \left[-\left(\frac{x^2}{2} + 2x\right) + \frac{x^3}{3} \right]_{-1}^2 \right| \\
 &= \left| \left(-6 + \frac{8}{3}\right) - \left(-\left(\frac{1}{2} + 2\right) - \frac{1}{3}\right) \right| = \frac{9}{2} \text{ sq. units.}
 \end{aligned}$$

[Article 11.1.1]

EXERCISE 11.1

1. (i) Find the area bounded by the curve $y = x^2$, the x -axis and the ordinates $x = 1$ and $x = 3$.
 (ii) Find the area of the region bounded by $y^2 = x - 2$ and the lines $x = 4$ and $x = 6$.
 (iii) Find the area of the region bounded by $x^2 = 4y$, $y = 2$, $y = 4$ and the y -axis in the first quadrant.
 (iv) Find the area of the region bounded by $x^2 = y - 3$ and the lines $y = 4$ and $y = 6$.
2. Using integration, find the area of the region bounded between the line $x = 2$ and the parabola $y^2 = 8x$.
3. Using integration, find the area of the region bounded by the line $2y = -x + 8$, x -axis and the lines $x = 2$ and $x = 4$.
4. Make a rough sketch of the graph of the function $f(x) = 9 - x^2$, $0 \leq x \leq 3$ and determine the area enclosed between the curve and the axes.
5. Draw a rough sketch of the curve $y = \sqrt{3x + 4}$ and find the area under the curve, above the x -axis and between $x = 0$ and $x = 4$.
6. Sketch the rough graph of $y = 4\sqrt{x-1}$, $1 \leq x \leq 3$ and compute the area between the curve, x -axis and the line $x = 3$.
7. Find the area enclosed between the curve $y = 2x - x^2$ and the x -axis.
8. Find the area of the region bounded by the curve $y^2 = 2y - x$ and the y -axis.
9. Find the area bounded by the curve $y = x^2 - 7x + 6$, the x -axis and the lines $x = 2$, $x = 6$.
10. Find the area of the region bounded by the curve $x = 4y - y^2$ and the y -axis.
 (I.S.C. 2012)
11. Sketch the graph of the curve $y = \sqrt{x} + 1$, $0 \leq x \leq 4$ and determine the area of the region enclosed by the curve, x -axis and the lines $x = 0$ and $x = 4$.
12. Find the area of the region bounded by the parabola $y^2 = 4ax$ and its latus-rectum.
13. (i) Find the area lying between the curve $y^2 = 4x$ and the line $y = 2x$.
 (ii) Find the area enclosed by the parabola $y^2 = 4ax$ and the chord $y = mx$.
14. Find the area lying in the first quadrant and bounded by the circle $x^2 + y^2 = 4$ and the lines $x = 0$ and $x = 2$.
15. Sketch the region $\{(x, y) ; 4x^2 + 9y^2 = 36\}$ and find its area, using integration.
16. Make a rough sketch of the curve $\frac{x^2}{4} + \frac{y^2}{9} = 1$ and find
 (i) the area under the curve and above the x -axis.
 (ii) the area enclosed by the curve.
17. Find the area of the region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.
18. (i) Find the area of the smaller part enclosed by the circle $x^2 + y^2 = 4$ and the line $x + y = 2$.
 (ii) Find the area of the smaller part of the circle $x^2 + y^2 = a^2$ cut off by the line $x = \frac{a}{\sqrt{2}}$.
19. Find the area of the region in the first quadrant enclosed by the x -axis, the line $y = x$ and the curve $x^2 + y^2 = 16$.

12. Find the area of the region enclosed by the curves $y = x^2$, $y = x^2 - 2x$ and the lines $x = 1$, $x = 3$.
13. Draw a rough sketch of the curves $y = \sin x$ and $y = \cos x$ as x varies from 0 to $\frac{\pi}{2}$ and find the area of the region enclosed by them and the x -axis.
14. Find the area enclosed by the curve $y = x^3$, the x -axis and the ordinates $x = -2$ and $x = 1$.
15. Find the area bounded by the curve $y = x^3$ and the line $y = x$.

ANSWERS

EXERCISE 11.1

1. (i) $\frac{26}{3}$ sq. units (ii) $\frac{8}{3}(4 - \sqrt{2})$ sq. units (iii) $\frac{8}{3}(4 - \sqrt{2})$ sq. units
 (iv) $\frac{4}{3}(3\sqrt{3} - 1)$ sq. units.
2. $\frac{32}{3}$ sq. units. 3. 5 sq. units. 4. 18 sq. units.
5. $\frac{112}{9}$ sq. units 6. $\frac{16\sqrt{2}}{3}$ sq. units. 7. $\frac{4}{3}$ sq. units.
8. $\frac{4}{3}$ sq. units. 9. $\frac{56}{3}$ sq. units. 10. $\frac{32}{3}$ sq. units. 11. $\frac{28}{3}$ sq. units.
12. $\frac{8}{3}a^2$ sq. units. 13. (i) $\frac{1}{3}$ sq. units (ii) $\frac{8a^2}{3m^3}$ sq. units. 14. π sq. units.
15. 6π sq. units. 16. (i) 3π sq. units (ii) 6π sq. units.
17. πab sq. units. 18. (i) $(\pi - 2)$ sq. units (ii) $\frac{a^2}{4}(\pi - 2)$ sq. units.
19. 2π sq. units. 20. $\frac{3}{2}(\pi - 2)$ sq. units. 21. (i) $\frac{1}{6}$ sq. units. (ii) $\frac{9}{8}$ sq. units.
22. 18 sq. units. 23. $(\pi - 2)$ sq. units 25. $\frac{2}{3}a^2$ sq. units.
26. (i) $\frac{16}{3}$ sq. units (ii) $\frac{16}{3}a^2$ sq. units.
27. (i) $\frac{23}{6}$ sq. units (ii) $\left(\frac{\pi}{4} - \frac{1}{2}\right)$ sq. units (iii) $\frac{1}{3}$ sq. units.
28. $\frac{16}{3}$ sq. units. 29. $\left(\frac{2\pi}{3} - \frac{\sqrt{3}}{2}\right)$ sq. units. 30. $\frac{13}{3}$ sq. units.
31. 4 sq. units. 32. $\frac{\pi}{4}$ sq. units. 33. 4 sq. units.

EXERCISE 11.2

1. 9; it represents the area below the graph, above the x -axis and bounded by the lines $x = -4$ and $x = 2$.
2. $\frac{16}{3}a$ sq. units. 3. (i) $\frac{3}{2}$ sq. units (ii) 6 sq. units.
4. $(8\pi - \sqrt{3}) : (4\pi + \sqrt{3})$. 5. 15 : 49.
6. $20\frac{5}{6}$ sq. units; $10\frac{2}{3}$ sq. units. 7. $\left(\frac{4 - \sqrt{2}}{\log 2} - \frac{5}{2}\log 2 + \frac{3}{2}\right)$ sq. units.