

Chapter 6: Measures of Dispersion

1. Dispersion refers to the variation of the items around an average.
2. Objectives of Dispersion
 - a) To determine the reliability of an average.
 - b) To compare the variability of two or more series.
 - c) It serves the basis of other statistical measures such as correlation etc.
 - d) It serves the basis of statistical quality control.

Properties of good measure of Dispersion

- a) It should be easy to understand.
- b) Easy to calculate.
- c) Rigidly defined
- d) Based on all observations.
- e) Should not be unduly affected by extreme values.

Measures of Dispersion may be either absolute measures or relative measure.

Absolute Measures of Dispersion are

- a) Range
- b) Quartile Deviation
- c) Mean Deviation
- d) Standard Deviation

Relative Measures of Dispersion are

- a) Coefficient of Range
- b) Coefficient of Quartile Deviation
- c) Coefficient of Mean Deviation
- d) Coefficient of Variation

Graphical method of dispersion

Lorenz Curve

Range

It is the difference between the largest and smallest value of distribution.

Computation of Range

Range = L – S

$$\text{Coefficient of Range} = \frac{L - S}{L + S}$$

Merits of Range

1. It is simple to understand and easy to calculate.
2. It is widely used in statistical quality control.

Demerits of Range

1. It is affected by extreme values in the series.
2. It cannot be calculated in case of open end series.
3. It is not based on all items.

Inter quartile range and quartile deviation

Inter quartile range is the difference between Upper Quartile (Q_3) and Lower Quartile Q_1 .

Quartile deviation is half of inter quartile range.

Computation of Inter quartile range and quartile deviation

Inter quartile Range = $Q_3 - Q_1$

Quartile Deviation Q.D = $\frac{Q_3 - Q_1}{2}$

Coefficient of Q.D = $\frac{Q_3 - Q_1}{Q_3 + Q_1}$

Merits of Q.D

1. Easy to compute
2. Less affected by extreme values.
3. Can be computed in open ended series.

Demerits of Q.D

1. Not based on all observations
2. It is influenced by change in sample and suffers from instability.

Mean Deviation

Mean Deviation is defined as the arithmetic average of the absolute deviations [ignoring signs] of various items from Mean or Median.

Computation of Mean Deviation

Individual Series

$$M.D = \frac{\sum |D|}{N}$$

Discrete/Continuous Series

$$M.D = \frac{\sum f |D|}{\sum f}$$

$$\text{Coefficient of M.D} = \frac{M.D}{\bar{X} \text{ or Median}}$$

Merits of Mean Deviation

1. Based on all observations.
2. It is less affected by extreme values.
3. Simple to understand and easy to calculate.

Demerits of Mean Deviation

1. It ignores \pm signs in deviations.
2. It is difficult to compute when deviations comes in fractions.

Standard Deviation: (σ)

It is defined as the root mean square deviation.

Features of Standard Deviation:

1. Value of its deviation is taken from Arithmetic Mean.
2. + and – signs of deviations taken from mean are not ignored.

Related Measures of Standard Deviation

Standard deviation = σ

Coefficient of standard deviation: $\frac{\sigma}{\bar{X}}$

Variance = σ^2

Coefficient of variation = $\frac{\sigma}{\bar{X}} \times 100$

Computation of σ

Individual Series

1. $\sigma = \sqrt{\frac{\sum x^2}{N}}$ Actual Mean Method
2. $\sigma = \sqrt{\frac{\sum d^2}{N} - \left(\frac{\sum d}{N}\right)^2}$ Assumed Mean Method

Discrete / Continuous Series:

- i. $\sigma = \sqrt{\frac{\sum fx^2}{\sum f}}$ Actual Mean Method
 $x = X - \bar{X}$
- ii. $\sigma = \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2}$ Assumed Mean Method
- iii. $\sigma = \sqrt{\frac{\sum fd^{12}}{\sum f} - \left(\frac{\sum fd^1}{\sum f}\right)^2}$ Step Deviation Method

Merits of Standard Deviation

- i. Rigidly defined
- ii. Based on all observations
- iii. Takes Algebraic signs in consideration
- iv. Amenable to further Algebraic treatment

Demerits

- i. Difficult to understand and compute.
- ii. Affected by extreme items.

Lorenz Curve

It is a graphical method of studying dispersion.

Lorenz curve is a cumulative percentage curve in which the percentage of frequency is combined with percentage of other items such as profit, income etc.

1 mark questions

1. What is range?
2. What is coefficient of variation?
3. What is Lorenz curve?

3 marks question

1. Mention any two merit and demerits each of Mean Deviation.

4 marks questions

1. Find out range and coefficient of range of the following series.

Size	5-10	10-15	15-20	20-25	25-30
Frequency	4	9	15	8	5

2. Find out Standard deviation (6) from the following data:

Class Interval	0-10	10-20	20-30	30-40	40-50
Frequency	10	03	02	01	04