(d) The reaction rate always remains constant

Ordinary Thinking

Objective Questions

Rate of a reaction

- **1.** The rate of a chemical reaction
 - [MP PMT 1973; CPMT 1982]
 - (a) Increases as the reaction proceeds
 - (b) Decreases as the reaction proceeds
 - (c) May increase or decrease during the reaction
 - (d) Remains constant as the reaction proceeds
- The rate of a reaction that not involve gases is not dependent on [CPMT 1988; AFMC 1995]
 (a) Pressure (b) Temperature
 - (c) Concentration (d) Catalyst
- **9.** The rate at which a substance reacts depends on its
 - [MP PMT 1987; BHU 1999; KCET 2005]
 - (a) Atomic weight (b) Equivalent weight
 - (c) Molecular weight (d) Active mass
- **10.** The rate law for the reaction
 - $RCl + NaOH(aq) \rightarrow ROH + NaCl$ is given by Rate
 - $= K_1[RCl]$. The rate of the reaction will be[IIT 1988]
 - (a) Doubled on doubling the concentration of sodium hydroxide
 - (b) Halved on reducing the concentration of alkyl halide to one half
 - (c) Decreased on increasing the temperature of the reaction
 - (d) Unaffected by increasing the temperature of the reaction
- 11. If doubling the concentration of a reactant `A' increases the rate 4 times and tripling the concentration of `A' increases the rate 9 times, the rate is proportional to [AIIMS 1991]
 - (a) Concentration of `A'
 - (b) Square of concentration of `A'
 - (c) Under root of the concentration of `A'
 - (d) Cube of concentration of `A'
- **12.** The rate of chemical reaction at constant temperature is proportional to
 - (a) The amount of products formed
 - (b) The product of masses of the reactants

(c) The product of the molar concentration of the reactants

(d) The mean free path of the reaction

- 13. The concentration of a reactant decreases from 0.2 M to 0.1 M in 10 minutes. The rate of the reaction is
 - (a) 0.01 M (b) 10^{-2}

(c) 0.01 mol $dm^{-3} min^{-1}$ (d) 1 mol $dm^{-3} min^{-1}$

- 14. When a reaction is progressing
 - (a) The rate of the reaction goes on increasing

(b) The concentration of the products goes on decreasing

(c) The concentration of the reactants goes on decreasing

15. In a catalytic conversion of N_2 to NH_3 by Haber's process, the rate of reaction was expressed as change in the concentration of ammonia per time is $40 \times 10^{-3} mol \ litre^{-1}s^{-1}$. If there are no side reaction, the rate of the reaction as expressed in terms of hydrogen is (in mol $\ litre^{-1}s^{-1}$)

- (a) 60×10^{-3} (b) 20×10^{-3}
- (c) 1.200 (d) 10.3×10^{-3}
- 16. If the concentration of the reactants is increased, the rate of reaction [MP PMT 1989]
 - (a) Remains unaffected (b) Increases

(c) Decreases (d) May increase or decrease

- **17.** Time required for completion of ionic reactions in comparison to molecular reactions is
 - (a) Maximum (b) Minimum
 - (c) Equal (d) None
- **18.** For reaction $2A + B \rightarrow$ products, the active mass of *B* is kept constant and that of *A* is doubled. The rate of reaction will then
 - (a) Increase 2 times (b) Increase 4 times
 - (c) Decrease 2 times (d) Decrease 4 times
- **19.** In a reaction $2A + B \rightarrow A_2B$, the reactant *A* will disappear at

[MP PET 1993]

- (a) Half the rate that *B* will decrease
- (b) The same rate that *B* will decrease
- (c) Twice the rate that *B* will decrease
- (d) The same rate that A_2B will form
- **20.** The rate of a gaseous reaction is given by the expression K[A][B]. If the volume of the reaction vessel is suddenly reduced to 1/4th of the initial volume, the reaction rate relating to original rate will be **[Roorkee 1992]**
 - (a) 1/10 (b) 1/8
 - (c) 8 (d) 16
- 21. A catalyst increases the rate of reaction because it [EAMCET 1992]
 - (a) Increases the activation energy
 - (b) Decreases the energy barrier for reaction
 - (c) Decreases the collision diameter
 - (d) Increases the temperature coefficient
- **22.** For the reaction $N_2(g)+3H_2(g) \rightarrow 2NH_3(g)$ under certain conditions of temperature and partial pressure of the reactants, the rate of formation of NH_3 is $0.001 \, kg \, h^{-1}$. The rate of conversion of H_2 under the same conditions is

[IIT 1994]

- (a) $1.82 \times 10^{-4} kg/hr$ (b) 0.0015 kg/hr
- (c) $1.52 \times 10^4 kg/hr$ (d) $1.82 \times 10^{-14} kg/hr$
- **23.** In the reaction $2A + B \rightarrow A_2B$, if the concentration of *A* is doubled and of *B* is halved, then the rate of the reaction will
 - (a) Increase by four times (b)Decrease by two times
 - (c) Increase by two times (d) Remain the same

24. The term $\left(-\frac{dc}{dt}\right)$ in a rate equation refers to the

[MP PMT 1996]

[AFMC 2002]

- (a) Concentration of the reactant
- (b) Decrease in concentration of the reactant with time
- (c) Increase in concentration of the reactant with time
 - (d) Velocity constant of the reaction
- The rate of a reaction depends upon the [Pb. PMT 1999] 25.
 - (a) Volume
 - (b) Force
 - (c) Pressure
 - (d) Concentration of reactant
- For a given reaction $3A + B \rightarrow C + D$ the rate of 26. reaction can be represented by [DCE 2000]

(a)
$$-\frac{1}{3}\frac{d[A]}{dt} = \frac{-d[B]}{dt} = \frac{+d[C]}{dt} = \frac{+d[D]}{dt}$$

(b) $-\frac{1}{3}\frac{d[A]}{dt} = \frac{d[C]}{dt} = K[A]^m[B]^n$
(c) $+\frac{1}{3}\frac{d[A]}{dt} = \frac{-d[C]}{dt} = K[A]^n[B]^m$

- (d) None of these **27.** For the reaction $N_2 + 3H_2 \rightarrow 2NH_3$
 - if $\frac{\Delta[NH_3]}{\Delta t} = 2 \times 10^{-4} \, mol \, l^{-1} s^{-1}$, the value of $\frac{-\Delta[H_2]}{\Delta t}$ would be
 - (a) $1 \times 10^{-4} mol \ l^{-1} s^{-1}$ (b) $3 \times 10^{-4} mol \ l^{-1}s^{-1}$
 - (c) $4 \times 10^{-4} mol \ l^{-1} s^{-1}$ (d) $6 \times 10^{-4} mol \ l^{-1} s^{-1}$
- 28. A gaseous hypothetical chemical equation 2A = 4B + C is carried out in a closed vessel. The concentration of B is found to increase by $5 \times 10^{-3} mol \ l^{-1}$ in 10 second. The rate of appearance of *B* is [AFMC 2001] (a) $5 \times 10^{-4} mol \ l^{-1} sec^{-1}$ (b) $5 \times 10^{-5} mol \ l^{-1} sec^{-1}$ (c) $6 \times 10^{-5} mol \ l^{-1} \text{ sec}^{-1}$ (d) $4 \times 10^{-4} mol l^{-1} sec^{-1}$
- The rate of a chemical reaction depends upon 29.
 - (a) Time

(b) Pressure (c) Concentration (d) All of these

The rate of disappearance of SO_2 in the reaction 30. $2SO_2 + O_2 \rightarrow 2SO_3$ is $1.28 \times 10^{-3} g/sec$ then the rate of formation of SO_3 is [JIPMER 2002]

> (a) $0.64 \times 10^{-3} g/sec$ (b) $0.80 \times 10^{-3} g / sec$

(c)
$$1.28 \times 10^{-3} g/sec$$
 (d) $1.60 \times 10^{-3} g/sec$

- When the concentration of A in the reaction 31. $A + B \Rightarrow AB$ is doubled, the rate of reaction will be [MP PM (a) Doubled (b) Decreased by half (c) Unchanged (d) Increased by four times
- The velocity of the chemical reaction doubles 32. rise of temperature. If the every $10^{\circ}C$ temperature is raised by $50^{\circ}C$, the velocity of the reaction increases to about

- (c) 20 times (d) 50 times
- An increase in temperature by $10^{\circ}C$, generally 33. increases the rate of a reaction by

	(a) 2 times	(b) 10 times
	(c) 9 times	(d) 100 times
34.	The temperature coeffic food deteriorates is 2. T	ient for reaction in which hen food deteriorates
	times as rapidly at 25°C	as it does at $5^{\circ}C$
	(a) Two	(b) Four
	(c) Six	(d) Twenty
35.	The rate of a reaction	is doubled for every 10°
	rise in temperature. The	e increase in reaction rate
)]	as a result of temperatu	are rise from 10° to 100°
	[KCET 1993; Kera	ala PET 2002; MP PET 2003]
	(a) 112	(b) 512
	(c) 400	(d) 614
36.	A catalyst increases reaction by	the rate of a chemical
	[MNR 1988]	; CPMT 1999; Pb. PMT 2000]
	(a) Increasing the activa	ation energy
	(b) Decreasing the activ	ation energy
	(c) Reacting with reacta	ints
	(d) Reacting with produ	cts
37.	Velocity constant of a re	action at 290 <i>K</i> was found
	to be 3.2×10^{-9} . At 310 k	(it will be about[KCET 1989, 91]
	(a) 1.28×10^{-2}	(b) 9.6×10^{-3}
	(c) 6.4×10^{-3}	(d) 3.2×10^{-4}
38.	The temperature coeffic	ient of a reaction is
[]	neaction rat	e at 25 ° C
	(b) Rate of the reaction	at 100 ° C
	(c) Ratio of the rate c	onstants at temperatures
	$35^{\circ}C$ and $25^{\circ}C$	
	(d) Ratio of the ra	ate constants at two
	temperatures differi	ng by 1°C
39.	The main function of a	catalyst in speeding up a
	(a) To increase the rate	of the forward reaction
	(b) To change the react	ion path so as to decrease
	the energy of activat	ion for the reaction
	(c) To reduce the temper	rature at which the reaction
	(d) To increase the oper	way of the melocules of the
	reactants	gy of the molecules of the
40.	The rate of a reaction	[CPMT 1973]
	(a) Increases with incre	ase in temperature
	(b) Decreases with incre	ease in temperature
	(c) Does not depend on	temperature
41	(u) Does not depend on which of the following	concentration
41.	relation to enzyme	
	(a) <i>pH</i> affects their func	tioning
рмт	2(b) Temperature affects	their functioning
1 141 1	(c) They always increas	e activation energy
	(d) Their reactions are s	specific

- **42.** A reaction is catalysed by 'X'. Here 'X' [MP PMT 2003]
 - (a) Decreases the rate constant of reaction
 - (b) Does not affect the equilibrium constant of reaction
 - (c) Decreases the enthalpy of reaction
 - (d) Decreases the activation energy
- Which reaction characteristics are changing by 43. the addition of a catalyst to a reaction at constant temperature

	(i) Activation energy(iii)	(ii) Equilibrium constant Reaction entropy (iv)	
		[DCE 2003]	
	(a) (i) Only	(b) (iii) only	5
	(c) (i) and I(ii) only	(d) All of these	
44.	The velocity constant o	of a reaction at 290 K was	
	found to be 3.2×10^{-3} . A	t 300 <i>K</i> it will be [MP PMT 2004]
	(a) 1.28×10^{-2}	(b) 6.4×10^{-3}	
	(c) 9.6×10^{-3}	(d) 3.2×10^{-4}	
4 5 .	In which of the following of farthest to completion	ng cases, does the reaction on	e
	(a) $K = 10^3$	(b) $K = 10^{-2}$	
	(c) $K = 10$	(d) $K = 1$	
46.	Rate of reaction	[Pb. CET 2004]	
	(a) Decreases with incr	ease in temperature	
	(b) Increases with incre	ease in temperature	
	(c) May increase or c temperature	lecrease with increase in	
	(d) Does not depend on	temperature	
47 •	For the reaction 2	$N_2 O_{5(g)} \to 4 N O_{2(g)} + O_{2(g)}$, if	
	concentration of NO_2 i	n 100 seconds is increased	-
	by $5.2 \times 10^{-3} m$. Then rate	e of reaction will be	1
		[Kerala CET 2005]	
	(a) $1.3 \times 10^{-5} m s^{-1}$	(b) $5 \times 10^{-4} m s^{-1}$	
	(c) $7.6 \times 10^{-4} m s^{-1}$	(d) $2 \times 10^{-3} m s^{-1}$	
	(e) $2.5 \times 10^{-5} m s^{-1}$		
48.	A first order reaction minutes then time requ	complete its 10% in 20 ired to complete its 19% is[Ken	ra
	(a) 30 minutes	(b) 40 minutes	
	(c) 50 minutes	(d) 38 minutes	

(e) 45 minutes

Rate law and Rate constant

1. Which of these does not influence the rate of reaction

[KCET 2005]

- (a) Nature of the reactants
- (b) Concentration of the reactants
- (c) Temperature of the reaction
- (d) Molecularity of the reaction
- 2. The rate law for reaction A + 2B = C + 2D will be (a) Rate = K[A][B] (b) Rate = K[A][2B]

(c) Rate =
$$K[A][B]^2$$
 (d) Rate = $K \frac{[C][D]^2}{[A][B]^2}$

3. In the reaction $2N_2O_5 \rightarrow 4NO_2 + O_2$, initial pressure is 500 *atm* and rate constant *K* is $3.38 \times 10^{-5} \text{ sec}^{-1}$. After 10 minutes the final pressure of N_2O_5 is **[Orissa JEE 2005]** (a) 490 atm (b) 250 atm

(c) 480 atm (d) 420 atm

4. The rate law for the reaction

Sucrose + Water $\xrightarrow{[H^+]}$ Glucose + Fructose is given by

- (a) Rate = K [sucrose] [water]
- (b) Rate = K [sucrose] [water]⁰

(c) Rate = K [sucrose]⁰ [water] Reaction enthalpy (d) Rate = K [sucrose]^{1/2} [water]^{1/2}

5. $A + 2B \to C + D$. If $-\frac{d[A]}{dt} = 5 \times 10^{-4} \mod l^{-1} s^{-1}$ 1, then $-\frac{d[B]}{dt}$ is [DPMT 2005]

(a) $2.5 \times 10^{-4} mol \ l^{-1}s^{-1}$ (b) $5.0 \times 10^{-4} mol \ l^{-1}s^{-1}$

(c) $2.5 \times 10 - 3mol \ l^{-1}s^{-1}$ (d) $1.0 \times 10^{-3} mol \ l^{-1}s^{-1}$

6. The data for the reaction $A + B \rightarrow C$ is [CBSE PMT 1994]

[UI	SEAT 200 Exp.	$[A]_0$	$[B]_0$	Initial rate
	(1)	0.012	0.035	0.10
	(2)	0.024	0.070	0.80
	(3)	0.024	0.035	0.10
	(4)	0.012	0.070	0.80

The rate law corresponds to the above data is

(a) Rate
$$= k[B]^3$$
 (b) Rate $= k[B]^4$

(c) Rate
$$= k[A][B]^3$$
 (d) Rate $= k[A]^2[B]^2$

7. The experimental data for the reaction $2A + B_2 \rightarrow 2AB$ is

[CBSE PMT 1997]

				L	-
	Exp.	$[A]_0$	$[B]_0$	Rate (mole s^{-1})	
	(1)	0.50	0.50	1.6×10^{-4}	
	(2)	0.50	1.00	3.2×10^{-4}	
Kerala CE'	г (2005]	1.00	1.00	3.2×10 ⁴	

The rate equation for the above data is

(a) Rate $= k[B_2]$	(b) Rate $= k [B_2]^2$
(c) Rate $= k[A]^2 [B]^2$	(d) Rate $= k[A]^2[B]$

8. The reaction $2NO(g) + O_2(g) = 2NO_2(g)$ is of first order. If volume of reaction vessel is reduced to 1/3, the rate of reaction would be

(a) 1/3 <i>times</i>	(b) 2/3 <i>times</i>
(c) 3 times	(d) 6 times

9. For a reaction $2A + B \rightarrow$ Products, doubling the initial concentration of both the reactants increases the rate by a factor of 8, and doubling the concentration of *B* alone doubles the rate. The rate law for the reaction is

[MP PET 2001]

(a)
$$\gamma = k[A][B]^2$$
 (b) $\gamma = k[A]^2[B]$

(c)
$$\gamma = k[A][B]$$
 (d) $\gamma = k[A]^2[B]^2$

10. For a reactions A + B → product, it was found that rate of reaction increases four times if concentration of 'A' is doubled, but the rate of reaction remains unaffected. If concentration of 'B' is doubled. Hence, the rate law for the reaction is [MP PET/PMT 1998; MP PMT 2003]

(a) rate =
$$k[A][B]$$
 (b) rate = $k[A]^2$

(c) rate = $k[A]^2[B]^1$ (d) rate = $k[A]^2[B]^2$

- **11.** Velocity constant K of a reaction is affected by
 - (a) Change in the concentration of the reactant

- (b) Change of temperature
- (c) Change in the concentration of the product
- (d) None of the above
- 12. Point out the wrong statement :

For a first order reaction

- (a) Time for half-change $(t_{1/2})$ is independent of initial concentration
- (b) Change in the concentration unit does not change the rate constant (*K*)
- (c) Time for half-change \times rate constant = 0.693
- (d) The unit of K is mole $^{-1}$ min $^{-1}$
- **13.** The rate constant of a reaction depends on

[CPMT 1989; DPMT 2001]

- (a) Temperature (b) Mass
- (c) Weight (d) Time
- 14. In a first order reaction the concentration of reactant decreases from $800 \text{ mol}/\text{dm}^3$ to $50 \text{ mol}/\text{dm}^3$ is $2 \times 10^2 \text{ sec}$. The rate constant of reaction in sec⁻¹ is

[IIT-JEE (Screening) 2003]

(a)
$$2 \times 10^4$$
 (b) 3.45×10^{-5}

- (c) 1.386×10^{-2} (d) 2×10^{-4}
- **15.** For a reaction $A \rightarrow B$, the rate of reaction quadrupled when the concentration of A is doubled. The rate expression of the reaction is $r = K(A)^n$. when the value of *n* is
 - (a) 1 (b) O

(c) 3 (d) 2

16. The velocity constant of a reaction is *K*. Which of the following statements is not true regarding *K*

(a) K is a constant for a reaction at a given temperature

(b) The value of *K* changes when the temperature changes

(c) *K* is the velocity of the reaction at unit concentrations of the reactant

(d) *K* is a constant for all reactions

- **17.** For the following reaction scheme (homogeneous), the rate constant has units : $A+B \xrightarrow{K} C$ [MP PET 1999]
 - (a) $\sec^{-1} mole$ (b) \sec^{-1}
 - (c) $\sec^{-1} litre mole^{-1}$ (d) sec
- 18. Which of the following oxides of nitrogen will be the most stable one [NCERT 1978]

(a)
$$2NO_2(g) \Rightarrow N_2(g) + 2O_2(g); K = 6.7 \times 10^{16} \text{ mol } l^{-1}$$

(b)
$$2NO(g) \Rightarrow N_2(g) + O_2(g); K = 2.2 \times 10^{30} \text{ mol } l^{-1}$$

(c)
$$2N_2O_5(g) \Rightarrow 2N_2(g) + 5O_2(g); K = 1.2 \times 10^{34} \text{ mol } l^{-5}$$

d)
$$2N_2O(g) \Rightarrow 2N_2(g) + O_2(g); K = 3.5 \times 10^{33} \text{ mol litre}^{-1}$$

- **19.** The rate of reaction is determined by slow step reaction. The step is called
 - (a) Reaction rate (b) Activation step
 - (c) Rate determining step (d) None of the above

20. The rate of the reaction

 $CCl_3CHO + NO \rightarrow CHCl_3 + NO + CO$ is given by Rate = $K[CCl_3CHO][NO]$. If concentration is expressed in moles/litre, the units of *K* are [MP PET 1993]

- (a) $litre^2 mole^{-2} \sec^{-1}$ (b) mole $litre^{-1} \sec^{-1}$
- (c) $litremole^{-1} \sec^{-1}$ (d) \sec^{-1}
- **21.** Rate constant for a reaction $H_2 + I_2 \rightarrow 2HI$ is 49, then rate constant for reaction $2HI \rightarrow H_2 + I_2$ is [Bihar MEI

(e) 63

22. The reaction

 N_2O_5 (in CCl_4 solution) $\rightarrow 2NO_2$ (solution) $+ \frac{1}{2}O_2(g)$

is of first order in N_2O_5 with rate constant $6.2 \times 10^{-1} s^{-1}$. What is the value of rate of reaction when $[N_2O_5] = 1.25$ mole l^{-1} [AFMC 1998]

(a) 7.75×10^{-1} mole $l^{-1}s^{-1}$ (b) 6.35×10^{-3} mole $l^{-1}s^{-1}$

(c)
$$5.15 \times 10^{-5}$$
 mole $l^{-1}s^{-1}$ (d) 3.85×10^{-1} mole $l^{-1}s^{-1}$

- 23. A reaction that is of the first order with respect to reactant *A* has a rate constant 6 min^{-1} . If we start with $[A] = 0.5 \text{ mol } l^{-1}$, when would [A] reach the value $0.05 \text{ mol } l^{-1}$ [KCET 2000] (a) 0.384 min (b) 0.15 min(c) 3 min (d) 3.84 min
- **24.** The rate constant for the reaction, $2N_2O_5 \rightarrow 4NO_2 + O_2$ is $3 \times 10^{-5} \text{ sec}^{-1}$. If the rate is $2.40 \times 10^{-5} \text{ mol litre}^{-1} \text{ sec}^{-1}$. Then the concentration of N_2O_5 (in mol litre⁻¹) is

[IIT Screening 2000]

(a) 1.4	(b) 1.2
(a) 1.4	(D) 1.2

- (c) 0.04 (d) 0.8
- **25.** For the reaction $2N_2O_5 \rightarrow 4NO_2 + O_2$ rate of reaction and rate constant are 1.02×10^{-4} and $3.4 \times 10^{-5} \text{ sec}^{-1}$ respectively. The concentration of N_2O_5 at that time will be

[BHU 2001]

- (a) 1.732 (b) 3
- (c) 1.02×10^{-4} (d) 3.4×10^{5}
- **26.** The rate law of the reaction $2N_2O_5 \rightarrow 4NO_2 + O_2$ is [MP PET 2002]

(a)
$$r = K[N_2O_5]$$
 (b) $r = K[N_2O_5]^2$
(c) $r = K[N_2O_5]^0$ (d) $r = K[NO_2]^4[O_2]$

27. If $R = K[NO]^2[O_2]$, rate constant may be increased by

[BHU 2003]

- (a) Increasing temperature
- (b) Decreasing temperature
- (c) Increasing concentration of O_2
- (d) Increasing concentration of NO
- **28.** The value of rate constant $A + B \rightarrow$ products depends on
 - (a) Concentration of A and B

[BHU 2003]

- (b) Pressure
- (c) Temperature
- (d) All of these
- 29. The rate constant of a reaction depends upon[BHU 2004](a) Extent of reaction
 - (b) Time of reaction
 - (c) Temperature of the system
 - (d) Concentration of the system
- **30.** The rate equation for the reaction $2A + B \rightarrow C$ is found to be: rate = k[A][B]. The correct statement in relation to this reaction is that the **[AIEEE 2004]**
 - (a) Rate of formation of *C* is twice the rate of disappearance of *A*
 - (b) $t_{1/2}$ is a constant
 - (c) Unit of k must be s^{-1}
 - (d) Value of k is independent of the initial concentrations of A and B
- **31.** The specific rate constant of a first order reaction depends on the

[IIT 1981, 83; DPMT 1991; Bihar MEE 1995; KCET 1998]

- (a) Concentration of the reactants
- (b) Concentration of the products
- (c) Time of reaction
- (d) Temperature of reaction
- **32.** If the concentration is expressed in moles per litre, the unit of the rate constant for a first order reaction is
 - [MNR 1986; MP PET 1994, 2000, 01;Bihar MEE 1996; CPMT 1997; MP PMT 1995, 96, 99;AFMC 2002]
 - (a) mole litre $^{-1} \sec^{-1}$ (b) mole litre $^{-1}$
 - (c) \sec^{-1} (d) $\operatorname{mole}^{-1} \operatorname{litre}^{-1} \operatorname{sec}^{-1}$
- 33. The dimension of rate constant of a second order reaction involves [NCERT 1982]
 - (a) Neither time nor concentration
 - (b) Only time
 - (c) Time and concentration
 - (d) Time and square of concentration
- **34.** The unit of rate constant of second order reaction is usually expressed as
 - [NCERT 1983, 84; MNR 1983; MP PMT 1994, 99]
 - (a) mole litre \sec^{-1} (b) mole $^{-1}$ litre $^{-1}$ sec $^{-1}$
 - (c) mole litre⁻¹ sec⁻¹ (d) mole⁻¹ litre sec⁻¹
- **35.** A zero order reaction is one whose rate is independent of

[NCERT 1981]

- (a) Temperature of the reaction
- (b) The concentrations of the reactants
- (c) The concentration of the products
- (d) The material of the vessel in which the reaction is carried out
- **36.** The unit of rate constant for a zero order reaction is

[NCERT 1981; MP PMT 2000; RPET 2000]

(a) litre \sec^{-1} (b) litre mole⁻¹ \sec^{-1}

(c) mole $litre^{-1} \sec^{-1}$ (d) mole \sec^{-1}

37. Which of the following rate laws has an overall order of 0.5 for reaction involving substances *x*, *y* and *z*

[AIIMS 1983]

(a) Rate $= K(C_x)(C_y)(C_z)$

- (b) Rate = $K(C_x)^{0.5} (C_y)^{0.5} (C_z)^{0.5}$
- (c) Rate = $K(C_x)^{1.5} (C_y)^{-1} (C_z)^0$

(d) Rate $= K(C_x)(C_z)^n / (C_y)^2$

38. The rates of a certain reaction (dc/dt) at different times are as follows

Time	e Rate (<i>m</i>	ole litre ⁻¹ sec ⁻¹)
0		2.8×10^{-2}
10		2.78×10^{-2}
20		2.81×10^{-2}
30		2.79×10^{-2}
The	reaction is	[NCERT 1978]
(a) Z	Zero order	(b) First order
(c) 5	Second order	(d) Third order

39. For a chemical reaction $A \rightarrow B$ it is found that the rate of reaction doubles, when the concentration of A is increased four times. The order in A for this reaction is

40. The following data are for the decomposition of ammonium nitrate in aqueous solution

Volume of N_2 in cc	Time (minutes)
6.25	10
9.50	15
11.42	20
13.65	25
35.05	Finally
The order of the react	ion is
(a) Zero	(b) One
(c) Two	(d) Three
mb - b - l - l - c - c - t -	

41. The hydrolysis of ethyl acetate is a reaction of [MP PMT 1987]

 $CH_{3}COOEt + H_{2}O \xrightarrow{H^{+}} CH_{3}COOH + EtOH$

- (a) First order (b) Second order
- (c) Third order (d) Zero order
- **42.** The rate of reaction between *A* and *B* increases by a factor of 100, when the concentration of *A* is increased 10 folds. The order of reaction with respect to *A* is

[CPMT 1985]

- (a) 10 (b) 1 (c) 4 (d) 2
- **43.** Which of the following is a first order reaction
 - [MP PMT 1987]
 - (a) $NH_4NO_2 \rightarrow N_2 + 2H_2O$
 - (b) $2HI \rightarrow H_2 + I_2$
 - (c) $2NO_2 \rightarrow 2NO + O_2$
 - (d) $2NO + O_2 \rightarrow 2NO_2$
- 44. The inversion of cane sugar is represented by $C_{12}H_{22}O_{11} + H_2O \rightarrow C_6H_{12}O_6 + C_6H_{12}O_6$

It is a reaction of

[AFMC 1982; MP PMT 1993; RPET 2000]

- (a) Second order (b) Unimolecular
- (c) Pseudo unimolecular (d) None of the three

45. Which one of the following formula represents a first order reaction [MP PMT 1999]

(a)
$$K = \frac{x}{t}$$
 (b) $K = \frac{1}{2t} \left[\frac{1}{(a-x)^2} - \frac{1}{a^2} \right]$
(c) $K = \frac{2.303}{t} \log_{10} \frac{a}{(a-x)}$ (d) $K = \frac{1}{t} \frac{x}{a(a-x)}$

46. The first order rate constant for the decomposition of N_2O_5 is $6.2 \times 10^{-4} \text{ sec}^{-1}$. The half life period for this decomposition in seconds is

[MNR 1991; MP PET 1997; UPSEAT 2000]

(a) 1117.7	(b) 111.7

- (c) 223.4 (d) 160.9
 47. A first order reaction which is 30% complete in 30 minutes has a half-life period of [AIIMS 1998]
 (a) 24.2 min (b) 58.2 min
 - (c) 102.2 min (d) 120.2 min
- **48.** The order of a reaction which has the rate expression $\frac{dc}{dt} = K[E]^{3/2}[D]^{3/2}$ is

(a) 3/2	(b) 3
(c) 2	(d) 0

- **49.** The reaction $2N_2O_5 = 2NO_2 + O_2$ follows first order kinetics. Hence, the molecularity of the reaction is
 - (a) Unimolecular
 - (b) Pseudo-unimolecular
 - (c) Bimolecular
 - (d) None of the above
- **50.** A reaction involving two different reactants

[KCET 1989; AIEEE 2005]

- (a) Can never be a second order reaction
- (b) Can never be a unimolecular reaction
- (c) Can never be a bimolecular reaction
- (d) Can never be a first order reaction

51. By "the overall order of a reaction", we mean

- (a) The number of concentration terms in the equation for the reaction
- (b) The sum of powers to which the concentration terms are raised in the velocity equation
- (c) The least number of molecules of the reactants needed for the reaction

(d) The number of reactants which take part in the reaction

- **52.** Catalyst decomposition of hydrogen peroxide is a order reaction
 - (a) First (b) Second
 - (c) Third (d) Zero

53. The half life of a first order reaction is

(a) Independent of the initial concentration of the reactant

- (b) Directly proportional to the initial concentration of the reactants
- (c) Inversely proportional to the initial concentration of the reactant
- (d) Directly proportional to the square of the initial concentration of the reactant

54. The decomposition of N_2O_5 is a first order reaction represented by $N_2O_5 \rightarrow N_2O_4 + \frac{1}{2}O_2$. After 15 *minutes* the volume of O_2 produced is 9 ml and at the end of the reaction 35 ml. The rate constant is equal to

[MP PET 1995]

(a)
$$\frac{1}{15} \ln \frac{35}{44}$$
 (b) $\frac{1}{15} \ln \frac{44}{26}$
(c) $\frac{1}{15} \ln \frac{44}{35}$ (d) $\frac{1}{15} \ln \frac{35}{26}$

55. The unit of specific reaction rate constant for a first order (if the concentration expressed in molarity) would be

[MNR 1988; UPSEAT 2000, 01]

(a) mole $litre^{-1}s^{-1}$	(b) mole litre ⁻¹
(c) mole s^{-1}	(d) s^{-1}

- **56.** A first order reaction requires 30 minutes for 50% completion. The time required to complete the reaction by 75% will be
 - (a) 45 minutes (b) 15 minutes
 - (c) 60 minutes (d) None of these
- **57.** Inversion of canesugar in dilute acid (conversion into glucose and fructose) is a
 - (a) Unimolecular reaction
 - (b) Bimolecular reaction
 - (c) Trimolecular reaction
 - (d) Pseudo-unimolecular reaction
- 58. The half life period of a first order reaction[AMU 2000]

(a)
$$\frac{0.693}{t}$$
 (b) $\frac{0.693}{K}$
(c) $\frac{2.303}{K}$ (d) $\frac{0.303}{K}$

- **59.** The order of a reaction is said to be 2 with respect to a reactant *X*, when **[DPMT 2000]**
 - (a) The rate of the reaction is proportional to [X]

 K_1

- (b) The rate of the reaction is proportional $[X]^2$
- (c) Two molecules of *X* are present in the stoichiometric equation
- (d) The reaction occurs in two steps
- **60.** Decay constant of a reaction is 1.1×10^{-9} / sec , then the half life of the reaction is
 - (a) 1.2×10^8 (b) 6.3×10^8
 - (c) 3.3×10^8 (d) 2.1×10^8
- **61.** If the half life period of a reaction is inversely proportional to the initial concentration, the order of the reaction is
 - (a) Zero (b) One

(c) Two	(d) Three
---------	-----------

62. Which one of the following statements is wrong

(a) Molecularity of a reaction is always a whole number

(b) Order and molecularity of a reaction need not be same

- (c) Order of a reaction may be zero
- (d) Order of a reaction depends upon the mechanism of the reaction

- **63.** The velocity constant of first order reaction is expressed in the units
 - (a) Concentration per unit time
 - (b) Time per unit concentration
 - (c) Per unit time
 - (d) Unit time per unit concentration
- **64.** For reation, $A + B \rightarrow$ products, it is found that the rate of the reaction is proportional to the concentration of *A*, but it is independent of the concentration of B, then
 - (a) The order of the reaction 2 and molecularity 1
 - (b) Molecularity of the reaction is 2 but order is 1
 - (c) Order is 2 and molecularity is 2
 - (d) Order of the reaction is 2 but molecularity is 0
- 65. For a zero order reaction
 - (a) The concentration of the reactant does not change during the reaction
 - (b) The concentration change only when the temperature changes
 - (c) The rate remains constant throughout
 - (d) The rate of the reaction is proportional to the concentration
- **66.** If 'a' is the initial concentration and 'n' is the order of the reaction and the half life period is 'T', then

[MH CET 2000]

(a)
$$T \propto a^{n-1}$$
 (b) $T \propto a^{n}$
(c) $T \propto \frac{1}{a^{n}}$ (d) $T \propto \frac{1}{a^{n-1}}$

- **67.** In presence of HCl, sucrose gets hydrolysed into glucose and fructose. The concentration of sucrose was found to reduce form 0.4 *M* to 0.2 *M* in 1 hour and 0.1 *M* in 2 hours. The order of the reaction is
 - (a) Zero (b) One
 - (c) Two (d) None of these
- **68.** The time of half change of a first order reaction in initial concentration
 - (a) Proportional to
 - (b) Inversely proportional to
 - (c) Independent of
 - (d) Equal to
- **69.** Half life period of a first order reaction is 138.6 minutes. The velocity constant of the reaction is
 - (a) 0.05 min^{-1} (b) 0.00005 min^{-1}

(c) 0.005 min^{-1} (d) 200 min^{-1}

- 70. An example of a pseudo -unimolecular reaction is(a) Dissociation of hydrogen iodide
 - (b) Hydrolysis of methyl acetate in dilute solution
 - (c) Dissociation of phosphorus pentachloride
 - (d) Decomposition of hydrogen peroxide
- **71.** About half life period of a first order reaction, which one of the following statements is generally false
 - (a) It is independent of initial concentration
 - (b) It is independent of temperature

(c) It decreases with the introduction of a catalyst

(d) It increases with increase of temperature

- **72.** Decomposition of nitrogen pentoxide is known to be a first order reaction 75 percent of the oxide had decomposed in the first 24 minutes. At the end of an hour, after the start of the reaction, the amount of oxide left will be
 - (a) Nil (b) About 1%
 - (c) About 2% (d) About 3%
- **73.** A reaction $2A \rightarrow$ products is found to follow zero oder kinetics, then

(a)
$$\frac{dx}{dt} = k[A]^2$$
 (b) $\frac{dx}{dt} = k[A]^6$

(c)
$$\frac{dx}{dt} = k[A]$$
 (d) $\frac{dx}{dt} = k[2A]$

74. The alkaline hydrolysis of ethyl acetate is represented by the equation

 $CH_3COOC_2H_5 + NaOH \rightarrow CH_3COONa + C_2H_5OH$

Experimentally it is found that for this reaction

$$\frac{dx}{dt} = k[CH_3COOC_2H_5][NaOH]$$

Then the reaction is

- (a) Bimolecular and of first order
- (b) Bimolecular and of second order
- (c) Pseudo-bimolecular
- (d) Pseudo-unimolecular
- **75.** For the reaction $2HI = H_2 + I_2$, the rate of the reaction is proportional to $[HI]^2$. This means that the reaction is

	[AMU 1985; MP PET 2000]
(a) Unimolecular	(b) Bimolecular
(c) Of first order	(d) Of second order

- 76. Inversion of sucrose is [AMU 1988; MP PET 2000](a) Zero order reaction (b) First order reaction
 - (c) Second order reaction (d)Third order reaction
- **77.** The one which is unimolecular reaction is

[MP PMT 1999; UPSEAT 2001]

[JIPMER 1999]

(a)
$$2HI \rightarrow H_2 + I_2$$
 (b) $N_2O_5 \rightarrow N_2O_4 + \frac{1}{2}O_2$

(c)
$$H_2 + Cl_2 \rightarrow 2HCl$$
 (d) $PCl_3 + Cl_2 \rightarrow PCl_5$

78. Integrated velocity equation for first order reaction is

(a)
$$[A]_{\rho} = [A]e^{-Kt}$$
 (b) $K = [A]_{\rho}e^{-A/t}$

(c)
$$Kt = 2.303 \log \frac{[A]_o}{[A]}$$
 (d) $\log \frac{[A]_o}{[A]} = -2.303 Kt$

- **79.** If the surface area of the reactants increases, then order of the reaction
 - (a) Increases
 - (b) Decreases
 - (c) Remain constant
 - (d) Sometimes increases and sometimes dereases
- **80.** Half life period $t_{1/2}$ for first order reaction is

(a)
$$K$$
 (b) $\frac{1.303 \log 2}{K}$
(c) $\frac{2.303 \log 2}{K}$ (d) $\frac{9}{K}$

81.	Molecularity	of reaction	of inversion	of sugar	is
-----	--------------	-------------	--------------	----------	----

(a) 3	(b) 2
(c) 1	(d) 0

82. For any reaction, if we plot a graph between time 't' and log(a-x), a simple line is obtained. The order of reaction is

(a) Zero	(b) One
(c) Two	(d) Three

- 83. Value of velocity constant for first order reaction is 3.46×10^{-3} min⁻¹, the time for half change is
 - (a) 100 minutes (b) 400 minutes
 - (c) 200 minutes (d) 346 minutes
- 84. The unit of the velocity constant in case of zero order reaction is [CPMT 1994]
 - (a) $Conc. \times time^{-1}$ (b) $Conc.^{-1} \times time$
 - (c) $Conc.^{-1} \times time^{-1}$ (d) $Conc. \times (time)^2$
- **85.** For the reaction $H_2(g) + Br_2(g) \rightarrow 2HBr(g)$, the experimental data suggest, rate $= K[H_2][Br_2]^{1/2}$. The molecularity and order of the reaction are respectively

[CPMT 1988; MP PET 1993]

(a)	$2, \frac{3}{2}$	(b)	$\frac{3}{2}, \frac{3}{2}$
(c)	1, 1	(d)	$1, \frac{1}{2}$

2

86. The incorrect order indicated against the rate of reaction $A+B \xrightarrow{K} C$ is [BHU 1990] Rate Order

RateOrder(a) $\frac{d[C]}{dt} = K[A]$ 1(b) $\frac{d[C]}{dt} = K[A][B]$ 2(c) $\frac{-d[A]}{dt} = K[A][B]^0$ 2(d) $\frac{-d[A]}{dt}K[A]$ 1

- **87.** Which of the following statements regarding the molecularity of a reaction is wrong[**CBSE PMT 1989**]
 - (a) It is the number of molecules of the reactants taking part in a single step chemical reaction
 - (b) It is calculated from the reaction mechanism
 - (c) It may be either a whole number or fractional
- (d) It depends on the rate determining step in the reaction

88. Diazonium salt decomposes as

 $C_6H_5N_2^+Cl^- \rightarrow C_6H_5Cl + N_2$

At $0^{\circ}C$, the evolution of N_2 becomes two times faster when the initial concentration of the salt is doubled. Therefore, it is [MNR 1994; UPSEAT 2002] (a) A first order reaction

- (b) A second order reaction
- (c) Independent of the initial concentration of the

(d) A zero order reaction

salt

89. In the reaction $A + B \rightarrow$ Products, if *B* is taken in excess, then it is an example of **[EAMCET 1992]**

- (a) Second order reaction
- (b) Zero order reaction
- (c) Pseudounimolecular reaction
- (d) First order reaction

90. The half life of a first order reaction is 69.35 sec . The value of the rate constant of the reaction is**[CBSE PMT**]

- (a) $1.0 s^{-1}$ (b) $0.1 s^{-1}$
- (c) $0.01 s^{-1}$ (d) $0.001 s^{-1}$

91. The half life for the reaction $N_2O_5 = 2NO_2 + \frac{1}{2}O_2$

in 24 hrs at 30 ^{o}C . Starting with 10 g of N_2O_5 how many grams of N_2O_5 will remain after a period of 96 hours

(a) 1.25 g (b) 0.63 g

```
(c) 1.77 g (d) 0.5 g
```

92. The half life of a first order reaction is 10 minutes. If initial amount is 0.08 mol/litre and concentration at some instant is 0.01 mol/litre, then t = [Roorkee 1990]
(a) 10 minutes (b) 30 minutes

(c) 20 minutes (d) 40 minutes

93. Half life period of second order reaction is[MP PMT 1994](a) Proportional to the initial concentration of reactants

(b) Independent of the initial concentration of reactants

- (c) Inversely proportional to initial concentration of reactants
- (d) Inversely proportional to square of initial concentration of reactants

94. The reaction $2H_2O_2 \rightarrow 2H_2O + O_2$ is a

[Manipal MEE 1995]

[KCET 1992]

- (a) Zero order reaction (b) First order reaction
- (c) Second order reaction (d)Third order reaction
- **95.** In a reaction involving hydrolysis of an organic chloride in presence of large excess of water $RCl + H_2O \rightarrow ROH + HCl$ [MP PET 1995]
 - (a) Molecularity is 2, order of reaction is also 2
 - (b) Molecularity is 2, order of reaction is 1
 - (c) Molecularity is 1, order of reaction is 2
 - (d) Molecularity is 1, order of reaction is also 1
- **96.** The thermal decomposition of a compound is of first order. If a sample of the compound decomposes 50% in 120 minutes, in what time will it undergo 90% decomposition

[MP PET 1996]

- (a) Nearly 240 minutes (b) Nearly 480 minutes
- (c) Nearly 450 minutes (d) Nearly 400 minutes
- **97.** The order of a reaction with rate equals $kC_A^{3/2} C_B^{-1/2}$ is

[MP PET 1996, 2001]

(a) 2 (b) 1
(c)
$$-\frac{1}{2}$$
 (d) $\frac{3}{2}$

- **98.** If the rate expression for a chemical reaction is [MP PMT 1996] given by Rate $= k[A]^m[B]^n$
 - (a) The order of the reaction is m
 - (b) The order of the reaction is *n*
 - (c) The order of the reaction is m+n
 - (d) The order of the reaction is m-n
- The half-life period of a first order reaction is 100 99. sec. The rate constant of the reaction is

[MP PMT 1997; MP PET 2001]

(a)
$$6.93 \times 10^{-3} \sec^{-1}$$
 (b) $6.93 \times 10^{-4} \sec^{-1}$

(c) 0.693 sec^{-1} (d) 69.3 sec^{-1}

100. For the first order reaction with rate constant k, which expression gives the half-life period ? (Initial concentration =a) [MP PET/PMT 1998]

(a)
$$\frac{1^2}{k}$$
 (b) $\frac{1}{ka}$
(c) $\frac{0.693}{k}$ (d) $\frac{3}{2ka^2}$

101. The rate constant of a first order reaction whose half-life is 480 seconds, is [MP PET 1999] (a) $2.88 \times 10^{-3} \text{ sec}^{-1}$ (b) 1.44×10^{-3} sec⁻¹

(a)
$$2.88 \times 10^{-1}$$
 (b) 1.44×10^{-3} sec⁻¹
(c) 1.44 sec^{-1} (d) 0.72×10^{-3} sec⁻¹

102. The conversion of $A \rightarrow B$ follows second order kinetics. Doubling the concentration of A will increase the rate of formation of B by a factor

[MP PET 1999; DCE 1999; KCET 2001; BCECE 2005]

(a)	1/4	(b) 2
(c)	1/2	(d) 4

- **103.** The reaction $2FeCl_3 + SnCl_2 \rightarrow 2FeCl_2 + SnCl_4$ is an example of [CBSE PMT 1996; MP PET 1999]
 - (a) First order reaction
 - (b) Second order reaction
 - (c) Third order reaction
 - (d) None of these
- **104.** If reaction between A and B to give C shows first order kinetics in A and second order in B, the rate equation can be written as [MP PET 1999]

,, .,

- (c) Rate $= k[A][B]^2$ (d) Rate $= k[A]^2[B]$
- 105. For a first order reaction, the half-life period is independent of

[CBSE PMT 1999]

- (a) Initial concentration
- (b) Cube root of initial concentration
- (c) First power of final concentration
- (d) Square root of final concentration
- **106.** Order of a reaction can have [DPMT 1996] (a) +*ve* values (b) Whole number values

(d) All of the above (c) Fractional values

107. The order of the reaction occurring by following mechanism should be [JIPMER 1997]

(i) $A_2 \rightarrow A + A$ (fast)

(ii) $A + B_2 \rightarrow AB + B$ (slow)

(iii) $A + B \rightarrow$ (fast)

(b) $3\frac{1}{2}$ (a) $1\frac{1}{2}$ (c) 2

(d) None of these

108. For the reaction $A \rightarrow B$, the rate law expression is : Rate =k[A]

Which of the following statements is incorrect

[Pb. PMT 1998]

(a) The reaction is said to follow first order kinetics

(b) The half life of the reaction will depend on the initial concentration of the reactant

(c) k is constant for the reaction at a constant temperature

- (d) The rate law provides a simple way of predicting the concentration of reactants and products at any time after the start of the reaction
- **109.** If initial concentration is reduced to its 1/4th in a zero order reaction, the time taken for half of the reaction to complete

[BHU 1998]

- (a) Remains same (b) Becomes 4 times
- (c) Becomes one-fourth (d) Doubles
- 110. For a reaction whose rate expression is :

Rate $= k[A]^{1/2}[B]^{3/2}$, the order would be[Pune CET 1998]

- (b) 2 (a) 1.5 (c) 3 (d) 1
- **111.** For the reaction $A \rightarrow B$, the rate increases by a factor of 2.25 when the concentration of A is increased by 1.5. What is the order of the reaction[KCET 1
 - (a) 3 (b) o
 - (c) 2 (d) 1
- **112.** For the reaction $H_2 + Cl_2 \xrightarrow{\text{Sunlight}} 2HCl$ taking place on water, the order of reaction is

[KCET 1998; AIIMS 2002; Pb. PMT 2002]

(a) 1	(b) 2

- (c) 3 (d) 0
- 113. For a first order reaction, rate constant is $0.6932 hr^{-1}$, then half-life for the reaction is [Bihar MEE 199
 - (a) 0.01 hr (b) 1*hr*
 - (c) 2hr (d) 10 hr
 - (e) 0.1*hr*
- **114.** The rate constant of a reaction is 0.69×10^{-1} min⁻¹ and the initial concentration is $0.2 mol l^{-1}$. The half-life period is

[AIIMS 1998]

(a) 400 sec	(b) 600 <i>sec</i>
(c) 800 sec	(d) 1200 sec

115. The rate constant of a first order reaction is 3×10^{-6} per second. If the initial concentration is 0.10 *m*, the initial rate of reaction is

[AFMC 1999 Pb. PMT 1999, 2000; BHU 1999;

AIIMS 1999; KCET 2000; DCE 2004]

- (a) $3 \times 10^{-5} m s^{-1}$ (b) $3 \times 10^{-6} m s^{-1}$
- (c) $3 \times 10^{-8} m s^{-1}$ (d) $3 \times 10^{-7} m s^{-1}$

- **116.** Certain bimolecular reactions which follow the first order kinetics are called
 - (a) First order reactions
 - (b) Unimolecular reactions
 - (c) Bimolecular reactions
 - (d) Pseudounimolecular reactions
- 117. The rate law of the reaction $A + 2B \rightarrow$ Product is given by $\frac{d[dB]}{dt} = k[B^2]$. If A is taken in excess, the
 - order of the reaction will be [AMU (Engg.) 1999] (a) 1 (b) 2
 - (c) 3 (d) 0
- **118.** For a first order reaction $A \rightarrow$ product, the rate of reaction at $[A] = 0.2 \mod l^{-1}$ is $1.0 \times 10^{-2} \mod l^{-1} \min^{-1}$. The half life period for the reaction is[Roorkee 1999]
 - (a) 832*s* (b) 440*s*
 - (c) 416 *s* (d) 13.86 *s*
- **119.** For the reaction $A + B \rightarrow$ products, doubling the concentration of A the rate of the reaction is doubled, but on doubling the concentration of B rate remains unaltered. The over all order of the reaction is [JIPMER 1999]
 - (a) 1 (b) 0
 - (c) 2 (d) 3
- **120.** Which among the following is a false statement

[KCET 1999]

- (a) Half life of a third order reaction is inversely proportional to the square of initial concentration of the reactant.
- (b) Molecularity of a reaction may be zero or fractional
- (c) For a first order reaction $t_{1/2} = \frac{0.693}{K}$
- (d) Rate of zero order reaction is independent of initial concentration of reactant
- 121. After how many seconds will the concentration of the reactants in a first order reaction be halved, if the decay constant is $1.155 \times 10^{-3} \text{ sec}^{-1}$ [CBSE PMT 2000]

00 <i>sec</i>	
2	200 sec

(c) 400 sec (d)600 sec

122. What is the order of a reaction which has a rate expression rate $= K[A]^{3/2}[B]^{-1}$ [DCE 2000]

(a) 3/2	(b) 1/2

(c) 0	(d) None of these

123. Which of the following expression is correct for first order reaction? (*CO*) refers to initial concentration of reactant

[DCE 2000]

(a)	$t_{1/2} \propto CO$	(b) $t_{1/2} \propto CO^{-1}$
(c)	$t_{1/2} \propto CO^{-2}$	(d) $t_{1/2} \propto CO^0$

124. For a reaction $2NO(g) + Cl_2(g) = 2NOCl(g)$. When concentration of Cl_2 is doubled, the rate of reaction becomes two times of the original. When the concentration of *NO* is doubled the rate becomes four times. What is the order of the reaction [MP PMT 2000]

- (a) 1 (b) 2
- (c) 3^[KCET (Med.) 1999] (d) 4
- **125.** The rate constant for a second order reaction is $8 \times 10^{-5} M^{-1} min^{-1}$. How long will it take a 1*M* solution to be reduced to 0.5 *M* [MH CET 2001]
 - (a) 8×10^{-5} min (b) 8.665×10^{3} min
 - (c) 4×10^{-5} min (d) 1.25×10^{4} min
- **126.** The rate for a first order reaction is $0.6932 \times 10^{-2} mol \ l^{-1} min^{-1}$ and the initial concentration of the reactants is 1*M*, $T_{1/2}$ is equal to **[JIPMER (Med.) 2001]**
 - (a) 6.932 min (b) 100 min
 - (c) 0.6932×10^{-3} min (d) 0.6932×10^{-2} min
- **127.** For a given reaction $t_{1/2} = \frac{1}{Ka}$. The order of the reaction is

[KCET 2001]

(a) 1	(b) O
(c) 3	(d) 2

128. 75% of a first order reaction is completed in 30 minutes. What is the time required for 93.75% of the reaction (in minutes) [KCET 2001]

(a) 45	(b) 120
(c) 90	(d) 60

129. A First order reaction is half completed in 45 minutes. How long does it need 99.9% of the reaction to be completed

[AIIMS 2001]

- (a) 5 hours (b) 7.5 hours
- (c) 10 hours (d) 20 hours
- **130.** A substance 'A' decomposes by a first order reaction starting initially with [A] =2.00m and after 200 min [A] = 0.15m. For this reaction what is the value of k [AIIMS 2001]

(a) $1.29 \times 10^{-2} \text{ min}^{-1}$	(b) $2.29 \times 10^{-2} \text{ min}^{-1}$
--	--

- (c) $3.29 \times 10^{-2} \text{ min}^{-1}$ (d) $4.40 \times 10^{-2} \text{ min}^{-1}$
- **131.** Which of the following statements about zero order reaction is not true
 - (a) Its unit is sec^{-1}
 - (b) The graph between log (reactant) versus rate of reaction is a straight line
 - (c) The rate of reaction increases with the decrease in concentration of reactants
 - (d) Rate of reaction is independent of concentration of reactants
- **132.** The given reaction $2NO + O_2 \rightarrow 2NO_2$ is an example of

[CBSE PMT 2001; JIPMER 2002]

(a) First order reaction (b) Second order reaction

- (c) Third order reaction (d) None of these
- 133. Order of a reaction is decided by [KCET 2002](a) Pressure
 - (b) Temperature
 - (c) Molecularity

(d) Relative concentration of reactants

134. From the following which is a second order reaction

[AMU 2002]

(a) $K = 5.47 \times 10^{-4} \text{ sec}^{-1}$

- (b) $K = 3.9 \times 10^{-3}$ mole lit sec⁻¹
- (c) $K = 3.94 \times 10^{-4}$ lit mole⁻¹ sec⁻¹
- (d) $K = 3.98 \times 10^{-5}$ lit mole⁻² sec⁻¹
- **135.** For the reaction $A + 2B \rightarrow C$, rate is given by R

 - (a) 3(b) 6
 - (c) 5 (d) 7
- 136. Units of rate constant of first and zero order reactions in terms of molarity M unit are respectively [AIEEE 2002]
 - (b) sec^{-1} , M (a) sec^{-1} , $M sec^{-1}$
 - (c) $M \sec^{-1}$, \sec^{-1} (d) M, sec^{-1}
- **137.** The reaction $2N_2O_5 = 2N_2O_4 + O_2$ is [MP PMT 2002]
 - (a) Bimolecular and second order
 - (b) Unimolecular and first order
 - (c) Bimolecular and first order
 - (d) Bimolecular and zero order
- **138.** The half-life period for a first order reaction is 693 seconds. The rate constants for this reaction would be

		[MP PET 2002]
	(a) $0.1 sec^{-1}$	(b) $0.01 sec^{-1}$
	(c) $0.001 sec^{-1}$	(d) $0.0001 sec^{-1}$
139.	For an elementary reac	tion, $2A + B \rightarrow C + D$ the
	molecularity is	[Kurukshetra CEE 2002]
	(a) Zero	(b) One
	(c) Two	(d) Three
140.	If the order of the react	ion $x + y \xrightarrow{hv} x y$ is zero,
	it means that the rate of	[Kurukshetra CEE 2002]
	(a) Reaction is independ	ent of temperature
	(b) Formation of activate	ed complex is zero
	(c) Reaction is independent reacting species	ent of the concentration of
	(d) Decomposition of act	ivated complex is zero
141.	For a first order rea	ction velocity constant,
	$K = 10^{-3} s^{-1}$. Two third life	e for it would be[MP PET 20
	(a) 1100 <i>s</i>	(b) 2200 <i>s</i>
	(c) 3300 <i>s</i>	(d) 4400 <i>s</i>
142.	In a reaction, the com- increased two times an increases in rate of reac- nine times respectively,	centration of reactant is nd three times then the ction were four times and order of reaction is
	(a) $7ama$	(h) 1

(a) Zero	(0) 1
(c) 2	(d) 3

143. For a chemical reaction....can never be a fraction [EAMCET 2003]

(a) Order	(b) Half-life
(c) Molecularity	(d) Rate constant

144. 75% of a first order reaction was completed in 32 minutes when was 50% of the reaction completed

[AMU 1999;	Kerala	(Med.)	2003]
------------	--------	--------	-------

- (a) 16 min. (b) 24 min. (c) 8 min. (d) 4 min.
- 145. The decomposition of N_2O_5 occurs as. $2N_2O_5 \rightarrow 4NO_2 + O_2$, and follows Ist order kinetics, hence

[BVP 2003]

- (a) The reaction is unimolecular
- (b) The reaction is bimolecular
- (c) $T_{1/2} \propto a^0$
- (d) None of these

= $[A][B]^2$ then the order of the reaction is [AIEEE 2002] 146. Which equation is correct for first order reactions

[MP PMT 2003]

- (a) $t_{1/2} \propto C^{-1}$ (b) $t_{1/2} \propto C$
- (c) $t_{1/2} \propto C^0$ (d) $t_{1/2} \propto C^{1/2}$
- **147.** For the reaction system $2NO(g) + O_2(g) \rightarrow 2NO_2(g)$ volume is suddenly produced to half its value by increasing the pressure on it. If the reaction is of first order with respect to O_2 and second order with respect to NO, the rate of reaction will [AIEEE 2003]
 - (a) Diminish to one fourth of its initial value
 - (b) Diminish to one eighth of its initial value
 - (c) Increase to eight times of its initial value
 - (d) Increase to four times of its initial value
- **148.** If the rate of the reaction is equal to the rate constant, the order of the reaction is [CBSE PMT 2003]

(a) 3	(b) O
(c) 1	(d) 2

- **149.** The reaction $A \rightarrow B$ follows first order kinetics. The time taken for 0.8 mole of A to produce 0.6 mole of B is 1 hour. What is the time taken for conversion of 0.9 mole of A to produce 0.675 mole of B [CBSE PMT 2003]
 - (a) 2 hours (b) 1 hour
 - (c) 0.5 hour (d) 0.25 hour
- 150. The unit of velocity constant in case of zero order reaction is

[MP PMT 2004]

- (a) Concentration \times Time⁻¹ (b) Concentrat ion $^{-1} \times \text{Time}^{-1}$
- (c) Concentrat ion \times Time² (d) Concentrat ion⁻¹ \times Time

151. Which one of the following is wrongly matched **2001; UPSEAT 2003**]

[KCET 2004]

- (a) Saponification of $CH_3COOC_2H_5$ Second order reaction
- (b) Hydrolysis of CH_3COOCH_3 Pseudo unimolecular [UPSEAT 2003]

reaction

- (c) Decomposition of H_2O_2 First order reaction
- (d) Combination of H_2 and Br_2 to give HBr Zero order

reaction

152. Which of the following is an example of pseudo unimolecular reaction [Pb. CET 2001]

- (a) $CH_3COOCH_3 + H_2O \xrightarrow{H^+} CH_3COOH + CH_3OH$
- (b) $CH_3COOCH_3 + H_2O \xrightarrow{OH^-} CH_3COOH + CH_3OH$
- (c) $2FeCl_3 + SnCl_2 \rightarrow SnCl_4 + 2FeCl_2$
- (d) $NaOH + HCl \rightarrow NaCl + H_2O$
- 153. Hydrolysis of DDT is a first order reaction, its half life is 10 years. Time to hydrolyse 10 g DDT to half is [BVP 2004]

(a) 100 years	(b) 50 years
(c) 5 years	(d) 10 years

154. In a first order reaction, the concentration of the reactant, decreases from 0.8 M to 0.4 M in 15 minutes. The time taken for the concentration to change from 0.1 *M* to 0.025 *M* is [AIEEE 2004] (a) 7 5 minutes (b) 15 minutes

(a) 7.5 minutes	(b) 15 minutes
(c) 30 minutes	(d) 60 minutes

155. In the first order reaction, the concentration of the reactant is reduced to 25% in one hour. The half life period of the reaction is [DCE 2004]

(a)	2 hr	(b) 4 hr
(c)	1/2 hr	(d) 1/4 hr

156. For a reaction, $X(g) \rightarrow Y(g) + Z(g)$ the half life period is 10 min. In What period of time would the concentration of X be reduced to 10% of original concentration [DCE 2004] (a) 20 min (b) 33 min

(c)	15 min	(d)	25	min

157. A first order reaction with respect to the reactant A has a rate constant of $6 \sec^{-1}$. If we start with [A] = 0.5 mol/litre, then in what time the concentration of A becomes 0.05 mol/litre

(a) 0.384 <i>sec</i>	(b) 0.214 sec
(c) 3.84 <i>sec</i>	(d) 0.402 <i>sec</i>

158. Order of radioactive disintegration reaction is [JEE Orissa 2004]

(a) Zero	(b) First
(c) Second	(d) Third

159. The rate of a first order reaction is $1.5 \times 10^{-2} molL^{-1} min^{-1}$ at 0.5 *M* concentration of the reactant. The half life of the reaction is[CBSE PMT 2004] (a) 8.73 min (b) 7.53 min

(c) $0.382 min$	(d) 22 1 min

160. A first order reaction was started with a decimolar solution of the reactant, 8 minutes and 20 seconds later its concentration was found to be M/100. So the rate of the reaction is [Kerala PMT 2004]

(a) $2.303 \times 10^{-5} \text{ sec}^{-1}$ (b)	b) $2.303 \times 10^{-4} \text{ sec}^{-1}$
---	--

(c)	4.606×10^{-3}	sec ⁻¹	(d)	2.606	$\times 10^{-5}$	sec ⁻¹

(e) $2.603 \times 10^{-4} \text{ sec}^{-1}$

161. Which is correct about zero order reaction

[JEE Orissa 2004]

[DCE 2004]

```
(a) Rate of reaction depends on decay constant
```

- (b) Rate of reaction is independent of concentration
- (c) Unit of rate constant is concentration⁻¹
- (d) Unit of rate constant is concentration⁻¹ time⁻¹

162. Decay of $_{92}U^{235}$ isorder reaction[JEE Orissa 2004]

(a) Zero	(b) First
(c) Second	(d) Third

- 163. The half-life of 2 sample are 0.1 and 0.4 seconds. Their respective concentration are 200 and 50 respectively. What is the order of the reaction [JEE Oriss (a) 0 (b) 2
 - (c) 1 (d) 4
- 164. The following statements(s) is(are) correct[IIT 1999] (a) A plot of $\log K_p$ versus 1/T is linear
 - (b) A plot of $\log[X]$ versus time is linear for a first order reaction $X \rightarrow P$
 - (c) A plot of $\log P$ versus 1/T is linear at constant volume
 - (d) A plot of *P* versus 1/V is linear at constant temperature
- 165. For a first order reaction [IIT 1998]
 - (a) The degree of dissociation is equal to $(1 e^{-kt})$
 - (b) A plot of reciprocal concentration of the reactant vs time gives a straight line
 - (c) The time taken for the completion of 75%reaction is thrice the $t_{1/2}$ of the reaction
 - (d) The pre-exponential factor in the Arrhenius equation has the dimension of time T^{-1}

166. For reaction a $A \rightarrow x P$, when [A] = 2.2 mM, the rate was found to be $2.4 \ mM \ s^{-1}$. On reducing concentration of A to half, the rate changes to $0.6 \ mM \ s^{-1}$. The order of reaction with respect to A is [AIIMS 2005]

- (a) 1.5 (b) 2.0
- (c) 2.5 (d) 3.0
- 167. Which one of the following statement for order of reaction is not correct [IIT 2005]
 - (a) Order can be determined experimentally.
 - (b) Order of reaction is equal to sum of the powers of concentration terms in differential rate law.
 - (c) It is not affected with the stoichiometric coefficient of the reactants.
 - (d) Order cannot be fractional.

168. *t*₁ can be taken as the time taken for the

concentration of a reactant to drop to $\frac{3}{4}$ of its initial value. If the rate constant for a first order reaction is K, the t_1 can be written as

[AIEEE 2005]

(a)	0.10 / K	(b)	0.29 / K
(c)	0.69 / K	(d)	0.75 / K

169. For a first order reaction $A \rightarrow B$ the reaction rate at reactant concentration of 0.01M is found to be $2.0 \times 10^{-5} mol \ L^{-1}s^1$. The half life period of the reaction is

[CBSE PMT 2005]

(a) 220 <i>s</i>	(b) 30 <i>s</i>
(c) 300 <i>s</i>	(d) 347 <i>s</i>

170. The rate of reaction between two reactants *A* and *B* decreases by a factor of 4 if the concentration of reactant B is doubled. The order of this reaction with respect to reactant *B* is

(a) – 1	(b) – 2
(C) 1	(d) 2

171. If a substance with half life 3 days is taken at other place in 12 days. What amount of substance is left now [AFMC 2005]

- (c) 1/16 (d) 1/32
- **172.** The half-life of a first order reaction having rate constant $K = 1.7 \times 10^{-5} s^{-1}$ is

(a) 12.1 <i>h</i>	(b) 9.7 h
(c) 11.3 <i>h</i>	(d) 1.8 h

- **173.** For the reaction $A+B \rightarrow C$, it is found that doubling the concentration of A increases the rate by 4 times, and doubling the concentration of B doubles the reaction rate. What is the overal order of the reaction. **[KCET 2005]**
 - (a) 4 (b) 3/2
 - (c) 3 (d) 1
- 174. Which of the following reactions end in finite time

		[DPMT	2005]
(a)	0 order	(b) 1st order	
(c)	2nd order	(d) 3rd order	

Collision theory, Energy of activation and Arrhenius equation

- 1. A large increase in the rate of a reaction for a rise in temperature is due to[EAMCET 1980; MP PET 1995]
 - (a) The decrease in the number of collisions

(b) The increase in the number of activated molecules

- (c) The shortening of the mean free path
- (d) The lowering of the activation energy
- **2.** Which of the following statements is not true according to collision theory of reaction rates
 - (a) Collision of molecules is a precondition for any reaction to occur

(b) All collisions result in the formation of the products

- (c) Only activated collisions result in the formation of the products
- (d) Molecules which have acquired the energy of activation can collide effectively
- **3.** According to the collision theory of chemical reactions
 - (a) A chemical reaction occurs with every molecular collision
 - (b) Rate is directly proportional to the number of collisions per second

(c) Reactions in the gas phase are always of zero order $% \left({{{\mathbf{r}}_{\mathbf{r}}}_{\mathbf{r}}} \right)$

(d) Reaction rates are of the order of molecular speeds

4. According to the collision theory of reaction rates, rate of reaction increases with temperature due to

[CBSE PMT 2005] (a) Greater number of collisions

(b) Greater velocity of the reacting molecules

(c) Greater number of molecules have activation energy

(d) None of the above

5. The reaction rate at a given temperature becomes slower, then

[MP PMT 1993; DPMT 2000]

- (a) The free energy of activation is higher
- (b) The free energy of activation is lower
- (c) The entropy changes
- (d) The initial concentration of the reactants remains constant
- **6.** A rise in temperature increases the velocity of a reaction. It is because it results in
 - (a) An increased number of molecular collisions

(b) An increased momentum of colliding molecules

- (c) An increase in the activation energy
- (d) A decrease in the activation energy
- 7. The number of collisions depend upon
 - (a) Pressure (b) Concentration
 - (c) Temperature (d) All the above
- **8.** If E_f and E_r are the activation energies of forward and reverse reactions and the reaction is known to be exothermic, then
 - (a) $E_f > E_r$
 - (b) $E_f < E_r$
 - (c) $E_f = E_r$
 - (d) No relation can be given between E_f and E_r as data are not sufficient
- **9.** According to Arrhenius theory, the activation energy is
 - (a) The energy it should possess so that it can enter into an effective collision
 - (b) The energy which the molecule should possess in order to undergo reaction
 - (c) The energy it has to acquire further so that it can enter into a effective collison
 - (d) The energy gained by the molecules on colliding with another molecule
- 10. The energy of activation is

(a) The energy associated with the activated molecules

(b) Threshold energy – energy of normal molecules

(c) Threshold energy + energy of normal molecules

- (d) Energy of products energy of reactants
- **11.** Which one of the following does not represent Arrhenius equation

(a)
$$k = Ae^{-E/RT}$$

(b) $\log_e k = \log_e A - \frac{E}{RT}$
(c) $\log_{10} k = \log_{10} A - \frac{E}{2.303 RT}$

(d) $k = AE^{-RT}$

- **12.** On increasing the temperature, the rate of the reaction increases because of
 - (a) Decrease in the number of collisions
 - (b) Decrease in the energy of activation
 - (c) Decrease in the number of activated molecules
 - (d) Increase in the number of effective collisions
- 13. Energy of activation of a reactant is reduced by(a) Increased temperature(b)Reduced temperature(c) Reduced pressure(d) Increased pressure
- 14. The minimum energy a molecule should possess in order to enter into a fruitful collision is known as

[Kurukshetra CEE 2002]

- (a) Reaction energy (b) Collision energy
- (c) Activation energy (d) Threshold energy
- **15.** Activation energy is
 - (a) The amount of energy to be added to the actual energy of a molecule so that the threshold energy is reached
 - (b) The amount of energy the molecule must contain so that it reacts
 - (c) The energy which a molecule should have in order to enter into an effective collision
 - (d) The average kinetic energy of the molecule
- **16.** The reason for almost doubling the rate of reaction on increasing the temperature of the reaction system by $10^{o^{C}}$ is

[J & K 2005]

- (a) The value of threshold energy increases
- (b) Collision frequency increases
- (c) The fraction of the molecule having energy equal to threshold energy or more increases(d) Activation on energy documents
- (d) Activation energy decreases
- **17.** The activation energy for a simple chemical reaction $A \rightarrow B$ is E_a in forward direction. The activation energy for reverse reaction
 - (a) Is always double of E_a
 - (b) Is negative of E_a
 - (c) Is always less than E_a
 - (d) Can be less than or more than E_a
- 18. Arrhenius equation is

(a)
$$\frac{d \ln K}{dT} = \Delta E^* / RT$$
 (b) $\frac{d \ln K}{dT} = \Delta E^* / RT^2$
(c) $\frac{d \ln K}{dT} = -\Delta E^* / RT^2$ (d) $\frac{d \ln K}{dT} = -\Delta E^* / RT$

- **19.** Activation energy of any reaction depends on(a) Temperature
 - (b) Nature of reactants
 - (c) Number of collisions per unit time
 - (d) Concentration of reactants
- **20.** Relation between rate constant and temperature by Arrhenius equation is

(a)
$$\log_e A = \log_e K + \frac{E_a}{RT}$$
 (b) $\log K = A \frac{E_a}{RT}$
(c) $\log_e K = \log_e A - \frac{E_a}{RT^2}$ (d) $\log A = RT \ln E_a - \ln K$

- **21.** An endothermic reaction $A \rightarrow B$ has an activation energy $15 \, kcal/mole$ and energy of reaction $5 \, kcal/mole$. The activation energy of the reaction $B \rightarrow A$ is [Pb. CET 1985]
 - (a) 20/15 kcal/mole (b) 15 kcal/mole
 - (c) 10 *kcal/mole* (d) None of these



23. The Arrhenius equation expressing the effect of temperature on the rate constant of a reaction is[MP PET : (a) k = e^{-E_a/RT}
(b) k = E_a/RT

(c)
$$k = \log_e \frac{E_a}{RT}$$
 (d) $k = Ae^{-E_a/RT}$

24. For a reaction, activation energy $(E_a) = 0$ and rate constant $(K) = 3.2 \times 10^6 s^{-1}$ at 300 K. What is the value of the rate constant at 310 K (a) $3.2 \times 10^{-12} s^{-1}$ (b) $3.2 \times 10^6 s^{-1}$

(a)
$$5.2 \times 10^{-1}$$
 (b) 5.2×10^{-5} (c) $6.4 \times 10^{12} s^{-1}$ (d) $6.4 \times 10^{6} s^{-1}$

25. Activation energy is given by the formula[DCE 1999]

(a)
$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

(b) $\log \frac{K_1}{K_2} = -\frac{E_a}{2.303 R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$
(c) $\log \frac{K_1}{K_2} = -\frac{E_a}{2.303 R} \left[\frac{T_1 - T_2}{T_1 T_2} \right]$

(d) None of these

26. A reaction having equal activation energies for forwards and prove for the second second

(c) Zero order (d) None of these

27. Collision theory is applicable to [MP PMT 2002](a) First order reactions (b) Zero order reactions

- (c) Bimolecular reactions (d)Intra molecular reactions
- **28.** A graph plotted between log *K* vs 1/*T* for calculating activation energy is shown by [**MP PET 2002**]



The rate constant of a reaction at temperature 29. 200K is 10 times less than the rate constant at 400 K. What is the activation energy (E_{a}) of the reaction (R = gas constant)

	[EAMCET 2003]
(a) 1842.4 <i>R</i>	(b) 921.2 <i>R</i>
(c) 460.6 R	(d) 230.3 <i>R</i>

In respect of the equation $k = Ae^{-E_a/RT}$ in chemical 30. kinetics, which one of the following statement is correct

[AIEEE 2003]

- (a) *k* is equilibrium constant
- (b) A is adsorption factor
- (c) E_a is energy of activation
- (d) R is Rydberg's constant
- The rate constant is doubled when temperature 31. increases from 27°C to 37°C. Activation energy in kJ is

[JEE Orissa	2004	[]
-------------	------	----

- (a) 34 (b) 54 (c) 100
- The activation energy of a reaction is zero. The 32. rate constant of this reaction

(d) 50

- (a) Increases with increase of temperature
- (b) Decreases with an increase of temperature
- (c) Decreases with decrease of temperature
- (d) Is independent of temperature
- The rate constant is given by the equation 33. $k = pze^{-E/RT}$. Which factor should register a decrease for the reaction to proceed more rapidly [MP PET/PMT 199
 - (a) T (b) Z
 - (c) E (d) p
- An endothermic reaction with high activation 34. energy for the forward reaction is given by the diagram: [AIIMS 2005]



- Consider an endothermic reaction $X \rightarrow Y$ with the 35. activation energies E_b and E_f for the backward and forward reactions, respectively, in general[AIEEE 2005]
 - (a) $E_h < E_f$
 - (b) $E_b > E_f$
 - (c) $E_b = E_f$
 - (d) There is no definite relation between E_{h} and E_{f}

36. Temperature dependent equation can be written as

[Orrisa JEE 2005]

(b) $\ln k = \ln A + e^{E_a / RT}$ (a) $\ln k = \ln A - e^{\frac{E_a}{RT}}$ (c) $\ln k = \ln A - e^{RT / E_a}$ (d) All of these

Photochemical reactions

- 1. The formation of starch in plants results from
 - (a) Photosynthesis (b) Photolysis (c) Flash photolysis (d) None of the above
- 2. The photolysis of water gives the substance
 - (a) $OH^- + H^+$ (b) $H_2 + OH^-$
 - (c) $H_2 + O_2$ (d) $H_2O + H_2O_2$
- The law of photochemical equivalence was given by 3.
 - (a) Drapper (b) Grauths
 - (c) Einstein (d) Labbert
- If 'I' is the intensity of absorbed light and C is the 4. concentration of AB for the photochemical process $AB + hv \rightarrow AB^*$, the rate of formation of AB^* is directly proportional to [IIT Screening 2001]

(a) <i>C</i>	(b) <i>I</i>
(c) I^2	(d) <i>C.I</i>

(c)
$$I^2$$



For the reaction $H_2(g) + I_2(g) = 2HI(g)$, the rate of 1. reaction is expressed as[CBSE PMT 1997; AIEEE 2002]

(a)
$$-\frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[HI]}{\Delta t}$$

(b) $\frac{\Delta[I_2]}{\Delta t} = \frac{\Delta[H_2]}{\Delta t} = \frac{\Delta[HI]}{2\Delta t}$
(c) $\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[HI]}{\Delta t}$

(d) None of these

If $3A \rightarrow 2B$, then the rate of reaction of $+\frac{d(B)}{dA}$ is 2. equal to

(b) $-\frac{1}{3}\frac{d(A)}{dt}$ (a) $+2\frac{d(A)}{dt}$ (c) $-\frac{2}{3}\frac{d(A)}{dt}$ (d) $-\frac{3}{2}\frac{d(A)}{dt}$

The differential rate law for the reaction 3. $H_2 + I_2 \rightarrow 2HI$ is [AIEEE 2002]

(a)
$$-\frac{d[H_2]}{dt} = -\frac{d[I_2]}{dt} = +\frac{1}{2}\frac{d[HI]}{dt}$$

[CBSE PMT 2002]

(b)
$$\frac{d[H_2]}{dt} = \frac{d[HI]}{dt} = \frac{1}{2} \frac{d[HI]}{dt}$$

(c) $\frac{1}{2} \frac{d[H_2]}{dt} = \frac{1}{2} \frac{d[I_2]}{dt} = -\frac{d[HI]}{dt}$
(d) $-2 \frac{d[H_2]}{dt} = -2 \frac{d[I_2]}{dt} = +\frac{d[HI]}{dt}$

4. The rate law for a reaction between the substances *A* and *B* is given by, rate $=k[A]^n[B]^m$. On doubling the concentration of *A* and halving the concentration of *B*, the ratio of the new rate to the earlier rate of the reaction will be as

(a)
$$\frac{1}{2^{(m+n)}}$$
 (b) $(m+n)$
(c) $(n-m)$ (d) $2^{(n-m)}$

5. If we plot a graph between log K and $\frac{1}{T}$ by Arrhenius equation, the slope is [UPSEAT 2001]

(a)
$$-\frac{E_a}{R}$$
 (b) $+\frac{E_a}{R}$
(c) $-\frac{E_a}{2.303 R}$ (d) $+\frac{E_a}{2.303 R}$

- 6. For an endothermic reaction, where ΔH represents the enthalpy of the reaction in kJ/mol, the minimum value for the energy of activation will be [IIT 1992] (a) Less than ΔH (b) Zero
 - (c) More than ΔH (d) Equal to ΔH
- 7. The rate constant (K') of one reaction is double of the rate constant (K'') of another reaction. Then the relationship between the corresponding activation energies of the two reactions (E_a'' and E_a''') will be

(a)
$$E_{a}^{'} > E_{a}^{''}$$
 (b) $E_{a}^{'} = E_{a}^{''}$

(c)
$$E_a' < E_a''$$
 (d) $E_a' = 4E_a$

8. The rate constant, the activation energy and the arrhenius parameter of a chemical reaction at $25 \,^{o}C$ are $3.0 \times 10^{-4} \, s^{-1}$, $104.4 \, kJ \, mol^{-1}$ and $6.0 \times 10^{14} \, s^{-1}$ respectively. The value of the rate constant as $T \to \infty$ is

(a)
$$2.0 \times 10^{18} s^{-1}$$
(b) $6.0 \times 10^{14} s^{-1}$ (c) Infinity(d) $3.6 \times 10^{30} s^{-1}$

- 9. The ΔH value of the reaction H₂ + Cl₂ = 2HCl is -44.12 kcal. If E₁ is the activation energy of the products, then for the above reaction[EAMCET 1997]
 (a) E₁ > E₂
 - (b) $E_1 < E_2$
 - (c) $E_1 = E_2$
 - (d) ΔH is not related to E_1 and E_2
 - (e) None is correct
- **10.** The temperature dependence of rate constant (k) of a chemical reaction is written in terms of Arrhenius equation, $K = A \cdot e^{-E^*/RT}$. Activation energy (E^*) of the reaction can be calculated by plotting **[CBSE PMT 2003]**

(a)
$$\log k vs \frac{1}{\log T}$$
 (b) $k vs T$

(c)
$$kvs\frac{1}{\log T}$$
 (d) $\log kvs\frac{1}{T}$

Activation energy of a chemical reaction can be determined by[CBSE PMT 1998; AFMC 1999; BHU 2000]
(a) Changing concentration of reactants

(b) Evaluating rate constant at standard temperature

(c) Evaluating rate constants at two different temperatures

(d) Evaluating velocities of reaction at two [AIEEE 2003] different temperatures

12. The activation energy for a reaction is
9.0 K cal/mol. The increase in the rate constant
when its temperature is increased from 298K to
308K is(a) 63%(b) 50%(c) 100%(d) 10%

13. Which of the following is the fastest reaction

(a)
$$C + \frac{1}{2}O_2 \xrightarrow{250^\circ C} CO$$
 (b) $C + \frac{1}{2}O_2 \xrightarrow{500^\circ C} CO$
(c) $C + \frac{1}{2}O_2 \xrightarrow{750^\circ C} CO$ (d) $C + \frac{1}{2}O_2 \xrightarrow{1000^\circ C} CO$

14. The rate constant k, for the reaction $N_2O_5(g) \rightarrow 2NO_2(g) + \frac{1}{2}O_2(g)$ is $2.3 \times 10^{-2} s^{-1}$. Which equation given below describes the change of $[N_2O_5]$ with time? $[N_2O_5]_0$ and $[N_2O_5]_r$ correspond to concentration of N_2O_5 initially and at time, *t* [AIIMS 2004]

a)
$$[N_2O_5]_t = [N_2O_5]_0 + kt$$

(b)
$$[N_2O_5]_0 = [N_2O_5]_t e^{kt}$$

(c)
$$\log_{10}[N_2O_5]_t = \log_{10}[N_2O_5]_0 - kt$$

(d)
$$\ln \frac{[N_2 O_5]_0}{[N_2 O_5]_t} = kt$$

- **15.** For the reaction $CH_3COOCH_3 + H_2O \xrightarrow{H^+} CH_3COOH + CH_3OH$ The progress of the process of reaction is followed by
 - (a) Finding the amount of methanol formed at different intervals
 - (b) Finding the amount of acetic acid formed at different intervals
 - (c) Using a voltmeter
 - (d) Using a polarimeter
- **16.** Half life of a reaction is found to be inversely proportional to the cube of its initial concentration. The order of reaction is

17. The integrated rate equation is $Rt = \log C_0 - \log C_t$. The straight line graph is obtained by plotting [AIEEE 200

(a) time v/s logC_t (b)
$$\frac{1}{\text{time}}$$
 v / s C_t

(c) time v/s C_t (d)
$$\frac{1}{\text{time}}$$
 v/s $\frac{1}{C_t}$

18. For which order reaction a straight line is obtained along with *x*-axis by plotting a graph between half life $(t_{1/2})$ and initial concentration 'a'

19. The reaction, $X \rightarrow$ product follows first order kinetics. In 40 minutes the concentration of *X* changes from 0.1 *M* to 0.025 *M* Then the rate of reaction when concentration of *X* is 0.01 *M*

(a)
$$1.73 \times 10^{-4} M \text{ min}^{-1}$$
 (b) $3.47 \times 10^{-5} M \text{ min}^{-1}$

(c) $3.47 \times 10^{-4} M \min^{-1}$ (d) $1.73 \times 10^{-5} M \min^{-1}$

20. A Substance undergoes first order decomposition. The decomposition follows two parallel first order reactions as

$$A \xrightarrow{k_1}_{k_2} K_1 = 1.26 \times 10^{-4} \text{ s}^{-1}$$

$$k_1 = 3.8 \times 10^{-5} \text{ s}^{-1}$$

The percentage distribution of B and C are

[Kerala PMT 2004]

5.

6.

- (a) 75% *B* and 25% *C* (b) 80% *B* and 20% *C*
- (c) 60% *B* and 40% *C* (d) 90% *B* and 10% *C*
- (e) 76.83% *B* and 23.17% *C*



Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.

1. Assertion : Instantaneous rate of reaction is equal to dx / dt.

Reason : It is the rate of reaction at any particular instant of time.

2. Assertion : Molecularity has no meaning for a complex reaction.

Reason : The overall molecularity of a complex reaction is equal to the molecularity of the slowest step.

3. Assertion : If in a zero order reaction, the concentration of the reactant is doubled, the half-life period is also doubled.

- Reason : For a zero order reaction, the rate of reaction is independent of initial concentration.
- **4.** Assertion : If the activation energy of a reaction is zero, temperature will have no effect on the rate constant.
 - Reason : Lower the activation energy, faster is the reaction.

- Assertion : According to steady state hypothesis, in a multistep reaction, the change in concentration with time for reactive intermediates is zero.
 - Reason : The intermediates are so reactive that after a brief initial period their concentrations rise from zero to a small value and remains constant for most of the duration of the reaction.
- Assertion : Half-life period of a reaction of first order is independent of initial concentration.
 - Reason : Half-life period for a first order reaction $t_{1/2} = \frac{2.303}{K} \log 2$.
- 7. Assertion : The photochemical reactions $H_2 + Cl_2 \rightarrow 2HCl$ and $H_2 + Br_2 \rightarrow 2HBr$ have equal quantum efficiencies.
 - Reason : Both the reactions proceed by similar mechanism.
- 8. Assertion : Vision is not a photochemical reaction.
 - Reason : Halogenation of alkenes is a photochemical reaction.
- **9.** Assertion : Glow worm shows chemiluminescence.

Reason : Glow worm emits light due to oxidation of protein, luciferin present in it.

10. Assertion : The rate of reaction is always negative.

Reason : Minus sign used in expressing the rate shows that concentration of product is decreasing.

11. Assertion : The kinetic of the reaction $mA + nB + pC \rightarrow m'X + n'Y + p'Z$ obeys the rate expression as $\frac{dx}{dt} = k[A]^m[B]^n$.

Reason

:

The rate of the reaction does not depend upon the concentration of *C*.



1	b	2	a	3	d	4	b	5	b
6	c	7	c	8	с	9	а	10	b
11	b	12	b	13	C	14	d	15	b
16	b	17	C	18	b	19	d	20	а
21	b	22	a	23	d	24	с	25	а
26	а	27	а	28	b	29	b	30	b
31	а	32	C	33	b	34	а	35	c
36	d	37	а	38	b	39	а	40	С
41	a	42	b						

Rate law and Rate constant

1	d	2	C	3	а	4	b	5	a
6	а	7	a	8	С	9	b	10	b
11	b	12	d	13	а	14	c	15	d
16	d	17	C	18	а	19	С	20	С
21	b	22	a	23	а	24	d	25	b
26	а	27	a	28	С	29	С	30	d
31	d	32	C	33	С	34	d	35	b
36	С	37	C	38	а	39	С	40	b
41	a	42	d	43	а	44	С	45	c
46	а	47	b	48	b	49	С	50	b
51	b	52	a	53	а	54	d	55	d
56	с	57	d	58	b	59	b	60	b
61	с	62	d	63	c	64	b	65	c
66	d	67	b	68	С	69	С	70	b
71	d	72	d	73	b	74	b	75	d
76	b	77	b	78	c	79	c	80	c
81	b	82	b	83	c	84	а	85	a
86	С	87	C	88	а	89	С	90	С
91	b	92	b	93	С	94	b	95	b
96	d	97	b	98	С	99	а	100	С
101	b	102	d	103	c	104	c	105	a
106	d	107	a	108	b	109	C	110	b
111	C	112	d	113	b	114	b	115	d
116	d	117	b	118	d	119	а	120	b
121	d	122	b	123	d	124	с	125	d
126	b	127	d	128	d	129	b	130	а
131	ac	132	с	133	d	134	с	135	а
136	а	137	с	138	С	139	d	140	с
141	C	142	c	143	C	144	а	145	C
146	c	147	c	148	b	149	b	150	a
151	d	152	а	153	d	154	C	155	C
156	b	157	a	158	b	159	d	160	c
161	b	162	b	163	b	164	abd	165	ad
166	b	167	d	168	b	169	d	170	b
171	C	172	c	173	с	174	а		

Collision theory, Energy of activation and Arrhenius equation

1	b	2	b	3	b	4	C	5	a
6	d	7	d	8	b	9	C	10	b
11	d	12	d	13	а	14	d	15	а
16	b	17	d	18	b	19	b	20	а
21	C	22	а	23	d	24	b	25	а
26	ab	27	C	28	b	29	b	30	C
31	b	32	d	33	C	34	C	35	а
36	а								

Photochemical reaction

1	а	2	а	3	c	4	b	

Critical Thinking Questions

1	а	2	с	3	а	4	d	5	c
6	c	7	c	8	b	9	а	10	d
11	c	12	а	13	d	14	d	15	b
16	d	17	а	18	b	19	C	20	е

Assertion & Reason

1	b	2	b	3	b	4	b	5	a
6	а	7	d	8	е	9	а	10	d
11	а								



Rate of a reaction

1	b	2	a	3	d	4	b	5	b
6	C	7	C	8	С	9	а	10	b
11	b	12	b	13	С	14	d	15	b
16	b	17	C	18	b	19	d	20	а
21	b	22	a	23	d	24	c	25	a
26	a	27	a	28	b	29	b	30	b
31	а	32	C	33	b	34	а	35	C
36	d	37	а	38	b	39	а	40	C
41	a	42	b						

Rate law and Rate constant

1	d	2	с	3	а	4	b	5	а
6	а	7	а	8	c	9	b	10	b
11	b	12	d	13	a	14	С	15	d
16	d	17	C	18	a	19	C	20	c
21	b	22	а	23	a	24	d	25	b
26	а	27	а	28	c	29	С	30	d
31	d	32	С	33	c	34	d	35	b
36	с	37	C	38	a	39	С	40	b
41	а	42	d	43	a	44	С	45	c
46	а	47	b	48	b	49	С	50	b
51	b	52	а	53	a	54	d	55	d
56	c	57	d	58	b	59	b	60	b
61	c	62	d	63	c	64	b	65	c
66	d	67	b	68	c	69	C	70	b
71	d	72	d	73	b	74	b	75	d
76	b	77	b	78	c	79	с	80	c
81	b	82	b	83	c	84	а	85	a
86	c	87	с	88	a	89	с	90	c
91	b	92	b	93	C	94	b	95	b
96	d	97	b	98	С	99	а	100	C
101	b	102	d	103	C	104	с	105	a
106	d	107	а	108	b	109	С	110	b
111	c	112	d	113	b	114	b	115	d

116	d	117	b	118	d	119	a	120	b
121	d	122	b	123	d	124	C	125	d
126	b	127	d	128	d	129	b	130	а
131	ac	132	С	133	d	134	C	135	а
136	а	137	C	138	С	139	d	140	C
141	С	142	C	143	с	144	a	145	c
146	С	147	С	148	b	149	b	150	а
151	d	152	a	153	d	154	C	155	c
156	b	157	a	158	b	159	d	160	C
161	b	162	b	163	b	164	abd	165	ad
166	b	167	d	168	b	169	d	170	b
171	C	172	C	173	C	174	a		

Collision theory, Energy of activation and Arrhenius equation

1	b	2	b	3	b	4	c	5	а
6	d	7	d	8	b	9	c	10	b
11	d	12	d	13	а	14	d	15	а
16	b	17	d	18	b	19	b	20	а
21	C	22	а	23	d	24	b	25	a
26	ab	27	С	28	b	29	b	30	C
31	b	32	d	33	c	34	c	35	a
36	а								

Photochemical reaction

1	а	2	а	3	C	4	b	

Critical Thinking Questions

1	а	2	C	3	а	4	d	5	C
6	с	7	С	8	b	9	а	10	d
11	С	12	а	13	d	14	d	15	b
16	d	17	а	18	b	19	с	20	е

Assertion & Reason

1	b	2	b	3	b	4	b	5	а
6	а	7	d	8	е	9	а	10	d
11	а								



Rate of a reaction

- 1. (b) Rate of reaction continuously decreases with time.
- (a) The rate of reaction depends upon conc. of reactant, surface area of reactant, temperature, presence of light and catalyst.
- **3.** (d) According to law of mass action.
- 4. (b) R = K[RCl], if [RCl] = 1/2, then rate = R/2.

5. (b)
$$2^2 = 4$$
, $3^2 = 9$

6. (c) The rate of chemical reaction ∞ The product of the molar conc. of the reactants (at constant *T*)

7. (c) Rate of reaction =
$$\frac{dx}{dt} = \left[\frac{0.2 - 0.1}{10}\right] = \frac{0.1}{10}$$

$$= 0.01 \ mol \ dm^{-3} \ min^{-1}$$

8. (c) As reaction progressing the concentration of the reactants decreases and the concentration of the product increases.

9. (a)
$$\frac{-d(N_2)}{dt} = -\frac{1}{3}\frac{d(H_2)}{dt} = \frac{1}{2}\frac{d(NH_3)}{dt} = \frac{3}{2} \times 40 \times 10^{-3}$$

= 60 × 10⁻³.

- 10. (b) Greater are the concentrations of the reactants, faster is the reaction. Conversely, as the concentrations of the reactants decreases, the rate of reaction also decreases.
- **11.** (b) Ionic reactions are very fast reactions i.e. take place instantaneously.
- **12.** (b) Rate = $K(A)^2(B)^1$ on doubling the active mass of *A* the rate of reaction increase 4 times.
- 13. (c) 'A' will disappear at twice the rate at which 'B' will decrease.
- 14. (d) When volume is reduced to $\frac{1}{4}$, concentrations become four times.

16. (b)
$$\frac{-dN_2}{dt} = \frac{-1}{3}\frac{dH_2}{dt} = \frac{1}{2}\frac{dNH_3}{dt}$$

 $\frac{dH_2}{dt} = \frac{3}{2} \times 0.001 = 0.0015 \, kg \, hr^{-1}.$

18. (b) $-\frac{dc}{dt}$ refers as decrease in concentration of the reactant with time.

19. (d) The rate of a reaction depends upon concentration of reactant.

20. (a)
$$-\frac{1}{3}\frac{d[A]}{dt} = -\frac{d[B]}{dt} = \frac{+d[C]}{dt} = \frac{+d(D)}{dt}$$

21. (b) $N_2 + 3H_2 = 2NH_3$

$$\frac{-\Delta[N_2]}{\Delta t} = -\frac{1}{3} \frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[NH_3]}{\Delta t}$$
$$\therefore \frac{\Delta[H_2]}{\Delta t} = \frac{3}{2} \times \frac{\Delta[NH_3]}{\Delta t} = \frac{3}{2} \times 2 \times 10^{-4}$$
$$= 3 \times 10^{-4} \ mol \ litre^{-1} sec^{-1}$$

22. (a) Increase in concentration of $B = 5 \times 10^{-3} mol l^{-1}$ Time = 10 sec

Rate of appearance of $B = \frac{\text{Increase of conc. B}}{\text{Time taken}}$

$$=\frac{5\times10^{-3}\,mol\ l^{-1}}{10\ \sec}=5\times10^{-4}\,mol\ l^{-1}\ Sec\ ^{-1}$$

24. (c) The rate of formation of SO_3 is $1.28 \times 10^{-3} \text{ g/sec}$.

26. (a)
$$\frac{K_t + 10}{K_t} = \frac{r_t + 10}{r_t} = 2$$
; For an increase of

temperature to $50^{\circ}C$, *i.e.* 5 times, the rate increases by 2^{5} times, *i.e.* 32 times.

28. (b)
$$\frac{K_t + 10}{K_t} = \frac{r_t + 10}{r_t} = 2$$
.

For an increase of temperature to $20^{\circ}C$ *i.e.* 2 times, the rate increase by 2^2 times, *i.e.* 4 times.

29. (b)
$$\frac{k_t + 10}{K_t} = \frac{r_t + 10}{r_t} = 2$$

For an increase of temperature to $90^{\circ}C$ *i.e.* 9 times, the rate increases by 2^{9} times *i.e.* 512.

- **30.** (b) Catalyst increases the rate by decreasing the activation energy.
- **31.** (a) For 10K rise in temperature, the rate of reaction nearly doubles.
- **32.** (c) Temperature coefficient $\frac{K_{35^{\circ}C}}{K_{25^{\circ}C}} = \frac{K_{308k}}{K_{298k}} = 2$ and 3 for most reactions.
- **33.** (b) Catalyst decrease energy of activation.
- **34.** (a) Thus both rate and rate constant *K* increase with temperature, $r = k(\text{reactant})^n$ and $k = Ae^{-E_a/RT}$
- **35.** (c) Enzymes does not always increase activation energy.

- **36.** (d) Catalyst reduce the activation energy for reaction and thus increase the rate of reaction.
- 37. (a) Catalyst affect only activation energy. It brings down the activation energy of reaction.
- **38.** (b) As we know that the velocity constant become double by increasing the temperature by $10^{\circ}C$ so if at 290 K, velocity constant = 3.2×10^{-3} then at 300 K, velocity constant = $2(K_{290}) = 2 \times 3.2 \times 10^{-3} = 6.4 \times 10^{-3}$.
- **39.** (a) Higher the value of rate constant so, faster the reaction rate.
- 40. (c) Rate of reaction may increase or decrease with increase in temperature. If reaction is exothermic, rate decreases with increasing temperature while that of endothermic reactions increase with increasing temperature.
- 41. (a) $2N_2O_{5(g)} \rightarrow 4NO_{2(g)} + O_{2(g)}$ Rate of reaction with respect to NO_2

$$=\frac{1}{4}\frac{d[NO_2]}{dt}=\frac{1}{4}\times\frac{5.2\times10^{-3}}{100}=1.3\times10^{-5}\,ms^{-1}$$

42. (b) In first phase,
$$K = \frac{2.303}{20} \log \frac{90}{100}$$
(i)

In second phase $K = \frac{2.303}{t} \log \frac{81}{100}$ (ii) From eq. (i) $\frac{2.303}{20} \log \frac{90}{100} = \frac{2.303}{t} \log \frac{81}{100}$

$$t = \frac{20(\log 81 - \log 100)}{(\log 90 - \log 100)}$$
$$= \frac{20(1.908 - 2)}{(1.954 - 2)} = \frac{20 \times (-0.092)}{(-0.046)} = 40 \text{ minutes}$$

Rate law and Rate constant

- (d) Molecularity of the reaction does not influence the rate of reaction.
- 2. (c) The rate law for an reaction can be given by; rate $= K(A)(B)^2$, *i.e.* the powers are raised which are given as coefficient of reactant.
- **3.** (a) $p_0 = 500 atm$

$$K = \frac{2.303}{t} \log_{10} \frac{p_0}{p_t}$$

3.38 × 10⁻⁵ = $\frac{2.303}{10 \times 60} \log \frac{500}{p_t}$
or 0.00880 = $\log \frac{500}{p_t} \Rightarrow \frac{500}{1.02} = 490 atm$

- **4.** (b) It is a pseudo-unimolecular reaction.
- $5. \qquad (a) \quad A+2B \to C+D$

$$\frac{-d[A]}{dt} = 5 \times 10^{-4}$$

$$-\frac{1}{2}\frac{d[B]}{dt} = \frac{5 \times 10^{-4}}{2} = 2.5 \times 10^{-4} \, mol^{-1} \, scc^{-1}$$

8. (c) For

reaction, $2NO_{(g)} + O_2(g) \rightarrow 2NO_2(g)$

When the volume of vessel change into $\frac{1}{3}$ then concentration of reactant become three

following

times.

The rate of reaction for first order reaction \propto concentration. So rate of reaction will increases three times.

9. (b) $2A + B \rightarrow$ Products

According to question : Rate of reaction of 'A' \propto [B] as increase in rate is double when [B] is doubled.

Rate of reaction \propto [A] [B] as increase in rate is 8 times when concentration of both reactant is doubled. It means that order of reaction is 3 and overall rate reaction should be $r = K[A]^2[B]$

10. (b) Let the rate of reaction depends on xth power of [A]. Then

$$r_1 = k[A]^x$$
 and $r_2 = k[2A]^x$
, $r_1 = [A]^x = 1 = (1)^2$

 $\therefore \frac{r_1}{r_2} = \frac{r_1}{[2A]^x} = \frac{1}{4} = \left(\frac{1}{2}\right) \quad (\because r_2 = 4r_1)$ $\therefore x = 2. \text{ As the reaction rate does not depend$

 $\therefore x = 2$. As the reaction rate does not depend upon the concentration of B. Hence, the correct rate law will be rate = $K[A]^2[B]^o$ or = $K[A]^2$

- (b) Velocity constant 'K' is characteristic constant of a reaction and depends only on temperature and catalyst.
- **12.** (d) Unit of K for Ist order reaction is $Time^{-1}$.
- **13.** (a) $K = Ae^{-E_a/RT}$ by this equation it is clear that rate constant of a reaction depends on temperature

14. (c)
$$k = \frac{2.303}{t} \log_{10} \frac{a}{a-x}$$
;

$$t = 2 \times 10^2$$
, $a = 800$, $a - x = 50$

$$k = \frac{2.303}{2 \times 10^2} \log_{10} \frac{800}{50} = \frac{2.303}{2 \times 10^2} \log_{10} 16$$
$$= \frac{2.303}{2 \times 10^2} \log_{10} 2^4 = \frac{2.303}{2 \times 10^4} \times 4 \times 0.301$$
$$= 1.38 \times 10^{-2} \ s^{-1}$$

- **15.** (d) Rate of reaction is quadrupled on doubling the concentration. Thus $r \propto [A]^2$.
- **16.** (d) *K* is not constant for all the reaction.
- **19.** (c) Rate determining step is a slow step by which rate of reaction can be determined.
- **20.** (c) It is a second order reaction and the unit of k for second order reaction is *litre* mol^{-1} sec⁻¹.

- **21.** (b) For reversible reaction rate constant is also reverse.
- **22.** (a) Rate = $K(N_2O_5) = 6.2 \times 10^{-1} \times 1.25$ = $7.75 \times 10^{-1} mol \ l^{-1} \ s^{-1}$
- 23. (a) We know that for first order kinetics

$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

(a-x) = 0.05 mol l⁻¹, 6 = $\frac{2.303}{t} \log \frac{0.5}{0.05}$
or $t = \frac{2.303}{6} \log \frac{0.5}{0.05} = \frac{2.303}{6} = 0.384$ min

 $K(N_2O_5)$

24. (d) Rate =

 $2.4 \times 10^{-5} = 3.0 \times 10^{-5} (N_2 O_5)$ or $(N_2 O_5) = 0.8 \ mol \ l^{-1}$

25. (b) R = K[A], $1.02 \times 10^{-4} = 3.4 \times 10^{-5}$, $[N_2O_5]$

or
$$(N_2O_5) = \frac{1.02 \times 10^{-4}}{3.4 \times 10^{-5}}, K = 3$$

- **26.** (a) Rate law for the reaction $2N_2O_5 \rightarrow 4NO_2 + O_2$ is $r = k[N_2O_5]$ first order reaction.
- **28.** (c) Rate constant depends on temperature only.
- **29.** (c) According to Arhenius concept Rate constant $K = A_e^{-E_a/RT}$, Hence rate constant depends only upon temperature of the system.
- **30.** (d) For $2A + B \rightarrow C$ Rate = K[A][B]

Value of rate constant $K = Ae^{-Ea/RT}$ here K is independent of the initial concentration of A and B.

- 31. (d) The specific rate constant of a first order reaction depends upon the temperature of reaction.
- **32.** (c) Unit of *K* for first order reaction $= sec^{-1}$
- **35.** (b) The rate of zero order reaction is not depend on the concentration of the reactants.
- **36.** (c) The unit of *K* for zero order reaction = mole litre⁻¹sec⁻¹.
- **37.** (c) Order of reaction is sum of the power raised on concentration terms to express rate expression.
- **38.** (a) The concentration of reactant does not change with time for zero order reaction (unit of *K* suggests zero order) since reactant is in excess.

39. (c)
$$r = K[A]^m$$
 also $2r = K[4A]^m$, $\frac{1}{2} = \left(\frac{1}{4}\right)^m$
 $\therefore m = \frac{1}{2}$

40. (b) $K = \frac{2.303}{t} \log \frac{V_{\infty}}{V_{\infty} - V_t}$ gives constant value of K.

With the given date. Hence it is 1st order.

42. (d) $r = K[A]^n$, $100 r = K[10 A]^n$

Thus
$$\frac{1}{100} = \left(\frac{1}{10}\right)^n$$
 or $n = 2$

- 43. (a) It is a standard example of first order because in that reaction rate of reaction affected by only one concentration term.
- **44.** (c) Inversion of cane sugar is a Pseudo unimolecular reaction.
- **45.** (c) For 1st order reaction

hence

$$K = \frac{2.303}{t} \log_{10} \frac{a}{(a-x)}$$

46. (a) $t_{1/2} = \frac{0.693}{k} = \frac{0.693}{6.2 \times 10^{-4}} = 1117.7$ sec.

17. (b)
$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

 $\frac{0.693}{T} = \frac{2.303}{t} \log \frac{100}{100-30}$
 $\therefore T = 58.2 \text{ min.}$

- **48.** (b) Order of reaction = $\frac{3}{2} + \frac{3}{2} = \frac{3+3}{2} = \frac{6}{2} = 3$
- (c) Two molecules are taking part in elementary step.
- **50.** (b) Because two molecules are taking part in elementary step.
- 51. (b) The overall order of a reaction is sum of powers T raised on concentration terms in order to write rate expression.
- **52.** (a) It is a first order reaction as is clear from rate law expression, $r = k(H_2O_2)$
- **53.** (a) For Ist order reaction half life is independent of concentration.

54. (d)
$$K = \frac{1}{t} \log e\left(\frac{a}{a-x}\right) = \frac{1}{15} \log e\left(\frac{35}{35-9}\right)$$

$$=\frac{1}{15}\log e\left(\frac{35}{26}\right)$$

56. (c)
$$k = \frac{0.693}{30} = 0.0231$$
 ; $t = \frac{2.303}{k} \log\left(\frac{100}{100 - 75}\right)$
 $t = \frac{2.303}{0.0231} \log 4 = 60$ minutes

57. (d) *Rate* =
$$K(sugar)(H_2O)^o$$

58. (b) Derive $t_{1/2}$ from $K_t = 2.303 \log \frac{a}{a-x}$, putting $t = \frac{1}{2}$ and $x = \frac{a}{2}$. Therefore it is $\frac{0.693}{K}$.

59. (b)
$$r \propto [X]^2$$
 or $r = k[X]^2$

60. (b) $t_{1/2} = \frac{0.693}{k}, \frac{0.693}{1.1 \times 10^{-9}} = 6.3 \times 10^8$ sec.

61. (c) $t_{1/2} \propto \frac{1}{(a)}$ for II order reaction.

- **62.** (d) Order of reaction is an experimental value, while molecularity is a theoretical value.
- **63.** (c) K for 1^{st} order reaction = per unit time *i.e.* $Time^{-1}$.
- **64.** (b) $R = K[A][B]^0$ so molecularity is two and order is two.
- **65.** (c) Rate of zero order reaction is independent of the concentration of the reactant and remains constant throughout the reaction.

66. (d)
$$t_{1/2} \propto \frac{1}{(a)^{n-1}}$$

67. (b) $t_{1/2} \propto \frac{1}{(C_o)^{n-1}}$: reaction is of first order.

68. (c) For Ist order reaction half life is independent of concentration.

69. (c)
$$K = \frac{0.693}{t_{1/2}}, K = \frac{0.693}{138.6 \text{ min}} = 0.005 \text{ min}^{-1}$$

70. (b) $CH_3COOCH_3 + H_2O \xrightarrow{H^+} CH_3COOH + CH_3OH$ It is a pseudo-unimolecular reaction.

71. (d)
$$t_{1/2} \propto \frac{1}{K}$$
 and $K \propto t$

- **73.** (b) For zero order reactions $\frac{dx}{dt} = K(A)^o$
- **74.** (b) It is an bimolecular and second order reaction.
- **75.** (d) $\frac{dx}{dt} = (HI)^2$, then order of reaction = 2
- **76.** (b) Rate = $K(\text{Sucrose})(H_2O)^o$
- 77. (b) Because in this reaction one molecule of N_2O_5 is used.
- **78.** (c) Integrated velocity equation for first order reaction is: $k = \frac{2.303}{t} \log \frac{(A)_o}{(A)}$

80. (c)
$$t_{1/2} = \frac{2.303 \log 2}{K} = \frac{0.693}{K}$$

- **81.** (b) Inversion of sugar is a pseudo-unimolecular reaction.
- **82.** (b) For Ist order reaction

$$\log(a-x) = \log a - \frac{kt}{2.303}$$

y = c + mx

83. (c)
$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{3.46 \times 10^{-3}} = 200 \text{ min}$$

84. (a)
$$\frac{1}{Time} \times (conc.) = \frac{moles \ l^{-1}}{Time} = moles \ l^{-1}time^{-1}$$
 for zero order reaction.

- **85.** (a) The order of reaction is $\frac{3}{2}$ and molecularity is
- **87.** (c) Molecularity can never be fractional.
- **88.** (a) As doubling the initial conc. doubles the rate of reaction, order = 1.
- **89.** (c) When *B* is in excess, it becomes a pseudo-unimolecular reaction.

90. (c)
$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{69.35} = 9.99 \times 10^{-3} = 0.01 \, s^{-1}$$

91. (b)
$$k = \frac{0.693}{24} hr^{-1} = \frac{2.303}{96} \log \frac{1}{a}$$

or
$$\log \frac{10}{a-x} = 1.2036$$
 or $1 - \log (a - x) =$

1.2036

С

or
$$\log(a - x) = -0.2036 = 1.7964$$

or
$$(a-x) =$$
antilog 1.7964 $= 0.6258 \ gm$

- **92.** (b) 0.08 mol l^{-1} to 0.01 mol l^{-1} involves 3 half-life. So the *t* is 30 minutes
- **93.** (c) $t_{1/2}$ of II order reaction is inversely proportional to initial concentration of reactants.
- **94.** (b) As $r = k(H_2O_2)$, it is a reaction of 1st order.

97. (b)
$$r = k(C_A)^{3/2}(C_B)^{-1/2}$$

Order $= \frac{3}{2} + \left(-\frac{1}{2}\right) = \frac{2}{2} = 1$

98. (c) If rate = $K(A)^m(B)^n$, then order of reaction = m + n.

99. (a)
$$k = \frac{0.693}{t_{1/2}} = \frac{0.693}{100 \text{ sec}} = 6.93 \times 10^{-3} \text{ sec}^{-1}$$

100. (c)
$$t_{1/2} = \frac{0.693}{k}$$

101. (b)
$$t_{1/2} = \frac{0.693}{k}, k = \frac{0.693}{480} = 1.44 \times 10^{-3} \text{ sec}^{-1}$$

102. (d) $r = k(A)^2$, when concentration is doubled $r = k(2A)^2 = k 4(A)^2$ the rate becomes 4 times.

103 (c)
$$r = K[FeCl_3]^2 [SnCl_2]^1$$
. Order $= 2 + 1 = 3$

105. (a) $t_{1/2}$ for I order reaction independent of initial concentration.

107. (a) The rate will be given by slowest step. Thus

$$r = K[A][B_2] \cdot K_c = \frac{[A][A]}{[A_2]}$$
 or $[A] = [K_c]^{1/2} [A_2]^{1/2}$
 $r = K \times [K_c]^{1/2} [A_2]^{1/2} [B_2] = K[A_2]^{1/2} [B]$. Thus order
is $0.5 + 1 = 1.5$

108. (b) For Ist order reaction half life is independent of concentration.

110. (b) Rate =
$$K[A]^{1/2}[B]^{3/2}$$

$$\therefore O.R. = \frac{1}{2} + \frac{3}{2} = \frac{4}{2} = 2$$

112. (d) The rate of this photochemical reaction is independent of the concentration, therefore, it is zero order reaction.

113. (b)
$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{0.6932 \ hr^{-1}} = 1 \ hr$$
.

114. (b) The unit of rate constant shows that reaction is of first order. For first order reaction, half life is independent of initial conc. of the reactant. Thus,

$$t_{1/2} = \frac{0.693}{0.69 \times 10^{-1}} = \frac{0.693 \times 60}{0.69 \times 10^{-1}} = 600 \text{ sec}$$

115. (d) *Given* : Rate constant of the first order reaction $(K) = 3 \times 10^{-6}$ per sec and initial concentration [A] = 0.10M. We know that initial rate constant

$$K[A] = 3 \times 10^{-6} \times 0.10 = 3 \times 10^{-7} ms^{-1}$$

- **116.** (d) It is the characteristic of pseudo-unimolecular reactions.
- **117.** (b) It is a second order reaction.
- **118.** (d) r = K [reactant]

$$\therefore K = \frac{1.0 \times 10^{-2}}{0.2} = 0.05$$
$$t_{1/2} = \frac{0.693}{0.05} = 13.86 s$$

- **119.** (a) For first order reactions rate is depend on the concentration of one reactant.
- **120.** (b) Molecularity of a reaction never become zero or fraction.

121. (d)
$$t = \frac{2.303}{k} \log \frac{a}{a-x}$$

 $t = \frac{2.303}{1.155 \times 10^{-3}} \log \frac{100}{100-50} = 600 \text{ sec}$

122. (b) Rate = $K[A]^{3/2}[B]^{-1}$

:
$$O.R. = \frac{3}{2} + (-1) = \frac{1}{2}$$

- **123.** (d) $t_{1/2} \propto (CO)^o$ *i.e.* half life for Ist order is independent of initial concentration.
- **124.** (c) $2NO_{(g)} + Cl_{2(g)} \rightarrow 2NOCl_{(g)}$

Rate =
$$K[NO]^2[Cl_2]^1$$
, $\therefore O.R. = 2 + 1 = 3$
125. (d) $8 \times 10^{-5} = \frac{1}{t} \left[\frac{1}{0.5} - \frac{1}{1} \right]$; $8 \times 10^{-5} = \frac{1}{t} [2 - 1]$

$$t = \frac{1}{8 \times 10^{-5}} = 0.125 \times 10^5 = 1.25 \times 10^4 \text{ min.}$$

126. (b)
$$r = k [\text{reactant }]^{-1} \therefore k = \frac{0.693 \times 10^{-2}}{1} \text{ also}$$

 $t_{1/2} = \frac{0.693}{k} = \frac{0.693}{0.693 \times 10^{-2}} = 100 \text{ min}$.

127. (d) $t_{1/2} = \frac{1}{Ka}$ for second order reactions.

128. (d)
$$k = \frac{2.303}{t} \log\left(\frac{a}{a-x}\right); \ k = \frac{2.303}{30} \log\left(\frac{100}{100-75}\right)$$

 $k = \frac{2.303}{t} \log\left(\frac{100}{100-93.75}\right)$ Put the value of K

from above equation we get the value of t therefore $\therefore t = 60 \text{ min}$.

129. (b)
$$k = \frac{0.693}{45} \min^{-1} = \frac{2.303}{t_{99.9\%}} \log \frac{a}{a - 0.999 a}$$
 or
 $t_{99.9\%} = \frac{2.303 \times 45}{0.693} \log 10^3 = 448 \min \approx 7.5 \ hrs$

130. (a) Given A(a) = 2.00 m, t = 200 min and a(a-x) = 0.15 m we know

$$k = \frac{2.303}{t} \log \frac{a}{a-x} = \frac{2.303}{200} \log \frac{2.00}{0.15}$$
$$= \frac{2.303}{200} \times (0.301 + 0.824) = 1.29 \times 10^{-2} \operatorname{min}^{-1}$$

- **131.** (a,c) The unit for zero order reaction is $mol \ litre^{-1}time^{-1}$
- **132.** (c) It is a third order reaction because Rate = $K[NO]^2[O_2]^1$ $\therefore O.R. = 2+1=3$
- **133.** (d) Order of a reaction is decided by relative concentration of reactants.
- **134.** (c) Unit of rate constant for second order reaction is $mol^{-1}litre \ time^{-1}$
- **135.** (a) $R = K[A]^2[B]$ order of reaction = 2 + 1 = 3
- **136.** (a) Units of 1^{st} rate constant order reaction are sec^{-1} and for zero order reaction, these are $mol \ litre^{-1} \ sec^{-1}$ i.e. $M \ sec^{-1}$.
- **137.** (c) This reaction is bimolecular and first order of reaction.

138. (c)
$$t_{1/2} = \frac{0.693}{k}$$
 Given $t_{1/2} = 693$ sec
 $693 = \frac{0.693}{k}$, $k = \frac{693 \times 10^{-3}}{693}$; $k = 10^{-3} = 0.001$ sec⁻¹.

139. (d) $2A + B \to C + B$ Rate = $k[A]^2[B]^1$

0 602

$$\therefore O.R. = 2 + 1 = 3$$
 and molecularity is $3[2A + B]$.

140. (c) In photochemical reactions the rate of reaction is independent of the concentration of reacting species.

141. (c) We know that
$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

 $10^{-3} = \frac{2.303}{t} \log \frac{a}{(a-\frac{2a}{3})}, 10^{-3} = \frac{2.303}{t} \log 3$
 $10^{-3} = \frac{2.303}{t} \times 0.4771, t = \frac{2.303 \times 0.4771}{10^{-3}} = 3300 \text{ sec}$
142. (c) $R = K(A)^2, R' = K(2A)^2, \therefore \frac{R'}{R} = 4$

$$R = K(A)^{2}, R' = K(3A)^{2}, \frac{R}{R} = 9$$

144. (a) $k = \frac{2.303}{t} \log\left(\frac{a}{a-x}\right)$
 $k = \frac{2.303}{32} \log\left(\frac{100}{100-75}\right) \quad \dots (i)$
 $k = \frac{2.303}{t} \log\left(\frac{100}{100-50}\right) \quad \dots (ii)$

from the two equation (i) and (ii), t = 16 minutes.

ית

146. (c) The relation between half – life period and initial concentration (c) for a nth order reaction is given by $t_{1/2} \propto \frac{1}{C^{n-1}}$ for first order reaction (n = 1). $t_{1/2} \propto \frac{1}{C^{1-1}}$ or $t_{1/2} \propto C^0$.

147. (c)
$$R = k[NO]^{2}[O_{2}], R' = k[2NO]^{2}[2O_{2}]$$

 $R' = k \times 4[NO]^{2}[O_{2}] = k \times 8[NO]^{2}[O_{2}]$
 $\frac{R'}{R} = \frac{k \times 8[NO]^{2}[O_{2}]}{k[NO]^{2}[O_{2}]} = 8$

148. (b) For zero order reaction r = k.

149. (b)
$$k = \frac{2.303}{t} \log \frac{0.8}{0.6} = 2.303 \log \frac{4}{3}$$

 $t = \frac{2.303}{k} \log \frac{0.9}{0.675} = \frac{2.303}{k} \log \frac{4}{3}$; $t = 1$ hour.

150. (a) For zero order reaction

Velocity constant
$$= \frac{dx}{dt} = \frac{\text{Concentrat ion}}{\text{Time}}$$

Unit = concentrat ion \times time⁻¹.

151. (d) $H_2 + Br_2 \approx 2HBr$ is a 1.5 order reaction

i.e.,
$$K = [H_2][Br_2]^{1/2}$$
.

152. (a) When in any chemical reaction, one of the reactant is present in large excess, then the second order reaction becomes first order reaction and is known as pseudo unimolecular reaction *e.g.*,

$$CH_{3}COOCH_{3} + H_{2}O \xrightarrow{H^{+}} CH_{3}COOH + CH_{3}OH$$

in this reaction molecularity is 2 but order of reaction is found to be first order experimentally, so it is an example of pseudo unimolecular reaction.

153. (d)
$$K = \frac{0.693}{t_{1/2}} = \frac{0.693}{10 \text{ years}}$$

If initial concentration a = 10 gm and final

concentration
$$x = \frac{a}{2} = 5 \ gm$$

then, $t = \frac{2.303}{K} \log \frac{a}{a-x} = \frac{2.303}{.693} \times 10 \times \log \frac{10}{5}$

 $=\frac{2.303\times10\times\log 2}{.693}=\frac{2.303\times10\times0.301}{0.693}=10 \text{ years }.$

154. (c) The concentration of the reactants decrease from 0.8 to 0.4 in 15 *min* i.e., $T_{1/2} = 15$ *min*, concentration from 0.1*m* to 0.025 will fall in 2 half lives so total time taken $= 2 \times T_{1/2} = 2 \times 15 = 30$ *min*.

155. (c)
$$K = \frac{2.303}{1 hr} \log \frac{100}{25} = \frac{2.303}{t} \log \frac{100}{50}$$

∴ $\log 4 = \frac{1}{t} \log 2$
∴ $2 \log 2 = \frac{1}{t} \log 2$; $t = \frac{1}{2} hr$.

156. (b)
$$x_{(g)} \rightarrow y_{(g)} + z_{(g)}$$

The reaction is a first order reaction hence, 0.693 - 2.303 = a = 0.693

$$K = \frac{0.093}{t_{1/2}} = \frac{2.303}{t} \log \frac{a}{a-x} = \frac{0.093}{10 \text{ min}}$$
$$= \frac{2.303}{t} \log \frac{a}{a/10} = \frac{0.693}{10} = \frac{2.303}{t} \log 10$$
$$\therefore t = \frac{2.303 \times 10}{.693} = 33 \text{ min} .$$

157. (a) For the first order reaction $t = \frac{2.303}{K} \log \frac{a}{a-x}$ Given: a = 0.5 mol / litre; a - x = 0.05, mol / litre $K = 6 \sec^{-1}$.

$$t = \frac{2.303}{6} \log \frac{0.5}{0.05} = \frac{2.303}{6} \log 10 = \frac{2.303}{6} = 0.384 \text{ sec} .$$

158. (b) The radioactive disintegration reactions are of first order because in this rate of disintegration depends on the concentration term of radioactive material only.

159. (d) Rate
$$\left(\frac{dx}{dt}\right) = K \cdot c$$
; $1.5 \times 10^{-2} = K \times 0.5$
For first order $K = \frac{1.5 \times 10^{-2}}{0.5} = 3 \times 10^{-2} \text{ min } ute^{-1}$
 $t_{1/2} = \frac{.693}{K} = \frac{.693}{3 \times 10^{-2}} = 23.1 \text{ min } ute$.

160. (c) For first order reaction
$$K = \frac{2.303}{t} \log \frac{a}{a-x}$$

Given:
$$a = \frac{1}{10} = .1m$$
; $a - x = \frac{1}{100} = .01m$; $t = 500$

sec

$$\therefore K = \frac{2.303}{500} \log \frac{.10}{.01} = \frac{2.303}{500} \log 10$$
$$= \frac{2.303}{500} = 0.004606 = 4.6 \times 10^{-3} \text{ sec}^{-1}.$$

- **161.** (b) For zero order reaction, rate of reaction is independent of concentration $R = K[\text{Reactant}]^0$
- 162. (b) The radioactive disintegration reactions are of first order because in this rate of disintegration depends on the concentration term of radioactive material only.

163. (b) $t_{1/2} \propto a^{1-n} \Rightarrow \frac{0.1}{0.4} = \frac{(200)^{1-n}}{(50)^{1-n}} \Rightarrow \frac{1}{4} = \left[\frac{4}{1}\right]^{1-n} = \left[\frac{1}{4}\right]^{n-1}$ $\Rightarrow \frac{1}{4^1} = \frac{1}{4^{n-1}} \therefore n-1 = 1; n = 2.$

164. (abd) By Vant's Hoff equation, $\frac{d \ln k}{dt} = \frac{\Delta H^o}{RT^2}$

or $\ln k_p = -\frac{\Delta H^o}{RT} + I$. Hence (a) is correct (b) is also correct as plot of log (X) vs time is linear. (c) is wrong because $p \propto T$ at constant volume. (d) is correct by Boyle's law.

165. (ad) (a) is correct because degree of dissociation =1-e^{-kt} at any time *t*. (b) is wrong because plot of log [A] vs *t* is a straight line (c) is wrong because time taken for 75% reaction is two half life. (d) is correct because in k = Ae^{-Ea/RT}, Ea/RT is dimensionless hence *A* has the unit of *K*.
166. (b) aA → xP Rate of reaction = [A]^a

Rate of reaction = $[A]^{*}$ Order of reaction = a $[A]_1 = 2.2 \ mM, \ r_1 = 2.4 \ mM \ s^{-1}$...(i)

$$[A]_2 = 2.2/2 \ mM, \ r_1 = 0.6 \ mM \ s^{-1} \ or, \ \frac{2.4}{4} \ ...(ii)$$

On reducing the concentration of *A* to half, the rate of reaction is decreased by four times. Rate of reaction = $[A]^2$ Order of reaction = 2.

167. (d) Order of a reaction can be fractional.

168. (b)
$$t_{1/4} = \frac{2.303}{K} \log \frac{1}{1 - \frac{1}{4}} = \frac{0.29}{K}$$

169.

(b)
$$R = k[B]^n$$
; $\frac{1}{2}R = k[2B]^n$; $4 = \left(\frac{1}{2}\right)^n$; $4 = 2^{-n}$;

170. (b)
$$R = k[B]^n$$
; $\frac{-}{4}R = k[2B]^n$; $4 = \left(\frac{-}{2}\right)$;
 $n = -2$.

171. (c) $T = t_{1/2} \times n$ i.e. $12 = 3 \times n \Rightarrow n = 4$

$$N = N_0 \left(\frac{1}{2}\right) \implies \frac{N}{N_0} = \left(\frac{1}{2}\right) = \frac{1}{16}$$

172. (c)
$$K = 1.7 \times 10^{-5} s^{-1}$$

 $t_{1/2} = \frac{0.693}{K} = \frac{0.693}{1.7} \times 10^{5} = 11.32h$
173. (c) $A + B \to C$

On doubling the concentration of A rate of reaction increases by four times. Rate $\propto [A]^2$

However on doubling the concentration of B, rate of reaction increases two times. Rate $\propto [B]$ Thus, overall order of reaction = 2 + 1= 3.

174. (a) In case of zero order reaction, the concentration of reactant decreases linearly with time, as its rate is independent of the concentration of the reactants.

Collision theory, Energy of activation and Arrhenius equation

- (b) The increase in collision frequency brings in an increase in effective collisions and thus rate of reaction increases.
- (b) All collisions are not effective and does not result in the formation of the products
- (b) The increase in collision frequency brings in an increase in effective collisions and thus rate of reaction increases.
- (c) The increase in collision frequency brings in an increase in effective collisions and thus rate of reaction increases.
- 5. (a) Slower reaction rate indicates higher energy of activation.
- (d) Energy of activation decreases, rate of reaction will increases.
- (d) Number of collision depend upon pressure, concentration and temperature.
- **9.** (c) The definition of activation energy.
- (b) Activation energy is the energy needed by reactant molecules to gain threshold energy level.
- **11.** (d) All other are different forms of Arrhenius equation.
- 12. (d) Increase in the rate of reaction is determined by the increase in the number of effective collisions.
- **13.** (a) Energy of activation reduced by increasing temperature.
- 14. (d) The definition of threshold energy.
- **15.** (a) The definition of activation energy.
- 16. (b) When the temperature is increased, heat energy is supplied which increases the kinetic energy of the reacting molecules. this will increase the number of collisions and ultimately the rate of reaction will be enhanced.
- 17. (d) In exothermic and endothermic reactions will be more and less than E_a respectively.
- **19.** (b) The value of activation energy for a chemical reaction primarily dependent on the nature of reacting species.
- **20.** (a) Arrhenius equation is $\log k = \log A \frac{E_a}{DT}$

- **22.** (a) A graph plotted between $\log k$ vs $\frac{1}{T}$ for calculating activation energy.
- **24.** (b) When $E_a = 0$ rate constant is independent of temperature.
- **25.** (a) It is modified form of Arrhenius equation.

(b) A graph plotted between $\log k \operatorname{Vs} \frac{1}{T}$ calculating activation energy is shown as from Arrhenius equation

$$\log k = \log A - \frac{E_A}{2.303 \ RT} \qquad \qquad \log k \boxed{1/T \rightarrow}$$

30. (c) It is Arrhenius equation.

28.

31. (b)
$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$$

If $\frac{K_2}{K_1} = 2$
 $\log 2 = \frac{E_a}{2.303 \times 8.314} \left[\frac{1}{300} - \frac{1}{310} \right]$
 $E_a = .3010 \times 2.303 \times 8.314 \left(\frac{300 \times 310}{10} \right)$
 $= 53598.59 \ Jmol^{-1} = 54 \ kJ$.

32. (d) When $E_a = 0$, the rate of reaction becomes independent of temperature.

(E_a = Energy of activation).

- **33.** (c) When *k* increases, rate of reaction also increases, $k = \frac{pz}{e^{E/RT}}$ for k to increase *p*, *z*, *T* should increase and *E* should decrease. ($e \approx 2.7$).
- **34.** (c) Endothermic reactions are those which involve absorption of heat. High activation energy means potential energy of product must be much greater than reactants.
- **35.** (a) For endothermic reaction $\Delta H = +ve$

Then from equation $\Delta H = E_{a_{F.R.}} - E_{a_{B.R.}}; E_{B.R.} < E_{F.R.}$

36. (a) Arrhenius suggested an equation which describes rate constant (*K*) as a function of temperature.

 $K = Ae^{-E_a/RT}$

 $\ln K = \ln A - e^{E_a/RT}$

Photochemical reactions

- **2.** (a) $H_2O \xrightarrow{\text{Photolysis}} OH^- + H^+$
- **3.** (c) Stark Einstein was given the law of photochemical equivalence.

 (b) In photochemical reaction the rate of formation of product is directly proportional to the intensity of absorbed light.

Critical Thinking Questions

1. (a) For the given reaction :

for

2.

8.

$$-\frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[HI]}{\Delta t}$$

(c)
$$-\frac{1}{3} \frac{d[A]}{dt} = \frac{1}{2} \frac{d[B]}{dt} \quad \therefore \frac{d[B]}{dt} = -\frac{2}{3} \frac{d[A]}{dt}$$

3. (a) Concentration of reactants decreases while concentration of product increases.

4. (d)
$$R = K(A)^n (B)^m$$

$$R' = K(2A)^{n} \left(\frac{B}{2}\right)^{m} = K(A)^{n} 2^{n} (B)^{m} 2^{-m}$$
$$= K(A)^{n} (B)^{m} 2^{n-m} , \frac{R'}{R} = \frac{K(A)^{n} (B)^{m} 2^{n-m}}{K(A)^{n} (B)^{m}} = 2^{n-m}$$

- 5. (c) $\ln K = \ln -\frac{E_a}{RT}$ is Arrhenius equation. Thus plots of $\ln K$ vs 1/T will give slope $= -E_a/RT$ or $-E_a/2.303 R$.
- **6.** (c) For a reaction E_a for forward reaction $= E_a$ for backward reaction $+ \Delta H$.
- 7. (c) As $K' > K'', E'_a < E''_a$ (Greater the rate constant, less is the activation energy).

(b)
$$T_2 = T(say), T = 25^{\circ} C = 298 K,$$

 $E_a = 104.4 \text{ kJ } mol^{-1} = 104.4 \times 10^3 \text{ J } mol^{-1}$
 $K_1 = 3 \times 10^{-4}, K_2 = ?,$
 $\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$
 $\log \frac{K_2}{3 \times 10^{-4}} = \frac{104.4 \times 10^3 \text{ J } mol^{-1}}{2.303 \times (8.314 \text{ J } k^{-1} \text{ mol}^{-1})}$
 $\left[\frac{1}{298 K} - \frac{1}{T} \right] \text{ As } T \to \infty, \frac{1}{T} \to 0$
 $\therefore \log \frac{K_2}{3 \times 10^{-4}} = \frac{104.4 \times 10^3 \text{ J } mol^{-1}}{2.303 \times 8.314 \times 298}$
 $\log \frac{K_2}{3 \times 10^{-4}} = 18.297, \frac{K_2}{3 \times 10^{-4}} = 1.98 \times 10^{18} \text{ or }$
 $K_2 = (1.98 \times 10^{18}) \times (3 \times 10^{-4}) = 6 \times 10^{14} s^{-1}$

9. (a) Because reaction is exothermic

10. (d)
$$k = Ae^{-E^0 / RT} \log_y K = \log_c A - \frac{E^0}{mx} RT \therefore \log K Vs \frac{1}{T}$$
.

11. (c)
$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

12. (a)
$$2.303 \log \frac{K_2}{K_1} = \frac{E_a}{R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

 $\log \frac{K_2}{K_1} = \frac{9.0 \times 10^3}{2.303 \times 2} \left[\frac{308 - 298}{308 \times 298} \right]$
 $\frac{K_2}{K_1} = 1.63$; $K_2 = 1.63 K_1$; $\frac{1.63 K_1 - K_1}{K_1} \times 100 = 63.0\%$

- 13. (d) Combustion is an exothermic process, which is expected to be favoured by low temperature, but it is not true. Combustion include burning of particles which takes place at higher temperature. Thus at high temperature due to combustion of more particles, the reaction proceeds at higher rate.
- 14. (d) Rate constant $= 2.3 \times 10^{-2} \text{ sec}^{-1}$ It means it is a first order reaction (because unit of rate constant is sec⁻¹)

For first order reaction
$$K = \frac{1}{t} \ln \frac{a}{a-x}$$

$$Kt = \ln \frac{a}{a-x} = \ln \frac{[N_2 O_5]_0}{[N_2 O_5]_t}$$

- **15.** (b) Because as reaction progresses the amount of acetic acid increases.
- 16. (d) Half life of a reaction is found to be inversely proportional to the cube of initial concentration. The order of reaction is 4.

17. (a) It is similar to
$$y = mx + c$$

18. (b) $r = K[A]^n$, $100 r = K[10A]^n$

Thus
$$\frac{1}{100} = \left(\frac{1}{10}\right)^n$$
 or $n = 2$

19. (c) Concentration will fall from 0.1 *M* concentration to 0.025 *M* concentration within 2 half lives. $2 \times T_{1/2} = 40 \text{ min}$

$$\therefore T_{1/2} = 20 min$$

2

%

Rate of reaction = K · c =
$$\frac{0.693}{T_{1/2}}$$
.c
= $\frac{0.693}{20} \times 10^{-2} M / \text{min} = 3.47 \times 10^{-4} M / \text{min}^{-1}$.

20. (e) % distribution of
$$B = \frac{K_1}{K_1 + K_2} \times 100$$

$$=\frac{1.26\times10^{-4}}{1.26\times10^{-4}+3.8\times10^{-4}}\times100$$

B% = 76.83%

6 Distribution of
$$C = \frac{K_2}{K_1 + K_2} \times 100$$

$$=\frac{5.8\times10}{1.26\times10^{-4}+3.8\times10^{-4}}\times100$$

C% = 23.17%

Assertion and Reason

- (b) Instantaneous rate of a reaction is equal to small change in concentration (*dx*) during a small interval of time (*dt*) at that particular instent of time divided by the time interval.
- 2. (b) Molecularity of a reaction can be defined only for an elementary reaction because complex reaction does not take place in one single step and it is almost impossible for all the total molecules of the reactants to be in a state of encounter simultaneously.
- 3. (b) For a zero order reaction, $t_{1/2} = [A_0]/2K$.
- **4.** (b) According to Arrhenius equation, $K = Ae^{-E_a/RT}$ when $E_a = 0, K = A$.
- 6. (a) For a first order reaction, $K = \frac{2.303}{t} \log \frac{a}{a-x}$, where *a* is initial concentration, and *x* is the amount reacted in time *t*. For half-life x = a/2, $t = t_{1/2}$ $K = \frac{2.303}{K} \log \frac{a}{a-a/2}$ $t_{1/2} = \frac{2.303}{K} \log 2 = \frac{0.693}{K}$.

7. (d)
$$H_2 + Cl_2 \rightarrow 2HCl$$
 has much higher quantum efficiency than $H_2 + Br_2 \rightarrow 2HBr$. The first step of secondary process of $H_2 + Cl_2 \rightarrow 2HCl$ is exothermic while the same for $H_2 + Br_2 \rightarrow 2HBr$ is endothermic.

- 8. (e) Vision is a fast photochemical reaction in which the compound, retinal, present in the eye undergoes isomerisation by absorbing a photon of light.
- **9.** (a) The emission of cold light during a chemical reaction is called chemiluminescence.
- 10. (d) The rate reaction is never negative. Minus sign used in expressing the rate only shows that the concentration of the reactant is decreasing.
- **11.** (a) Rate expression $\frac{dx}{dt} = K[A]^m [B]^n$

Shows that the total order of reactions is m+n+O = m+n as the rate of reaction is independent of concentration of *C*, *i.e.* the order with respect to *C* is zero. This is the reason that *C* does not figure in the rate expression.