

Chapter 5



UNIT VIII: Biotechnology

Plant Tissue Culture



Learning Objectives

The learner will be able to

- ❖ Perceive the concepts of tissue culture.
- ❖ Cognize the steps of tissue culture techniques and its types.
- ❖ Understand the protoplast culture in detail.
- ❖ Elicit the list of secondary metabolites obtained through cell suspension culture.
- ❖ Learn plant regeneration pathway.
- ❖ Appreciate the uses of micro propagation, somatic hybridization, shoot meristem culture and germplasm conservation.
- ❖ Acquire the knowledge of patenting Biosafety and Bioethics.



Chapter outline

- 5.1 Basic concepts in tissue culture
- 5.2 Plant tissue culture techniques and types
- 5.3 Plant regeneration pathway
- 5.4 Applications of plant tissue culture
- 5.5 Conservation of plant genetic resources
- 5.6 Intellectual Property Right (IPR), Biosafety and Bioethics
- 5.7 Future of Biotechnology



Growing plant protoplasts, cells, tissues or organs away from their natural or normal environment, under artificial condition, is known as Tissue Culture. It is also known as *in vitro* (*In vitro* is a Latin word, it means that - in glass or in test-tube) growth of plant protoplasts, cells, tissues and organs. A single explant can be multiplied into several thousand plants in a short duration and space under controlled conditions.



Gottlieb
Haberlandt

Tissue culture techniques are often used for commercial production of plants as well as for plant research. Plant tissue culture serves as an indispensable tool for regeneration of transgenic plants. Apart from this some of the main applications of Plant tissue culture are clonal propagation of elite varieties, conservation of endangered plants, production of virus-free plants, germplasm preservation, industrial production of secondary metabolites. etc., In this chapter let us discuss the history , techniques, types , applications of plant tissue culture and get awareness on ethical issues.

Gottlieb Haberlandt (1902) the German Botanist proposed the concept **Totipotency** and he was also the first person to culture plant cells in artificial conditions using the mesophyll cells of *Lamium purpureum* in culture medium and obtained cell proliferation. He is regarded as the father of tissue culture.

5.1 Basic concepts of Tissue Culture

Basic concepts of plant tissue culture are totipotency, differentiation, dedifferentiation and redifferentiation.

Totipotency

The property of live plant cells that they have the genetic potential when cultured in nutrient medium to give rise to a complete individual plant.

Differentiation

The process of biochemical and structural changes by which cells become specialized in form and function.




Figure 5.1: Totipotency

Redifferentiation

The further differentiation of already differentiated cell into another type of cell. For example, when the component cells of callus have the ability to form a whole plant in a nutrient medium, the phenomenon is called redifferentiation.

Dedifferentiation

The phenomenon of the reversion of mature cells to the meristematic state leading to the formation of callus is called dedifferentiation. These two phenomena of redifferentiation and dedifferentiation are the inherent capacities of living plant cells or tissue. This is described as totipotency.

5.2 Plant Tissue Culture (PTC)

Plant tissue culture is used to describe the *in vitro* and aseptic growth of any plant part on a tissue culture medium. This technology is based on three fundamental principles:

- The plant part or explant must be selected and isolated from the rest of plant body.
- The explant must be maintained in controlled physically (environmental) and chemically defined (nutrient medium) conditions.

Explant: The tissue taken from a selected plant transferred to a culture medium often to establish a new plant.

5.2.1 Laboratory Facilities for PTC

For PTC, the laboratory must have the following facilities:



Figure 5.2: Tissue culture lab

- Washing facility for glassware and ovens for drying glassware.
- Medium preparation room with autoclave, electronic balance and pH meter.
- Transfer area sterile room with laminar air-flow bench and a positive pressure ventilation unit called High Efficiency Particulate Air (HEPA) filter to maintain aseptic condition.
- Culture facility: Growing the explant inoculated into culture tubes at 22-28° C with illumination of light 2400 lux, with a photoperiod of 8-16 hours and a relative humidity of about 60%.

5.2.2 Technique Involved in PTC

1. Sterilization:

Sterilization is the technique employed to get rid of microbes such as bacteria and fungi in the culture medium, vessels and explants.

i. Maintenance of Aseptic Environment:

During in vitro tissue culture maintenance of aseptic environmental condition should be followed, i.e., sterilization of glassware, forceps, scalpels, and all accessories in wet steam sterilization by autoclaving at 15 psi (121°C) for 15 to 30 minutes or dipping in 70% ethanol followed by flaming and cooling.

ii. **Sterilization of culture room:** Floor and walls are washed first with detergent and then with 2% sodium hypochlorite or 95% ethanol. The cabinet of laminar airflow is sterilized by clearing the work surface with 95% ethanol and then exposure of UV radiation for 15 minutes.

iii. **Sterilization of Nutrient Media:** Culture media are dispensed in glass containers, plugged with non-absorbent cotton or sealed with plastic closures and then sterilized using autoclave at 15 psi (121°C) for 15 to 30 minutes. The plant extracts, vitamins, amino acids and hormones are sterilized by passing through Millipore filter with 0.2 mm pore diameter and then added to sterilized culture medium inside Laminar Airflow Chamber under sterile condition.

iv. **Sterilization of Explants:** The plant materials to be used for tissue culture should be surface sterilized by first exposing the material in running tap water and then treating it in surface sterilization agents like 0.1% mercuric chloride, 70% ethanol under aseptic condition inside the Laminar Air Flow Chamber.

2. Media Preparation

The success of tissue culture lies in the composition of the growth medium, plant growth regulators and culture conditions such as temperature, pH, light and humidity. No single medium is capable of maintaining optimum growth of all plant tissues. Suitable nutrient medium as per the principle of tissue culture is prepared and used.

MS nutrient medium (Murashige and Skoog 1962) is commonly used. It has carbon sources, with suitable vitamins and hormones. The media formulations available for plant tissue culture other than MS are B5 medium (Gamborg et al. 1968), White medium (White 1943), Nitsch's medium (Nitsch & Nitsch 1969). A medium may be solid or semisolid or liquid. For solidification, a gelling agent such as agar is added.

Agar: A complex mucilaginous polysaccharide obtained from marine algae (sea weeds) used as solidifying agent in media preparation.




Figure 5.3: Basic steps in Plant tissue culture technology



3. Culture condition

pH

The pH of medium is normally adjusted between 5.6 to 6.0 for the best result.

Temperature

The cultures should be incubated normally at constant temperature of $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for optimal growth.

Humidity and Light Intensity

The cultures require 50-60% relative humidity and 16 hours of photoperiod by the illumination of cool white fluorescent tubes of approximately 1000 lux.

Aeration

Aeration to the culture can be provided by shaking the flasks or tubes of liquid culture on automatic shaker or aeration of the medium by passing with filter-sterilized air.

4. Induction of Callus

Explant of 1-2 cm sterile segment selected from leaf, stem, tuber or root is inoculated (transferring the explants to sterile glass tube containing nutrient medium)



Figure 5.4:

Induction of callus

in the MS nutrient medium supplemented with auxins and incubated at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in an alternate light and dark period of 12 hours to induce cell division and soon the upper surface of explant develops into callus. Callus is a mass of unorganized growth of plant cells or tissues in *in vitro* culture medium.

5. Embryogenesis

The callus cells undergoes differentiation and produces somatic embryos, known as **Embryoids**. The embryoids are sub-cultured to produce plantlets.



Figure 5.5:

Embryogenesis

6. Hardening

The plantlets developed *in vitro* require a hardening period and so are transferred to greenhouse or hardening chamber and then to normal environmental conditions.

Hardening is the gradual exposure of *in vitro* developed plantlets in humid chambers in diffused light for acclimatization so as to enable them to grow under normal field conditions.

5.2.3 Types of Plant tissue cultures

Based on the type of explants other plant tissue culture types are

1. Organ culture
2. Meristem culture
3. Protoplast culture
4. Cell suspension culture.

1. Organ culture




Figure 5.6: Organ Culture

The culture of embryos, anthers, ovaries, roots, shoots or other organs of plants on culture media.

2. Meristem Culture:

The culture of any plant meristematic tissue on culture media.






Figure 5.7: Meristem Culture

3. Protoplast Culture:

Protoplasts are cells without a cell wall, but bound by a cell membrane or plasma membrane. Using protoplasts, it is possible to regenerate

whole plants from single cells and also develop somatic hybrids. The steps involved in protoplast culture.

i. **Isolation of protoplast:** Small bits of plant tissue like leaf tissue are used for isolation of protoplast. The leaf tissue is immersed in 0.5% Macrozyme and 2% Onozuka cellulase enzymes dissolved in 13% sorbitol or mannitol at pH 5.4. It is then incubated over-night at 25°C. After a gentle teasing of cells, protoplasts are obtained, and these are then transferred to 20% sucrose solution to retain their viability. They are then centrifuged to get pure protoplasts as different from debris of cell walls.

ii. **Fusion of protoplast:** It is done through the use of a suitable fusogen. This is normally PEG (Polyethylene Glycol). The isolated protoplast are incubated in 25 to 30% concentration of PEG with Ca^{++} ions and the protoplast shows agglutination (the formation of clumps of cells) and fusion.

iii. **Culture of protoplast:** MS liquid medium is used with some modification in droplet, plating or micro-drop array techniques. Protoplast viability is tested with fluorescein diacetate before the culture. The cultures are incubated in continuous light 1000-2000 lux at 25°C. The cell wall formation occurs within 24-48 hours and the first division of new cells occurs between 2-7 days of culture.

iv. **Selection of somatic hybrid cells:** The fusion product of protoplasts without nucleus of different cells is called a cybrid. Following this nuclear fusion take place. This process is called somatic hybridization.

4. Cell Suspension Culture

The growing of cells including the culture of single cells or small




Figure 5.8: Protoplast Culture



aggregates of cells *in vitro* in liquid medium is known as cell suspension culture. The cell suspension is prepared by transferring a portion of callus to the liquid medium and agitated using rotary shaker instrument. The cells are separated from the callus tissue and used for cell suspension culture.

Production of Secondary Metabolites

Cell suspension culture can be useful for the production of secondary metabolites like alkaloids, flavonoids, terpenoids, phenolic compounds and recombinant proteins. Secondary metabolites are chemical compounds that are not required by the plant for normal growth and development but are produced in the plant as 'byproducts' of cell metabolism. For Example: Biosynthesis and isolation of indole alkaloids from *Catharanthus roseus* plant cell culture.

The process of production of secondary metabolites can be scaled up and automated using bio-reactors for commercial production. Many strategies such as biotransformation, elicitation and immobilization have been used to make cell suspension cultures more efficient in the production of secondary metabolites. Few examples of industrially important plant secondary metabolites are listed below in the table:

Secondary metabolites	Plant source	Uses
Digoxin	<i>Digitalis purpurea</i>	Cardiac tonic
Codeine	<i>Papaver somniferum</i>	Analgesic
Capsaicin	<i>Capsicum annum</i>	Rheumatic pain treatment
Vincristine	<i>Catharanthus roseus</i>	Anti-carcinogenic
Quinine	<i>Cinchona officinalis</i>	Antimalarial

Table 5.1: Secondary metabolites and its plant resources

5.3 Plant Regeneration Pathway

From the explants, plants can be regenerated by somatic embryogenesis or organogenesis.




Figure 5.9: Flow chart of Plant regeneration pathway




Figure 5.10: Plant Regeneration Pathway



5.3.1 Somatic Embryogenesis

Somatic embryogenesis is the formation of embryos from the callus tissue directly and these embryos are called **Embryoids** or from the *in vitro* cells directly form pre-embryonic cells which differentiate into embryoids.


Applications

- Somatic embryogenesis provides potential plantlets which after hardening period can establish into plants.
- Somatic embryoids can be used for the production of synthetic seeds.
- Somatic embryogenesis is now reported in many plants such as *Allium sativum*, *Hordeum vulgare*, *Oryza sativa*, *Zea mays* and this possible in any plant.

Synthetic seeds are produced by encapsulation of embryoids in agarose gel or calcium alginate.

5.3.2 Organogenesis

The morphological changes occur in the callus leading to the formation of shoot and roots is called organogenesis.



- Organogenesis can be induced *in vitro* by introducing plant growth regulators in the MS medium.
- Auxin and cytokinins induce shoot and root formation.

5.4 Applications of Plant Tissue Culture

Plant tissue culture techniques have several applications such as:

- i. Improved hybrids production through somatic hybridization.
- ii. Somatic embryoids can be encapsulated into synthetic seeds (synseeds). These encapsulated seeds or synthetic seeds help

in conservation of plant biodiversity.

- iii. Production of disease resistant plants through meristem and shoot tip culture.
- iv. Production of stress resistant plants like herbicide tolerant, heat tolerant plants.
- v. Micropropagation technique to obtain large numbers of plantlets of both crop and tree species useful in forestry within a short span of time and all through the year.
- vi. Production of secondary metabolites from cell culture utilized in pharmaceutical, cosmetic and food industries.

Somaclonal variations: Somatic variations found in plants regenerated *in vitro* (i.e. variations found in leaf, stem, root, tuber or propagule)

Gametoclonal variations: Gametophytic variations found in plants regenerated *in vitro* gametic origin (i.e. variations found in gametes and gametophytes)

5.4.1 Micropropagation of Banana

Micropropagation of plants at industrial level maintains high standards of homogeneity in plants like pineapple, banana, strawberry and potato.



Figure 5.11: Micropropagation of Banana

5.4.2 Artificial Seed

Artificial seeds or synthetic seeds (synseeds) are produced by using embryoids (somatic embryos) obtained through *in vitro* culture. They may even be derived from single cells from any part of the plant that later divide to form cell mass containing dense cytoplasm, large nucleus, starch grains, proteins, and oils etc., To prepare the artificial seeds different

inert materials are used for coating the somatic embryoids like agrose and sodium alginate.




Figure 5.12: Artificial seeds

Advantages of Artificial seeds

Artificial seeds have many advantages over the true seeds

- Millions of artificial seeds can be produced at any time at low cost.
- They provide an easy method to produce genetically engineered plants with desirable traits.
- It is easy to test the genotype of plants.
- They can potentially stored for long time under cryopreservation method.
- Artificial seeds produce identical plants
- The period of dormancy of artificial seeds is greatly reduced, hence growth is faster with a shortened life cycle.

5.4.3 Virus-free plants




Figure 5.13: Shoot tip - Apical Meristem

The field grown plants like perennial crops, usually are infected by variety of pathogens like fungi, bacteria, mycoplasma, viruses which cause considerable economic losses. Chemical methods can be used to control fungal and bacterial pathogens, but not viruses generally.

Shoot meristem tip culture is the method

to produce virus-free plants, because the shoot meristem tip is always free from viruses.

5.5 Conservation of plant genetic resources

5.5.1 Germplasm Conservation

Germplasm conservation refers to the conservation of living genetic resources like pollen, seeds or tissue of plant material maintained for the purpose of selective plant breeding, preservation in live condition and used for many research works.

Germplasm conservation resources is a part of collection of seeds and pollen that are stored in seed or pollen banks, so as to maintain their viability and fertility for any later use such as hybridization and crop improvement.

Germplasm conservation may also involve a gene bank, DNA bank of elite breeding lines of plant resources for the maintenance of biological diversity and also for food security.



Figure 5.14: Seed bank

5.5.2 Cryopreservation (–195°C)

Cryopreservation, also known as Cryo-conservation, is a process by which protoplasts, cells, tissues, organelles, organs, extracellular matrix, enzymes or any other biological materials are subjected to preservation by cooling to very low temperature of –196°C using liquid nitrogen. At this extreme low temperature any enzymatic or chemical activity of the biological material will be totally stopped and this leads to preservation of material in dormant status. Later these materials can be activated by bringing to room temperature slowly for any experimental work.



Figure 5.15: Cryopreservation

Protective agents like dimethyl sulphoxide, glycerol or sucrose are added before cryopreservation process. These protective agents are called cryoprotectants, since they protect the cells, or tissues from the stress of freezing temperature.

5.6 Intellectual Property Right (IPR)

Intellectual property right (IPR) is a category of rights that includes intangible creation of the human intellect, and primarily consists of copyrights, patents, and trademarks. It also includes other types of rights, such as trade secrets, publicity rights, moral rights, and rights against unfair competition.



Figure 5.16: IPR in India

- In biotechnology, the transformed microorganisms and plants and technologies for the production of commercial products are exclusively the property of the discoverer.
- The discoverer has the full rights on his property. It should not be neglected by the others without legal permission.
- The right of discoverer must be protected and it does by certain laws framed by a country.
- The IPR is protected by different ways like patents, copyrights, trade secrets and trademarks, designs and geographical indications.

5.6.1 Patents

- It is a special right to the discoverer/inventor that has been granted by the government

through legislation for trading new articles.

- A patent is a personal property which can be licensed or sold by the person or organisation just like any other property.
- Patent terms give the inventor the rights to exclude others from making, using or selling his invention.

5.6.2 Biosafety and Bioethics

Advances in biotechnology and their applications deals with genetic manipulation.

Biosafety

Biosafety is the prevention of large-scale loss of biological integrity, focusing both on ecology and human health. These prevention mechanisms include conduction of regular reviews of the biosafety in laboratory settings, as well as strict guidelines to follow. Many laboratories handling biohazards employ an ongoing risk management assessment and enforcement process for biosafety. Failures to follow such protocols can lead to increased risk of exposure to biohazards or pathogens.

Bioethics - Ethical, Legal and Social Implications (ELSI)

Bioethics refers to the study of ethical issues emerging from advances in biology and medicine. It is also a moral discernment as it relates to medical policy and practice. Bioethicists are concerned with the ethical questions that arise in the relationships among life sciences, biotechnology and medicine. It includes the study of values relating to primary care and other branches of medicine.

The scope of bioethics is directly related to biotechnology, including cloning, gene therapy, life extension, human genetic engineering, astroethics life in space, and manipulation of basic biology through altered DNA, RNA and proteins. These developments in biotechnology will affect future evolution, and may require new principles, such as biotic ethics, that values life and its basic biological characters and structures. The Ethical, Legal, and Social Implications



(ELSI) program was founded in 1990 as an integral part of the Human Genome Project. The mission of the ELSI program was to identify and address issues raised by genomic research that would affect individuals, families, and society. A percentage of the Human Genome Project budget at the National Institutes of Health and the U.S. Department of Energy was devoted to ELSI research.

Genetic Engineering Appraisal Committee (GEAC)

GEAC is an apex body under Ministry of Environment, Forests and Climate change for regulating manufacturing, use, import, export and storage of hazardous microbes or genetically modified organisms (GMOs) and cells in the country. It was established as an apex body to accord approval of activities involving large scale use of hazardous microorganisms and recombinants in research and industrial production. The GEAC is also responsible for approval of proposals relating to release of genetically engineered organisms and products into the environment including experimental field trials.

5.7 Future of Biotechnology

Biotechnology has become a comprehensive scientific venture from the point of academic and commercial angles, within a short time with the sequencing of human genome and genome of some important organisms. The future developments in biotechnology will be exciting. Thus the development in biotechnology will lead to a new scientific revolution that would change the lives and future of people. Like industrial and computer revolution, biotechnological revolution will also promise major changes in many aspects of modern life.

Summary

Tissue culture is the in vitro aseptic culture of cells, tissues or organs into whole plants under controlled nutritional and environmental conditions. A German physiologist Gottlieb Haberlandt in 1902 for the first time attempted

to culture plant cells in artificial medium, hence he was regarded as father of Tissue culture. Tissue culture mainly based on the concepts totipotency, differentiation, redifferentiation and dedifferentiation. Plant tissue culture technique involves selection of explants, sterilization, media preparation, maintaining culture condition, callus formation, embryogenesis or organogenesis and hardening. Based on the explants chosen the types of tissue culture are organ culture, meristem culture, protoplast culture and cell suspension culture. From the explants, plants can be regenerated by somatic embryogenesis or organogenesis is said to be plant regeneration pathway. Some of the main applications of tissue culture are production of somatic hybrids, artificial seeds, disease resistant and stress resistant plants, germplasm conservation, micropropagation and production of secondary metabolites. Intellectual Property Right (IPR) is primarily aimed at patents, copyrights, trade secret and trademark given to the discoverer / inventor for the commercial production of transformed micro organisms or plants. Biosafety is the prevention mechanism to protect harmful incidents due to biohazards or pathogens. Bioethics dealt with ethical issue emerging from biotechnological advancement. ELSI program addresses issues related to genenomic research. GEAC (Genetic Engineering Appraisal Committee) is a regulatory authority for release of genetically modified products or organisms into the environment.

Evaluation

Choose the correct answer from the given option:

1. Totipotency refers to
 - a) capacity to generate genetically identical plants.
 - b) capacity to generate a whole plant from any plant cell / explant.





- c) capacity to generate hybrid protoplasts.
d) recovery of healthy plants from diseased plants.
2. Micro propagation involves
- vegetative multiplication of plants by using micro-organisms.
 - vegetative multiplication of plants by using small explants.
 - vegetative multiplication of plants by using microspores.
 - Non-vegetative multiplication of plants by using microspores and megaspores.


3. Match the following :

Column A	Column B
1) Totipotency	A) Reversion of mature cells into meristem
2) Dedifferentiation	B) Biochemical and structural changes of cells
3) Explant	C) Properties of living cells develops into entire plant
4) Differentiation	D) Selected plant tissue transferred to culture medium

- | | | | | |
|----|---|---|---|---|
| | 1 | 2 | 3 | 4 |
| a) | C | A | D | B |
| b) | A | C | B | D |
| c) | B | A | D | C |
| d) | D | B | C | A |
4. The time duration for sterilization process by using autoclave is _____ minutes and the temperature is _____
- 10 to 30 minutes and 125° C
 - 15 to 30 minutes and 121° C
 - 15 to 20 minutes and 125° C
 - 10 to 20 minutes and 121° C
5. Which of the following statement is correct
- Agar is not extracted from marine algae such as seaweeds.
 - Callus undergoes differentiation and produces somatic embryoids.
 - Surface sterilization of explants is done by using mercuric bromide
 - P^H of the culture medium is 5.0 to 6.0

6. Select the incorrect statement from given statement
- A tonic used for cardiac arrest is obtained from *Digitalis purpurea*
 - Medicine used to treat Rheumatic pain is extracted from *Capsicum annum*
 - An anti malarial drug is isolated from *Cinchona officinalis*.
 - Anti-carcinogenic property is not seen in *Catharanthus roseus*.
7. Virus free plants are developed from
- Organ culture
 - Meristem culture
 - Protoplast culture
 - Cell suspension culture
8. The prevention of large scale loss of biological integrity
- Biopatent
 - Bioethics
 - Biosafety
 - Biofuel
9. Cryopreservation means it is a process to preserve plant cells, tissues or organs
- at very low temperature by using ether.
 - at very high temperature by using liquid nitrogen
 - at very low temperature of -196 by using liquid nitrogen
 - at very low temperature by using liquid nitrogen
10. Solidifying agent used in plant tissue culture is
- Nicotinic acid
 - Cobaltous chloride
 - EDTA
 - Agar

11. What is the name of the process given below? Write its 4 types.



12. How will you avoid the growing of microbes in nutrient medium during culture process? What are the techniques used to remove the



microbes?

13. Write the various steps involved in cell suspension culture.
14. What do you mean Embryoids? Write its application.
15. Give the examples for micro propagation performed plants .
16. Explain the basic concepts involved in plant tissue culture.
17. Based on the material used, how will you classify the culture technology? Explain it.
18. Give an account on Cryopreservation.
19. What do you know about Germplasm conservation. Describe it.
20. Write the protocol for artificial seed preparation.

Glossary

Aseptic condition: Preparation of materials free from microbes in *in vitro* cultures.

Cell Culture: Growing of cells *in vitro*, including the culture of single cells or small aggregates of cells in a liquid medium.

Chemically defined medium: A nutritive medium used for culturing cells or tissue; each chemical of this medium is known and defined;

Cybrid: Cytoplasmic hybrid obtained by the fusion of cytoplasm of cells of different parental sources; a term applied to the fusion of cytoplasms of two different protoplasts;

Organogenesis: The process of initiation and development of shoot or root though *in vitro* culture particularly from callus

APPENDIX

Composition of MS (Murashige and Skoog) Medium

Macronutrients:

Ammonium nitrate (NH_4NO_3)	1650.0 mg/l
Potassium nitrate (KNO_3)	1900.0 mg/l
Calcium chloride ($\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$)	440.0 mg/l
Magnesium sulphate ($\text{MgSO}_4 \cdot 6\text{H}_2\text{O}$)	370.0 mg/l
Potassium dihydrogen phosphate (KH_2PO_4)	170.0 mg/l

Micronutrients:

Manganese sulphate ($\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$)	22.3 mg/l
Zinc sulphate ($\text{ZnSO}_4 \cdot 4\text{H}_2\text{O}$)	8.6 mg/l
Boric acid (H_3BO_3)	6.2 mg/l
Potassium iodide (KI)	0.83 mg/l

Minor nutrient:

Sodium molybdate ($\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$)	0.250 mg/l
Cupric sulphate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$)	0.025 mg/l
Cobaltous chloride ($\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$)	0.025 mg/l

Iron stock

Na EDTA	37.25 mg/l
Ferrous Sulphate ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$)	27.85 mg/l

Vitamins

Glycine	2.0 mg/l
Nicotinic acid	0.5 mg/l
Pyridoxin HCl	0.5 mg/l
Thiamine HCl	0.1 mg/l

Growth Hormones

IAA	1.30 mg/l
Kinetin	0.4–10.0 mg/l
Myo-inositol	100.0 mg/l
Sucrose	30.0 g/l

Solidifying Agent

Agar	8.0 g/l
------	---------



Milestones in Plant Tissue Culture

Haberlandt (1902)

cultured plant cells in artificial condition called *in vitro* (inside glass) in culture medium (Knop's salt solution) containing glucose and peptone and developed callus (unorganized growth of cells and tissue) and proposed the concept Totipotency, it means the development of whole plant from isolated cells or tissue in *in vitro* condition.

P.R. White (1934)

developed root cultures, used Knop's solution along with three vitamins like pyridoxine, thiamine and nicotinic acid

F.C. Steward (1948)

used coconut water in plant tissue culture work and obtained cell proliferation from carrot explants (Cellular totipotency).

Morel and Martin (1952, 1955)

developed virus-free *Dahlia* and potato plants using shoot meristem culture.

Murashige and Skoog (1962)

formulated tissue culture medium, a landmark in plant tissue culture and it is the most frequently used medium for all kinds of tissue culture work.

Kanta *et al.* (1962)

produced test-tube fertilization in flowering plants.

Yamada *et al.* (1963)

produced *calli* and free cells in tissue culture of *Tradescantia reflexa*.

Guha and Maheshwari (1964)

developed *in vitro* production of haploid embryos from anthers of *Datura*.

Vasil and Hildbrandt (1965)

achieved differentiation of tobacco plants from single, isolated cells in micro propagation.

Takebe *et al.* (1971)

regenerated tobacco plants from isolated mesophyll protoplasts.

Carlson

and co-workers obtained protoplast fusion between *Nicotiana glauca* and *Nicotiana longsdorffii* and developed first interspecific somatic hybrid in 1971.

Melchers and co-workers in 1978

developed intergenic hybrid between potato and tomato called pomato.

Chilton (1983)

produced transformed tobacco plants from single cell transformation and gene insertion.

Horsh *et al.* (1984)

developed transgenic tobacco by *Agrobacterium* mediated gene transfer.

Knop's solution: Nutrient solution used in growth experiments of plants which contains:

Calcium nitrate 3.0 g

Potassium nitrate 1.0 g

Sucrose 50.0 g (optimal)

Magnesium sulfate 1.0 g

Dibasic Potassium phosphate 1.0 g

Deionized water 1000.0 ml