


Tips & Tricks


✍ **Badecker reaction** : This reaction involves the following chemical change.



✍ **Everitt's salt** : It is $K_2[Fe(CN)_6]$ obtained by reduction of prussian blue.

✍ **Masking** : Masking is the process in which a substance without physical separation of it is so transformed that it does not enter into a particular reaction e.g., masking of Cu^{2+} by CN^- ion.

✍ **Macrocyclic effect** : This term refers to the greater thermodynamic stability of a complex with a cyclic polydentate ligand when compared to the complex formed with a non-cyclic ligand. e.g., $Zn(II)$ complex with ligand;



✍ **Prussian blue** and **Turnbull's blue** is pot. ferric ferrocyanide. However colour of Turnbull's blue is less intense than prussian blue. Decrease in colour is due to the presence in it of a white compound of the formula $K_2\{Fe[Fe(CN)_6]\}$ named as potassium ferrous ferrocyanide.

✍ Crystal field theory explains why certain geometries are more favoured than other by certain metals in terms of crystal field stabilization energies.

✍ The colour of complexes are explained in terms of electronic transitions between the various d orbitals of different energies.

✍ In octahedral complexes, the energy of the d_{z^2} and $d_{x^2-y^2}$ orbitals increases much more than the energy of the d_{xy}, d_{yz}, d_{xz} orbitals. The potential energy increases as a result of repulsive interactions. The greater the repulsion, greater is the increases in energy.

✍ Ligands with larger groups from unstable rings than the ligands with smaller groups. This is ascribable to steric hinderance produced by a larger group.

✍ When Ca^{2+} or Mg^{2+} forms complexes with EDTA, the pH of the solution decreases.

✍ The copper sulphate solution e.g., turns deep

blue when excess of ammonia is added. This is also due to the formation of $[Cu(NH_3)_4]SO_4$

✍ Co-ordination compound with a general formula MA_4 , MA_3B or $MABCD$ do not show any geometric isomerism.

✍ **Flexidentate character** : polydentate ligand are said to have flexidentate character if they do not use all its donar atoms to get coordinated to the metal ion e.g., EDTA generally act as a hexadentate ligand but it can also act as a pentadentate and tetradentate ligand.

✍ Octahedral complex of the type $[M(ABCDEF)]$ have 15 different geometrical isomers with a pair of enantiomers. Although a few geometrical isomer have been prepared but none has been resolved e.g., $[Pt(Py)(NH_3)(NO_2)(Cl)(Br)I]$.

Ordinary Thinking

Objective Questions

Basic Terms

- In $K_4Fe(CN)_6$
 - (CN) are linked with primary valency
 - (CN) are linked with secondary valency
 - K are linked with secondary valency
 - K are linked with non-ionic valency
- The co-ordination number of copper in cuprammonium sulphate is
 - 2
 - 6
 - 4
 - 4
- Which of the following acts as a bidentate ligand in complex formation
 - Acetate
 - Oxalate
 - Thiocyanate
 - EDTA
- The co-ordination number of cobalt in the complex $[Co(en)_2Br_2]Cl_2$ is
 - 2
 - 6
 - 5
 - 4
- Which of the following ligands forms a chelate

[MP PET/PMT 1998]

 - Acetate
 - Oxalate
 - Cyanide
 - Ammonia
- According to Werner's theory

[MP PMT 2000, 02]

 - Primary valency can be ionized
 - Secondary valency can be ionized
 - Primary and secondary valencies both cannot be ionized

908 Co-ordination Chemistry

- (d) Only primary valency cannot be ionized
7. Which of the following is not true for ligand-metal complex [MP PET 1993]
- Larger the ligand, the more stable is the metal-ligand bond
 - Highly charged ligand forms strong bond
 - Larger the permanent dipole moment of ligand, the more stable is the bond
 - Greater the ionization potential of central metal, the stronger is the bond
8. What is the co-ordination number of the metal in $[Co(en)_2Cl_2]^+$
- 4
 - 5
 - 6
 - 3
9. Bidentate ligand is
- CN^-
 - Ethylene diammine
 - SCN^-
 - EDTA
- (en)
10. The coordination number of Pt in $[Pt(NH_3)_4Cl_2]^{++}$ ion is [MP PET 1995]
- 2
 - 4
 - 6
 - 8
11. Which is the example of hexadentate ligand
- 2, 2'-dipyridyl
 - Dimethyl glyoxime
 - Aminodiacetate ion
 - Ethylene diammine tetra acetate ion [EDTA]
12. The coordination number of a metal in coordination compounds is [MP PET 1996; KCET (Engg./Med.) 1999]
- Same as primary valency
 - Sum of primary and secondary valencies
 - Same as secondary valency
 - None of these
13. Ligand in a complex salt are [KCET 1992]
- Anions linked by coordinate bonds to a central metal atom or ion
 - Cations linked by coordinate bonds to a central metal atom or ion
 - Molecules linked by coordinate bonds to a central metal atom or ion
 - Ions or molecules linked by coordinate bonds to a central metal atom or ion
14. A group of atoms can function as a ligand only when [KCET 1989; DCE 1999; MP PMT 2000]
- It is a small molecule
 - It has an unshared electron pair
 - It is a negatively charged ion
 - It is a positively charged ion
15. Which of the following complexes show six coordination number [RPET 2000]
- $[Zn(CN)_4]^{2-}$
 - $[Cr(H_2O)_6]^{3+}$
 - $[Cu(CN)_4]^{2-}$
 - $[Ni(NH_3)_4]^{2+}$
16. The number of ions formed when cuprammonium sulphate is dissolved in water is [KCET 1993]
- 1
 - 2
 - 4
 - Zero
17. The coordination number of Cu in complex $[Cu(H_2O)_4]^{++}$ is
- 4
 - 3
 - 2
 - 1
18. The primary valence of the metal ion in the coordination compound $K_2[Ni(CN)_4]$ is
- Four
 - Zero
 - Two
 - Six
19. The metal which does not form a polynuclear carbonyl is
- Mn
 - Co
 - Cr
 - Fe
20. Which one of the following forms with an excess of CN^- (Cyanide) a complex having coordination number two [AIIMS 2004]
- Cu^+
 - Ag^+
 - Ni^{2+}
 - Fe^{2+}
21. According to Lewis the ligands are [MP PMT 2002]
- Acidic in nature
 - Basic in nature
 - Neither acidic nor basic
 - Some are acidic and others are basic
22. The coordination number of a central metal atom in a complex is determined by
- The number of ligands around a metal ion bonded by sigma and pi-bonds both.
 - The number around a metal ion bonded by pi-bonds
 - The number of ligands around a metal ion bonded by sigma bonds
 - The number of only anionic ligands bonded to the metal ion
23. In the extraction of which of the following, complex ion forms [MP PET 1989]
- Cu
 - Ag
 - Fe
 - Na
24. Potassium ferrocyanide is a [AFMC 2000]
- Normal salt
 - Mixed salt
 - Double salt
 - Complex salt
25. A monodentate ligand has
- One co-ordinate site
 - Two co-ordinate sites
 - Any number of co-ordinate sites
 - No capacity to co-ordinate

26. EDTA has coordination number [AFMC 2004] (a) 3 (b) 4 (c) 5 (d) 6
27. Among the properties (a) reducing (b) oxidising (c) complexing, the set of properties shown by CN^- ion towards metal species is (a) c, a (b) b, c (c) a, b (d) a, b, c
28. That ion or molecule which forms a complex compound with transitional metal ion is called (a) Recipient (b) Ligand (c) Coordinate ion (d) No special name
29. Coordination number of Zn in ZnS (zinc blende) is [Orissa JEE 2004] (a) 6 (b) 4 (c) 8 (d) 12
30. Wilkinson's catalyst used as a homogeneous catalyst in the hydrogenation of alkenes contains (a) Iron (b) Aluminium (c) Rhodium (d) Cobalt
31. Given the molecular formula of the hexa coordinated complexes (A) $CoCl_3 \cdot 6NH_3$ (B) $CoCl_3 \cdot 5NH_3$ (C) $CoCl_3 \cdot 4NH_3$. If the number of coordinated NH_3 molecules in A, B and C respectively are 6, 5 and 4, the primary valency in (A), (B) and (C) are: [DCE 2003] (a) 6, 5, 4 (b) 3, 2, 1 (c) 0, 1, 2 (d) 3, 3, 3
32. Generally, a group of atoms can function as a ligand if [MP PET 1996] (a) They are positively charged ions (b) They are free radicals (c) They are either neutral molecules or negatively charged ions (d) None of these
33. The ligand in potassium ferricyanide is (a) K^+ (b) CN^- (c) Fe^{3+} (d) $(CN)_6$
34. Co-ordination number of aluminum is [MHCET 2004] (a) 8 (b) 6 (c) 12 (d) 4
35. In $K_4Fe(CN)_6$, Fe is in the form of (a) An atom (b) An ion (c) Cationic complex (d) Anionic complex
36. Which of the following ligands is expected to bidentate [CBSE PMT 1994] (a) Br (b) $C_2O_4^{2-}$ (c) CH_3NH_2 (d) $CH_3C \equiv N$
37. In the compound lithium tetrahydroaluminate, the ligand is (a) H^+ (b) H^- (c) H (d) None of these
38. Which of the following is the odd one out [MP PET 1996] (a) Potassium ferrocyanide (b) Ferrous ammonium sulphate (c) Potassium ferricyanide (d) Tetrammine copper (II) sulphate
39. The basic ligand is (a) NH_3 (b) CN^- (c) F^- (d) All
40. The negative ligand is (a) Aqua (b) Sulphato (c) Carboxyl (d) Nitro sodium
41. Which has yellow colour (a) Potassium cobaltinitrite (b) Potassium hexanitro cobaltate (III) (c) Fischer's salt (d) All the above
42. Ligands, in complex compounds [MP PMT 2003] (a) Accept e^- -pair (b) Donate e^- -pair (c) Neither accept e^- -pair nor donate (d) All of these happen
43. Which of the following is a common donor atom in ligands [BHU 2001] (a) Arsenic (b) Nitrogen (c) Oxygen (d) Both (b) and (c)
44. Trunbull's blue is a compound [KCET 1993] (a) Ferricyanide (b) Ferrous ferricyanide (c) Ferrous cyanide (d) Ferriferrocyanide
45. Tollen's reagent is [KCET 1990] (a) $[Ag(NH_3)_2]^+$ (b) Ag_2O (c) $[Cu(OH)_4]^{2-}$ (d) Cu_2O
46. Finely divided iron combines with CO to give [UPSEAT 2002] (a) $Fe(CO)_5$ (b) $Fe_2(CO)_9$ (c) $Fe_2(CO)_{12}$ (d) $Fe(CO)_6$
47. In a complex, the highest possible coordination number is (a) 6 (b) 12 (c) 4 (d) 8
48. The number of neutral molecules or negative groups attached to the central metal atom in a complex ion is called (a) Atomic number (b) Effective atomic number (c) Coordination number (d) Primary valency
49. EDTA combines with cations to form

910 Co-ordination Chemistry

- (a) Ion-exchange resins (b) Chelates
(c) Clathrates (d) Polymers
50. An example of a double salt is [MP PET 2001]
(a) Bleaching powder (b) Hypo
(c) $K_4[Fe(CN)_6]$ (d) Potash alum
51. In complex compounds, metal ligand bond is
(a) Coordinate bond (b) Hydrogen bond
(c) Ionic bond (d) Covalent bond
52. Ammonia forms the complex ion $[Cu(NH_3)_4]^{2+}$ with copper ions in alkaline solutions but not in acidic solution. What is the reason for it [AIEEE 2003]
(a) In acidic solutions hydration protects copper ions
(b) In acidic solutions protons coordinate with ammonia molecules forming NH_4^+ ions and NH_3 molecules are not available
(c) In alkaline solutions insoluble $Cu(OH)_2$ is precipitated which is soluble in excess of any alkali
(d) Copper hydroxide is an amphoteric substance
53. Zeigler–Natta catalyst is used for which type of reaction
(a) Hydrogenation (b) Polymerization
(c) Oxidation (d) Reduction
54. Which of the following is not considered as an organometallic compound.
(a) Cis-platina (b) Ferrocene
(c) Zeise's salt (d) Grignard reagent
55. Which one is organometallic compound [MP PMT 2004]
(a) Lithium methoxide (b) Lithium dimethyl amide
(c) Lithium acetate (d) Methyl lithium
56. An aqueous solution of potash alum gives [UPSEAT 2004]
(a) Two types of ions (b) Only one type of ion
(c) Four types of ions (d) Three types of ions
57. Carnallite in solution in H_2O , shows the properties of [DCE 2003]
(a) K^+, Mg^{2+}, Cl^- (b) $K^+, Cl^-, SO_4^{2-}, Br^-$
(c) K^+, Mg^{2+}, CO_3^{2-} (d) K^+, Mg^{2+}, Cl^-, Br^-
58. What is the co-ordination number of cobalt in $Co(NH_3)_3Cl_3$ [MP PET 1994]
(a) 3 (b) 4
(c) 5 (d) 6
59. The formula of alum is [Pb. CET 2002]
(a) $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$
(b) $K_4[Fe(CN)_6]$
(c) $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 6H_2O$
(d) $Na_2CO_3 \cdot 10H_2O$
60. Number of ions present in $K_4[Fe(CN)_6]$ [Pb. CET 2000]
(a) 2 (b) 10
(c) 3 (d) 5
61. CH_3MgI is an organometallic compound due to [DCE 2002]
(a) $Mg-I$ bond (b) $C-I$ bond
(c) $C-Mg$ bond (d) $C-H$ bond
62. What is the EAN of nickel in $Ni(CO)_4$ [BVP 2003]
(a) 34 (b) 35
(c) 32 (d) 36

Nomenclature, Oxidation state and EAN number

1. How many ions are produced in aqueous solution of $[Co(H_2O)_6]Cl_2$ [RPMT 2002]
(a) 2 (b) 3
(c) 4 (d) 6
2. IUPAC name of $[Pt(NH_3)_3(Br)(NO_2)Cl]Cl$ is [CBSE PMT 1998]
(a) Triamminechlorobromonitroplatinum (IV) chloride
(b) Triamminebromonitrochloroplatinum (IV) chloride
(c) Triamminebromochloronitroplatinum (IV) chloride
(d) Triamminenitrochlorobromoplatinum (IV) chloride
3. Oxidation state of nitrogen is incorrectly given for [UPSEAT 2000, 01]
- | Compound | Oxidation state |
|--------------------------|-----------------|
| (a) $[Co(NH_3)_5Cl]Cl_2$ | 0 |
| (b) NH_2OH | -1 |
| (c) $(N_2H_5)_2SO_4$ | +2 |
| (d) Mg_3N_2 | -3 |
4. The formula of dichloro bis (urea) copper (II) is [CBSE PMT 1997]
(a) $[Cu\{O=C(NH_2)_2\}Cl_2]$
(b) $[CuCl_2\{O=C(NH_2)_2\}_2]$
(c) $[Cu\{O=C(NH_2)_2\}Cl]Cl$
(d) $[CuCl_2]\{O=C(NH_2)_2H_2\}$
5. The IUPAC name of the complex $[Pt(NH_3)_2Cl_2]$ is
(a) Platinum (II) diammino dichloride
(b) Diammino dichloro platinate (IV)
(c) Bis (ammino) dichloro platinum (IV)
(d) Dichloro diammine platinum (II)
6. Correct formula of diammine silver (I) chloride is

- (a) $Ag(NH_3)Cl$ (b) $Ag(NH_2)Cl$
 (c) $[Ag(NH_3)_2]Cl$ (d) $[Ag(NH_2)_2]Cl$
7. The formula of sodium nitroprusside is [AIIMS 1992]
 (a) $Na_4[Fe(CN)_5NOS]$ (b) $Na_2[Fe(CN)_5NO]$
 (c) $NaFe[Fe(CN)_6]$ (d) $Na_2[Fe(CN)_6NO_2]$
8. The correct name of $[Pt(NH_3)_4Cl_2][PtCl_4]$ is [MP PET 2003]
 (a) Tetraammine dichloro platinum (iv) tetrachloro platinate (ii)
 (b) Dichloro tetra ammine platinum (iv) tetrachloro platinate (ii)
 (c) Tetrachloro platinum (ii) tetraammine platinate (iv)
 (d) Tetrachloro platinum (ii) dichloro tetraammine platinate (iv)
9. Correct formula of potassium ferrocyanide is [CBSE PMT 1988]
 (a) $K_4[Fe(CN)_6]$ (b) $K_2[Fe(CN)_6] \cdot H_2O$
 (c) $K_3[Fe(CN)_6]$ (d) None of these
10. The IUPAC name of $[Co(NH_3)_3(NO_2)_3]$ is
 (a) Trinitrotriammine cobalt (III)
 (b) Trinitrotriammine cobalt (II)
 (c) Trinitrotriammine cobalt (III) ion
 (d) Trinitritriammine cobaltate (III)
11. In $K_4[Fe(CN)_6]$, the E.A.N. of Fe is [DCE 2000]
 (a) 33 (b) 35
 (c) 36 (d) 26
12. Which of the following pairs is not correctly matched [MP PET 1993]
 (a) Effective atomic number of Pt in $[PtCl_6]^{2-} = 84$
 (b) Absorption peak for $[Cr^{III}(NH_3)_6]^{+3} = 21680 \text{ cm}^{-1}$
 (c) Crystal field stabilization energy of d^2 in weak ligand field $= (-0.8\Delta_0)$
 (d) Example of weak ligand field for d^5 configuration $= [Mn^{II}F_6]^{-4}$
13. The oxidation number of chromium in sodium tetrafluoro oxochromate complex is
 (a) II (b) IV
 (c) VI (d) III
14. The IUPAC name of $K_4[Fe(CN)_6]$ is [CBSE PMT 1990; MP PET 1992; MP PMT 1995, 97; Kurukshetra CET 2002]
 (a) Potassium hexacyanoferrate (II)
 (b) Potassium ferrocyanide
 (c) Tetrapotassium hexacyanoferrate (II)
 (d) Tetrapotassium ferrous hexacyanide (II)
15. The IUPAC name of $[Ni(CO)_4]$ is [RPET 1999]
 (a) Tetra carbonyl nickel (II)
 (b) Tetra carbonyl nickel (0)
 (c) Tetra carbonyl nickelate (II)
 (d) Tetra carbonyl nickelate (0)
16. The correct nomenclature for $Fe_4[Fe(CN)_6]_3$ is [MP PMT 1994]
 (a) Ferroso-ferric cyanide
 (b) Ferric-ferrous hexacyanate
 (c) Iron (III) hexacyanoferrate (II)
 (d) Hexacyanoferrate (III-II)
17. The IUPAC name of compound $Na_3[Co(ONO)_6]$ will be [MP PMT 2000]
 (a) Hexanitritocobalt (III) sodium
 (b) Sodium cobalt nitrite
 (c) Sodium hexanitrocobaltate (III)
 (d) Sodium hexanitritocobaltate (III)
18. In which of the following complexes oxidation state of metal is zero [MP PET 1997]
 (a) $[Pt(NH_3)_2Cl_2]$ (b) $[Cr(CO)_6]$
 (c) $[Cr(NH_3)_3Cl_3]$ (d) $[Cr(en)_2Cl_2]$
19. The oxidation number of Cr in $[Cr(NH_3)_6]Cl_3$ is [CBSE PMT 2001]
 (a) 8 (b) 6
 (c) 4 (d) 3
20. In $[Ni(NH_3)_4]SO_4$, the E.A.N. of Ni is
 (a) 34 (b) 35
 (c) 36 (d) 37
21. IUPAC name of $[Co(ONO)(NH_3)_5Cl_2]$ is [AMU 2002]
 (a) Pentaammine nitro cobalt (III) chloride
 (b) Pentaammine nitrito cobalt (III) chloride
 (c) Pentaammine nitroso cobalt (III) chloride
 (d) Pentaammine oxo-nitro cobalt (III) chloride
22. The oxidation number of Pt in $[Pt(C_2H_4)Cl_3]^-$ is [MNR 1993]
 (a) +1 (b) +2
 (c) +3 (d) +4
23. The oxidation state of cobalt in the complex compound $[Co(NH_3)_6]Cl_3$ is
 (a) +3 (b) +6
 (c) +5 (d) +2
24. The correct IUPAC name of potassium cuprochloride is
 (a) Potassium copper (I) tetrachloride
 (b) Potassium tetrachlorocuprate (I)
 (c) Tetrachloropotassium cuprate (I)

912 Co-ordination Chemistry

- (d) Tetrachlorocopper (I) potassiate
25. The effective atomic number of cobalt in the complex $[Co(NH_3)_6]^{3+}$ is [MP PET 2003]
 (a) 36 (b) 33
 (c) 24 (d) 30
26. IUPAC name of $K_3Fe(CN)_6$ is [MP PMT 1993; MP PET 1997]
 (a) Potassium ferrocyanide (II)
 (b) Potassium hexaferrocyanate (III)
 (c) Potassium ferrohexacyanate (II)
 (d) Potassium hexacyanoferrate (III)
27. The EAN of iron in potassium ferricyanide is [Pb. CET 2000]
 (a) 18 (b) 54
 (c) 35 (d) 23
28. In the coordination compound, $K_4[Ni(CN)_4]$ oxidation state of nickel is
 (a) -1 (b) 0
 (c) +1 (d) +2
29. According to IUPAC nomenclature sodium nitroprusside is named is
 (a) Sodium pentacyanonitrosyl ferrate (III)
 (b) Sodium nitroferrocyanide
 (c) Sodium nitroferrocyanide
 (d) Sodium pentacyanonitrosyl ferrate (II)
30. Pick out the complex compound in which the central metal atom obeys EAN rule strictly [KCET 2003]
 (a) $K_4[Fe(CN)_6]$ (b) $K_3[Fe(CN)_6]$
 (c) $[Cr(H_2O)_6]Cl_3$ (d) $[Cu(NH_3)_4]SO_4$
31. Which of the following is wrong statement [BHU 2003]
 (a) $Ni(CO)_4$ has oxidation number + 4 for Ni
 (b) $Ni(CO)_4$ has zero oxidation number for Ni
 (c) Ni is metal
 (d) CO is gas
32. Oxidation state of Fe in $K_3[Fe(CN)_6]$ [RPMT 2002]
 (a) 2 (b) 3
 (c) 0 (d) None of these
33. Which complexes have zero oxidation state
 (a) Carbonyl (b) Ferrocyanide
 (c) Amine (d) Cyanide
34. The proper name for $K_2[PtCl_6]$ is [MH CET 2002]
 (a) Potassium platinum hexachloride
 (b) Potassium hexachloro platinum IV
 (c) Potassium hexachloro platinate IV
 (d) Potassium hexachloro platinum
35. IUPAC name of $K_3[Al(C_2O_4)_3]$ is called [MP PMT 1993, 02, 03]
 (a) Potassium alumino oxalato
 (b) Potassium aluminium (III) trioxalate
 (c) Potassium trioxalato aluminate (III)
 (d) Potassium trioxalato aluminate (IV)
36. The I.U.P.A.C. name of $K_3[Ir(C_2O_4)_3]$ is [MP PMT 2001]
 (a) Potassium tri oxalato iridium (III)
 (b) Potassium tri oxalato iridate (III)
 (c) Potassium tris (oxalato) iridium (III)
 (d) Potassium tris (oxalato) iridate (III)
37. The charge on $[Ag(CN)_2]^-$ complex is [AIIMS 2001]
 (a) -1 (b) +1
 (c) +2 (d) +3
38. The IUPAC name of $[Co(NH_3)_6]Cl_3$ is [IIT-JEE 1994]
 (a) Hexammine cobalt (III) chloride
 (b) Hexammine cobalt (II) chloride
 (c) Triammine cobalt (III) trichloride
 (d) None of these
39. IUPAC name of $[Co(NH_3)_3(H_2O)_2Cl]Cl_2$ is [AIEEE 2003] [MP PET 1994]
 (a) Diaquachlorodiammine cobalt (III) chloride
 (b) Triamminediaquachloro cobalt (III) chloride
 (c) Chlorodiamminediaqua cobalt (III) chloride
 (d) Diamminediaquachloro cobalt (II) chloride
40. Dichloro diammine platinum (II) complex has the formula [MP PMT 1997]
 (a) $Pt[Cl_2(NH_3)_2]$ (b) $Pt[R.(NH_3)_2]Cl_2$
 (c) $[PtCl_2(NH_3)_2]$ (d) $[Pt.R.(NH_3)_2]Cl_2$
41. The formula of potassiumdicyano bis (oxalato) nickelate (II) is
 (a) $K_4[Ni(CN)_2(Ox)_2]$ (b) $K_3[Ni_2[Ni_2(CN)_2(Ox)_2]]$
 (c) $K[Ni(CN)(Ox)_2]$ (d) $K_2[Ni(CN)_2(Ox)_2]$
42. The value of x which appears in the complex $[Ni(CN)_4]^x$ is
 (a) + 2 (b) - 2
 (c) 0 (d) 4
43. Pick the correct name of $[Co(NH_3)_5Cl]Cl_2$ [AMU 2001]
 (a) Chloropentammine cobalt (III)
 (b) Pentammine cobalt (III) chloride
 (c) Chloropentammine cobalt (III) chloride
 (d) Chloropentammine cobalt (II) chloride
44. The valency of cuprammonium ion is
 (a) + 4 (b) + 2
 (c) - 2 (d) - 4
45. In which of the following compounds transition metal has zero oxidation state
 (a) CrO_5 (b) $NH_2.NH_2$
 (c) $NOClO_4$ (d) $[Fe(CO)_5]$
46. The complex chlorocompound diaquatriammine cobalt (III) chloride is represented as [CBSE PMT 2002]
 (a) $[Co(NH_3)_3(H_2O)_3]Cl_2$

- (b) $[Co(NH_2)_3(H_2O)_2]Cl_2$
 (c) $[CoCl(NH_3)_3(H_2O)_2]Cl_3$
 (d) $[CoCl(NH_3)_3(H_2O)_2]Cl_2$
47. The complex compound $[Co(NH_3)_3NO_2ClCN]$ is named as
 (a) Chlorocyanonitrotriammine cobalt (III)
 (b) Nitrochlorocyanotriammine cobalt (III)
 (c) Cyanonitrochlorotriammine cobalt (III)
 (d) Triamminenitrochlorocyano cobalt (III)
48. The oxidation number of Pt in $[Pt(C_2H_4)Cl_3]^-$ is [UPSEAT 1999, 01]
 (a) +1 (b) +2
 (c) +3 (d) +4
49. What is the structural formula of lithium tetrahydridoaluminate [MP PMT 2003]
 (a) $Al[LiH_4]$ (b) $Al_2[LiH_4]_3$
 (c) $Li[AlH_4]$ (d) $Li[AlH_4]_2$
50. IUPAC name for $K[Ag(CN)_2]$ is
 (a) Potassium argentocyanide
 (b) Potassium silver cyanide
 (c) Potassium dicyanoargentate (I)
 (d) Potassium dicyanosilver (II)
51. The oxidation state of Co in $[Co(H_2O)_5Cl]^{2+}$ is
 (a) +2 (b) +3
 (c) +1 (d) +4
52. The chemical formula of diammine silver (I) chloride is [BHU 2004]
 (a) $[Ag(NH_3)]Cl$ (b) $[Ag(NH_3)_2]Cl$
 (c) $[Ag(NH_3)_2]Cl$ (d) $[Ag(NH_4)_2]Cl$
53. IUPAC name of $[Co(NH_3)_5NO_2]Cl_2$ [Pb. CET 2000]
 (a) Pentamminenitrocobalt (III) chloride
 (b) Pentamminenitrosocobalt (III) chloride
 (c) Pentamminenitrocobalt (II) chloride
 (d) None of these
54. The pair of the compounds in which both the metals are in the highest possible oxidation state is [IIT-JEE (Screening) 2004]
 (a) $[Fe(CN)_6]^{3-}$, $[Co(CN)_6]^{3-}$
 (b) CrO_2Cl_2 , MnO_4^-
 (c) TiO_3 , MnO_2
 (d) $[Co(CN)_6]^{3-}$, MnO_3
55. The IUPAC name of $[Cr(NH_3)_6]^{3+}$ is [Pb. CET 2001]
 (a) Hexamminechromium (VI) ion
 (b) Hexamminechromium (III) ion
 (c) Hexamminechromium (II) ion
 (d) Hexamminechloride
56. The IUPAC name of $K_2[Cr(CN)_2O_2(O)_2(NH_3)]$ is [DCE 2003]
 (a) Potassium ammine dicyano dioxoperoxochromate (VI)
 (b) Potassium ammine cyano peroxo dioxochromate (VI) [MP PMT 1996]
 (c) Potassium ammine cyano peroxo dioxochromium (VI)
 (d) Potassium ammine cyano peroxo dioxochromatic (IV)
57. The IUPAC name of the coordination compound $K_3[Fe(CN)_6]$ is [AIEEE 2005]
 (a) Potassium hexacyanoferrate (II)
 (b) Potassium hexacyanoferrate (III)
 (c) Potassium hexacyanoiron (II)
 (d) Tripotassium hexacyanoiron (II)
58. Which compound is zero valent metal complex [KCET 2000]
 (a) $[Cu(NH_3)_4]SO_4$ (b) $[Pt(NH_3)_2Cl_2]$
 (c) $[Ni(CO)_4]$ (d) $K_3[Fe(CN)_6]$

Isomerism and magnetic properties

1. Which one of the following octahedral complexes will not show geometric isomerism (A and B are monodentate ligands) [CBSE PMT 2003]
 (a) $[MA_5B]$ (b) $[MA_2B_4]$
 (c) $[MA_3B_3]$ (d) $[MA_4B_2]$
2. The number of unpaired electrons in the complex ion $[CoF_6]^{3-}$ is (Atomic no. of Co = 27) [CBSE PMT 2003]
 (a) Zero (b) 2
 (c) 3 (d) 4
3. Which would exhibit co-ordination isomerism
 (a) $[Cr(NH_3)_6][Co(CN)_6]$ (b) $[Co(en)_2Cl_2]$
 (c) $[Cr(NH_3)_6]Cl_3$ (d) $[Cr(en)_2Cl_2]^+$
4. $[Co(NH_3)_5NO_2]Cl_2$ and $[Co(NH_3)_5(ONO)]Cl_2$ are related to each other as
 (a) Geometrical isomers (b) Optical isomers
 (c) Linkage isomers (d) Coordination isomers
5. $[Co(NH_3)_5Br]SO_4$ and $[Co(NH_3)_5SO_4]Br$ are examples of which type of isomerism [MP PMT 1993, 94, 95; MP PET 1997; MP PET/PMT 1998]
 (a) Linkage (b) Geometrical
 (c) Ionization (d) Optical
6. $[Co(NH_3)_4Cl_2]NO_2$ and $[Co(NH_3)_4Cl.NO_2]Cl$ are isomers [MP PMT 1993; MP PET 1995, 2001]
 (a) Geometrical (b) Optical
 (c) Linkage (d) Ionization
7. Which would exhibit ionisation isomerism [MP PET 1997]
 (a) $[Cr(NH_3)_6]Cl_3$ (b) $[Co(NH_3)_5Br]SO_4$

914 Co-ordination Chemistry

- (c) $[Cr(en)_2Cl_2]$ (d) $[Cr(en)_3Cl_3]$ (a) $[Cu(NH_3)_4]Cl_2$ (b) $[Ag(NH_3)_2]Cl$
8. $[Ti(H_2O)_6]^{+3}$ is paramagnetic in nature due to [RPMT 2002] (c) NO (d) NO_2
- (a) One unpaired e^- (b) Two unpaired e^- 17. The number of geometrical isomers for $[Pt(NH_3)_2Cl_2]$ is [CBSE PMT 1995]
- (c) Three unpaired e^- (d) No unpaired e^- (a) Two (b) One
9. Coordination isomerism is caused by the interchange of ligands between the [UPSEAT 2002] (c) Three (d) Four
- (a) *Cis* and *Trans* structure (b) Complex cation and complex anion
- (c) Inner sphere and outer sphere 18. The pair of complex compounds $[Cr(H_2O)_6Cl_3]$ and $[Cr(H_2O)_5Cl]Cl_2 \cdot H_2O$ are an example of [MP PMT 1997]
- (d) Low oxidation and higher oxidation states (a) Linkage isomerism (b) Ionisation isomerism
10. Which one of the following will not show geometrical isomerism [MP PMT 2002] (c) Coordination isomerism (d) Hydrate isomerism
- (a) $[Cr(NH_3)_4Cl_2]Cl$ (b) $[Co(en)_2Cl_2]Cl$ 19. The number of geometrical isomers of the complex $[Co(NO_2)_2(NH_3)_2]$ is [CBSE PMT 1997]
- (c) $[Co(NH_3)_5NO_2]Cl_2$ (d) $[Pt(NH_3)_2Cl_2]$ (a) 2 (b) 3
11. Paramagnetic co-ordination compounds contain electrons (c) 4 (d) 0
- (a) No 20. The type of isomerism present in nitropentamine chromium (III) chloride is
- (b) Both paired and unpaired (a) Optical (b) Linkage
- (c) Paired (c) Ionization (d) Polymerisation
- (d) Unpaired 21. Which of the following compounds exhibits linkage isomerism [MP PMT 2001]
12. Which of the following isomeric pairs shows ionization isomerism [MP PET 1993]
- (a) $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$ (a) $[Co(en)_3]Cl_3$ (b) $[Co(NH_3)_6][Cr(CN)_6]$
- (b) $[Cr(H_2O)_6]Cl_3$ and $[Cr(H_2O)_5Cl]Cl_2 \cdot H_2O$ (c) $[Co(en)_2NO_2Cl]Br$ (d) $[Co(NH_3)_5Cl]Br_2$
- (c) $[Pt(NH_3)_2Cl_2]$ and $[Pt(NH_3)_4][PtCl_4]$ 22. Pick out from the following complex compounds, a poor electrolytic conductor in solution [MP PMT 1994]
- (d) $[Co(NH_3)_5Br]SO_4$ and $[Co(NH_3)_5SO_4]Br$ (a) $K_2[PtCl_6]$ (b) $[Co(NH_3)_3(NO_2)_3]$
13. Among the following ions which one has the highest paramagnetism [IIT 1993; UPSEAT 2002] (c) $K_4[Fe(CN)_6]$ (d) $[Cu(NH_3)_4]SO_4$
- (a) $[Cr(H_2O)_6]^{3+}$ (b) $[Fe(H_2O)_6]^{2+}$ 23. The possible number of optical isomers in $[Co(en)_2Cl_2]^+$ are
- (c) $[Cu(H_2O)_6]^{2+}$ (d) $[Zn(H_2O)_6]^{2+}$ (a) 2 (b) 3
14. Amongst $Ni(CO)_4$, $[Ni(CN)_4]^{2-}$ and $[NiCl_4]^{2-}$ [IIT 1991] (c) 4 (d) 6
- (a) $Ni(CO)_4$ and $[NiCl_4]^{2-}$ are diamagnetic and $[Ni(CN)_4]^{2-}$ is paramagnetic 24. Magnetic moment of $[Cu(NH_3)_4]^{2+}$ ion is [RPET 2003]
- (b) $[NiCl_4]^{2-}$ and $[Ni(CN)_4]^{2-}$ are diamagnetic and $Ni(CO)_4$ is paramagnetic (a) 1.414 (b) 1.73
- (c) $Ni(CO)_4$ and $[Ni(CN)_4]^{2-}$ are diamagnetic and $[NiCl_4]^{2-}$ is paramagnetic (c) 2.23 (d) 2.38
- (d) $Ni(CO)_4$ is diamagnetic and $[NiCl_4]^{2-}$ and $[Ni(CN)_4]^{2-}$ are paramagnetic 25. What is true for $[Fe(CN)_6]^{3-}$ and $[FeF_6]^{3-}$ [RPET 1999]
15. $[Co(NH_3)_4Cl_2]^+$ exhibits (a) Both are paramagnetic
- (a) Geometrical isomerism (b) Optical isomerism (b) Only $[Fe(CN)_6]^{3-}$ is paramagnetic
- (c) Bonding isomerism (d) Ionisation isomerism (c) Only $[FeF_6]^{3-}$ is paramagnetic
16. The compound which does not show paramagnetism is (d) Both are diamagnetic
- (a) $[Ni(CO)_4]$ (b) $[Co(NH_3)_6]^{3+}$ 26. Which of the following is paramagnetic [AFMC 1997]
- (c) $[Ni(CN)_4]^{2-}$ (d) $[NiCl_4]^{2-}$ (a) $[Ni(CO)_4]$ (b) $[Co(NH_3)_6]^{3+}$
27. The total number of possible isomers for the complex compound $[Cu^{II}(NH_3)_4][Pt^{II}Cl_4]$ are [CBSE PMT 1998; DPMT 2004; J & K CET 2005]
- (a) 3 (b) 4
- (c) 5 (d) 6

[IIT 1992]

28. Which one of the following shows maximum paramagnetic character [AIIMS 1998]
 (a) $[Cr(H_2O)_6]^{3+}$ (b) $[Fe(CN)_6]^{4-}$
 (c) $[Fe(CN)_6]^{3-}$ (d) $[Cu(H_2O)_6]^{2+}$
29. The complexes $[Co(NH_3)_6][Cr(C_2O_4)_3]$ and $[Cr(NH_3)_6][Co(C_2O_4)_3]$ [AMU 2002]
 (a) Linkage isomerism (b) Geometrical isomerism
 (c) Coordination isomerism (d) Ionisation isomerism
30. Which of the following exhibits highest molar conductivity [MP PET 1994]
 (a) $[Co(NH_3)_6]Cl_3$ (b) $[Co(NH_3)_5Cl]Cl_2$
 (c) $[Co(NH_3)_4Cl_2]Cl$ (d) $[Co(NH_3)_3Cl_3]$
31. Which of the following compounds is colourless [MP PET 1994]
 (a) $Cu_2(CH_3COO)_4 \cdot 2H_2O$ (b) Cu_2Cl_2
 (c) $CuSO_4 \cdot 5H_2O$ (d) $[Cu(NH_3)_4]SO_4 \cdot 4H_2O$
32. The type of magnetism exhibited by $[Mn(H_2O)_6]^{2+}$ ion is [IIT 1994]
 (a) Paramagnetism (b) Diamagnetism
 (c) Both (a) and (b) (d) None of these
33. The number of isomers possible for square planar complex $K_2[PdClBr_2(SCN)]$ is [MP PET 1994]
 (a) 2 (b) 3
 (c) 4 (d) 6
34. The number of unpaired electrons in $Ni(CO)_4$ is [AIIMS 1997]
 (a) Zero (b) One
 (c) Three (d) Five
35. If the magnetic moment of $[Ag(CN)_2]^{-1}$ is zero, then the number of unpaired electrons will be [MP PET 1995]
 (a) 1 (b) 2
 (c) 3 (d) Zero
36. $[Pt(NH_3)_4Cl_2]Br_2$ and $[Pt(NH_3)_4Br_2]Cl_2$ are related to each other as [MP PET 1996; AFMC 2000; CBSE PMT 2001]
 (a) Optical isomers (b) Coordinate isomers
 (c) Ionization isomers (d) Linkage isomers
37. Which of the following complex will show geometrical as well as optical isomerism (*en* = ethylene diamine) [KCET 1996]
 (a) $Pt(NH_3)_2Cl_2$ (b) $[Pt(NH_3)Cl_4]$
 (c) $[Pt(en)_3]^{4+}$ (d) $[Pt(en)_2]Cl_2$
38. Which of the following complexes is diamagnetic [RPMT 1997]
 (a) $Ni(CO)_4$ (b) $NiCl_4^{2-}$
 (c) $Ni(Br)_4^{2-}$ (d) $NiCl_2 \cdot 4H_2O$
39. Which one has the highest paramagnetism [AMU 2001]
 (a) $Ni(CO)_4$ (b) $[Ni(NH_3)_4]Cl_2$
 (c) $[Ni(NH_3)_6]Cl_2$ (d) $[Cu(NH_3)_4]Cl_2$
40. Which of the following coordination compounds would exhibit optical isomerism
 (a) *trans*-dicyanobis (ethylenediamine) chromium (III) chloride
 (b) *tris*-(ethylenediamine) cobalt (III) bromide
 (c) pentaamminenitrocobalt (III) iodide
 (d) diamminedichloroplatinum (II)
41. Which of the following does not have optical isomer [AIIMS 2004]
 (a) $[Co(NH_3)_3Cl_3]$ (b) $[Co(en)_3]Cl_3$
 (c) $[Co(en)_2Cl_2]Cl$ (d) $[Co(en)(NH_3)_2Cl_2]Cl$
42. Change in composition of co-ordination sphere yields which types of isomers
 (a) Optical (b) Geometrical
 (c) Ionisation (d) None of these
43. Types of isomerism shown by $[Cr(NH_3)_5NO_2]Cl_2$ is
 (a) Optical (b) Ionisation
 (c) Geometrical (d) Linkage
44. Which of the following will not give a precipitate with $AgNO_3$ [MP PET 2003]
 (a) $[Co(NH_3)_3Cl_3]$ (b) $[Co(NH_3)_4Cl_2]Cl$
 (c) $[Co(NH_3)_5Cl]Cl_2$ (d) $[Co(NH_3)_6]Cl_3$
45. How many ions are produced from $[Co(NH_3)_6]Cl_3$ in solution [RPET 1999]
 (a) 6 (b) 4
 (c) 3 (d) 2
46. The colour of $CoCl_3 \cdot 5NH_3 \cdot H_2O$ is [Kerala (Med.) 2002]
 (a) Orange yellow (b) Orange
 (c) Green (d) Violet
 (e) Pink
47. Which one of the following is expected to be a paramagnetic complex [MP PMT 1991, 2000]
 (a) $[Ni(H_2O)_6]^{2+}$ (b) $[Ni(CO)_4]$
 (c) $[Zn(NH_3)_4]^{2+}$ (d) $[Co(NH_3)_6]^{+3}$
48. Which one of the following will give a white precipitate with $AgNO_3$ in aqueous medium [MP PMT 1994]
 (a) $[Co(NH_3)_5Cl](NO_2)_2$ (b) $[Pt(NH_3)_2Cl_2]$
 (c) $[Pt(en)Cl_2]$ (d) $[Pt(NH_3)_4]Cl_2$
49. How many ions will be produced in solution from one molecule of chloropentammine cobalt (III) chloride
 (a) 1 (b) 2
 (c) 3 (d) 4

916 Co-ordination Chemistry

50. Which of the following complex will give white precipitate with $BaCl_2$ (aq.)
 (a) $[Co(NH_3)_4SO_4]NO_2$ (b) $[Cr(NH_3)_5SO_4]Cl$
 (c) $[Cr(NH_3)_5Cl]SO_4$ (d) Both (b) and (c)
51. The number of precipitable halide ions in the sample $[Pt(NH_3)Cl_2Br]Cl$ will be
 (a) 2 (b) 3
 (c) 4 (d) 1
52. The colour of tetrammine copper (II) sulphate is
 (a) Blue (b) Red
 (c) Violet (d) Green
53. The number of ions per mole of a complex $[CoCl_2.5NH_3]Cl_2$ in aqueous solution will be
 [MP PMT 2001]
 (a) Nine (b) Four
 (c) Three (d) Two
54. How many unpaired electrons are present in the central metal ion of $[CoCl_4]^{2-}$ [Orissa JEE 2005]
 (a) 3 (b) 4
 (c) 5 (d) 2
55. What is the magnetic moment of $K_3[FeF_6]$
 [Orissa JEE 2005]
 (a) 5.91 BM (b) 4.89 BM
 (c) 3.87 BM (d) 6.92 BM
56. (i) $K_4[Fe(CN)_6]$ (ii) $K_3[Cr(CN)_6]$ (iii) $K_3[Fe(CN)_6]$
 (iv) $K_2[Ni(CN)_4]$
 Choose the complex which is paramagnetic
 [Kerala CET 2005]
 (a) (i), (ii) and (iii) (b) (i), (iii) and (iv)
 (c) (i), (iii) and (iv) (d) (i), (ii) and (iv)
 (e) (ii) and (iv)
57. In coordination compound $[Co(en)_2Cl_2]Cl$ which is false
 [Kerala CET 2005]
 (a) Show geometrical Isomerism
 (b) Show optical Isomerism
 (c) Show ionic Isomerism
 (d) A octahedral complex
 (e) A cationic complex
- (b) sp^3, sp^3, sp^3d and sp^2
 [JIPMER 1997]
 (c) dsp^2, sp^2, sp^3 and dsp^3
 (d) dsp^2, sp^3, sp^2 and dsp^3
3. One mole of the complex compound $Co(NH_3)_5Cl_3$, gives 3 moles of ions on dissolution in water. One mole of the same complex reacts with two moles of $AgNO_3$ solution to yield two moles of $AgCl(s)$. The structure of the complex is
 [AIEEE 2003]
 (a) $[Co(NH_3)_5Cl]Cl_2$
 (b) $[Co(NH_3)_3Cl_3].2NH_3$
 (c) $[Co(NH_3)_4Cl_2]Cl.NH_3$
 (d) $[Co(NH_3)_4Cl]Cl_2.NH_3$
4. Cuprammonium ion $[Cu(NH_3)_4]^{2+}$ is
 [MP PMT 1997; KCET 2002]
 (a) Tetrahedral (b) Square planar
 (c) Triangular bipyramid (d) Octahedral
5. In the complex $[SbF_5]^{2-}, sp^3d$ hybridisation is present. Geometry of the complex is [Pb. PMT 2000]
 (a) Square pyramidal (b) Square bipyramidal
 (c) Tetrahedral (d) Square
6. The type of hybridization involved in the metal ion of $[Ni(H_2O)_6]^{2+}$ complex is
 (a) d^3sp^2 (b) sp^3d^2
 (c) sp^3 (d) dsp^2
7. In the formation of $K_4Fe(CN)_6$, the hybridisation involved is
 (a) sp^2 (b) d^2sp^3
 (c) d^3sp^2 (d) d^4p
8. Which of the following is not true for metal carbonyls
 [MP PET 1993]
 (a) The oxidation state of the metal in the carbonyls is zero
 (b) The secondary carbonyls are obtained from photo-decomposition
 (c) Metal carbonyls are single bonded species
 (d) $d\pi - p\pi$ overlap is observed in metal carbonyls
9. An octahedral complex is formed, when hybrid orbitals of the following type are involved [DCE 2003]
 (a) sp^3 (b) dsp^2
 (c) sp^3d^2 (d) sp^2d
10. Which one is an example of octahedral complex
 [MP PET 2000]
 (a) FeF_6^{3-} (b) $Zn(NH_3)_4^{2+}$
 (c) $Ni(CN)_4^{2-}$ (d) $Cu(NH_3)_4^{2+}$
11. Which of the following complexes has a square planar geometry

Hybridisation and Geometry

1. The correct structural formula of zeise's salt is
 (a) $K^+[PtCl_3 - \eta^2 - (C_2H_4)]^-$
 (b) $K_2[PtCl_3 - \eta^2 - C_2H_4]$
 (c) $K^+[PtCl_2 - \eta^2 - (C_2H_4)]Cl^-$
 (d) $K^+[PtCl_3(C_2H_4)]^-$
2. The correct order of hybridisations of central atom in $NH_3, [PtCl_4]^{2-}, PCl_5$ and BCl_3 is [MP PMT 2003]
 (a) dsp^2, dsp^3, sp^2 and sp^3

- (a) $Ag(NH_3)_2^+$ (b) $Cu(en)_2^{2+}$
(c) $[MnCl_4]^{2-}$ (d) $Ni(CO)_4$
12. The shape of $[Fe(CN)_6]^{4-}$ ion is
(a) Hexagonal (b) Pyrimidal
(c) Octahedral (d) Octagonal
13. What is the shape of $Fe(CO)_5$ [CBSE PMT 2000]
(a) Linear (b) Tetrahedral
(c) Square planar (d) Trigonal bipyramidal
14. What type of hybridization is involved in $[Fe(CN)_6]^{3-}$ [AMU 1999]
(a) d^2sp^3 (b) dsp^2
(c) sp^3d^2 (d) dsp^3
15. The example of dsp^2 hybridisation is [MP PET 1999; AIIMS 2001]
(a) $Fe(CN)_6^{3-}$ (b) $Ni(CN)_4^{2-}$
(c) $Zn(NH_3)_4^{2+}$ (d) FeF_6^{3-}
16. The shape of $[Cu(NH_3)_4]^{2+}$ is square planar, Cu^{2+} in this complex is [NCERT 1989; RPET 1999]
(a) sp^3 hybridised (b) dsp^2 hybridised
(c) sp^3d hybridised (d) sp^3d^2 hybridised
17. The geometry of $Ni(CO)_4$ and $Ni(PPh_3)_2Cl_2$ are [IIT-JEE 1999; DCE 2002]
(a) Both square planar
(b) Tetrahedral and square planar respectively
(c) Both tetrahedral
(d) Square planar and tetrahedral respectively
18. Which complex has square planar structure [JIPMER 2002]
(a) $Ni(CO)_4$ (b) $[NiCl_4]^{2-}$
(c) $[Ni(H_2O)_6]^{2+}$ (d) $[Cu(NH_3)_4]^{2+}$
19. $[Pt(NH_3)_4]Cl_2$ is [DCE 2001]
(a) Square planar (b) Tetrahedral
(c) Pyramidal (d) Pentagonal
20. A complex involving dsp^2 hybridization has
(a) A square planar geometry
(b) A tetrahedral geometry
(c) An octahedral geometry
(d) Trigonal planar geometry
21. A tetrahedral complex ion is formed due to hybridization
(a) sp^2 (b) sp^3
(c) dsp^2 (d) d^2sp^3
22. Back bonding is involved in which of the organometallic compounds
(a) $[(CH_3)_3Al]_2$ (b) $Mg^{2+}(C_5H_5^-)_2$
(c) $R-Mg-X$ (d) $[C_5H_5)_2Fe]$
23. d^2sp^3 hybridisation leads to
(a) Hexagonal shape (b) Trigonal bipyrimidal
(c) Octahedral shape (d) Tetrahedral shape
24. Chromium hexacarbonyl is an octahedral compound involving
(a) sp^3d^2 (b) dsp^2
(c) d^2sp^3 (d) d^3sp^2 orbitals
25. $[CoF_6]^{3-}$ is formed by hybridization
(a) d^2sp^3 (b) d^3sp^2
(c) d^2sp^3 (d) sp^3d^2
26. The species havoing tetrahedral shape is [IIT-JEE (Screening) 2004]
(a) $[PdCl_4]^{2-}$ (b) $[Ni(CN)_4]^{2-}$
(c) $[Pd(CN)_4]^{2-}$ (d) $[NiCl_4]^{2-}$
27. Among $[Ni(CO)_4]$, $[Ni(CN)_4]^{2-}$, $[NiCl_4]^{2-}$ species, the hybridization states at the Ni atom are, respectively [CBSE PMT 2004; MP PMT 1992; BHU 1995; AFMC 1997]
(a) sp^3 , sp^3 , dsp^2 (d) dsp^2 , sp^3 , sp^3
(c) sp^3 , dsp^2 , dsp^2 (d) sp^3 , dsp^2 , sp^3
(At. no. of Ni = 28)
28. The bond in $K_4[Fe(CN)_6]$ are: [MP PET 2004]
(a) All ionic
(b) All covalent
(c) Ionic and covalent
(d) Ionic, covalent and co-ordiante covalent
29. Hybridization of Fe in $K_3Fe(CN)_6$ is [DCE 2002]
(a) sp^3 (b) dsp^3
(c) sp^3d^2 (d) d^2sp^3
30. The complex ion which has no 'd' electrons in the central metal atom is [IIT-JEE Screening 2001]
(a) $[MnO_4]^-$ (b) $[Co(NH_3)_6]^{3+}$
(c) $[Fe(CN)_6]^{3-}$ (d) $[Cr(H_2O)_6]^{3+}$
31. Which of the following statement is correct
(a) $[Cu(NH_3)_6]^{2+}$ is a colourless ion
(b) $[Zn(H_2O)_6]^{2+}$ ion is blue coloured
(c) $[Ni(CN)_4]^{2-}$ ion has a tetrahedral shape
(d) Nickel dimethyl glyoxides is red in colour
32. Which of the following shall form an octahedral complex [DCE 2001]
(a) d^4 (low spin) (b) d^8 (high spin)
(c) d^6 (low spin) (d) None of these
33. Which one of the following is a strong field ligand
(a) CN^- (b) NO_2^-
(c) en (d) NH_3
34. The strongest ligand in the following is [MP PET 1995]
(a) CN^- (b) Br^-
(c) HO^- (d) F^-
35. The neutral ligand is

918 Co-ordination Chemistry

- (a) Chloro (b) Hydroxo
(c) Ammine (d) Oxalato
36. The ligands which can get attached to the central metal ion through more than one atom are called
(a) Ambident ligands (b) Polydentate ligands
(c) Chelate ligands (d) Neutral ligands
37. A strong ligand gives a complex which is generally called
(a) High spin (b) High energy
(c) Low spin (d) Stable
38. CN^- is a strong field ligand. This is due to the fact that
[CBSE PMT 2004]
(a) It can accept electron from metal species
(b) It forms high spin complexes with metal species
(c) It carries negative charge.
(d) It is a pseudohalide
39. Considering H_2O as a weak field ligand, the number of unpaired electrons in $[Mn(H_2O)_6]^{2+}$ will be (At. No. of Mn = 25)
[CBSE PMT 2004]
(a) Two (b) Four
(c) Three (d) Five
40. Which of the following is a π - complex
(a) Trimethyl aluminium (b) Ferrocene
(c) Diethyl zinc (d) Nickel carbonyl
41. Which of the following is a π - acid ligand
[KCET 1996; AIIMS 2003]
(a) NH_3 (b) CO
(c) F^- (d) Ethylene diamine
42. The value of the 'spin only' magnetic moment for one of the following configurations is 2.84 BM. the correct one
[AIEEE 2005]
(a) d^4 (in strong ligand field)
(b) d^4 (in weak ligand field)
(c) d^3 (in weak as well as in strong fields)
(d) d^5 (in strong ligand field)
43. The geometry of $Ni(CO)_4$ and $Ni(PPh_3)_2Cl_2$ are
[BHU 2005]
(a) Both square planar
(b) Tetrahedral and square planar respectively
(c) Both tetrahedral
(d) Square planar and tetrahedral respectively
- (a) $[Fe(OH)_3]^{3-}$ (b) $[Fe(Cl)_6]^{3-}$
(c) $[Fe(CN)_6]^{3-}$ (d) $[Fe(H_2O)_6]^{3+}$
3. The most stable complex among the following is
[MP PMT 2002]
(a) $K_3[Al(C_2O_4)_3]$ (b) $[Pt(en)_2]Cl_2$
(c) $Ag(NH_3)_2Cl$ (d) $K_2[Ni(EDTA)]$
4. Which of the following factors tends to increase the stability of metal ion complexes
(a) Higher ionic radius of the metal ion
(b) Higher charge/size ratio of the metal ion
(c) Lower ionisation potential of the metal ion
(d) Lower basicity of the ligand
5. $CuSO_4$ reacts with KCN solution and forms: [DPMT 2004]
(a) $K_3[Cu(CN)_4]$ (b) $Cu(CN)$
(c) $Cu(CN)_2$ (d) $K_4[Cu(CN)_6]$
6. A co-ordination complex compound of cobalt has the molecular formulae containing five ammonia molecules, one nitro group and two chlorine atoms for one cobalt atom. One mole of this compound produces three mole ions in an aqueous solution on reacting with excess of $AgNO_3$, $AgCl$ precipitate. The ionic formula for this complex would be:
[DPMT 2004; Kerala PMT 2004]
(a) $[Co(NH_3)_5(NO_2)]Cl_2$
(b) $[Co(NH_3)_5Cl][Cl(NO_2)]$
(c) $[Co(NH_3)_4(NO_2)Cl][(NH_3)Cl]$
(d) $[Co(NH_3)_5][(NO_2)_2Cl_2]$
7. In any ferric salt, on adding potassium ferrocyanide, a prussian blue colour is obtained, which is
[BIT 1992; BHU 2002]
(a) $K_3Fe(CN)_6$ (b) $KFe[Fe(CN)_6]$
(c) $FeSO_4 \cdot Fe(CN)_6$ (d) $Fe_4[Fe(CN)_6]_3$
8. Prussian blue is formed when [CBSE PMT 1990]
(a) Ferrous sulphate reacts with $FeCl_3$
(b) Ferric sulphate reacts with $K_4[Fe(CN)_6]$
(c) Ferrous ammonium sulphate reacts with $FeCl_3$
(d) Ammonium sulphate reacts with $FeCl_3$
9. Complex salt can be made by the combination of $[Co^{III}(NH_3)_5Cl]^x$ with [RPMT 2000, AFMC 2002]
(a) PO_4^{3-} (b) Cl^-
(c) $2Cl^-$ (d) $2K^+$
10. Which reagent can be used to identify nickel ion
(a) Resorcinol
(b) Dimethyl glyoxime [DMG]
(c) Diphenyl benzidine
(d) Potassium ferrocyanide
11. Dimethyl glyoxime forms a coloured complex with

Complexes and complex stability

1. Which of the following complexes has the highest stability constant at 298 K
(a) $[CdCl_4]^{2-}$ (b) $[CdBr_4]^{2-}$
(c) $[CdI_4]^{2-}$ (d) $[Cd(CN)_4]^{2-}$
2. The most stable ion is [AIEEE 2002]

- (a) Ag (b) Ni
(c) Cr (d) Zn
12. Silver chloride dissolves in excess of NH_4OH . The cation present in this solution is [EAMCET 1998]
(a) Ag^+ (b) $[Ag(NH_3)_2]^+$
(c) $[Ag(NH_3)_4]^+$ (d) $[Ag(NH_3)_6]^+$
13. Silver sulphide dissolved in a solution of sodium cyanide to form the complex [AMU 1999]
(a) $Na[Ag(CN)_2]$ (b) $Na_3[Ag(CN)_4]$
(c) $Na_5[Ag(CN)_6]$ (d) $Na_2[Ag(CN)_2]$
14. Which one will give Fe^{3+} ions in solution
(a) $[Fe(CN)_6]^{3-}$
(b) $Fe_2(SO_4)_3$
(c) $[Fe(CN)_6]^{4-}$
(d) $NH_4(SO_4)_2 \cdot FeSO_4 \cdot 6H_2O$
15. The cation that does not form an amine complex with excess of ammonia is
(a) Cd^{2+} (b) Al^{3+}
(c) Cu^{2+} (d) Ag^+

Application of organometallics

1. Ziegler-Natta catalyst is an organometallic compound of which metal
(a) Iron (b) Zirconium
(c) Rhodium (d) Titanium
2. In the case of small cuts, bleeding is stopped by applying potash alum. Here alum acts as [KCET (Med.) 2001]
(a) Fungicide (b) Disinfectant
(c) Germicide (d) Coagulating agent
3. The formula of ferrocene is
(a) $[Fe(CN)_6]^{4-}$ (b) $[Fe(CN)_6]^{3-}$
(c) $[Fe(CO)_5]$ (d) $[(C_5H_5)_2Fe]$
4. Which of the following is formed when *n*-butyl lithium reacts with tin (II) chloride [AFMC 2001]
(a) $LiBr$ (b) Et_4Pb
(c) $(C_4H_9)_4Sn$ (d) $(C_2H_5)_4Pb$
5. Which of the following organo-silicon compound on hydrolysis will give a three dimensional silicone [Orissa JEE 2003]
(a) R_3SiCl (b) $RSiCl_3$
(c) $SiCl_4$ (d) R_2SiCl_2
6. Which one is not an organometallic compound [J & K CET 2005; Pb. CET 2003]
(a) $RMgX$ (b) C_2H_5ONa
(c) $(CH_3)_4Sn$ (d) KC_4H_9
7. The complex used as an anticancer agent is [AIIMS 2003]
(a) $trans-[Co(NH_3)_3Cl_3]$ (b) $cis-[PtCl_2(NH_3)_2]$
(c) $cis-K_2[PtCl_2Br_2]$ (d) Na_2CO_3
8. The compound that is not olefinic organometallic is
(a) $K[C_2H_4PtCl_3] \cdot 3H_2O$ (b) $Be(CH_2)_2$
(c) $(C_2H_4PtCl_3)_2$ (d) $C_4H_4Fe(CO)_3$
9. Among the following, which is not the π -bonded organometallic compound [CBSE PMT 2003]
(a) $(CH_3)_4Sn$ (b) $K[PtCl_3(\eta^2-C_2H_4)]$
(c) $Fe(\eta^5-C_5H_5)_2$ (d) $Cr(\eta^6-C_6H_6)_2$
10. Wilkinson's catalyst is used in
(a) Polymerization (b) Condensation
(c) Halogenation (d) Hydrogenation
11. What is the use of tetraethyl lead
(a) As a catalyst in addition reaction of alkenes
(b) As a catalyst in polymerization reaction of alkenes
(c) For reducing knocking
(d) For creating knocking
12. Which of the following is an organo-metallic compound [MP PMT 2001]
(a) Lithium ethoxide (b) Ethyl lithium
(c) Lithium acetate (d) Lithium carbide
13. Which of the following is an organometallic compound [J & K CET 2005] [AIIMS 1997]
(a) $Ti(C_2H_5)_4$ (b) $Ti(OC_2H_5)_4$
(c) $Ti(OCOCH_3)_4$ (d) $Ti(OC_6H_5)_4$
14. Which of the following is not an organometallic compound [MP PET 1996; BHU 2002]
(a) Ethyl magnesium bromide
(b) Tetraethyl lead
(c) Sodium ethoxide
(d) Trimethyl aluminium
15. An organometallic compound amongst the following is
(a) Ferrocene (b) CaC_2
(c) Tetraethyl lead (TEL) (d) All of these
16. Which of the following does not have a metal carbon bond [CBSE PMT 2004]
(a) $K[Pt(C_2H_4)Cl_3]$ (b) $Ni(CO)_4$
(c) $Al(OC_2H_5)_3$ (d) C_2H_5MgBr
17. Coordination compounds have great importance in biological systems. In this context which of the following statements is incorrect
(a) Cyanocobalamin is B_{12} and contains cobalt.
(b) Haemoglobin is the red pigment of blood and contains iron.
(c) Chlorophylls are green pigments in plants and contains calcium.

- (d) Carboxypeptidase-A is an enzyme and contains zinc.
18. Ziegler-Natta catalyst is [Pb. CET 2004]
 (a) $(Ph_3P)_3RhCl$ (b) $K[PtCl_3(C_2H_4)]$
 (c) $[Al_2(C_2H_5)_6]+TiCl_4$ (d) $[Fe(C_2H_5)_2]$
19. Dimethyl glyoxime gives a red precipitate with Ni^{2+} , which is used for its detection. To get this precipitate readily the best pH range is [AIIMS 2004]
 (a) < 1 (b) 2-3
 (c) 3-4 (d) 9-11
20. The π -bonded organo metallic compound which has ethene as one of its component is [J & K CET 2005]
 (a) Zeise's salt (b) Ferrocene
 (c) Dibenzene chromium (d) Tetraethyl tin.
7. On hydrolysis $(Me)_2SiCl_2$ will produce [IIT-JEE 2003]
 (a) $(Me)_2Si(OH)_2$
 (b) $(Me)_2Si = O$
 (c) $[-O-(Me)_2Si-O-]_n-$
 (d) $Me_2SiCl(OH)$
8. Which of the following represents a chelating ligand [JIPMER 2002]
 (a) H_2O (b) OH^-
 (c) DMG (d) Cl^-
9. $CuSO_4$ reacts with KCN solution and forms [MP PMT 1992; IIT 1996; UPSEAT 2001, 02]
 (a) $Cu(CN)_2$ (b) $Cu(CN)$
 (c) $K_2[Cu(CN)_4]$ (d) $K_3[Cu(CN)_4]$
10. In the process of extraction of gold,
 Roasted gold ore + $CN^- + H_2O \xrightarrow{O_2} [X] + OH^-$
 $[X] + Zn \rightarrow [Y] + Au$
 Identify the complexes [X] and [Y]
 (a) $X = [Au(CN)_2]^-$, $Y = [Zn(CN)_4]^{2-}$
 (b) $X = [Au(CN)_4]^{3-}$, $Y = [Zn(CN)_4]^{2-}$
 (c) $X = [Au(CN)_2]^-$, $Y = Zn(CN)_6]^{4-}$
 (d) $X = [Au(CN)_4]^-$, $Y = [Zn(CN)_4]^{2-}$ [BHU 1999]
11. A solution of potassium ferrocyanide would contain ions
 (a) 2 (b) 3
 (c) 4 (d) 5
12. Which one of the following complexes is an outer orbital complex [AIEEE 2004]
 (a) $[Co(NH_3)_6]^{3+}$ (b) $[Mn(CN)_6]^{4-}$
 (c) $[Fe(CN)_6]^{4-}$ (d) $[Ni(NH_3)_6]^{2+}$ [IIT JEE 2003]
 Atomic nos : Mn = 25, Fe = 26, Co = 27, Ni = 28
13. Which one of the following has largest number of isomers? [AIEEE 2004]
 (a) $[Ir(PP_3)_2H(CO)]^{2+}$ (b) $[Co(NH_3)_5Cl]^{2+}$
 (c) $[Ru(NH_3)_4Cl_2]^+$ (d) $[Co(en)_2Cl_2]^+$
 (R = alkyl group; en = ethylenediamine)
14. Which kind of isomerism is exhibited by octahedral $Co(NH_3)_4Br_2Cl$?
 (a) Geometrical and Ionization
 (b) Geometrical and Optical
 (c) Optical and Ionization
 (d) Geometrical only
15. Which one of the following is expected to exhibit optical isomerism (en = ethylenediamine)
 (a) cis - $[Pt(NH_3)_2Cl_2]$ (b) trans - $[Co(en)_2Cl_2]$
1. The coordination number and oxidation state of Cr in $K_3[Cr(C_2O_4)_3]$ are, respectively [CBSE PMT 1995]
 (a) 4 and +2 (b) 6 and +3
 (c) 3 and +3 (d) 3 and 0
2. The complex compounds which result from the coordination of carbon monoxide are known as
 (a) Electronic (b) Carbonyls
 (c) Carbonates (d) Carbon permono
3. Mixture X = 0.02 mol of $[Co(NH_3)_5SO_4]Br$ and 0.02 mol of $[Co(NH_3)_5Br]SO_4$ was prepared in 2 litre of solution
 1 litre of mixture X + excess $AgNO_3 \rightarrow Y$.
 1 litre of mixture X + excess $BaCl_2 \rightarrow Z$
 Number of moles of Y and Z are
 (a) 0.01, 0.01 (b) 0.02, 0.01
 (c) 0.01, 0.02 (d) 0.02, 0.02
4. Which of the following organometallic compound is σ and π bonded [MH CET 2001; PCET 2002]
 (a) $Fe(CH_3)_3$
 (b) $[Fe(\eta^5-C_5H_5)_2]$
 (c) $[Co(CO)_5NH_3]^{2+}$
 (d) $K[PtCl_3(\eta^2-C_2H_4)]$
5. In $[Ni(NH_3)_4]SO_4$, the valency and coordination number of Ni will be respectively
 (a) 3 and 6 (b) 2 and 4
 (c) 4 and 2 (d) 4 and 4
6. Co-ordination number of Fe in the complexes $[Fe(CN)_6]^{4-}$, $[Fe(CN)_6]^{3-}$ and $[FeCl_4]^-$ would be respectively [MP PET 2003]
 (a) 2, 3, 3 (b) 6, 6, 4
 (c) 6, 3, 3 (d) 6, 4, 6

Critical Thinking

Objective Questions

(c) *trans* - $[Pt(NH_3)_2Cl_2]$ (d) *cis* - $[Co(en)_2Cl_2]$ 16. $[EDTA]^{4-}$ is a: [UPSEAT 2004]


- (a) Monodentate ligand (b) Bidentate ligand
(c) Quadridentate ligand (d) Hexadentate ligand

17. Which of the following statements is incorrect ?

[KCET 2004]

- (a) In $K_3[Fe(CN)_6]$, the ligand has satisfied only the secondary valency of ferric ion.
(b) In $K_3[Fe(CN)_6]$, the ligand has satisfied both primary and secondary valencies of ferric ion.
(c) In $K_4[Fe(CN)_6]$, the ligand has satisfied both primary and secondary valencies of ferrous ion.
(d) In $[Cu(NH_3)_4]SO_4$, the ligand has satisfied only the secondary valency of copper.

18. Which of the following is considered as an anticancer species. [CBSE PMT 2004]

19. An aqueous solution of $CoCl_2$ on addition of excess of concentrated HCl turns blue due to formation of

[AIIMS 2005]

- (a) $[Co(H_2O)_4Cl_2]$
(b) $[Co(H_2O)2Cl_4]^{2-}$
(c) $[CoCl_4]^{2-}$
(d) $[Co(H_2O)2Cl_2]$

20. The correct order for the wavelength of absorption in the visible region is

- (a) $[Ni(NO_2)_6]^{4-} < [Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+}$
(b) $[Ni(NO_2)_6]^{4-} < [Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+}$
(c) $[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
(d) $[Ni(NH_3)_6]^{2+} < [Ni(H_2O)_6]^{2+} < [Ni(NO_2)_6]^{4-}$

21. In which of the following pairs both the complexes show optical isomerism

- (a) *cis*- $[Cr(C_2O_4)_2Cl_2]^{3-}$, *cis*- $[Co(NH_3)_4Cl_2]$
(b) $[Co(en)_3]Cl_3$, *cis*- $[Co(en)_2Cl_2]Cl$
(c) $[PtCl(dien)]Cl$, $[NiCl_2Br_2]^{2-}$
(d) $[Co(NO_3)_3(NH_3)_3]$, *cis*- $[Pt(en)_2Cl_2]$

22. Which of the following compounds shows optical isomerism

[AIEEE 2005; CBSE PMT 2005]

- (a) $[Cu(NH_3)_4]^{2+}$ (b) $[ZnCl_4]^{2-}$
(c) $[Cr(C_2O_4)_3]^{3-}$ (d) $[Co(CN)_6]^{3-}$

Assertion & Reason

For AIEEE Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
(c) If assertion is true but reason is false.
(d) If the assertion and reason both are false.
(e) If assertion is false but reason is true.

1. Assertion : Potassium ferrocyanide and potassium ferricyanide both are diamagnetic.

Reason : Both have unpaired electrons.

2. Assertion : NF_3 is a weaker ligand than $N(CH_3)_3$

Reason : NF_3 ionizes to give F^- ions in aqueous solution.

3. Assertion : The $[Ni(en)_3]Cl_2$ (*en* = ethylene diamine) has lower stability than $[Ni(NH_3)_6]Cl_2$


Reason : In $[Ni(en)_3]Cl_2$ the geometry of *Ni* is trigonal bipyramidal

4. Assertion : The ligands nitro and nitrito are called ambidentate ligands.

Reason : These ligands give linkage isomers.

5. Assertion : Geometrical isomerism is also called *cis-trans* isomerism.

Reason : Tetrahedral complexes show geometrical isomerism.

6. Assertion :  is named as tetrakis (ethylene diamine) μ -hydroxo-imido dicobalt (III) ion.

Reason : In naming polynuclear complexes i.e., containing two or more metal atoms joined by bridging ligands, the word μ is added with hyphen before the name of such ligands.

7. Assertion : $[Fe(CN)_6]^{3-}$ is weakly paramagnetic while $[Fe(CN)_6]^{4-}$ is diamagnetic.

Reason : $[Fe(CN)_6]^{3-}$ has +3 oxidation state while $[Fe(CN)_6]^{4-}$ has +2 oxidation state.

8. Assertion : H_2N-NH_2 is a chelating ligand.

Reason : A chelating ligand must possess two or more lone pairs at such a distance

922 Co-ordination Chemistry

that it may form suitable strain free rings at the metal ion.

9. Assertion : $[Ti(H_2O)_6]^{3+}$ is coloured while $[Sc(H_2O)_6]^{3+}$ is colourless.

Reason : $d-d$ transition is not possible in $[Sc(H_2O)_6]^{3+}$.

10. Assertion : All the octahedral complexes of Ni^{2+} must be outer orbital complexes.

Reason : Outer orbital octahedral complexes are given by weak ligands.

11. Assertion : Potassium ferrocyanide is diamagnetic whereas potassium ferricyanide is paramagnetic.

Reason : Crystal field splitting in ferrocyanide ion is greater than that of ferricyanide ion.

[AIIMS 2005]

Answers

Basic Terms

1	b	2	c	3	b	4	b	5	d
6	a	7	b	8	c	9	b	10	c
11	d	12	c	13	d	14	b	15	b
16	b	17	a	18	c	19	c	20	b
21	b	22	c	23	b	24	d	25	a
26	d	27	a	28	b	29	b	30	c
31	b	32	c	33	b	34	b	35	d
36	d	37	b	38	b	39	d	40	b
41	d	42	b	43	d	44	b	45	a
46	a	47	d	48	c	49	b	50	d
51	a	52	b	53	d	54	a	55	d
56	d	57	a	58	d	59	a	60	d
61	c	62	d						

Nomenclature, oxidation State and EAN number

1	b	2	c	3	c	4	b	5	d
6	c	7	b	8	a	9	a	10	a
11	c	12	a	13	b	14	a	15	b
16	c	17	d	18	b	19	d	20	a
21	b	22	b	23	a	24	b	25	a
26	d	27	c	28	b	29	a	30	a
31	a	32	b	33	a	34	b	35	c
36	b	37	a	38	a	39	b	40	c
41	a	42	b	43	c	44	b	45	b
46	d	47	a	48	b	49	c	50	c

51	b	52	b	53	c	54	d	55	b
56	a	57	b	58	c				

Isomerism and Magnetic properties

1	a	2	d	3	a	4	c	5	c
6	d	7	b	8	a	9	a	10	c
11	d	12	d	13	b	14	c	15	a
16	b	17	a	18	d	19	a	20	b
21	c	22	b	23	b	24	a	25	a
26	d	27	d	28	a	29	c	30	a
31	b	32	a	33	c	34	a	35	d
36	c	37	d	38	a	39	d	40	b
41	a	42	c	43	d	44	a	45	b
46	e	47	a	48	d	49	c	50	c
51	d	52	a	53	c	54	a	55	a
56	c	57	c						

Hybridisation and Geometry

1	a	2	b	3	a	4	b	5	a
6	b	7	b	8	d	9	c	10	a
11	b	12	c	13	d	14	a	15	b
16	b	17	d	18	d	19	a	20	a
21	b	22	d	23	c	24	c	25	c
26	d	27	d	28	d	29	d	30	a
31	d	32	c	33	a	34	a	35	c
36	a	37	c	38	d	39	d	40	b
41	b	42	a	43	c				

Complexes and complex stability

1	d	2	d	3	d	4	b	5	a
6	a	7	d	8	b	9	c	10	b
11	b	12	b	13	a	14	b	15	b

Application of organometallics

1	d	2	d	3	d	4	c	5	b
6	b	7	b	8	a	9	a	10	d
11	c	12	b	13	a	14	c	15	d
16	c	17	c	18	c	19	d	20	a

Critical Thinking Questions


1	b	2	b	3	a	4	b	5	b
6	b	7	c	8	c	9	d	10	a
11	d	12	d	13	d	14	a	15	d
16	d	17	ad	18	a	19	c	20	a
21	b	22	c						


Assertion & Reason

1	d	2	c	3	d	4	a	5	c
6	e	7	b	8	e	9	a	10	b
11	c								

Answers and Solutions

Basic terms

- (b) (CN) are linked with secondary valency.
- (c) In Cuprammonium sulphate $[Cu(NH_3)_4]SO_4$ co-ordination no. of Cu is 4.
- (b) As it makes use of its two atoms to form two co-ordinate covalent bonds with the central metal ion.
- (b) $[Co(en)_2Br_2]Cl_2$
C.N. of Co = $2 \times$ number of bidentate ligand + $1 \times$ number of monodentate ligand = $2 \times 2 + 1 \times 2 = 6$.
- (b) The charge does not decide the formation of bond but availability of lone pair decide the formation of Co-ordinate bond.
- (c) In $[Co(en)_2Cl_2]^+$
No. of monodentate ligand = 2
No. of bidentate ligand = 2
Co-ordination no. of the metal = $2 + 2(2) = 6$.
- (c)
- (d)
 
- (c) According to modern view primary valency of complex compound is its oxidation number while secondary valency is the co-ordination number.
- (b) Co-ordination number is equal to total number of ligands in a complex.
- (b) Cuprammonium salt- $[Cu(NH_3)_4]SO_4$
 $[Cu(NH_3)_4]SO_4 \rightleftharpoons [Cu(NH_3)_4]^{2+} + SO_4^{2-}$
So, it will give two ions in water.
- (a) The co-ordination no. = no. of ligands attached.
- (c) Primary valencies are also known as oxidation state.
 $K_2[Ni(CN)_4], 2 + x - 4 = 0 \Rightarrow x = +2$
- (b) Ag^+ has two coordination number forms complex with excess of CN^- $Ag(CN)_2^-$.
- (b) According to Lewis the ligands are basic in nature.
- (c) The number of atoms of the ligand that are directly bounded to the central metal atom or ion by coordinate bond is known as the coordination number of the metal atom or ion
coordination number of metal = number of σ bonds formed by metal with ligand.

- (b) $Ag_2S + 4NaCN \rightleftharpoons 2Na[Ag(CN)_2] + Na_2S$
Sodium dicyano argentate
 $2Na[Ag(CN)_2] + Zn \rightarrow Na_2[Zn(CN)_4] + 2Ag \downarrow$
Sodium tetracyano zincate (ppt)
 - (d) In $K_4Fe(CN)_6$, the species retains its identity in solid as well as in solution state.
 - (d) The number of atom of the ligand that are directly bound to the central metal atom or ion by coordinate bonds is known as the coordinate number of the metal or ion. It is actually the number of chemical bonds which the ligand form with the central metal atom or ion.
- 
- (a) CN^- ions acts both as reducing agent as well as good complexing agent.
 - (b) ZnS structure shows the coordination number of Zn is 4.
 - (c) Wilkinson's catalyst is $Rh(PPh_3)_3Cl$.
 - (b) The complexes can be written as follows
 $[Co(NH_3)_6]Cl_3$ (A) $[Co(NH_3)_5Cl]Cl_2$ (B) $[Co(NH_3)_4Cl_2]Cl$ (C)
 Hence, number of primary valencies are 3, 2 and 1 respectively.
 - (c) Ligand must have capacity to donate lone pair of electrons to form co-ordinate bond.
 - (b) $K_3[Fe(CN)_6]$ because in it CN^- donats a lone pair of electron.
 - (b) Al has 6 co-ordination number.
 - (d) $K_4[Fe(CN)_6] \rightarrow 4K^+ + [Fe(CN)_6]^{4-}$
Complex has negative charge so it is anionic complex.
 - (d) Co-ordination number is equal to total number of ligands in a complex.
 - (b) Lithium tetrahydroaluminate is $Li[Al(H)_4]$.
 - (b) Negative ligands end in -O eg. SO_4^{2-} (sulphato).
 - (b) Turnbull's blue is $K\{Fe^{II}[Fe^{III}(CN)_6]\}$.
 - (a) $Fe + 5CO \xrightarrow[\text{Pressure}]{\text{Heat}} [Fe(CO)_5]$
Iron pentacarbonyl.
 - (c) When a ligand attaches to the central metal atom through two or more atoms to form a ring like structure is known as chelates and the ligand is called chelating ligand.

53. (b) eg. $n \text{CH}_3 - \text{CH} = \text{CH}_2 \xrightarrow[(\text{C}_2\text{H}_5)_3\text{Al}]{\text{TiCl}_4+} \begin{array}{c} (-\text{CH} - \text{CH}_2)_n \\ | \\ \text{CH}_3 \end{array}$
55. (d) CH_3Li is the organometallic compound in which lithium bonded with carbon and organometallic are those in which metal-carbon bond found.
56. (d) Potash alum is a mixed salt of K_2SO_4 and $\text{Al}_2(\text{SO}_4)_3$ and on dissolving, it gives all three ion Al^{3+} , K^+ and SO_4^{2-} of which it is made.
57. (a) Carnallite is a double salt with molecular formula $\text{KCl} \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$. It gives K^+ , Mg^{2+} and Cl^- ions in solution.
59. (a) General formula for alum is $\text{M}_2\text{SO}_4 \cdot \text{R}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$
 M = mono valent cation (K^+ , Na^+ )
 R = Trivalent cation (Al^{+3} , Fe^{+3})
Hence, $\text{K}_2\text{SO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$ represent an alum.
60. (d) $\text{K}_4[\text{Fe}(\text{CN})_6]$ is a complex salt. On ionisation it will dissociate in 4K^+ and $[\text{Fe}(\text{CN})_6]^{4-}$ ion. Hence, in $\text{K}_4[\text{Fe}(\text{CN})_6]$ five ions are present.
61. (c) Due to $\text{C}-\text{Mg}$ bond.
62. (d) $\text{EAN} = (\text{At. number} - 0.5 + 2 \times \text{C.N.})$
Hence, EAN of Ni in $\text{Ni}(\text{CO})_4$ is
 $= 28 - 0 + 2 \times 4 = 36$
20. (a) $\text{EAN} = (\text{atomic no.}) - (\text{oxidation state}) + (2 \times \text{number of Ligands}) = 28 - 2 + 2 \times 4 = 34$
21. (b) Follow IUPAC rule.
22. (b) $[\text{Pt}(\text{C}_2\text{H}_4)_2\text{Cl}_3]^-$
 $x + 0 + 3x(-1) = -1$, $x - 3 = -1$, $x = +2$.
23. (a) $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3 \rightarrow [\text{Co}(\text{NH}_3)_6]^{3+} + 3\text{Cl}^-$
 $x + 6(0) = +3 \Rightarrow x = +3$.
24. (b) Follow IUPAC rule.
25. (a) $\text{EAN} = \text{at. no. of central atom} - \text{oxidation state} + 2 \times (\text{no. of ligands}) = 27 - 3 + 2 \times 6 = 36$.
26. (d) Follow IUPAC rules for nomenclature.
27. (c) $\text{EAN of a central metal ion} = (\text{atomic no. of central atom}) - \text{oxidation state} + \text{no. of ligands} \times 2 = 26 - 3 + (6 \times 2) = 23 + 12 = 35$
28. (b) $+1 \times 4 + x - 1 \times 4 = 0$
 $4 + x - 4 = 0 \Rightarrow x = 0$ for Ni .
29. (a) Follow IUPAC rule.
30. (a) In complex $\text{K}_4[\text{Fe}(\text{CN})_6]$ the Fe obey EAN rule strictly.
31. (a) $\text{Ni}(\text{CO})_4$ has O.N. zero for Ni .
32. (b) $3 \times (+1) + x + 6 \times (-1) = 0$ or $x = 6 - 3 = +3$ Oxidation state of $\text{Fe} = +3$.
33. (a) Complexes containing carbonyl ligand (CO) have zero oxidation state.
34. (b) Follow IUPAC rule.
35. (c) Follow IUPAC rule.
36. (b) Follow IUPAC rule.
38. (a) Follow IUPAC rule.
39. (b) Follow IUPAC rule.
41. (a) Follow IUPAC rule.
42. (b) $[\text{Ni}(\text{CN})_4]^{x-}$, ($\text{Ni} = +2$) ($\text{CN} = -1$)
 $x = 2 + 4(-1) \Rightarrow x = -2$.
44. (b) $[\text{Cu}(\text{NH}_3)_4]^{2+}$ so, copper has + 2 valency.
47. (a) Follow IUPAC rule.
50. (c) Follow IUPAC rule.
51. (b) $[\text{Co}(\text{H}_2\text{O})_5\text{Cl}]^{+2}$
 $x + 0 + (-1) = +2$
 $x - 1 = 2 \Rightarrow x = +3$.
52. (b) It is clear from the chemical formulae that Ag is central metal atom and ligands are 2 ammonia molecule.
Hence, compound is $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$.


Nomenclature, Oxidation State & EAN number

1. (b) $[\text{Co}(\text{H}_2\text{O})_6]\text{Cl}_2 = [\text{Co}(\text{H}_2\text{O})_6]^{+2} + 2\text{Cl}^-$.
5. (d) Follow IUPAC rule.
7. (b) Follow IUPAC rule.
9. (a) Follow IUPAC rule.
10. (a) Follow IUPAC rule.
11. (c) $\text{EAN} = \text{Atomic number} - \text{Oxidation state} + 2 \times \text{number of Ligands} = 26 - 2 + 2(6) = 36$.
12. (a) EAN of Pt in $[\text{PtCl}_6]^{2-} = 80$
13. (b) $\text{Na}_2[\text{CrF}_4\text{O}]$
 $x + 4(-1) + (-2) = -2$
 $x - 6 = -2 \Rightarrow x = +4$.
14. (a) Follow IUPAC rule.
16. (c) Follow IUPAC rule.
17. (d) Follow IUPAC rule.
18. (b) The oxidation state of metal in metal carbonyl is zero.
19. (d) $x + 6 \times (0) + 3 \times (-1) = 0$

53. (a) In the compounds $[Co(NH_3)_5NO_2]Cl_2$, the oxidation state of cobalt is +3 and here 5 NH_3 ligand, a NO_2 ligand are attached to the central atom. therefore its name is pentaamminecobalt (III) chloride.
54. (b) CrO_2Cl_2 , MnO_4^{+7} .
55. (b) In the given ion $[Cr(NH_3)_6]^{3+}$, the oxidation state of chromium is +3 and here 6 NH_3 ligand are attached to the central metal atom. Therefore its IUPAC name is hexamminechromium (III) ion.
56. (a) It is potassium ammine dicyano dioxo peroxo-chromate (VI)
57. (b) Potassium hexa cyanoferrate (III).

Isomerism and Magnetic Properties

1. (a) Octahedral complexes of the type $[MA_4B_2]$, $[MA_2B_4]$, $[MA_3B_3]$ exhibit geometrical isomerism.
2. (d) The number of unpaired electrons in the Complex ion $[CoF_6]^{3-}$ is 4.




Here more than one atom function as donor, as oxygen in first one and nitrogen in second, so they show linkage isomerism


5. (c) The two given compounds have same composition but in solution both will give different ions. The isomerism is known as ionisation isomerism.
6. (d) Both produce different ions in solution state-
 $[Co(NH_3)_4Cl_2]NO_2 \rightleftharpoons [Co(NH_3)_4Cl_2]^+ NO_2^-$
 $[Co(NH_3)_4Cl.NO_2]Cl \rightleftharpoons [Co(NH_3)_4Cl.NO_2]^+ + Cl^-$
7. (b) The compound which has same composition but give different ions in solution, show ionization. So $[Co(NH_3)_5Br]SO_4$ is ionization isomer.
 $[Co(NH_3)_5Br]SO_4 \rightleftharpoons [Co(NH_3)_5Br]^{2+} + SO_4^{2-}$
 $[Co(NH_3)_5SO_4]Br \rightleftharpoons [Co(NH_3)_5SO_4]^+ + Br^-$
9. (a) Co-ordination isomerism is caused by the interchange of ligands between cis and trans structure.
10. (c) $[Co(NH_3)_5NO_2]Cl_2$ will not show geometrical isomerism because this complex showed 4 and 6 co-ordination number.

13. (b) $[Fe(H_2O)_6]^{2+}$ has four unpaired electrons, $[Cr(H_2O)_6]^{3+}$, $[Cu(H_2O)_6]^{2+}$ and $[Zn(H_2O)_6]^{2+}$ have 3, 1, 0 unpaired electrons respectively.
14. (c) The electronic configuration of Ni in $[Ni(CN)_4]^{2-}$, $[NiCl_4]^{2-}$ and $Ni(CO)_4$ are as following


Ni^+ in $[Ni(CN)_4]^{2-}$ -



Ni^{2+} in $[NiCl_4]^{2-}$ -




Ni in $[Ni(CO)_4]$ -




CO and CN^- are strong ligands so they induces pairing of electrons so their complexes are diamagnetic while Cl^- is a weak ligand so it does not induce the pairing of electrons so its complex is paramagnetic.

15. (a) $[Co(NH_3)_4Cl_2]^+$ is the Ma_4b_2 and Ma_2b_3 type complex.



16. (b) In $[Ag(NH_3)_2]Cl$, Ag^+ contains d^{10} configuration. All others contain unpaired electrons.


17. (a) $Pt(NH_3)_2Cl_2$.



19. (a)
-

21. (c) $[Co(en)_2NO_2Cl]Br$; $[Co(en)_2ONOC]Br$
22. (b) Because it will not give any ions in solution.

23. (b) $[Co(en)_2Cl_2]^+$ have three optical isomers which are.




24. (a) Magnetic moment of $[Cu(NH_3)_4]^{2+}$ ion is 1.414 due to the presence of one unpaired electron.
25. (a) Due to the presence of one unpaired electron, both are slightly paramagnetic.
27. (d) 1. $[Cu(NH_3)_4].[PtCl_4]$
 2. $[Cu(NH_3)_3Cl].[PtCl_3(NH_3)]$
 3. $[Cu(NH_3)_2Cl_2].[PtCl_2(NH_3)_2]_{cis}$
 4. $[Cu(NH_3)_2Cl_2].[PtCl_2(NH_3)_2]_{Trans}$
 5. $[Cu(NH_3)Cl_3].[PtCl(NH_3)_3]$
 6. $[Pt(NH_3)_4Cl].[CuCl_4]$
29. (c) Co-ordination isomerism is possible when both +ve and -ve ions of a salt are complex ions and the two isomers differ in the distribution of ligands in the cation and the anion.
30. (a) On ionisation it gives maximum number of (four) ions.
31. (b) The anhydrous complex of Cu^+ do not involve d-d transition and are thus colourless.
32. (a) Mn^{2+}

↑↓	↑↓	↑	↑	↑	
$3d^7$					$4s^0$


In presence of H_2O which is a weak ligand no pairing occurs which results in unpaired electrons left in the compound, due to which it shows paramagnetism.

33. (c) Geometrical isomers (cis and trans) and linkage isomers ($-SCN$ and $-NCS$).
34. (a) Due to presence of strong ligand all the e^- get paired. So no. of unpaired electrons are 0.
36. (c) Both have same composition but give different ions in the solution.
37. (d) $[Pt(en)_2]Cl_2$






38. (a) CO is a strong ligand so induce pairing of electrons and hence $Ni(CO)_4$ is diamagnetic.

40. (b) $[Co(en)_3]^{3+}$






41. (a) $[Co(NH_3)_3Cl_3]$ does not have optical isomers because it is of formula MA_3B_3 which does not show optical isomerism.
42. (c) Change in composition of co-ordination sphere yield ionization isomers.
 $[Cr(H_2O)_6]Cl_3$ and $[CrCl_3(H_2O)_3].3H_2O$
43. (d) NO_2 is ambident and can be linked either to N-side as ($-NO_2$) or to O-side as ($-ONO$).
44. (a) As $[Co(NH_3)_3Cl_3]$ does not losses any Cl^- ions in the solution so it will not give ppt. with $AgNO_3$.
45. (b) $[Co(NH_3)_6]Cl_3 = [Co(NH_3)_6]^{3+} + 3Cl^-$.
46. (e) $CoCl_3.5NH_3.H_2O$ is pink in colour.
47. (a) The configuration of Ni^{2+} has two unpaired electron so it is paramagnetic.
48. (d) $[Pt(NH_3)_4]Cl_2 \rightarrow [Pt(NH_3)_4]^{2+} + 2Cl^-$
 As it gives Cl^- ions in solution so it will give white ppt. of $AgCl$ with $AgNO_3$.
49. (c) $[Co(NH_3)_5Cl]Cl_2 = [Co(NH_3)_5Cl]^{2+} + 2Cl^-$
50. (c) $[Cr(NH_3)_5Cl]SO_4 \rightarrow [Cr(NH_3)_5Cl]^{2+} + SO_4^{2-}$




926 Co-ordination Chemistry



Cl^- ion is precipitable.





Number of unpaired electrons = 3.




Number of unpaired electrons = 5

$$\text{Magnetic moment} = \sqrt{n(n+2)} = \sqrt{5(5+2)}$$


$$= \sqrt{35} = 5.91 \text{ BM}.$$




d^2sp^3 ,
















d^2sp^3 ,




d^2sp^3 ,















Hybridisation and Geometry




In presence of CN^-



Hence, Octahedral geometry.






Electronic configuration of $Fe = [Ar]4s^2 3d^6$

Electronic configuration of $Fe^{+3} = [Ar]3d^5$

Number of ligand (coordination number) = 6

Nature of ligand is strong field.



Hybridization of Fe is $d^2 sp^3$.



32. (c) The compounds which show d^6 - configuration are octahedral complexes.

33. (a) CN^- ligand has strong field ligand because of higher value of Δ .

35. (c) Ammine (NH_3) is neutral ligand.


37. (c) A strong field ligand produces low spin complexes.

38. (d) Cyanide ion is strong field ligand because it is a pseudohalide ion pseudohalide ions are stronger coordinating ligand & they have the ability to form σ bond (from the pseudohalide to the metal) and π bond (from the metal to pseudohalide).



In presence of weak ligand field, there will be no pairing of electrons. So it will form a high spin complex. i.e. the number of unpaired electrons = 5.

40. (b) The complexes, in which, the metal and ligand form a bond that involves the π -electrons of the ligand are known as π -complexes e.g. Ferrocene



41. (b) CO has strong $M \rightarrow L$ π -bonding ability so it can accept electron pairs from metal ion so it is a π -acid ligand.

42. (a) d^4 (in strong ligand field).

43. (c) Both $Ni(CO)_4$ and $Ni(PPh_3)_2Cl_2$ are tetrahedral.

Complexes and complex stability


1. (d) The cyano and ammine complexes are far more stable than those formed by halide ions. This is due to the fact that NH_3 and CN^- are strong Lewis bases.

4. (b) Greater the charge on central metal ion greater is the stability of complex.




5. (a) $CuSO_4$ on reaction with KCN gives $K_3[Cu(CN)_4]$







6. (a) The most probable complex which gives three moles ions in aqueous solution may be $[Co(NH_3)_5NO_2]Cl_2$ because it gives two chlorine atoms on ionisation.



7. (d) $3K_4[Fe(CN)_6] + 4FeCl_3 \rightarrow Fe_4[Fe(CN)_6]_3 + 12KCl$
 pot. ferrocyanide (Ferri ferrocyanide)
 (Prussian blue)



10. (b) Ni reacts with dimethylglyoxime to give red ppt. of nickel-dimethyl glyoxime complex.




15. (b) Aluminum is a p -block element and does not form complex compounds.

Application of Organometallics

1. (d) $(CH_3CH_2)_3Al + TiCl_4$ is the Ziegler-Natta catalyst.

2. (d) Alum acts as coagulating agent.




6. (b) As there is no direct bonding between the metal atom and the carbon atom.


9. (a) $(CH_3)_4Sn$ is a σ -bonded organometallic compound.

10. (d) Wilkinson's catalyst is used as homogeneous catalyst in the hydrogenation of alkenes as -


928 Co-ordination Chemistry




12. (b) C_2H_5Li is an organo-metallic compound.
13. (a) Because there is direct bonding of metal ion with carbon.
14. (c) Organometallic compounds are those compounds in which carbon atom is directly linked to metal. But in sodium ethoxide as oxygen is attached to sodium metal so it is not an organometallic compound.
16. (c) $Al(OC_2H_5)_3$ contains bonding through O and thus it does not have metal-carbon bond.
17. (c) Chlorophyll are green pigment in plant & contain magnesium instead of calcium.
19. (d) This reaction carried out in alkaline pH i.e., 9-11
20. (a)



Zeise's salt



Ferrocene



dibenzene chromium


$(C_2H_5)_4Sn$
tetraethyl
tin

Critical Thinking Questions


1. (b) Its coordination number will be 6 because it is bonded with three bidentate ligands.
Oxi. No. of Cr in $K_3[Cr(C_2O_4)_3]$ is
 $x + 3(-2) + 3(+1) = 0 \Rightarrow x = +3$
2. (b) e.g. $Fe(CO)_5, Ni(CO)_4$ etc.
3. (a) $[Co(NH_3)_5SO_4]Br + AgNO_3 \rightarrow$
0.02 mole
 $[Co(NH_3)_5SO_4]NO_3 + AgBr$
0.02 mole (y)
 $[Co(NH_3)_5Br]SO_4 + BaCl_2 \rightarrow$
0.02 mole
 $[Co(NH_3)_5Br]Cl_2 + BaSO_4$
0.02 mole (z)
On using one lit. solution, we will get 0.01 mole y and 0.01 mole z.
4. (b) $[Fe(\eta^5-C_5H_5)_2]$ is the organometallic compound which has σ and π bonds present.
5. (b) $[Ni(NH_3)_4]SO_4$
 $x + 0 + (-2) = 0 \Rightarrow x = +2$ is valency and 4 is C.N. of Ni .
6. (b) Co-ordination number is equal to total number of ligands in a complex.
7. (c) $n \left[\begin{array}{c} Cl \\ \diagup \quad \diagdown \\ Si \\ \diagdown \quad \diagup \\ CH_3 \end{array} \right] + 2nH_2O \rightarrow \left(\begin{array}{c} CH_3 \\ | \\ -O-Si-O- \\ | \\ CH_3 \end{array} \right)_n$
8. (c) Because it is a polydentate ligand which binds the central atom nickel forming a ring like structure.
9. (d) $4KCN + CuSO_4 \rightarrow K_3[Cu(CN)_4]$
Potassium tetracyano cuprate
10. (a) Roasted + $CN^- + H_2O \xrightarrow{O_2} [Au(CN)_2]^- + OH^-$
gold ore
 $[Au(CN)_2]^- + Zn \rightarrow [Zn(CN)_4]^{2-}$.
11. (d) Potassium ferrocyanide $K_4[Fe(CN)_6]$ will ionize as $K_4[Fe(CN)_6] \rightleftharpoons 4K^+ + [Fe(CN)_6]^{4-}$
So, it will give five ions in solution.
12. (d) Complex ion Hybridization of central atom

$[Fe(CN)_6]^{4-}$	d^2sp^3 (inner)
$[Mn(CN)_6]^{4-}$	d^2sp^3 (inner)
$[Co(NH_3)_6]^{3+}$	d^2sp^3 (inner)
$[Ni(NH_3)_6]^{2+}$	sp^3d^2 (outer)
13. (d) $[Co(en)_2Cl_2]^+$ shows geometrical as well as optical isomerism.
14. (a)

15. (d)



16. (d)



Ethylenediamine tetraacetate ion (EDTA)

They have six donor atoms. poly dentate ligands have flexidentate character. It is not necessary that all the donor atoms present in the polydentate ligands should form co-ordinate bonds with central metal atom or ion.

17. (a,d) In $K_3[Fe(CN)_6]$ the ligand are negative which is present in coordination spheres shows a dual behaviour. It may satisfied both primary & secondary valencies while neutral ligand satisfied only secondary valencies.




18. (a) Cis Isomer of $[Pt(NH_3)_2Cl_2]$ is used as an anticancer drugs for treating several type of malignant tumours. When it is injected into the blood stream the more reactive Cl groups are lost so the Pt atom bonds to a N atom in guanosine (a part of DNA) This molecule can bond to two different guanosine units & by bridging between them it upsets the normal reproduction of DNA.

19. (c) $CoCl_2$ is a weak Lewis acid, reacting with chloride ion to produce salt containing the tetrahedral $[CoCl_4]^{2-}$ ion. $CoCl_2$ is blue when anhydrous, and a deep magenta colour when hydrated, for this reason it is widely used as an indicator for water.

20. (a) The absorption of energy or the observation of colour in a complex transition compound depends on the charge of the metal ion and

the nature of the ligands attached. The same metal ion with different ligands shows different absorption depending upon the type of ligand. The presence of weak field ligands make the central metal ion to absorb low energies i.e., of higher wavelength. The field strength of ligands can be obtained from spectrochemical series. i.e.

(weak field) $I^- < Br^- < S^{2-} < Cl^- < NO_3^- < F^- < OH^- < H_2O < NH_3 < NO_2^- < CN^- < CO$ (strong field)

21. (b) $[Co(en)_3]Cl_3$ cis- $[Co(en)_2Cl_2]Cl$ 

Assertion and Reason

- (d) Potassium ferrocyanide and potassium ferricyanide both are not diamagnetic because both do not have paired electrons. Assertion and reason both are false.
- (c) It is correct statement that NF_3 is a weaker ligand than $N(CH_3)_3$, the reason is that fluorine is highly electronegative therefore, it with draw electrons from nitrogen atom. Hence, the lone pair of nitrogen atom cannot be ligated. While $N(CH_3)_3$ is a strong ligand because CH_3 has electron releasing group.

3. (d) Both assertion and reason are false. $[Ni(en)_3]Cl_2$ is a chelating compound and chelated complexes are more stable than similar complexes involves breaking of two bonds rather than one. In $[Ni(en)_3]Cl_2$, Ni with d^8 configuration shows octahedral geometry. Six electrons will occupy the t_{2g} orbitals and two electrons will occupy the e_g orbitals.

4. (a) Both assertion and reason are true and reason is the correct explanation of assertion. When a monodentate ligand has two possible donor atoms and attached in two ways to the central metal atom are called ambidentate ligands.

5. (c) Assertion is true but reason is false. Tetrahedral complexes do not show geometrical isomerism because the relative position of the atoms with respect to each other will be the same.

6. (e) Assertion is false but reason is true.

$\left[(en)_2 Co \begin{array}{c} \nearrow NH \\ \searrow OH \end{array} Co(en)_2 \right]^{3+}$ is named as tetrakis (ethylene diamine)- μ -hydroxo- μ -imido dicobalt (III) ion. For more than one bridging group the word μ is repeated before each bridging group.

7. (b) Both assertion and reason are true but reason is not the correct explanation of assertion. $[Fe(CN)_6]^{3-}$ has EAN equal to 35 and thus possesses unpaired electron to show paramagnetic nature while $[Fe(CN)_6]^{4-}$


possesses no unpaired electron and thus shows diamagnetic nature.

8. (e) Assertion is false but reason is true. H_2N-NH_2 does not act as chelating ligand. The coordination by hydrazine leads to a three member highly unstable strained ring and thus it does not act as chelating agent.

9. (a) Both assertion and reason are true and reason is the correct explanation of assertion. $[Sc(H_2O)_6]^{3+}$ has no unpaired electron in its d subshell and thus $d-d$ transition is not possible whereas $[Ti(H_2O)_6]^{3+}$ has one unpaired electron in its d subshell which gives rise to $d-d$ transition to impart colour.

10. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.

Ni^{2+} configuration



During rearrangement only one $3d$ -orbital may be made available by pairing the electrons. Thus, inner d^2sp^3 hybridization is not possible, so, only sp^3d^2 (outer) hybridization can occur.

11. (c) $K_4[Fe(CN)_6]$ and $K_3[Fe(CN)_6]$ both are low spin complex due to strong ligand field. That is why it is false that crystal field splitting in ferrocyanide ion is greater than ferricyanide ion.