





# Alcohols, Phenols and Ethers

## NCERT Intext Questions

**Q. 1.** Classify the following as primary, secondary and tertiary alcohols:



**Ans.** (i) Primary

(ii) Primary

(iii) Primary


(iv) Secondary

(v) Secondary


(vi) Tertiary

**Q. 2.** Identify allylic alcohols in the above examples.

**Ans.** (ii)  $\text{H}_2\text{C} = \text{CH} - \text{CH}_2\text{OH}$





Q. 3. Name the following compounds according to IUPAC system.





- Ans. (i) 3-Chloromethyl-2-isopropylpentan-1-ol (ii) 2, 5-Dimethylhexane-1,3-diol  
 (iii) 3-Bromocyclohexanol (iv) Hex-1-en-3-ol  
 (v) 2-Bromo-3-methylbut-2-en-1-ol


Q. 4. Show how are the following alcohols prepared by the reaction of a suitable Grignard reagent on methanal?




Q. 5. Write structures of the products of the following reactions.




(ii)  $\text{NaBH}_4$  is a weak reducing agent, it reduces the aldehydes/ketones and not the esters.





**Q. 8.** *Ortho* and *para*-nitrophenols are more acidic than phenol. Draw the resonance structures of the corresponding phenoxide ions.

**Ans.** The resonance structures of *o*- and *p*-nitrophenoxide ions and phenoxide ion are given as follows:



It is clear from the above structures that due to  $-R$ -effect of  $\text{NO}_2$  group, *o*- and *p*-nitrophenoxide ions are more stable than phenoxide ion. Consequently, *o*- and *p*-nitrophenols are more acidic than phenol.




**Q. 9.** Write the equations involved in the following reactions:

(i) Reimer-Tiemann reaction



(ii) Kolbe's reaction

**Ans.** Refer to Points to remember 15.

**Q. 10.** Write the reactions of Williamson synthesis of 2-ethoxy-3-methylpentane starting from ethanol and 3-methylpentan-2-ol. [HOTS]




**Q. 11.** Which of the following is an appropriate set of reactants for the preparation of 1-methoxy-4-nitrobenzene and why?





**Ans.** Chemically both sets are equally probable. In set (i), the Br atom is activated by the electron withdrawing effect of  $-\text{NO}_2$  group. Therefore, nucleophilic attack by  $\text{CH}_3\text{O}^-\text{Na}^+$  followed by elimination of NaBr gives desired ether, while in set (ii) nucleophilic attack by 4-nitrophenoxide ion on methyl bromide gives the desired ether.






**Q. 12. Predict the products of the following reactions:**












[CBSE (F) 2012]



## NCERT Exercises

**Q. 1. Write the IUPAC names of the following compounds:**





**Ans.** (i) 2, 2, 4-Trimethylpentan-3-ol

(iii) Butane-2, 3-diol

(v) 2-Methylphenol

(vii) 2, 5-Dimethylphenol

(ix) 1-Methoxy-2-methylpropane

(xi) 1-Phenoxyheptane

(ii) 5-Ethylheptane-2, 4-diol

(iv) Propane-1, 2, 3-triol

(vi) 4-Methylphenol

(viii) 2, 6-Dimethylphenol

(x) Ethoxybenzene

(xii) 2-Ethoxybutane

**Q. 2.** Write the structures of the compounds whose IUPAC names are as follows:

(i) 2-Methylbutan-2-ol

(iii) 3, 5-Dimethylhexane-1, 3, 5-triol

(v) 1-Ethoxypropane

(vii) Cyclohexylmethanol

(ix) Cyclopent-3-en-1-ol










(ii) 1-Phenylpropan-2-ol

(iv) 2, 3-Diethylphenol

(vi) 2-Ethoxy-3-methylpentane

(viii) 3-Cyclohexylpentan-3-ol




(x) 4-Chloro-3-ethylbutan-1-ol







**Q. 3.** (i) Draw the structures of all isomeric alcohols of molecular formula  $\text{C}_5\text{H}_{12}\text{O}$  and give their IUPAC names.

(ii) Classify the isomers in Q. 3(i) into primary, secondary and tertiary alcohols.


**Ans.** Eight isomers are possible. These are:





**Q. 4. Explain why propanol has higher boiling point than that of the hydrocarbon, butane?**


**Ans.** The molecules of butane are held together by weak van der Waal's forces of attraction while those of propanol are held together by stronger intermolecular hydrogen bonds.



Hence, the boiling point of propanol is much higher than that of butane.

**Q. 5. Alcohols are comparatively more soluble in water than hydrocarbons of comparable molecular masses. Explain this fact.**


**Ans.** Alcohols can form H-bonds with water and break the H-bonds already existing between water molecules. Hence, they are soluble in water.



In contrast, hydrocarbons cannot form H-bonds with water and hence are insoluble in water.

**Q. 6. What is meant by hydroboration-oxidation reaction? Illustrate it with an example.**


**Ans.** The addition of diborane to alkenes to form trialkylboranes followed by their oxidation with alkaline hydrogen peroxide to form alcohols is called hydroboration-oxidation reaction.



The alcohols obtained by this process is contrary to the Markovnikov's rule.



**Q. 7. Give the structures and IUPAC names of monohydric phenols of molecular formula,  $\text{C}_7\text{H}_8\text{O}$ .**

**Ans.** The three isomers are given as follows:




**Q. 8. While separating a mixture of *ortho* and *para*-nitrophenols by steam distillation, name the isomer which will be steam volatile. Give reasons.**


**Ans.** *o*-Nitrophenol is steam volatile as it exists as discrete molecules due to intramolecular H-bonding and hence can be separated by steam distillation from *p*-nitrophenol which is less volatile as it exists as associated molecules because of intermolecular H-bonding.



**Q. 9. Give the equations of reactions for the preparation of phenol from cumene. [CBSE 2020 (56/3/2)]**





**Q. 10. Write chemical reaction for the preparation of phenol from chlorobenzene.**




**Q. 11. Write the mechanism of hydration of ethene to form ethanol. [CBSE 2019 (56/4/2), 2020 (56/5/1)]**


**Ans.** Direct addition of  $\text{H}_2\text{O}$  to ethene in presence of an acid does not occur. Ethene is first passed through conc.  $\text{H}_2\text{SO}_4$ , when ethyl hydrogen sulphate is formed.



Ethyl hydrogen sulphate when boiled with water undergoes hydrolysis to form ethanol.




**Q. 12. You are given benzene, conc.  $\text{H}_2\text{SO}_4$  and NaOH. Write the equations for the preparation of phenol using these reagents.**




**Q. 13. Show how will you synthesise:**

- 1-phenylethanol from a suitable alkene?
- cyclohexylmethanol using an alkyl halide by an  $\text{S}_{\text{N}}2$  reaction?
- pentan-1-ol using a suitable alkyl halide?


**Ans.** (i) Addition of  $\text{H}_2\text{O}$  to ethenylbenzene in presence of dil.  $\text{H}_2\text{SO}_4$  gives 1-phenylethanol.



(ii) Hydrolysis of cyclohexylmethyl bromide by aqueous NaOH gives cyclohexylmethanol.




(iii) Hydrolysis of 1-bromopentane by aqueous NaOH gives pentan-1-ol.



**Q. 14. Give two reactions that show the acidic nature of phenol. Compare acidity of phenol with that of ethanol.**

**Ans.** The reactions showing acidic nature of phenol are:


(i) **Reaction with sodium:** Phenol reacts with active metals like sodium to liberate H<sub>2</sub> gas.



(ii) **Reaction with NaOH:** Phenol dissolves in NaOH to form sodium phenoxide and water.




**Comparison of acidic character of phenol and ethanol:** Phenol is more acidic than ethanol. This is due to the reason that phenoxide ion left after the loss of a proton from phenol is stabilised by resonance while ethoxide ion left after loss of a proton from ethanol is not stabilised by resonance.




**Q. 15. Explain why is *ortho*-nitrophenol more acidic than *ortho*-methoxyphenol.**

[HOTS]

**Ans.** Due to strong –R and –I-effect of the NO<sub>2</sub> group, electron density in the O—H bond decreases and hence the loss of a proton becomes easy.



Moreover, after the loss of a proton, the *o*-nitrophenoxide ion left behind is stabilised by resonance.



This makes *o*-nitrophenol a stronger acid.

In contrast, due to +R effect, OCH<sub>3</sub> group increases the electron-density in the O—H bond thereby making the loss of proton difficult.



A small amount of *p*-hydroxybenzaldehyde is also formed.



**Q. 18. Explain the following with an example:**

- (i) Kolbe's reaction (ii) Reimer-Tiemann reaction  
(iii) Williamson ether synthesis (iv) Unsymmetrical ether

**Ans.**


- (i) Refer to Points to remember 15 (i).  
(ii) Refer to Points to remember 15 (ii).  
(iii) Refer to Points to remember 15(v).  
(iv) **Unsymmetrical ether:** If the alkyl or aryl groups attached to the oxygen atom are different, ethers are called unsymmetrical ethers. For example, ethyl methyl ether, methyl phenyl ether, etc.

**Q. 19. Write the mechanism of acid catalysed dehydration of ethanol to yield ethene.**


[CBSE 2019 (56/4/1)][CBSE 2023 (56/1/1)]

**Ans.** The mechanism of dehydration of ethanol involves the following steps:


**Step 1: Formation of protonated alcohol**



**Step 2: Formation of carbocation:** It is the slowest step and hence, the rate determining step of the reaction.



**Step 3: Formation of ethane by elimination of a proton**





The acid used in step 1 is released in step 3. To drive the equilibrium to the right, ethene is removed as it is formed.


**Q. 20. How are the following conversions carried out?**

- (i) Propene  $\longrightarrow$  Propan-2-ol  
(ii) Benzyl chloride  $\longrightarrow$  Benzyl alcohol  
(iii) Ethyl magnesium chloride  $\longrightarrow$  Propan-1-ol  
(iv) Methyl magnesium bromide  $\longrightarrow$  2-Methylpropan-2-ol

**Ans.**

- (i) Propan-2-ol can be prepared from propene by hydration as shown below:





**Q. 21. Name the reagents used in the following reactions:**

- Oxidation of primary alcohol to carboxylic acid
- Oxidation of a primary alcohol to aldehyde
- Bromination of phenol to 2, 4, 6-tribromophenol
- Benzyl alcohol to benzoic acid
- Dehydration of propan-2-ol to propene
- Butan-2-one to butan-2-ol


[CBSE (F) 2014]

**Ans.**

- Acidified potassium dichromate solution ( $\text{K}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4$ ) or alkaline potassium permanganate solution ( $\text{KMnO}_4/\text{OH}^-$ ) followed by hydrolysis with dil.  $\text{H}_2\text{SO}_4$ .
- Pyridinium chlorochromate (PCC),  $\text{C}_5\text{H}_5\text{NHCICrO}_3^+$  in  $\text{CH}_2\text{Cl}_2$
- Aqueous bromine ( $\text{Br}_2/\text{H}_2\text{O}$ )
- Acidified or alkaline potassium permanganate.
- Conc.  $\text{H}_2\text{SO}_4$  at 433–443 K.
- Lithium aluminium hydride ( $\text{LiAlH}_4$ ) or sodium borohydride ( $\text{NaBH}_4$ )



**Q. 22. Give reason for the higher boiling point of ethanol in comparison to methoxymethane.**

**Ans.** Ethanol undergoes intermolecular H-bonding due to the presence of a hydrogen attached to the electronegative oxygen atom. As a result, ethanol exists as associated molecules.



Consequently, a large amount of energy is required to break these hydrogen bonds. Therefore, the boiling point of ethanol is higher than that of methoxymethane which does not form H-bonds.

**Q. 23. Give the IUPAC names of the following ethers:**

- $\text{C}_2\text{H}_5\text{OCH}_2-\underset{\text{CH}_3}{\text{CH}}-\text{CH}_3$
- $\text{CH}_3\text{OCH}_2\text{CH}_2\text{Cl}$
- $\text{O}_2\text{N}-\text{C}_6\text{H}_4-\text{OCH}_3$  (p)
- $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_3$
- 
- 

**Ans.**

- 1-Ethoxy-2-methylpropane
- 2-Chloro-1-methoxyethane
- 4-Nitroanisole
- 1-Methoxypropane
- 4-Ethoxy-1, 1-dimethylcyclohexane
- Ethoxybenzene



COc1ccccc1.O=[N+]([O-])O.OS(=O)(=O)(O)O>>COc1ccccc1[N+](=O)[O-].COc1ccc([N+](=O)[O-])cc1

Anisole

conc. HNO<sub>3</sub>  
+ conc. H<sub>2</sub>SO<sub>4</sub>

*o*-Nitroanisole  
(Minor)



+

*p*-Nitroanisole  
(Major)





**Q. 30. Write mechanism of the reaction of HI with methoxymethane.**

**Ans.** With equimolar amounts of HI and methoxymethane, a mixture of methyl alcohol and methyl iodide are formed by the following mechanism:




If excess of HI is used, methyl alcohol formed in step II is also converted into methyl iodide by the following mechanism:




**Q. 31. Write equations of the following reactions:**

- (i) Friedel-Crafts reaction—alkylation of anisole
- (ii) Nitration of anisole
- (iii) Bromination of anisole in ethanoic acid medium
- (iv) Friedel-Crafts acetylation of anisole.


**Ans.** (i) Friedel-Crafts reaction (Alkylation):




(ii) **Nitration of anisole:** Anisole reacts with a mixture of concentrated sulphuric and nitric acids to yield a mixture of ortho and para nitroanisole.



(iii) **Halogenation:**



(iv) Friedel-Crafts acetylation of anisole:




**Q. 32.** Show how would you synthesise the following alcohols from appropriate alkenes?



[HOTS]





**Ans.** As addition and elimination are opposite to each other, the general strategy is to first dehydrate alcohol to give either a single alkene or a mixture of alkenes. If a mixture of alkenes is possible, then find out which of the alkenes will give the desired product.




Addition of  $\text{H}_2\text{O}$  to both these alkenes gives 1-methylcyclohexan-1-ol.



Addition of  $\text{H}_2\text{O}$  to 4-methylhept-3-ene in presence of an acid gives the desired alcohol.




Addition of  $\text{H}_2\text{O}$  to pent-1-ene gives pentan-2-ol.







Now, addition of  $\text{H}_2\text{O}$  to pent-2-ene gives a mixture of pentan-2-ol, pentan-3-ol.




Hence, the desired alkene is pent-1-ene and not pent-2-ene.



Now addition of  $\text{H}_2\text{O}$  to both 2-cyclohexylbut-2-ene and 2-cyclohexylidenobutane in presence of an acid gives 2-cyclohexylbutan-2-ol.



**Q. 33.** When 3-methylbutan-2-ol is treated with  $\text{HBr}$ , the following reaction takes place.




Give a mechanism for this reaction.


(Hint: The secondary carbocation formed in step II rearranges to a more stable tertiary carbocation by a hydride ion shift from 3rd carbon atom.)

**Ans.** **Mechanism:** The reaction takes place through the following mechanism:


**Step I:** Formation of protonated alcohol.




**Step II:** Formation of carbocation.



2° carbocation being less stable undergoes 1, 2 hydride shift to form more stable 3° carbocation.



**Step III:** Attack of nucleophile



## Multiple Choice Questions

Choose and write the correct option(s) in the following questions.

1. The C—O—H bond angle in alcohol is [CBSE 2022 (56/3/4)]

(a) slightly greater than  $109^{\circ}28'$ . (b) slightly less than  $109^{\circ}28'$ .  
(c) slightly greater than  $120^{\circ}$ . (d) slightly less than  $120^{\circ}$ .

2. How many alcohols with molecular formula  $C_4H_{10}O$  are chiral in nature? [NCERT Exemplar]

(a) 1 (b) 2 (c) 3 (d) 4

3. IUPAC name of *m*-cresol is \_\_\_\_\_. [NCERT Exemplar]

(a) 3-methylphenol (b) 3-chlorophenol (c) 3-methoxyphenol (d) benzene-1, 3-diol

4. The correct IUPAC name of  $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{C}-\text{CH}_2\text{CH}_3 \\ | \\ \text{OH} \end{array}$  is [CBSE 2020 (56/2/1)]

(a) *tert*-butyl alcohol (b) 2, 2-Dimethylpropanol  
(c) 2-Methylbutan-2-ol (d) 3-Methylbutan-3-ol

5. IUPAC name of the compound  $\begin{array}{c} \text{CH}_3-\text{CH}-\text{OCH}_3 \\ | \\ \text{CH}_3 \end{array}$  is \_\_\_\_\_. [NCERT Exemplar]

(a) 1-methoxy-1-methylethane (b) 2-methoxy-2-methylethane  
(c) 2-methoxypropane (d) isopropylmethyl ether





6. The C—O bond length in phenol is less than that in methanol due to

(a) partial double bond character of oxygen with aromatic ring  
(b)  $sp^2$  hybridised carbon to which oxygen attached  
(c)  $sp^3$  hybridised carbon  
(d) both (a) and (b)

7. Which of the following are not used to convert  $RCHO$  into  $RCH_2OH$ ?

(a)  $H_2/Pd$  (b)  $LiAlH_4$   
(c)  $NaBH_4$  (d) Reaction with  $RMgX$  followed by hydrolysis

8. Which of the following reactions will not yield phenol?



9. Which of the following alcohols will not undergo oxidation?


(a) Butanol (b) Butan-2-ol  
(c) 2-Methylbutan-2-ol (d) 3-Methylbutan-2-ol

10. Which of the following species can act as the strongest base?

(a)  $^{\ominus}OH$

(b)  $^{\ominus}OR$


(c)  $^{\ominus}OC_6H_5$




11. Phenol is less acidic than \_\_\_\_\_. [NCERT Exemplar]  
 (a) ethanol (b) *o*-nitrophenol (c) *o*-methylphenol (d) *o*-methoxyphenol
12. Which is the correct order of acid strength of the following?  
 (a)  $C_6H_5OH > H_2O > ROH > HC \equiv CH$  (b)  $C_6H_5OH > ROH > H_2O > HC \equiv CH$   
 (c)  $C_6H_5OH > HC \equiv CH > H_2O > ROH$  (d)  $C_6H_5OH > H_2O > HC \equiv CH > ROH$
13. Mark the correct order of decreasing acid strength of the following compounds. [NCERT Exemplar]



(i)




(ii)



(iii)






(iv)



(v)





- (a) (v) > (iv) > (ii) > (i) > (iii) (b) (ii) > (iv) > (i) > (iii) > (v)  
 (c) (iv) > (v) > (iii) > (ii) > (i) (d) (v) > (iv) > (iii) > (ii) > (i)
14. During dehydration of alcohols to alkenes by heating with concentrated  $H_2SO_4$ , the initiation step is: [CBSE Sample Paper 2022]  
 (a) protonation of alcohol molecule (b) formation of carbocation  
 (c) elimination of water (d) formation of an ester
15. Which of the following reactions are feasible? [CBSE 2023 (56/2/1)]  
 (a)  $CH_3CH_2Br + Na^+ O^-C(CH_3)_3 \longrightarrow CH_3CH_2-O-C(CH_3)_3$   
 (b)  $(CH_3)_3C-Cl + Na^+ O^-CH_2CH_3 \longrightarrow CH_3CH_2-O-C(CH_3)_3$   
 (c) Both (a) and (b)  
 (d) Neither (a) nor (b)
16. The synthesis of alkyl fluoride is best obtained from: [CBSE 2023 (56/1/1)]  
 (a) Free radicals (b) Swarts reaction  
 (c) Sandmeyer reaction (d) Finkelstein reaction
17. The compound that reacts fastest with Lucas reagent at room temperature is  
 (a) butan-1-ol (b) butan-2-ol  
 (c) 2-methylpropan-1-ol (d) 2-methylpropan-2-ol
18.  $CH_3CH_2OH$  can be converted into  $CH_3CHO$  by \_\_\_\_\_. [NCERT Exemplar]  
 (a) catalytic hydrogenation (b) treatment with  $LiAlH_4$   
 (c) treatment with pyridinium chlorochromate (d) treatment with  $KMnO_4$
19. An alcohol on oxidation is found to give  $CH_3COOH$  and  $CH_3CH_2COOH$ . The structure of the alcohol is:  
 (a)  $CH_3CH_2CH_2OH$  (b)  $(CH_3)_2C(OH)CH_2-CH_3$   
 (c)  $CH_3(CH_2)_3CH_2OH$  (d)  $CH_3CHOHCH_2CH_2CH_3$
20. Which of the following compounds will react with sodium hydroxide solution in water?



The electrophile involved in the above reaction is:



22. Monochlorination of toluene in sunlight followed by hydrolysis with *aq.* NaOH yields.



[NCERT Exemplar]

23. Which of the following compounds is oxidised to methyl ethyl ketone?

- (a) 2-Propanol (b) 1-Butanol (c) 2-Butanol (d) *tert.* Butyl alcohol

24. The reaction of  $\text{CH}_3-\text{C}(\text{CH}_3)_2-\text{Br}$  with  $\text{NaOCH}_3$  forms the product

- (a)  $\text{CH}_3-\text{O}-\text{C}(\text{CH}_3)_2-\text{Br}$  (b)  $\text{CH}_3-\text{C}(\text{CH}_3)=\text{CH}_2$  (c)  $\text{CH}_3-\text{C}(\text{CH}_3)_2-\text{ONa}$  (d)  $\text{CH}_3-\text{CH}(\text{CH}_3)-\text{CH}=\text{CH}_2$

25. Which of the following analogue is correct?

- (a) *o*-nitrophenol :  $pK_a$  value (7.2) :: *m*-nitrophenol :  $pK_a$  value (7.1)  
 (b) Benzene-1, 2-diol : Catechol :: Benzene-1, 4-diol : Quinol  
 (c)  $\text{C}_6\text{H}_5(\text{OH})\text{CHO}$  : Salicylaldehyde ::  $\text{C}_6\text{H}_5(\text{OH})\text{COO}^-\text{Na}^+$  : Salicylic acid  
 (d) *n*-pentane : B.P. (309.1 K) :: Ethoxy ethane : B.P. (390 K)

26. Williamson's synthesis of preparing dimethyl ether is an:

[CBSE Sample Paper 2022]

- (a)  $\text{S}_{\text{N}}1$  reaction (b) Elimination reaction  
 (c)  $\text{S}_{\text{N}}2$  reaction (d) Nucleophilic addition reaction

27. Match the species in column I with the corresponding boiling point in column II and choose the correct answer using the codes given below:

Column-I	Column-II
(i) $\text{CH}_3\text{CH}_2\text{OH}$	A. 373 K
(ii) $\text{CH}_3-\text{O}-\text{CH}_3$	B. 348 K
(iii) $\text{H}_2\text{O}$	C. 351 K
(iv) $\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_3$	D. 248 K

- (a) (i) — C, (ii) — D, (iii) — A, (iv) — B (b) (i) — A, (ii) — B, (iii) — C, (iv) — D  
 (c) (i) — B, (ii) — D, (iii) — A, (iv) — C (d) (i) — D, (ii) — B, (iii) — A, (iv) — C

## Answers

1. (b) 2. (a) 3. (a) 4. (c) 5. (c) 6. (d) 7. (d) 8. (d) 9. (c) 10. (b)  
 11. (b) 12. (a) 13. (b) 14. (a) 15. (a) 16. (b) 17. (d) 18. (c) 19. (d) 20. (a)  
 21. (d) 22. (d) 23. (c) 24. (b) 25. (b) 26. (b) 27. (a)


## Assertion-Reason Questions

In the following questions, two statements are given—one labeled Assertion (A) and the other labeled Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below:

- (a) Both Assertion (A) and Reason (R) are correct statements, and Reason (R) is the correct explanation of the Assertion (A).  
 (b) Both Assertion (A) and Reason (R) are correct statements, but Reason (R) is not the correct explanation of the Assertion (A).  
 (c) Assertion (A) is correct, but Reason (R) is incorrect statement.  
 (d) Assertion (A) is incorrect, but Reason (R) is correct statement.



1. Assertion (A) : IUPAC name of the compound



Reason (R) : In IUPAC nomenclature, ether is regarded as hydrocarbon derivative in which a hydrogen atom is replaced by —OR or —OAr group [where R = alkyl group and Ar = aryl group].

2. Assertion (A) : The boiling point of ethanol is higher than methoxymethane.

Reason (R) : There is intramolecular hydrogen bonding in ethanol.

3. Assertion (A) :  $\text{C}_2\text{H}_5\text{OH}$  is a weaker base than phenol but is a stronger nucleophile than phenol.

Reason (R) : In phenol the lone pair of electrons on oxygen is withdrawn towards the ring due to resonance. [CBSE Sample Paper 2020]

4. Assertion (A) : *o*-Nitrophenol is less soluble in water than the *m*- and *p*-isomers.

Reason (R) : *m*- and *p*- Nitrophenols exist as associated molecules.

5. Assertion (A) : *o*-nitrophenol is a weaker acid than *p*-nitrophenol.

Reason (R) : Intramolecular hydrogen bonding makes *ortho* isomer weaker than *para* isomer.

[CBSE 2020 (56/2/1)]

6. Assertion (A) : *p*-nitrophenol is more acidic than phenol.

Reason (R) : Nitro group helps in the stabilisation of the phenoxide ion by dispersal of negative charge due to resonance.

7. Assertion (A) : In Lucas test, 3° alcohols react immediately.

Reason (R) : An equimolar mixture of anhyd.  $\text{ZnCl}_2$  and conc.  $\text{HCl}$  is called Lucas reagent.

8. Assertion (A) : Phenol on oxidation with chromic acid gives benzoquinone.

Reason (R) : Pure phenol is colourless but turn pink due to oxidation to phenoquinone.

9. Assertion (A) : Phenol is more reactive than benzene towards electrophilic substitution reaction.

Reason (R) : In the case of phenol, the intermediate carbocation is more resonance stabilized.

10. Assertion (A) : Phenols give *o*- and *p*-nitrophenol on nitration with conc.  $\text{HNO}_3$  and  $\text{H}_2\text{SO}_4$  mixture.

Reason (R) : —OH group in phenol is *o*-, *p*-directing.

11. Assertion (A) : Like bromination of benzene, bromination of phenol is also carried out in the presence of Lewis acid.

Reason (R) : Lewis acid polarises the bromine molecule.

12. Assertion (A) : The reaction of phenol with acid chloride to form ether occurs in the presence of pyridine.

Reason (R) : Pyridine is a base so it neutralises the  $\text{HCl}$  formed and proceeds the reaction.

13. Assertion (A) : The C—O—H bond angle in alcohols is slightly less than the tetrahedral angle.

Reason (R) : This is due to the repulsive interaction between the two lone electron pairs on oxygen.

[CBSE 2020 (56/3/1)]

14. Assertion (A) : Methoxy ethane reacts with HI to give ethanol and iodomethane

Reason (R) : Reaction of ether with HI follows  $\text{S}_{\text{N}}2$  mechanism [CBSE Sample Paper 2021]

15. Assertion (A) :  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$  gives  $(\text{CH}_3)_3\text{C}-\text{I}$  and  $\text{CH}_3\text{OH}$  on treatment with HI.

Reason (R) : The reaction occurs by  $\text{S}_{\text{N}}1$  mechanism.

[CBSE 2020 (56/4/1)]

## Answers

- |         |         |         |         |         |        |        |        |        |         |
|---------|---------|---------|---------|---------|--------|--------|--------|--------|---------|
| 1. (d)  | 2. (c)  | 3. (d)  | 4. (b)  | 5. (a)  | 6. (a) | 7. (b) | 8. (b) | 9. (a) | 10. (d) |
| 11. (d) | 12. (a) | 13. (a) | 14. (a) | 15. (a) |        |        |        |        |         |


## Passage-based/Case-based/ Source-based Questions

Read the given passages and answer the questions that follow.

### PASSAGE-1

Alcohols and Phenols are acidic in nature. Electron withdrawing groups in phenol increase its acidic strength and electron donating groups decrease it. Alcohols undergo nucleophilic substitution with hydrogen halides to give alkyl halides. On oxidation primary alcohols yield aldehydes with mild oxidising agents and carboxylic acids with strong oxidising agents while secondary alcohols yield ketones. The presence of —OH groups in phenols activates the ring towards electrophilic substitution. Various important products are obtained from phenol like salicylaldehyde, salicylic acid, picric acid etc.

1. Give the structure of an alcohol which is resistant to oxidation?
2. Name any one group that increases the acidic character of phenol?
3. Consider the following reaction:




Identify X and Y.

OR


*p*-nitrophenol is a stronger acid than phenol while *p*-cresol is a weaker acid. Why?

### Answers



3° alcohols do not undergo oxidation because they do not have α-hydrogens.

2. NO<sub>2</sub> is electron withdrawing group, due to which it increases the stability of phenoxide ion.



OR

This is because —NO<sub>2</sub> decreases electron density on oxygen of O—H group making *p*-nitrophenol a stronger acid. While —CH<sub>3</sub> group increases electron density.

### PASSAGE-2

Williamson synthesis is one of the best methods for the preparation of ethers. It involves the treatment of an alkyl halide with a suitable sodium alkoxide. Williamson's synthesis involves nucleophilic substitution of the halogen atom of an alkyl halide by an alkoxide group as shown below:



When alkyl halide used in the reaction is primary, Williamson's synthesis proceeds via S<sub>N</sub>2 mechanism leading to the formation of an ether. This method is a versatile method for the synthesis of both symmetrical and unsymmetrical ethers.



1. Name the alkyl halide and sodium alkoxide used to synthesise tert-butyl ethyl ether.
2. What happens when benzyl ethyl ether reacts with HI?
3. Why is Williamson's synthesis not applicable when the alkyl halide used is tertiary?

OR

$(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$  on reaction with HI gives  $(\text{CH}_3)_3\text{C}-\text{I}$  and  $\text{CH}_3-\text{OH}$  as the main products and not  $(\text{CH}_3)_3\text{C}-\text{OH}$  and  $\text{CH}_3-\text{I}$ . Give reason.

## Answers

1. Ethyl bromide and sodium *tert*-butoxide.
2. Benzyl iodide and ethyl alcohol will be formed after the reaction of benzyl ethyl ether with HI.
3. This is because  $3^\circ$  alkyl halides are highly susceptible to dehydrohalogenation in the presence of sodium alkoxides and elimination reaction occurs.


OR

The reaction between  $(\text{CH}_3)_3\text{COCH}_3$  and HI follows  $\text{S}_\text{N}1$  mechanism. For an  $\text{S}_\text{N}1$  reaction, the formation of product is controlled by stability of the carbocation formed in the slowest step. Since *tert*-butyl carbonium ion  $(\text{CH}_3)_3\text{C}^+$  formed after the cleavage of C—O bond in the slowest step is more stable than methyl carbonium ion  $(\text{CH}_3)^+$  therefore  $(\text{CH}_3)_3\text{C}-\text{I}$  and  $\text{CH}_3\text{OH}$  are the main products.

## CONCEPTUAL QUESTIONS

Q. 1. Write the IUPAC name of the following compound:


[CBSE (AI) 2017]




Ans. 2-Bromo-3-methylbut-2-en-1-ol

Q. 2. Write the IUPAC name of the following compound:

[CBSE (AI) 2017]




Ans.



Q. 3. Write the IUPAC name of the following compound:


[CBSE Chennai 2015]



Ans. 1-methoxypropan-2-ol


Q. 4. How would you convert ethanol to ethene?

[CBSE (AI) 2011]



Q. 5. How will you synthesise the following alcohol from appropriate alkene:

[CBSE 2019 (56/4/1)]



**Q. 6. Why *ortho*-nitrophenol is steam volatile while *para*-nitrophenol is not?** [CBSE 2023 (56/2/1)]

**Ans.** Due to intramolecular H-bonding *o*-nitrophenol exists as a discrete molecule whereas due to intermolecular H-bonding *p*-nitrophenol exists as associated molecules. As a result of this *p*-nitrophenol has higher boiling point than *o*-nitrophenol.

**Q. 7. Of the two hydroxy organic compounds ROH and R'OH, the first one is basic and the other is acidic in behaviour. How is R different from R'?**

**Ans.** R is an alkyl group whereas R' is an aryl group.

**Q. 8. Arrange the following compounds in the increasing order of their acid strengths:**

4-nitrophenol, phenol, 2, 4, 6-trinitrophenol

[CBSE (F) 2013]

**Ans.** Phenol < 4-nitrophenol < 2,4,6-trinitrophenol

**Q. 9. Phenol is an acid but does not react with sodium bicarbonate solution. Why?**

[HOTS]

**Ans.** Phenol is a weaker acid than carbonic acid ( $\text{H}_2\text{CO}_3$ ) and hence does not liberate  $\text{CO}_2$  from sodium bicarbonate.

**Q. 10. Suggest a reagent for conversion of ethanol to ethanoic acid.**

[NCERT Exemplar]


**Ans.** (i)  $\text{CrO}_3 - \text{H}_2\text{SO}_4$   
(ii)  $\text{KMnO}_4/\text{OH}^-$   
(iii)  $\text{H}_3\text{O}^+$   
(iv) Acidified  $\text{KMnO}_4$  or  $\text{K}_2\text{Cr}_2\text{O}_7$

**Q. 11. How is toluene obtained from phenol?**

[CBSE Delhi 2013C]



**Q. 12. Suggest a reagent for the following conversion:**



[NCERT Exemplar]

**Q. 13. What is denatured alcohol?**

[NCERT Exemplar]

**Ans.** Alcohol is made unfit for drinking by mixing some copper sulphate and pyridine in it. This is called denatured alcohol.

**Q. 14. Explain why sodium metal can be used for drying diethyl ether but not ethyl alcohol.**

[HOTS]

**Ans.** Due to presence of an active hydrogen atom, ethyl alcohol reacts with sodium metal.



Diethyl ether, on the other hand, does not have replaceable hydrogen atom therefore does not react with sodium metal hence can be dried by metallic sodium.

**Q. 15. What is the order of dehydration of primary, secondary and tertiary alcohols?**

**Ans.** Tertiary alcohols > Secondary alcohols > Primary alcohols

**Q. 16. Name the reagents used in the following reactions:**

[CBSE (F) 2014]

**Friedel-Crafts alkylation of anisole**

**Ans.** Anhydrous aluminium chloride (anhyd.  $\text{AlCl}_3$ )



Phenol is stronger acid than alcohol because the phenoxide ion left-after the release of a proton is stabilised by resonance but alkoxide ion is not, moreover, the +I effect of alkyl group destabilizes the alkoxide ion by intensifying negative charge on oxygen atom.

- (ii) In alcohols with increase of branching in carbon chain surface area decreases which results in decrease in vander Waals forces and hence decrease in boiling point.

**Q. 5. Account for the following:**

- (i) Phenol does not react with  $\text{NaHCO}_3$  whereas carboxylic acids react.  
 (ii) Phenol is more easily nitrated than benzene.

[HOTS]



Such acid-base reaction takes place only if the acid formed is weaker than the reacting acid. In other words, phenol is not a strong enough acid to react with  $\text{NaHCO}_3$ .

- (ii) Nitration involves attack of electrophile nitronium ion ( $\text{NO}_2^+$ ) on benzene ring. Due to +R effect of  $-\text{OH}$  group electron density on benzene increases. Therefore, phenol is more easily nitrated as compared to benzene.

**Q. 6. Answer the following questions:**

- (i) Dipole moment of phenol is smaller than that of methanol. Why?  
 (ii) In Kolbe's reaction, instead of phenol, phenoxide ion is treated with carbon dioxide. Why?

[NCERT Exemplar]


- Ans.** (i) In phenol,  $\text{C}-\text{O}$  bond is less polar due to electron-withdrawing effect of benzene ring whereas in methanol,  $\text{C}-\text{O}$  bond is more polar due to electron-releasing effect of  $-\text{CH}_3$  group.  
 (ii) Phenoxide ion is more reactive than phenol towards electrophilic aromatic substitution and hence undergoes electrophilic substitution with carbon dioxide which is a weak electrophile.

**Q. 7. Write the mechanism of the following reaction:**



[CBSE Delhi 2016, 2019 (56/4/3), 2020 (56/3/2)] [HOTS]


- Ans.** The formation of ether is a nucleophilic bimolecular reaction ( $\text{S}_{\text{N}}2$ ) involving the attack of alcohol molecule on protonated alcohol as shown below:



**Q. 8. For the pair phenol and cyclohexanol, answer the following:**

- (i) Why is phenol more acidic than cyclohexanol?  
 (ii) Give one chemical test to distinguish between the two.

[CBSE 2023 (56/5/2)]






Due to higher electronegativity of  $sp^2$  hybridised carbon of phenol to which  $-\text{OH}$  groups is attached, electron density decreases on oxygen.

This increases the polarity of  $\text{O}-\text{H}$  bond and results in an increase in ionisation of phenols than that of cyclohexanol where  $-\text{OH}$  group is attached to  $sp^3$  hybridised carbon atom.

- (ii) Phenol gives a violet colouration with  $\text{FeCl}_3$  solution while cyclohexanol does not.




**Q. 9. What happens when**

(i) Phenol reacts with Conc.  $\text{HNO}_3$ ?

(ii) Ethyl chloride reacts with  $\text{NaOC}_2\text{H}_5$ ?

Write the chemical equations involved in the above reactions.


[CBSE 2019 (56/4/3)]



**Q. 10. How do you convert**

(i) phenol to toluene

(ii) formaldehyde to ethanol ? [CBSE North 2016]



## Short Answer Questions-II


Each of the following questions are of 3 marks.

**Q. 1. Explain the mechanism of the following reactions:**


- Addition of Grignard's reagent to the carbonyl group of a compound forming an adduct followed by hydrolysis.
- Acid catalysed dehydration of an alcohol forming an alkene.
- Acid catalysed hydration of an alkene forming an alcohol.

[CBSE 2020 (56/1/2)]


**Ans.** (i) **Step I:** Nucleophilic addition of Grignard reagent to carbonyl group.



**Step II:** Hydrolysis




(ii) Refer to NCERT Exercises, Q. 19.




**Mechanism:**


**Step I:** Protonation of alkene to form carbocation by electrophilic attack of  $\text{H}_3\text{O}^+$ .



**Step II:** Nucleophilic attack of water on carbocation.



**Step III:** Deprotonation to form an alcohol






**Q. 2. How do you convert the following:**

(i) Phenol to anisole

(ii) Propan-2-ol to 2-methylpropan-2-ol

(iii) Aniline to phenol

[CBSE Delhi 2015]






Q. 3. Arrange the following compounds in decreasing order of acidity.

[NCERT Exemplar]



Ans. (i)  $\text{H}_2\text{O} > \text{ROH} > \text{HC} \equiv \text{CH}$



Q. 4. Give one chemical test to distinguish between the following:

[CBSE 2019 (56/5/1)]

(i) Phenol and propan-1-ol (ii) Ethanol and dimethyl ether

(iii) propan-1-ol and 2-Methyl-propan-2-ol

Ans. (i) Add neutral  $\text{FeCl}_3$  to both the compounds. Phenol gives violet colouration while, 1 - propanol does not. 1



(ii) Add  $\text{I}_2/\text{NaOH(aq)}$  to both the compounds, Ethanol gives yellow precipitate while ether does not 1

(iii) Add  $\text{HCl}$  and  $\text{ZnCl}_2$  to both the compounds, 2-methyl-2-propanol gives turbidity immediately while 1-propanol does not 1


(Or any other suitable chemical test)

[CBSE Marking Scheme 2019 (56/5/1)]

Q. 5. An organic compound 'A' having molecular formula  $\text{C}_3\text{H}_6$  on treatment with aqueous  $\text{H}_2\text{SO}_4$  gives 'B' which on treatment with  $\text{HCl}/\text{ZnCl}_2$  gives 'C'. The compound C on treatment with ethanolic  $\text{KOH}$  gives back the compound 'A'. Identify the compounds A, B, C. [HOTS]



Q. 6. A compound 'A' is optically active. On mild oxidation, it gives a compound 'B' but on vigorous oxidation gives another compound 'C'. C along with D is also formed from B by reaction with iodine and alkali. Deduce the structures of A, B, C, D. [HOTS]



**Q. 7. Account for the following:**







- (i) Rectified spirit cannot be converted into absolute alcohol by simple distillation.
- (ii) Diethyl ether does not react with sodium.
- (iii) Phenols do not undergo substitution of the  $\text{—OH}$  group like alcohols.

**Ans.** (i) Rectified spirit containing 95% ethyl alcohol and 5% water forms an azeotropic mixture which distils at a constant temperature of 351.13 K.

- (ii) Since diethyl ether does not contain an active hydrogen attached to oxygen like alcohols and phenols, it does not react with sodium.
- (iii) The C—O bond in phenols has some double bond character due to resonance and hence cannot be easily cleaved by nucleophile. In contrast, the C—O bond in alcohols is a pure single bond and hence can be easily cleaved by nucleophile.

**Q. 8. Write the main product(s) in each of the following reactions:**



[CBSE Delhi 2016]




**Q. 9. How will you bring the following conversions?**

- (i) Phenol to benzyl alcohol                      (ii) Phenol to *m*-bromophenol  
(iii) Phenol to aspirin.

[HOTS]





Q. 10. (i) Why is the C—O bond length in phenols less than that in methanol?

(ii) Arrange the following in order of increasing boiling point:

Ethoxyethane, Butanal, Butanol, *n*-butane

(iii) How can phenol be prepared from anisole? Give reaction.

[CBSE 2023 (56/5/2)]


Ans.

(i) This is due to

— partial double bond character on account of the conjugation of unshared electron pairs of oxygen with the aromatic ring and

—  $sp^2$  hybridised state of carbon to which oxygen is attached.

(ii) *n*-Butane < ethoxyethane < butanal < butanol




Q. 11. (i) Write the mechanism of the following reaction:






(ii) Write the equation involved in the acetylation of Salicylic acid.

[CBSE Delhi 2015]



Ans.




Q. 12. Give the major products that are formed by heating each of the following ethers with HI.



[HOTS]



Q. 13. The following is not an appropriate reaction for the preparation of *tert*-butyl ethyl ether:




- What would be the major product of the given reaction?
- Write a suitable reaction for the preparation of *tert*-butyl ethyl ether, specifying the names of reagents used.


Justify your answer in both cases.

[CBSE Sample Paper 2016] [HOTS]

- Ans. (i) The major product of the given reaction is 2-methylprop-1-ene. It is because sodium ethoxide ( $\text{CH}_3\text{CH}_2\text{O}^-\text{Na}^+$ ) is a strong nucleophile as well as a strong base. Thus, elimination reaction predominates over substitution reaction.



- (ii) To prepare tertiary butyl ether, the alkyl halide should be primary, i.e., ethyl chloride and the nucleophile should be tertiary, i.e., tertiary butoxide ion. It is because the reaction occurs by  $\text{S}_\text{N}2$  mechanism and primary alkyl halides are most reactive in  $\text{S}_\text{N}2$  reactions.




Q. 14. How do you convert the following: (Any three)

(i) Phenol to picric acid




(ii) Propanone to 2-Methylpropan-2-ol

(iii) Phenol to anisole

(iv) Propene to Propan-1-ol [CBSE 2023 (56/1/1)]









(Any three)

Q. 15. Write the structures of the main products in the following reactions:




[CBSE 2018]


Ans.




Q. 16. Write the products of the following reactions:



Ans.






**Q. 17. What happens when**

- (i)  $(\text{CH}_3)_3\text{C}-\text{OH}$  is treated with Cu at 573 K,
- (ii) Anisole is treated with  $\text{CH}_3\text{Cl}/\text{anhydrous AlCl}_3$ ,
- (iii) Phenol is treated with Zn dust?

Write chemical equations in support of your answer.

[CBSE (F) 2017]




**Q. 18. What happens when**

- (i) Salicylic acid is treated with  $(\text{CH}_3\text{CO})_2\text{O}/\text{H}^+$ ?
- (ii) Phenol is oxidised with  $\text{Na}_2\text{Cr}_2\text{O}_7/\text{H}^+$ ?
- (iii) Anisole is treated with  $\text{CH}_3\text{Cl}/\text{anhydrous AlCl}_3$ ?

Write chemical equation in support of your answer.

[CBSE 2019 (56/3/1)]



## Long Answer Questions

Each of the following questions are of 5 marks.


Q. 1. (i) Write the formula of reagents used in the following reactions:

- (a) Bromination of phenol to 2, 4, 6-tribromophenol  
 (b) Hydroboration of propene and then oxidation to propanol.

(ii) Arrange the following compound groups in the increasing order of their property indicated:

- (a) *p*-nitrophenol, ethanol, phenol (acidic character)  
 (b) Propanol, propane, propanal (boiling point)

(iii) Write the mechanism (using curved arrow notation) of the following reaction:




Ans. (i) (a)  $\text{Br}_2(\text{aq})$

(b)  $\text{B}_2\text{H}_6$ ,  $\text{H}_2\text{O}$ ,  $\text{H}_2\text{O}_2$  and  $\text{OH}^-$

(ii) (a) Ethanol < Phenol < *p*-nitrophenol

(b) Propane < Propanal < Propanol







Q. 2. (i) How can you obtain phenol from (a) Cumene, (b) Benzene sulphonic acid, (c) Benzene diazonium chloride?

(ii) Write the structure of the major product obtained from dinitration of 3-methylphenol.

(iii) Write the reaction involved in Kolbe's reaction.


[CBSE 2019 (56/2/1)]

Ans.




[CBSE Marking Scheme 2019 (56/2/1)]

- Q. 3. An aromatic compound 'A' on treatment with  $\text{CHCl}_3/\text{KOH}$  gives two compounds 'B' and 'C'. Both B and C give the same product 'D' when distilled with zinc dust. Oxidation of D gives E having molecular formula  $\text{C}_7\text{H}_6\text{O}_2$ . The sodium salt of E on heating with sodalime gives F which may also be obtained by distilling A with zinc dust. Identify A to F. [HOTS]




Reactions involved:



- Q. 4. (i) Give equations of the following reactions:
- Phenol is treated with conc.  $\text{HNO}_3$ .
  - Propene is treated with  $\text{B}_2\text{H}_6$  followed by  $\text{H}_2\text{O}_2/\text{OH}^-$ .
  - Sodium *t*-butoxide is treated with  $\text{CH}_3\text{Cl}$ .
- (ii) How will you distinguish between butan-1-ol and butan-2-ol?
- (iii) Arrange the following in increasing order of acidity:


Phenol, ethanol, water

[CBSE 2019 (56/2/1)]





- (ii) Butan-2-ol when warmed with  $I_2$  in NaOH (NaOI) gives yellow precipitate of iodoform while butan-1-ol does not respond to this test.



- (iii) Ethanol < water < phenol

Q. 5. (i) What happens when



- (a) phenol reacts with bromine water?  
 (b) ethanol reacts with  $\text{CH}_3\text{COCl}$ /pyridine?  
 (c) anisole reacts with HI?

Write the chemical equations involved in the above reactions.


- (ii) Distinguish between:

- (a) Ethanol and phenol  
 (b) Propan-2-ol and 2-methylpropan-2-ol


[CBSE (F) 2016]



- (ii) (a) Phenol gives violet colouration with  $\text{FeCl}_3$  solution but ethanol does not.



- (b) Propan-2-ol when warmed with  $I_2$  in NaOH gives yellow precipitate of iodoform while 2-methylpropan-2-ol does not respond to this test.




Q. 6. (i) Write equations of the following reactions:



- (a) Bromine in  $\text{CS}_2$  with phenol  
 (b) Treating phenol with chloroform in the presence of aq. NaOH  
 (c) Anisole reacts with HI

- (ii) Distinguish between


- (a) Ethanol and Diethyl ether (b) Propanol and *t*-butyl alcohol

[CBSE South 2016]






- (ii) (a) Ethanol when warmed with  $I_2$  in NaOH gives yellow precipitate of iodoform while diethyl ether does not.



- (b) *tert*-butyl alcohol on treatment with the Lucas reagent immediately gives turbidity while propanol does not produce turbidity at room temperature.



## Questions for Practice

Choose and write the correct answer for each of the following.

1. The order of reactivity of the alkenes, (i)  $(CH_3)_2C=CH_2$ , (ii)  $CH_3CH=CH_2$ , (iii)  $CH_2=CH_2$ , when subjected to acid catalysed hydration is:

- (a) (i) > (iii) > (ii)                      (b) (i) > (ii) > (iii)  
(c) (ii) > (i) > (iii)                      (d) (iii) > (ii) > (i)

2. Alcohols are soluble in water because

- (a) alcohol forms ionic bond with water.  
(b) alcohol forms hydrogen bond with water.  
(c) alcohol forms covalent bond with water.  
(d) alcohol forms interstitial compound with water.

3. Mark the correct increasing order of reactivity of the following compounds with HBr/HCl.



(i)




(ii)



(iii)


- (a) (i) < (ii) < (iii)                      (b) (ii) < (i) < (iii)  
(c) (ii) < (iii) < (i)                      (d) (iii) < (ii) < (i)

4. In the reaction given below, X is:



- (a) 2-methylpent-2-ene                      (b) 2-methylpentane  
(c) 2-methylbut-2-ene                      (d) neopentane

5. What is the correct order of reactivity of alcohols in the following reaction?



[NCERT Exemplar]

(a)  $1^\circ > 2^\circ > 3^\circ$

(b)  $1^\circ < 2^\circ < 3^\circ$

(c)  $3^\circ > 2^\circ > 1^\circ$

(d)  $3^\circ > 1^\circ > 2^\circ$

In the following questions, two statements are given—one labeled Assertion (A) and the other labeled Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below:

(a) Both Assertion (A) and Reason (R) are correct statements, and Reason (R) is the correct explanation of the Assertion (A).

(b) Both Assertion (A) and Reason (R) are correct statements, but Reason (R) is not the correct explanation of the Assertion (A).

(c) Assertion (A) is correct, but Reason (R) is incorrect statement.

(d) Assertion (A) is incorrect, but Reason (R) is correct statement.

6. Assertion (A) : Phenol is more reactive than benzene towards electrophilic substitution reaction.

Reason (R) : In the case of phenol, the intermediate carbocation is more resonance stabilized.

7. Assertion (A) : Addition reaction of water to but-1-ene in acidic medium yields butan-2-ol.

Reason (R) : Addition of water in acidic medium proceeds through the formation of primary carbocation.

8. Assertion (A) : Ethers behave as bases in the presence of mineral acids.

Reason (R) : Due to the presence of lone pairs of electrons on oxygen.

9. Assertion (A) : The bond angles in alcohols is slightly less than  $109^\circ 28'$ .

Reason (R) : Lone pair – lone pair repulsion decreases the bond angles.

10. Assertion (A) : Alkyl aryl ethers are cleaved at the alkyl-oxygen bond.

Reason (R) : More stable aryl oxygen bond responsible for this.

Answer the following questions:

11. (i) While separating a mixture of ortho and para-nitrophenols by steam distillation, name the isomer which will be steam volatile. Give reason.

(ii) Give a chemical test to distinguish between ethyl alcohol and methyl alcohol.

12. Write the chemical equation involved in the following reactions: [CBSE 2023 (56/2/2)]


(i) Reimer-Tiemann reaction

(ii) Acetylation of Salicylic acid

13. (i) Arrange the following compounds in the increasing order of their acid strength:

*p*-cresol, *p*-nitrophenol, phenol

(ii) Write the mechanism (using curved arrow notation) of the following reaction:



[CBSE (AI) 2017]

14. Give reasons for the following:

(i) Anisole reacts with HI to give phenol and methyl iodide and not iodobenzene and methyl alcohol.

(ii) *o*-Nitrophenol is less soluble in water than *p*-nitrophenol.



15. Write the structures of the products when Butan-2-ol reacts with the following:

(i)  $\text{CrO}_3$


(ii)  $\text{SOCl}_2$

[CBSE (AI) 2017]

16. Alcohols react both as nucleophiles as well as electrophiles. Write one reaction of each type and describe its mechanism.
17. Which of the following is an appropriate set of reactant for the preparation of 1-methoxy-4-nitrobenzene and why? [CBSE 2023 (56/4/2)]




18. (i) What happens when phenol reacts with  
 (a) Conc. HNO<sub>3</sub>, and  
 (b) CHCl<sub>3</sub> in presence of aqueous NaOH followed by acidification?
- (ii) Why does the reaction of CH<sub>3</sub>ONa with (CH<sub>3</sub>)<sub>3</sub>C—Br give 2-methylpropene and not (CH<sub>3</sub>)<sub>3</sub>C—OCH<sub>3</sub>? [CBSE 2023 (56/4/2)]
19. (i) Write the mechanism of the following reaction: [CBSE 2023 (56/2/1)]



- (ii) Why *ortho*-nitrophenol is steam volatile while *para*-nitrophenol is not.
20. A compound 'A' having molecular formula C<sub>4</sub>H<sub>10</sub>O is found to be soluble in concentrated sulphuric acid. It does not react with sodium metal or potassium permanganate. On heating with excess of HI, it gives a single alkyl halide. Deduce the structure of compound A and explain all the reactions. [HOTS]
21. How would you obtain the following?  
 (i) Benzoquinone from phenol [CBSE 2020 (56/1/2)]  
 (ii) 2-Methylpropan-2-ol from methylmagnesium bromide  
 (iii) Propan-2-ol from propene [CBSE (AI) 2011, (F) 2011]

22. Answer the following questions:

- (i) Why are ethers insoluble in water?  
 (ii) Complete the reaction equation:





- (iii) How will you know whether a given OH group is alcoholic or phenolic in nature?
23. (i) Arrange the following sets of compounds in order of their increasing boiling points:  
 (a) Pentan-1-ol, butan-1-ol, butan-2-ol, ethanol, propan-1-ol, methanol  
 (b) Pentan-1-ol, *n*-butane, pentanal, ethoxyethane.
- (ii) Arrange the following compounds in increasing order of acidity and give a suitable explanation.  
 Phenol, *o*-nitrophenol, *o*-cresol [NCERT Exemplar]



24. Name the reagents which are used in the following conversions:

- (i) A primary alcohol to aldehyde
- (ii) Butan-2-one to butan-2-ol
- (iii) Benzyl alcohol to benzoic acid

25. (i) Write the product(s) in the following reactions:





(ii) Give simple chemical tests to distinguish between the following pairs of compounds:


- (a) Ethanol and Phenol
- (b) Pentan-2-ol and pentan-3-ol

26. (i) Write mechanism of the reaction of HI with methoxymethane.

(ii) Identify A and B in the following reactions:



(iii) Give the structure and the IUPAC name of the major product obtained in the following reaction:



[CBSE Sample Paper 2015]

## Answers

1. (b)    2. (b)    3. (c)    4. (c)    5. (c)    6. (a)    7. (c)    8. (a)    9. (a)    10. (a)

