


7. p-Block Elements

Group 15 elements:




- The valence shell electronic configuration is $ns^2 np^3$.
- Nitrogen differs in chemical properties from other elements of the group due to its small size, high electronegativity, high ionisation enthalpy and non-availability of d -orbitals.
- They exhibit two oxidation states, +3 and +5. Heavier elements exhibit mainly +3 oxidation state due to inert pair effect.

The main use of nitrogen is in the manufacture of ammonia


Ammonia

- On a small scale, ammonia is obtained from ammonium salts, which decompose when treated with caustic soda or lime. It forms metal salt, water, and ammonia gas.
- Ammonia can also be prepared by treating metal nitrides with warm water.
- It has trigonal pyramidal structure with nitrogen atom at the apex.





- **Forms**
 - Dry ammonia gas (gaseous ammonia)
 - Liquid ammonia (liquified ammonia)
 - Liquor ammonia fortis (saturated solution of ammonia in water)
 - Laboratory bench reagent (dilute solution of liquor ammonia)
- On large scale, ammonia is obtained by **Haber's process**.
 - Raw material: Mixture of hydrogen and nitrogen gases in the ratio 3:1
 - Pressure: 200 atm to 900 atm pressure
 - Temperature: 450 – 500°C
 - Catalyst: Finely divided iron
 - Promoter: molybdenum or Al_2O_3
- **Properties:**
 - It is a colourless non-poisonous gas with a characteristic pungent odour.
 - It is lighter than air and extremely soluble in water because of hydrogen bonding.



- It can be liquefied when cooled to 10°C under pressure of 6 atm. It forms white crystals on cooling.
- It has basic nature because of the presence of lone pair of electrons.
- It acts as a reducing agent.
- Inhaling this gas causes irritation to the eyes and respiratory system.
- **Uses:**
 - Due to high dielectric constant, ammonia is a good solvent for ionic compounds.
 - It is used as a cleaning agent for removing grease in dry cleaning.
 - It is used in the manufacturing of artificial silk.
 - It is used as laboratory reagent.
- Nitric acid (HNO_3)



1. Preparation: Ostwald's process




- **Detection of the presence of nitrate:**
(Brown ring test)



- Phosphorus exists as P_4 in elemental form.
- **Allotropic forms of phosphorus:**
 1. White phosphorus
 2. Red phosphorus
 3. Black phosphorus (α -block phosphorus and β -block phosphorus)

- Phosphorus forms two types of halides, PX_3 ($\text{X} = \text{F}, \text{Cl}, \text{Br}, \text{I}$) and PX_5 ($\text{X} = \text{F}, \text{Cl}, \text{Br}$).
- The structure of PCl_5 is trigonal bipyramidal



- Phosphorus forms a number of oxoacids such as ortho-phosphoric acid (H_3PO_4), ortho-phosphorus acid (H_3PO_3), hypo-phosphorus acid (H_3PO_2).
- The oxoacids containing P – H bond are strong reducing agents.


Group 16 elements: (known as chalcogens)

Oxygen
Sulphur
Selenium
Tellurium
Polonium

- The valence shell electronic configuration is $ns^2 np^4$.

Like nitrogen, oxygen differs from other elements of the group due to its small size and high electronegativity

1. Preparation:



1. Three stable isotopes – ^{16}O , ^{17}O , ^{18}O

Uses

- In normal respiration and combustion
- As an oxidant (in liquid state) for propelling rockets
- In oxyacetylene welding
- In the manufacture of many metals (particularly steel)
- Oxygen cylinders are used in hospitals, high altitude flying and mountaineering.

1.

Acidic oxides – Combine with water to give an acid

Example – SO_2 , Cl_2O_7 , CO_2 , N_2O_5

Basic oxides – Combine with water to give bases

Examples – Na_2O , CaO , BaO

Amphoteric oxides – Show the characteristics of both acidic as well as basic oxides

React with both acids and alkalies

Example – Al_2O_3

Neutral oxides – Neither acidic nor basic

Examples – CO, NO, N₂O

Ozone (O₃) is an allotropic form of oxygen. It is a powerful oxidising agent.

- **Sulphur –**

Allotropic forms of sulphur:

1. Rhombic sulphur (α – sulphur)
2. Monoclinic sulphur (β – sulphur)

Both rhombic and monoclinic sulphur exist as S₈ molecules.

Oxides of sulphur – SO₂, SO₃

Sulphuric Acid

- Concentrated sulphuric acid is known as oil of vitriol. It occurs in free state in hot water of sulphur springs. In combined state, it occurs as mineral sulphates.
- Sulphuric acid is prepared by contact process. It involves burning of a pure and dry mixture of two parts of sulphur or sulphide ores and one part of air in the presence of vanadium pentoxide or platinised asbestos as catalyst.
- Chemical reactions of H₂SO₄ are because of its
 1. low volatility
 2. strong acidic character
 3. strong affinity for water
 4. ability to act as an oxidising agent
- Dilute sulphuric acid reacts with active metals, metal oxides, metal hydroxides, metal carbonates, metal sulphites to form their respective metal sulphates and acid sulphates.
- Because of low volatility, it can be used for the manufacture of more volatile acids from their corresponding salts.
- It is a strong dehydrating agent. Because of its strong affinity for water, sulphuric acid removes water from hydrated salts and organic compounds.
- Concentrated sulphuric acid is a moderately strong oxidising agent and can oxidise both metals and non-metals.

Fluorine
Chlorine
Bromine
Iodine
Astatine

- The valence shell electronic configuration is $ns^2 np^5$.
- They have very high electronegativity.
- The common oxidation state is -1 . However, $+1$, $+3$, $+5$ and $+7$ oxidation states are also exhibited.
- Fluorine show anomalous properties in the group due to its very small size.

- Chlorine has an atomic number 17 and an atomic mass of 35.5 u.
- It does not occur in free state as it is highly reactive in nature.


Manufacture of chlorine:

- **Laboratory methods of preparation of chlorine**


1. By the oxidation of conc. HCl and manganese dioxide (MnO_2)



2. By the action of HCl on $KMnO_4$



- **Deacon's process:**



Physical properties of chlorine:

- It is a greenish yellow gas.
- It has a pungent smell.
- It has a slight sour taste.
- It is fairly soluble in water.
- It is 2.5 times heavier than air.
- It is poisonous in nature. When inhaled, it causes severe headache accompanied by cough and breathlessness.

Chemical properties of chlorine:

- Chlorine gas is non-combustible.
- Chlorine reacts with water to form hypochlorous acid.
- It reacts with burning sodium to form sodium chloride.
- When white phosphorus comes in contact with chlorine, it melts and spontaneously catches fire to form dense white fumes.
- It has strong affinity for hydrogen.

- It reacts with slaked lime to give bleaching powder.
- HOCl releases nascent oxygen, which is responsible for oxidising and bleaching action.
- Bleaching effect of Cl₂ is permanent. It bleaches vegetable or organic matter in the presence of moisture.

Hydrogen Chloride

- In laboratory, hydrogen chloride gas is prepared by heating sodium chloride with concentrated sulphuric acid.
- It is also prepared by burning hydrogen gas in the atmosphere of chlorine gas or by exposing hydrogen gas and chlorine gas to diffused sunlight.
- It is colourless and pungent-smelling with sour taste and a very irritating odour.
- It is extremely soluble in water.
- Hydrogen chloride is neither combustible nor does it support combustion.
- On heating at above 500°C, it dissociates into hydrogen and chlorine.
- On mixing with ammonia gas, it forms dense white fumes due to formation of ammonium chloride.
- Aqueous solution of hydrogen chloride is called **hydrochloric acid**.
- It is prepared by dissolving hydrogen chloride in water.
- It reacts with metals to form respective chlorides and hydrogen gas.
- **Aqua regia** is a mixture of 3 parts of concentrated hydrochloric acid and 1 part of concentrated nitric acid. It is a very corrosive acid and is the only known acid that can dissolve gold.
- Halogens form a number of oxoacids.

Halic (I) acid (Hypohalous acid)	HO_F (Hypo-fluorous acid)	HOCl (Hypochlorous acid)	HOBr (Hypobromous acid)	HOI (Hypoiodous acid)
Halic (III) acid (Halous acid)	— —	HOCIO (Chlorous acid)	— —	— —
Halic (V) acid (Halic acid)	— —	HOCIO ₂ (Chloric acid)	HOBrO ₂ (Bromic acid)	HOIO ₂ (Iodic acid)
Halic (VII) acid (Perhalic acid)	— —	HOCIO ₃ (Perchloric acid)	HOBrO ₃ (Perbromic acid)	HOIO ₃ (Periodic acid)

•

- Halogens form a number of inter-halogen compounds (compounds formed by two different halogens).


Type	Formula	Structure
XX'_3	ClF_3	Bent T-shaped
	BrF_3	Bent T-shaped
	IF_3	Bent T-shaped
	ICl_3	Bent T-shaped
XX'_5	IF_5	Square pyramidal
	BrF_5	Square pyramidal
	ClF_5	Square pyramidal
XX'_7	IF_7	Pentagonal bipyramidal

Helium
Neon
Argon
Krypton
Xenon
Radon

- The valence shell electronic configuration is $ns^2 np^6$. (Exception: Helium $\rightarrow 1s^2$)
- **Physical Properties**
 - Monoatomic
 - Colourless, odourless, and tasteless
 - Sparingly soluble in water
 - Low melting and boiling points.
- **Xenon-Fluorine Compounds**



(Xe in excess)



(1:5 ratio)



(1:20 ratio)

Structure

- $\text{XeF}_2 \longrightarrow$ Linear
- $\text{XeF}_4 \longrightarrow$ Square planar
- $\text{XeF}_6 \longrightarrow$ Distorted octahedral

Xenon-Oxygen Compounds

XeO_3 has a pyramidal

XeOF_4 has a square pyramidal