

# Principle of Mathematical Induction

## Short Answer Type Questions

**Q. 1** Give an example of a statement  $P(n)$  which is true for all  $n \geq 4$  but  $P(1)$ ,  $P(2)$  and  $P(3)$  are not true. Justify your answer.

**Sol.** Let the statement  $P(n)$ :  $3n < n!$

For  $n = 1$ ,  $3 \times 1 < 1!$  [false]

For  $n = 2$ ,  $3 \times 2 < 2!$   $\Rightarrow 6 < 2$  [false]

For  $n = 3$ ,  $3 \times 3 < 3!$   $\Rightarrow 9 < 6$  [false]

For  $n = 4$ ,  $3 \times 4 < 4!$   $\Rightarrow 12 < 24$  [true]

For  $n = 5$ ,  $3 \times 5 < 5!$   $\Rightarrow 15 < 5 \times 4 \times 3 \times 2 \times 1 \Rightarrow 15 < 120$  [true]

**Q. 2** Give an example of a statement  $P(n)$  which is true for all  $n$ . Justify your answer.

**Sol.** Consider the statement

$$P(n): 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\text{For } n = 1, \quad 1 = \frac{1(1+1)(2 \times 1 + 1)}{6}$$

$$\Rightarrow \quad 1 = \frac{2(3)}{6}$$

$$\Rightarrow \quad 1 = 1$$

$$\text{For } n = 2, \quad 1 + 2^2 = \frac{2(2+1)(4+1)}{6}$$

$$\Rightarrow \quad 5 = \frac{30}{6} \Rightarrow 5 = 5$$

$$\text{For } n = 3, \quad 1 + 2^2 + 3^2 = \frac{3(3+1)(7)}{6}$$

$$\Rightarrow \quad 1 + 4 + 9 = \frac{3 \times 4 \times 7}{6}$$

$$\Rightarrow \quad 14 = 14$$

Hence, the given statement is true for all  $n$ .

Prove each of the statements in the following questions from by the Principle of Mathematical Induction.

**Q. 3**  $4^n - 1$  is divisible by 3, for each natural number  $n$ .

**Thinking Process**

In step I put  $n=1$ , the obtained result should be a divisible by 3. In step II put  $n=k$  and take  $P(k)$  equal to multiple of 3 with non-zero constant say  $q$ . In step III put  $n=k+1$ , in the statement and solve till it becomes a multiple of 3.

**Sol.** Let  $P(n) : 4^n - 1$  is divisible by 3 for each natural number  $n$ .

Step I Now, we observe that  $P(1)$  is true.

$$P(1) = 4^1 - 1 = 3$$

It is clear that 3 is divisible by 3.

Hence,  $P(1)$  is true.

Step II Assume that,  $P(n)$  is true for  $n = k$

$P(k) : 4^k - 1$  is divisible by 3

$$x4^k - 1 = 3q$$

Step III Now, to prove that  $P(k+1)$  is true.

$$\begin{aligned} P(k+1) : 4^{k+1} - 1 &= 4^k \cdot 4 - 1 \\ &= 4^k \cdot 3 + 4^k - 1 \\ &= 3 \cdot 4^k + 3q & [\because 4^k - 1 = 3q] \\ &= 3(4^k + q) \end{aligned}$$

Thus,  $P(k+1)$  is true whenever  $P(k)$  is true.

Hence, by the principle of mathematical induction  $P(n)$  is true for all natural number  $n$ .

**Q. 4**  $2^{3n} - 1$  is divisible by 7, for all natural numbers  $n$ .

**Sol.** Let  $P(n) : 2^{3n} - 1$  is divisible by 7

Step I We observe that  $P(1)$  is true.

$$P(1) : 2^{3 \times 1} - 1 = 2^3 - 1 = 8 - 1 = 7$$

It is clear that  $P(1)$  is true.

Step II Now, assume that  $P(n)$  is true for  $n = k$ ,

$P(k) : 2^{3k} - 1$  is divisible by 7.

$$\Rightarrow 2^{3k} - 1 = 7q$$

Step III Now, to prove  $P(k+1)$  is true.

$$\begin{aligned} P(k+1) : 2^{3(k+1)} - 1 &= 2^{3k} \cdot 2^3 - 1 \\ &= 2^{3k} \cdot 8 - 1 \\ &= 2^{3k} (7 + 1) - 1 \\ &= 7 \cdot 2^{3k} + 2^{3k} - 1 \\ &= 7 \cdot 2^{3k} + 7q & [\text{from step II}] \\ &= 7(2^{3k} + q) \end{aligned}$$

Hence,  $P(k+1) :$  is true whenever  $P(k)$  is true.

So, by the principle of mathematical induction  $P(n)$  is true for all natural number  $n$ .

**Q. 5**  $n^3 - 7n + 3$  is divisible by 3, for all natural numbers  $n$ .

**Sol.** Let  $P(n) : n^3 - 7n + 3$  is divisible by 3, for all natural number  $n$ .

**Step I** We observe that  $P(1)$  is true.

$$\begin{aligned} P(1) &= (1)^3 - 7(1) + 3 \\ &= 1 - 7 + 3 \\ &= -3, \text{ which is divisible by 3.} \end{aligned}$$

Hence,  $P(1)$  is true.

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$\therefore P(k) = k^3 - 7k + 3 = 3q$$

**Step III** To prove  $P(k + 1)$  is true

$$\begin{aligned} P(k + 1) &: (k + 1)^3 - 7(k + 1) + 3 \\ &= k^3 + 1 + 3k(k + 1) - 7k - 7 + 3 \\ &= k^3 - 7k + 3 + 3k(k + 1) - 6 \\ &= 3q + 3[k(k + 1) - 2] \end{aligned}$$

Hence,  $P(k + 1)$  is true whenever  $P(k)$  is true.

[from step II]

So, by the principle of mathematical induction  $P(n) :$  is true for all natural number  $n$ .

**Q. 6**  $3^{2n} - 1$  is divisible by 8, for all natural numbers  $n$ .

**Sol.** Let  $P(n) : 3^{2n} - 1$  is divisible by 8, for all natural numbers.

**Step I** We observe that  $P(1)$  is true.

$$\begin{aligned} P(1) : 3^{2(1)} - 1 &= 3^2 - 1 \\ &= 9 - 1 = 8, \text{ which is divisible by 8.} \end{aligned}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : 3^{2k} - 1 = 8q$$

**Step III** Now, to prove  $P(k + 1)$  is true.

$$\begin{aligned} P(k + 1) &: 3^{2(k + 1)} - 1 \\ &= 3^{2k} \cdot 3^2 - 1 \\ &= 3^{2k} \cdot (8 + 1) - 1 \\ &= 8 \cdot 3^{2k} + 3^{2k} - 1 \\ &= 8 \cdot 3^{2k} + 8q \\ &= 8(3^{2k} + q) \end{aligned}$$

[from step II]

Hence,  $P(k + 1)$  is true whenever  $P(k)$  is true.

So, by the principle of mathematical induction  $P(n)$  is true for all natural numbers  $n$ .

**Q. 7** For any natural numbers  $n$ ,  $7^n - 2^n$  is divisible by 5.

**Sol.** Consider the given statement is

$P(n) : 7^n - 2^n$  is divisible by 5, for any natural number  $n$ .

**Step I** We observe that  $P(1)$  is true.

$$P(1) = 7^1 - 2^1 = 5, \text{ which is divisible by 5.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) = 7^k - 2^k = 5q$$

**Step III** Now, to prove  $P(k + 1)$  is true,

$$\begin{aligned} P(k + 1) &: 7^{k + 1} - 2^{k + 1} \\ &= 7^k \cdot 7 - 2^k \cdot 2 \end{aligned}$$

$$\begin{aligned}
 &= 7^k \cdot (5 + 2) - 2^k \cdot 2 \\
 &= 7^k \cdot 5 + 2 \cdot 7^k - 2^k \cdot 2 \\
 &= 5 \cdot 7^k + 2(7^k - 2^k) \\
 &= 5 \cdot 7^k + 2(5q) \\
 &= 5(7^k + 2q), \text{ which is divisible by 5.} \quad [\text{from step II}]
 \end{aligned}$$

So,  $P(k + 1)$  is true whenever  $P(k)$  is true.

Hence, by the principle of mathematical induction  $P(n)$  is true for any natural number  $n$ .

**Q. 8** For any natural numbers  $n$ ,  $x^n - y^n$  is divisible by  $x - y$ , where  $x$  and  $y$  are any integers with  $x \neq y$ .

**Sol.** Let  $P(n) : x^n - y^n$  is divisible by  $x - y$ , where  $x$  and  $y$  are any integers with  $x \neq y$ .

**Step I** We observe that  $P(1)$  is true.

$$P(1) : x^1 - y^1 = x - y$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : x^k - y^k \text{ is divisible by } (x - y).$$

$\therefore$

$$x^k - y^k = q(x - y)$$

**Step III** Now, to prove  $P(k + 1)$  is true.

$$\begin{aligned}
 P(k + 1) : x^{k+1} - y^{k+1} \\
 &= x^k \cdot x - y^k \cdot y \\
 &= x^k \cdot x - x^k \cdot y + x^k \cdot y - y^k \cdot y \\
 &= x^k(x - y) + y(x^k - y^k) \\
 &= x^k(x - y) + yq(x - y) \\
 &= (x - y)[x^k + yq], \text{ which is divisible by } (x - y). \quad [\text{from step II}]
 \end{aligned}$$

Hence,  $P(k + 1)$  is true whenever  $P(k)$  is true. So, by the principle of mathematical induction  $P(n)$  is true for any natural number  $n$ .

**Q. 9**  $n^3 - n$  is divisible by 6, for each natural number  $n \geq 2$ .

### Thinking Process

In step I put  $n=2$ , the obtained result should be divisible by 6. Then, follow the same process as in question no. 4.

**Sol.** Let  $P(n) : n^3 - n$  is divisible by 6, for each natural number  $n \geq 2$ .

**Step I** We observe that  $P(2)$  is true.  $P(2) : (2)^3 - 2$

$$\Rightarrow 8 - 2 = 6, \text{ which is divisible by 6.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : k^3 - k \text{ is divisible by 6.}$$

$\therefore$

$$k^3 - k = 6q$$

**Step III** To prove  $P(k + 1)$  is true

$$\begin{aligned}
 P(k + 1) : (k + 1)^3 - (k + 1) \\
 &= k^3 + 1 + 3k(k + 1) - (k + 1) \\
 &= k^3 + 1 + 3k^2 + 3k - k - 1 \\
 &= k^3 - k + 3k^2 + 3k \\
 &= 6q + 3k(k + 1) \quad [\text{from step II}]
 \end{aligned}$$

We know that,  $3k(k + 1)$  is divisible by 6 for each natural number  $n = k$ .

So,  $P(k + 1)$  is true. Hence, by the principle of mathematical induction  $P(n)$  is true.

**Q. 10**  $n(n^2 + 5)$  is divisible by 6, for each natural number  $n$ .

**Sol.** Let  $P(n) : n(n^2 + 5)$  is divisible by 6, for each natural number  $n$ .

**Step I** We observe that  $P(1)$  is true.

$$P(1) : 1(1^2 + 5) = 6, \text{ which is divisible by 6.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : k(k^2 + 5) \text{ is divisible by 6.}$$

$\therefore$

$$k(k^2 + 5) = 6q$$

**Step III** Now, to prove  $P(k + 1)$  is true, we have

$$\begin{aligned} P(k + 1) &: (k + 1)[(k + 1)^2 + 5] \\ &= (k + 1)[k^2 + 2k + 1 + 5] \\ &= (k + 1)[k^2 + 2k + 6] \\ &= k^3 + 2k^2 + 6k + k^2 + 2k + 6 \\ &= k^3 + 3k^2 + 8k + 6 \\ &= k^3 + 5k + 3k^2 + 3k + 6 \\ &= k(k^2 + 5) + 3(k^2 + k + 2) \\ &= (6q) + 3(k^2 + k + 2) \end{aligned}$$

We know that,  $k^2 + k + 2$  is divisible by 2, where,  $k$  is even or odd.

Since,  $P(k + 1) : 6q + 3(k^2 + k + 2)$  is divisible by 6. So,  $P(k + 1)$  is true whenever  $P(k)$  is true.

Hence, by the principle of mathematical induction  $P(n)$  is true.

**Q. 11**  $n^2 < 2^n$ , for all natural numbers  $n \geq 5$ .

**Sol.** Consider the given statement

$$P(n) : n^2 < 2^n \text{ for all natural numbers } n \geq 5.$$

**Step I** We observe that  $P(5)$  is true

$$\begin{aligned} P(5) &: 5^2 < 2^5 \\ &= 25 < 32 \end{aligned}$$

Hence,  $P(5)$  is true.

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) = k^2 < 2^k \text{ is true.}$$

**Step III** Now, to prove  $P(k + 1)$  is true, we have to show that

$$P(k + 1) : (k + 1)^2 < 2^{k+1}$$

Now,

$$\begin{aligned} k^2 < 2^k &= k^2 + 2k + 1 < 2^k + 2k + 1 \\ &= (k + 1)^2 < 2^k + 2k + 1 \end{aligned} \quad \dots(i)$$

Now,  $(2k + 1) < 2^k$

$$\begin{aligned} &= 2^k + 2k + 1 < 2^k + 2^k \\ &= 2^k + 2k + 1 < 2 \cdot 2^k \\ &= 2^k + 2k + 1 < 2^{k+1} \end{aligned} \quad \dots(ii)$$

From Eqs. (i) and (ii), we get  $(k + 1)^2 < 2^{k+1}$

So,  $P(k + 1)$  is true, whenever  $P(k)$  is true. Hence, by the principle of mathematical induction  $P(n)$  is true for all natural numbers  $n \geq 5$ .

**Q. 12**  $2n < (n + 2)!$  for all natural numbers  $n$ .

**Sol.** Consider the statement

$P(n) : 2n < (n + 2)!$  for all natural number  $n$ .

**Step I** We observe that,  $P(1)$  is true.  $P(1) : 2(1) < (1 + 2)!$

$$\Rightarrow 2 < 3! \Rightarrow 2 < 3 \times 2 \times 1 \Rightarrow 2 < 6$$

Hence,  $P(1)$  is true.

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ ,

$P(k) : 2k < (k + 2)!$  is true.

**Step III** To prove  $P(k + 1)$  is true, we have to show that

$$P(k + 1) : 2(k + 1) < (k + 1 + 2)!$$

Now,

$$2k < (k + 2)!$$

$$2k + 2 < (k + 2)! + 2$$

$$2(k + 1) < (k + 2)! + 2 \quad \dots(i)$$

Also,

$$(k + 2)! + 2 < (k + 3)! \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$2(k + 1) < (k + 1 + 2)!$$

So,  $P(k + 1)$  is true, whenever  $P(k)$  is true.

Hence, by principle of mathematical induction  $P(n)$  is true.

**Q. 13**  $\sqrt{n} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{n}}$ , for all natural numbers  $n \geq 2$ .

**Sol.** Consider the statement

$P(n) : \sqrt{n} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{n}}$ , for all natural numbers  $n \geq 2$ .

**Step I** We observe that  $P(2)$  is true.

$$P(2) : \sqrt{2} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}}, \text{ which is true.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : \sqrt{k} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{k}} \text{ is true.}$$

**Step III** To prove  $P(k + 1)$  is true, we have to show that

$$P(k + 1) : \sqrt{k + 1} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{k + 1}} \text{ is true.}$$

Given that,

$$\sqrt{k} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{k}}$$

$$\Rightarrow \sqrt{k} + \frac{1}{\sqrt{k + 1}} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{k}} + \frac{1}{\sqrt{k + 1}}$$

$$\Rightarrow \frac{(\sqrt{k})(\sqrt{k + 1}) + 1}{\sqrt{k + 1}} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{k}} + \frac{1}{\sqrt{k + 1}} \quad \dots(i)$$

$$\text{If } \sqrt{k + 1} < \frac{\sqrt{k}\sqrt{k + 1} + 1}{\sqrt{k + 1}}$$

$$\Rightarrow k + 1 < \sqrt{k}\sqrt{k + 1} + 1$$

$$\Rightarrow k < \sqrt{k(k + 1)} \Rightarrow \sqrt{k} < \sqrt{k + 1} \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$\sqrt{k + 1} < \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{k + 1}}$$

So,  $P(k + 1)$  is true, whenever  $P(k)$  is true. Hence,  $P(n)$  is true.

**Q. 14**  $2 + 4 + 6 + \dots + 2n = n^2 + n$ , for all natural numbers  $n$ .

**Sol.** Let  $P(n) : 2 + 4 + 6 + \dots + 2n = n^2 + n$

For all natural numbers  $n$ .

**Step I** We observe that  $P(1)$  is true.

$$P(1) : 2 = 1^2 + 1$$

$$2 = 2 \text{ which is true.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$\therefore P(k) : 2 + 4 + 6 + \dots + 2k = k^2 + k$$

**Step III** To prove that  $P(k + 1)$  is true.

$$\begin{aligned} P(k + 1) : 2 + 4 + 6 + 8 + \dots + 2k + 2(k + 1) \\ &= k^2 + k + 2(k + 1) \\ &= k^2 + k + 2k + 2 \\ &= k^2 + 2k + 1 + k + 1 \\ &= (k + 1)^2 + k + 1 \end{aligned}$$

So,  $P(k + 1)$  is true, whenever  $P(k)$  is true.

Hence,  $P(n)$  is true.

**Q. 15**  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$  for all natural numbers  $n$ .

**Sol.** Consider the given statement

$$P(n) : 1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1, \text{ for all natural numbers } n$$

**Step I** We observe that  $P(0)$  is true.

$$P(1) : 1 = 2^{0+1} - 1$$

$$1 = 2^1 - 1$$

$$1 = 2 - 1$$

$$1 = 1, \text{ which is true.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$\text{So, } P(k) : 1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1 \text{ is true.}$$

**Step III** Now, to prove  $P(k + 1)$  is true.

$$\begin{aligned} P(k + 1) : 1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} \\ &= 2^{k+1} - 1 + 2^{k+1} \\ &= 2 \cdot 2^{k+1} - 1 \\ &= 2^{k+2} - 1 \\ &= 2^{(k+1)+1} - 1 \end{aligned}$$

So,  $P(k + 1)$  is true, whenever  $P(k)$  is true.

Hence,  $P(n)$  is true.

**Q. 16**  $1 + 5 + 9 + \dots + (4n - 3) = n(2n - 1)$ , for all natural numbers  $n$ .

**Sol.** Let  $P(n) : 1 + 5 + 9 + \dots + (4n - 3) = n(2n - 1)$ , for all natural numbers  $n$ .

**Step I** We observe that  $P(1)$  is true.

$$P(1) : 1 = 1(2 \times 1 - 1), 1 = 2 - 1 \text{ and } 1 = 1, \text{ which is true.}$$

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$\text{So, } P(k) : 1 + 5 + 9 + \dots + (4k - 3) = k(2k - 1) \text{ is true.}$$

**Step III** Now, to prove  $P(k+1)$  is true.

$$\begin{aligned}
 P(k+1) &: 1 + 5 + 9 + \dots + (4k-3) + 4(k+1) - 3 \\
 &= k(2k-1) + 4(k+1) - 3 \\
 &= 2k^2 - k + 4k + 4 - 3 \\
 &= 2k^2 + 3k + 1 \\
 &= 2k^2 + 2k + k + 1 \\
 &= 2k(k+1) + 1(k+1) \\
 &= (k+1)(2k+1) \\
 &= (k+1)[2k+1+1-1] \\
 &= (k+1)[2(k+1)-1]
 \end{aligned}$$

So,  $P(k+1)$  is true, whenever  $p(k)$  is true, hence  $P(n)$  is true.

## Long Answer Type Questions

Use the Principle of Mathematical Induction in the following questions.

**Q. 17** A sequence  $a_1, a_2, a_3, \dots$  is defined by letting  $a_1 = 3$  and  $a_k = 7a_{k-1}$ , for all natural numbers  $k \geq 2$ . Show that  $a_n = 3 \cdot 7^{n-1}$  for all natural numbers.

**Sol.** A sequence  $a_1, a_2, a_3, \dots$  is defined by letting  $a_1 = 3$  and  $a_k = 7a_{k-1}$ , for all natural numbers  $k \geq 2$ .

Let  $P(n) : a_n = 3 \cdot 7^{n-1}$  for all natural numbers.

**Step I** We observe  $P(2)$  is true.

For  $n=2$ ,  $a_2 = 3 \cdot 7^{2-1} = 3 \cdot 7^1 = 21$  is true.

As  $a_1 = 3, a_k = 7a_{k-1}$

$\Rightarrow a_2 = 7 \cdot a_{2-1} = 7 \cdot a_1$

$\Rightarrow a_2 = 7 \times 3 = 21$  [ $\because a_1 = 3$ ]

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : a_k = 3 \cdot 7^{k-1}$$

**Step III** Now, to prove  $P(k+1)$  is true, we have to show that

$$P(k+1) : a_{k+1} = 3 \cdot 7^{k+1-1}$$

$$\begin{aligned}
 a_{k+1} &= 7 \cdot a_{k+1-1} = 7 \cdot a_k \\
 &= 7 \cdot 3 \cdot 7^{k-1} = 3 \cdot 7^{k-1+1}
 \end{aligned}$$

So,  $P(k+1)$  is true, whenever  $p(k)$  is true. Hence,  $P(n)$  is true.

**Q. 18** A sequence  $b_0, b_1, b_2, \dots$  is defined by letting  $b_0 = 5$  and  $b_k = 4 + b_{k-1}$ , for all natural numbers  $k$ . Show that  $b_n = 5 + 4n$ , for all natural number  $n$  using mathematical induction.

**Sol.** Consider the given statement,

$P(n) : b_n = 5 + 4n$ , for all natural numbers given that  $b_0 = 5$  and  $b_k = 4 + b_{k-1}$

**Step I**  $P(1)$  is true.

$$P(1) : b_1 = 5 + 4 \times 1 = 9$$



As  $b_0 = 5, b_1 = 4 + b_0 = 4 + 5 = 9$

Hence,  $P(1)$  is true.

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : b_k = 5 + 4k$$

**Step III** Now, to prove  $P(k + 1)$  is true, we have to show that

$$\therefore P(k + 1) : b_{k+1} = 5 + 4(k + 1)$$

$$b_{k+1} = 4 + b_{k+1-1}$$

$$= 4 + b_k$$

$$= 4 + 5 + 4k = 5 + 4(k + 1)$$

So, by the mathematical induction  $P(k + 1)$  is true whenever  $P(k)$  is true, hence  $P(n)$  is true.

**Q. 19** A sequence  $d_1, d_2, d_3, \dots$  is defined by letting  $d_1 = 2$  and  $d_k = \frac{d_{k-1}}{k}$ , for all natural numbers,  $k \geq 2$ . Show that  $d_n = \frac{2}{n!}$ , for all  $n \in N$ .

**Sol.** Let  $P(n) : d_n = \frac{2}{n!}, \forall n \in N$ , to prove  $P(2)$  is true.

$$\text{Step I} \quad P(2) : d_2 = \frac{2}{2!} = \frac{2}{2 \times 1} = 1$$

As, given

$$d_1 = 2$$

$$\Rightarrow d_k = \frac{d_{k-1}}{k}$$

$$\Rightarrow d_2 = \frac{d_1}{2} = \frac{2}{2} = 1$$

Hence,  $P(2)$  is true.

**Step II** Now, assume that  $P(k)$  is true.

$$P(k) : d_k = \frac{2}{k!}$$

**Step III** Now, to prove that  $P(k + 1)$  is true, we have to show that  $P(k + 1) : d_{k+1} = \frac{2}{(k + 1)!}$

$$\begin{aligned} d_{k+1} &= \frac{d_{k+1-1}}{k} = \frac{d_k}{k} \\ &= \frac{2}{k!k} = \frac{2}{(k + 1)!} \end{aligned}$$

So,  $P(k + 1)$  is true. Hence,  $P(n)$  is true.

**Q. 20** Prove that for all  $n \in N$

$$\cos \alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos[\alpha + (n - 1)\beta]$$

$$= \frac{\cos\left[\alpha + \left(\frac{n-1}{2}\right)\beta\right] \sin\left(\frac{n\beta}{2}\right)}{\sin\frac{\beta}{2}}$$

### Thinking Process

To prove this, use the formula  $2 \cos A \sin B = \sin(A + B) - \sin(A - B)$  and

$$\sin A - \sin B = 2 \cos\left(\frac{A+B}{2}\right) \cdot \sin\left(\frac{A-B}{2}\right)$$

**Sol.** Let  $P(n) : \cos\alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos[\alpha + (n-1)\beta]$

$$= \frac{\cos\left[\alpha + \left(\frac{n-1}{2}\right)\beta\right] \sin\left(\frac{n\beta}{2}\right)}{\sin\frac{\beta}{2}}$$

**Step I** We observe that  $P(1)$

$$P(1) : \cos\alpha = \frac{\cos\left[\alpha + \left(\frac{1-1}{2}\right)\beta\right] \sin\frac{\beta}{2}}{\sin\frac{\beta}{2}} = \frac{\cos(\alpha + 0) \sin\frac{\beta}{2}}{\sin\frac{\beta}{2}}$$

$$\cos\alpha = \cos\alpha$$

Hence,  $P(1)$  is true.

**Step II** Now, assume that  $P(n)$  is true for  $n = k$ .

$$P(k) : \cos\alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos[\alpha + (k-1)\beta]$$

$$= \frac{\cos\left[\alpha + \left(\frac{k-1}{2}\right)\beta\right] \sin\frac{k\beta}{2}}{\sin\frac{\beta}{2}}$$

**Step III** Now, to prove  $P(k+1)$  is true, we have to show that

$$P(k+1) : \cos\alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos[\alpha + (k-1)\beta] + \cos(\alpha + k\beta) = \frac{\cos\left(\alpha + \frac{k\beta}{2}\right) \sin(k+1)\frac{\beta}{2}}{\sin\frac{\beta}{2}}$$

$$\begin{aligned} \text{LHS} &= \cos\alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos[\alpha + (k-1)\beta] + \cos(\alpha + k\beta) \\ &= \frac{\cos\left[\alpha + \left(\frac{k-1}{2}\right)\beta\right] \sin\frac{k\beta}{2}}{\sin\frac{\beta}{2}} + \cos(\alpha + k\beta) \\ &= \frac{\cos\left[\alpha + \left(\frac{k-1}{2}\right)\beta\right] \sin\frac{k\beta}{2} + \cos(\alpha + k\beta) \sin\frac{\beta}{2}}{\sin\frac{\beta}{2}} \\ &= \frac{\sin\left(\alpha + \frac{k\beta}{2} - \frac{\beta}{2} + \frac{k\beta}{2}\right) - \sin\left(\alpha + \frac{k\beta}{2} - \frac{\beta}{2} - \frac{k\beta}{2}\right) + \sin\left(\alpha + k\beta + \frac{\beta}{2}\right) - \sin\left(\alpha + k\beta - \frac{\beta}{2}\right)}{2\sin\frac{\beta}{2}} \\ &= \frac{\sin\left(\alpha + k\beta + \frac{\beta}{2}\right) - \sin\left(\alpha - \frac{\beta}{2}\right)}{2\sin\frac{\beta}{2}} \\ &= \frac{2\cos\frac{1}{2}\left(\alpha + \frac{\beta}{2} + k\beta + \alpha - \frac{\beta}{2}\right) \sin\frac{1}{2}\left(\alpha + \frac{\beta}{2} + k\beta - \alpha + \frac{\beta}{2}\right)}{2\sin\frac{\beta}{2}} \\ &= \frac{\cos\left(\frac{2\alpha + k\beta}{2}\right) \sin\left(\frac{k\beta + \beta}{2}\right)}{\sin\frac{\beta}{2}} = \frac{\cos\left(\alpha + \frac{k\beta}{2}\right) \sin(k+1)\frac{\beta}{2}}{\sin\frac{\beta}{2}} = \text{RHS} \end{aligned}$$

So,  $P(k+1)$  is true. Hence,  $P(n)$  is true.

**Q. 21** Prove that  $\cos \theta \cos 2\theta \cos 2^2\theta \dots \cos 2^{n-1}\theta = \frac{\sin 2^n \theta}{2^n \sin \theta}, \forall n \in N.$

**Sol.** Let  $P(n) : \cos \theta \cos 2\theta \dots \cos 2^{n-1}\theta = \frac{\sin 2^n \theta}{2^n \sin \theta}$

$$\begin{aligned} \text{Step I For } n = 1, P(1) : \cos \theta &= \frac{\sin 2^1 \theta}{2^1 \sin \theta} \\ &= \frac{\sin 2\theta}{2 \sin \theta} = \frac{2 \sin \theta \cos \theta}{2 \sin \theta} = \cos \theta \end{aligned}$$

which is true.

Step II Assume that  $P(n)$  is true, for  $n = k$ .

$$P(k) : \cos \theta \cdot \cos 2\theta \cdot \cos 2^2\theta \dots \cos 2^{k-1}\theta = \frac{\sin 2^k \theta}{2^k \sin \theta} \text{ is true.}$$

Step III To prove  $P(k+1)$  is true.

$$\begin{aligned} P(k+1) : \cos \theta \cdot \cos 2\theta \cdot \cos 2^2\theta \dots \cos 2^{k-1}\theta \cdot \cos 2^k\theta \\ &= \frac{\sin 2^k \theta}{2^k \sin \theta} \cdot \cos 2^k\theta \\ &= \frac{2 \sin 2^k\theta \cdot \cos 2^k\theta}{2 \cdot 2^k \sin \theta} \\ &= \frac{\sin 2 \cdot 2^k\theta}{2^{k+1} \sin \theta} = \frac{\sin 2^{(k+1)}\theta}{2^{k+1} \sin \theta} \end{aligned}$$

which is true.

So,  $P(k+1)$  is true. Hence,  $P(n)$  is true.

**Q. 22** Prove that,  $\sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta = \frac{\sin n\theta \sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}},$   
for all  $n \in N.$

**Thinking Process**

To use the formula of  $2 \sin A \sin B = \cos(A-B) - \cos(A+B)$  and

$$\cos A - \cos B = 2 \sin \frac{A+B}{2} \cdot \sin \frac{B-A}{2} \text{ also } \cos(-\theta) = \cos \theta.$$

**Sol.** Consider the given statement

$$\begin{aligned} P(n) : \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta \\ &= \frac{\sin \frac{n\theta}{2} \sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}}, \text{ for all } n \in N \end{aligned}$$

Step I We observe that  $P(1)$  is

$$P(1) : \sin \theta = \frac{\sin \frac{\theta}{2} \cdot \sin \frac{(1+1)\theta}{2}}{\sin \frac{\theta}{2}} = \frac{\sin \frac{\theta}{2} \cdot \sin \theta}{\sin \frac{\theta}{2}}$$

$$\sin \theta = \sin \theta$$

Hence,  $P(1)$  is true.

**Step II** Assume that  $P(n)$  is true, for  $n = k$ .

$$P(k) : \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin k\theta \\ = \frac{\sin \frac{k\theta}{2} \sin \left( \frac{k+1}{2} \theta \right)}{\sin \frac{\theta}{2}} \text{ is true.}$$

**Step III** Now, to prove  $P(k+1)$  is true.

$$P(k+1) : \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin k\theta + \sin (k+1)\theta \\ = \frac{\sin \frac{(k+1)\theta}{2} \sin \left( \frac{k+1+1}{2} \theta \right)}{\sin \frac{\theta}{2}}$$

$$\begin{aligned} \text{LHS} &= \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin k\theta + \sin (k+1)\theta \\ &= \frac{\sin \frac{k\theta}{2} \sin \left( \frac{k+1}{2} \theta \right)}{\sin \frac{\theta}{2}} + \sin (k+1)\theta = \frac{\sin \frac{k\theta}{2} \sin \left( \frac{k+1}{2} \theta \right) + \sin (k+1)\theta \cdot \sin \frac{\theta}{2}}{\sin \frac{\theta}{2}} \\ &= \frac{\cos \left[ \frac{k\theta}{2} - \left( \frac{k+1}{2} \theta \right) \right] - \cos \left[ \frac{k\theta}{2} + \left( \frac{k+1}{2} \theta \right) \right] + \cos \left[ (k+1)\theta - \frac{\theta}{2} \right] - \cos \left[ (k+1)\theta + \frac{\theta}{2} \right]}{2 \sin \frac{\theta}{2}} \\ &= \frac{\cos \frac{\theta}{2} - \cos \left( k\theta + \frac{\theta}{2} \right) + \cos \left( k\theta + \frac{\theta}{2} \right) - \cos \left( k\theta + \frac{3\theta}{2} \right)}{2 \sin \frac{\theta}{2}} \\ &= \frac{\cos \frac{\theta}{2} - \cos \left( k\theta + \frac{3\theta}{2} \right)}{2 \sin \frac{\theta}{2}} = \frac{2 \sin \frac{1}{2} \left( \frac{\theta}{2} + k\theta + \frac{3\theta}{2} \right) \cdot \sin \frac{1}{2} \left( k\theta + \frac{3\theta}{2} - \frac{\theta}{2} \right)}{2 \sin \frac{\theta}{2}} \\ &= \frac{\sin \left( \frac{k\theta + 2\theta}{2} \right) \cdot \sin \left( \frac{k\theta + \theta}{2} \right)}{\sin \frac{\theta}{2}} = \frac{\sin (k+1)\theta \cdot \sin (k+1)\theta}{\sin \frac{\theta}{2}} \end{aligned}$$

So,  $P(k+1)$  is true, whenever  $P(k)$  is true. Hence,  $P(n)$  is true.

**Q. 23** Show that  $\frac{n^5}{5} + \frac{n^3}{3} + \frac{7n}{15}$  is a natural number, for all  $n \in \mathbb{N}$ .

### Thinking Process

Here, use the formula  $(a+b)^5 = a^5 + 5ab^4 + 10a^2b^3 + 10a^3b^2 + 5a^4b + b^5$

and  $(a+b)^3 = a^3 + b^3 + 3ab(a+b)$

**Sol.** Consider the given statement

$$P(n) : \frac{n^5}{5} + \frac{n^3}{3} + \frac{7n}{15} \text{ is a natural number, for all } n \in \mathbb{N}.$$

**Step I** We observe that  $P(1)$  is true.

$$P(1) : \frac{(1)^5}{5} + \frac{1^3}{3} + \frac{7(1)}{15} = \frac{3+5+7}{15} = \frac{15}{15} = 1, \text{ which is a natural number. Hence, } P(1) \text{ is true.}$$

**Step II** Assume that  $P(n)$  is true, for  $n = k$ .

$$P(k) : \frac{k^5}{5} + \frac{k^3}{3} + \frac{7k}{15} \text{ is natural number.}$$

**Step III** Now, to prove  $P(k+1)$  is true.

$$\begin{aligned}
 & \frac{(k+1)^5}{5} + \frac{(k+1)^3}{3} + \frac{7(k+1)}{15} \\
 &= \frac{k^5 + 5k^4 + 10k^3 + 10k^2 + 5k + 1}{5} + \frac{k^3 + 1 + 3k(k+1)}{3} + \frac{7k+7}{15} \\
 &= \frac{k^5 + 5k^4 + 10k^3 + 10k^2 + 5k + 1}{5} + \frac{k^3 + 1 + 3k^2 + 3k}{3} + \frac{7k+7}{15} \\
 &= \frac{k^5}{5} + \frac{k^3}{3} + \frac{7k}{15} + \frac{5k^4 + 10k^3 + 10k^2 + 5k + 1}{5} + \frac{3k^2 + 3k + 1}{3} + \frac{7k+7}{15} \\
 &= \frac{k^5}{5} + \frac{k^3}{3} + \frac{7k}{15} + k^4 + 2k^3 + 2k^2 + k + k^2 + k + \frac{1}{5} + \frac{1}{3} + \frac{7}{15} \\
 &= \frac{k^5}{5} + \frac{k^3}{3} + \frac{7k}{15} + k^4 + 2k^3 + 3k^2 + 2k + 1, \text{ which is a natural number}
 \end{aligned}$$

So,  $P(k+1)$  is true, whenever  $P(k)$  is true. Hence,  $P(n)$  is true.

**Q. 24** Prove that  $\frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} > \frac{13}{24}$ , for all natural numbers  $n > 1$ .

**Sol.** Consider the given statement

$$P(n): \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} > \frac{13}{24}, \text{ for all natural numbers } n > 1.$$

**Step I** We observe that,  $P(2)$  is true,

$$\begin{aligned}
 P(2): \frac{1}{2+1} + \frac{1}{2+2} &> \frac{13}{24} \\
 \frac{1}{3} + \frac{1}{4} &> \frac{13}{24} \\
 \frac{4+3}{12} &> \frac{13}{24} \\
 \frac{7}{12} &> \frac{13}{24}, \text{ which is true.}
 \end{aligned}$$

Hence,  $P(2)$  is true.

**Step II** Now, we assume that  $P(n)$  is true,

For  $n = k$ ,

$$P(k): \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k} > \frac{13}{24}.$$

**Step III** Now, to prove  $P(k+1)$  is true, we have to show that

$$\begin{aligned}
 P(k+1): \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k} + \frac{1}{2(k+1)} &> \frac{13}{24} \\
 \text{Given, } \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k} &> \frac{13}{24} \\
 \frac{1}{k+1} + \frac{1}{k+2} + \frac{1}{2k} + \frac{1}{2(k+1)} &> \frac{13}{24} + \frac{1}{2(k+1)} \\
 \frac{13}{24} + \frac{1}{2(k+1)} &> \frac{13}{24} \\
 \therefore \frac{1}{k+1} + \frac{1}{k+2} + \dots + \frac{1}{2k} + \frac{1}{2(k+1)} &> \frac{13}{24}
 \end{aligned}$$

So,  $P(k+1)$  is true, whenever  $P(k)$  is true. Hence,  $P(n)$  is true.

**Q. 25** Prove that number of subsets of a set containing  $n$  distinct elements is  $2^n$ , for all  $n \in N$ .

**Sol.** Let  $P(n)$ : Number of subset of a set containing  $n$  distinct elements is  $2^n$ , for all  $n \in N$ .

*Step I* We observe that  $P(1)$  is true, for  $n = 1$ .

Number of subsets of a set contain 1 element is  $2^1 = 2$ , which is true.

*Step II* Assume that  $P(n)$  is true for  $n = k$ .

$P(k)$ : Number of subsets of a set containing  $k$  distinct elements is  $2^k$ , which is true.

*Step III* To prove  $P(k + 1)$  is true, we have to show that

$P(k + 1)$ : Number of subsets of a set containing  $(k + 1)$  distinct elements is  $2^{k+1}$ .

We know that, with the addition of one element in the set, the number of subsets become double.

$\therefore$  Number of subsets of a set containing  $(k + 1)$  distinct elements  $= 2 \times 2^k = 2^{k+1}$ .

So,  $P(k + 1)$  is true. Hence,  $P(n)$  is true.

## Objective Type Questions

**Q. 26** If  $10^n + 3 \cdot 4^{n+2} + k$  is divisible by 9, for all  $n \in N$ , then the least positive integral value of  $k$  is

(a) 5

(b) 3

(c) 7

(d) 1

**Sol. (a)** Let  $P(n)$ :  $10^n + 3 \cdot 4^{n+2} + k$  is divisible by 9, for all  $n \in N$ .

For  $n = 1$ , the given statement is also true  $10^1 + 3 \cdot 4^{1+2} + k$  is divisible by 9.

$$\begin{aligned} \therefore &= 10 + 3 \cdot 64 + k = 10 + 192 + k \\ &= 202 + k \end{aligned}$$

If  $(202 + k)$  is divisible by 9, then the least value of  $k$  must be 5.

$$\therefore 202 + 5 = 207 \text{ is divisible by 9}$$

$$\Rightarrow \frac{207}{9} = 23$$

Hence, the least value of  $k$  is 5.

**Q. 27** For all  $n \in N$ ,  $3 \cdot 5^{2n+1} + 2^{3n+1}$  is divisible by

(a) 19

(b) 17

(c) 23

(d) 25

**Sol. (b, c)**

Given that,  $3 \cdot 5^{2n+1} + 2^{3n+1}$

For  $n = 1$ ,

$$\begin{aligned} &3 \cdot 5^{2(1)+1} + 2^{3(1)+1} \\ &= 3 \cdot 5^3 + 2^4 \\ &= 3 \times 125 + 16 = 375 + 16 = 391 \end{aligned}$$

Now,  $391 = 17 \times 23$

which is divisible by both 17 and 23.

**Q. 28** If  $x^n - 1$  is divisible by  $x - k$ , then the least positive integral value of  $k$  is

- (a) 1 (b) 2 (c) 3 (d) 4

**Sol.** Let  $P(n) : x^n - 1$  is divisible by  $(x - k)$ .  
 For  $n = 1$ ,  $x^1 - 1$  is divisible by  $(x - k)$ .  
 Since, if  $x - 1$  is divisible by  $x - k$ . Then, the least possible integral value of  $k$  is 1.

## Fillers

**Q. 29** If  $P(n) : 2n < n!$ ,  $n \in N$ , then  $P(n)$  is true for all  $n \geq \dots\dots\dots$ .

**Sol.** Given that,  $P(n) : 2n < n!$ ,  $n \in N$

For $n = 1$ ,	$2 < 1!$	[false]
For $n = 2$ ,	$2 \times 2 < 2! \quad 4 < 2$	[false]
For $n = 3$ ,	$2 \times 3 < 3!$	
	$6 < 3!$	
	$6 < 3 \times 2 \times 1$	
	$(6 < 6)$	[false]
For $n = 4$ ,	$2 \times 4 < 4!$	
	$8 < 4 \times 3 \times 2 \times 1$	
	$(8 < 24)$	[true]
For $n = 5$ ,	$2 \times 5 < 5!$	
	$10 < 5 \times 4 \times 3 \times 2 \times 1$	
	$(10 < 120)$	[true]

Hence,  $P(n)$  is for all  $n \geq 4$ .

## True/False

**Q. 30** Let  $P(n)$  be a statement and let  $P(k) \Rightarrow P(k + 1)$ , for some natural number  $k$ , then  $P(n)$  is true for all  $n \in N$ .

**Sol. False**  
 The given statement is false because  $P(1)$  is true has not been proved.