

Chapter : 4. TRIANGLES

Exercise : 4A

Question: 1 A

D and E are point

Solution:

Given: AD = 3.6 cm, AB = 10 cm and AE = 4.5 cm.

Applying Thale's Theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow EC = \frac{AE}{AD} \times DB$$

$$\Rightarrow EC = \frac{4.5}{3.6} \times DB \quad [\because DB = AB - AD \Rightarrow DB = 10 - 3.6 = 6.4]$$

$$\Rightarrow EC = \frac{4.5}{3.6} \times 6.4$$

$$\Rightarrow EC = 8$$

Now, AC = AE + EC

$$\Rightarrow AC = 4.5 + 8 = 12.5$$

Hence, EC = 8 cm and AC = 12.5 cm

Question: 1 B

D and E are point

Solution:

Given: AB = 13.3 cm, AC = 11.9 cm and EC = 5.1 cm.

Applying Thale's Theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Since we need to find DB first, we add 1 on both sides

$$\frac{AD}{DB} + 1 = \frac{AE}{EC} + 1$$

$$\Rightarrow \frac{AD+DB}{DB} = \frac{AE+EC}{EC}$$

$$\Rightarrow \frac{AB}{DB} = \frac{AC}{EC}$$

$$\Rightarrow DB = \frac{AB \times EC}{AC}$$

$$\Rightarrow DB = \frac{13.3 \times 5.1}{11.9}$$

$$\Rightarrow DB = 5.7$$

AD is given by,

$$AD = AB - DB$$

$$\Rightarrow AD = 13.3 - 5.7$$

$$\Rightarrow AD = 7.6 \text{ cm}$$

Hence, AD is 7.6

Question: 1 C

D and E are point

Solution:

Given: $AD/DB = 4/7$ or $AD = 4$ cm, $DB = 7$ cm, and $AC = 6.6$

Applying Thale's Theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

We have AE at RHS but we need AC, as the value of AC is given. So by adding 1 to both sides of the equation, we can get the desired result

$$\Rightarrow \frac{AD}{DB} + 1 = \frac{AE}{EC} + 1$$

$$\Rightarrow \frac{AD+DB}{DB} = \frac{AE+EC}{EC}$$

$$\Rightarrow \frac{4+7}{7} = \frac{AC}{EC}$$

$$\Rightarrow \frac{11}{7} = \frac{6.6}{EC}$$

$$\Rightarrow EC = \frac{6.6 \times 7}{11}$$

$$\Rightarrow EC = 4.2$$

AE is given by,

$$AE = AC - EC$$

$$\Rightarrow AE = 6.6 - 4.2$$

$$\Rightarrow AE = 2.4$$

Hence, AE is 2.4 cm.

Question: 1 D

D and E are point

Solution:

Given: $AD/AB = 8/15$ or $AD = 8$ cm, $AB = 15$ cm, and $EC = 3.5$ cm

By applying Thale's Theorem,

$$\frac{AD}{AB} = \frac{AE}{AC}$$

$$\Rightarrow \frac{AD}{AB} = \frac{AE}{AE+EC}$$

$$\Rightarrow \frac{8}{15} = \frac{AE}{AE+3.5}$$

$$\Rightarrow 8 \times (AE + 3.5) = 15 \times AE$$

$$\Rightarrow 8 \times AE + 28 = 15 \times AE$$

$$\Rightarrow 15 \times AE - 8 \times AE = 28$$

$$\Rightarrow 7 \times AE = 28$$

$$\Rightarrow AE = 28/7 = 4$$

Hence, AE is 4 cm.

Question: 2 A

D and E are point

Solution:

Given: AD = x cm,

DB = (x - 2) cm,

AE = (x + 2) cm and,

EC = (x - 1) cm

By applying Thale's Theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow \frac{x}{x-2} = \frac{x+2}{x-1}$$

$$\Rightarrow x(x - 1) = (x + 2)(x - 2)$$

$$\Rightarrow x^2 - x = x^2 - 4$$

$$\Rightarrow x = 4$$

Thus, x = 4 cm

Question: 2 B

D and E are point

Solution:

Given: AD = 4 cm, DB = (x - 4) cm, AE = 8 cm and EC = (3x - 19) cm

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow \frac{4}{x-4} = \frac{8}{3x-19}$$

$$\Rightarrow 4(3x - 19) = 8(x - 4)$$

$$\Rightarrow 12x - 76 = 8x - 32$$

$$\Rightarrow 12x - 8x = 76 - 32$$

$$\Rightarrow 4x = 44$$

$$\Rightarrow x = 44/4 = 11$$

Thus, x = 11 cm

Question: 2 C

D and E are point

Solution:

Given: AD = (7x - 4) cm, AE = (5x - 2), DB = (3x + 4) cm and EC = 3x cm

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow \frac{7x-4}{3x+4} = \frac{5x-2}{3x}$$

$$\Rightarrow 3x(7x - 4) = (5x - 2)(3x + 4)$$

$$\Rightarrow 21x^2 - 12x = 15x^2 + 20x - 6x - 8$$

$$\Rightarrow 21x^2 - 12x = 15x^2 + 14x - 8$$

$$\Rightarrow 21x^2 - 15x^2 - 12x - 14x + 8 = 0$$

$$\Rightarrow 6x^2 - 26x + 8 = 0$$

$$\Rightarrow 2 \times (3x^2 - 13x + 4) = 0 \text{ [Simplifying the equation]}$$

$$\Rightarrow 3x^2 - 13x + 4 = 0$$

$$\Rightarrow 3x^2 - 12x - x + 4 = 0$$

$$\Rightarrow 3x(x - 4) - (x - 4) = 0$$

$$\Rightarrow (3x - 1)(x - 4) = 0$$

$$\Rightarrow (3x - 1) = 0 \text{ or } (x - 4) = 0$$

$$\Rightarrow x = 1/3 \text{ or } x = 4$$

Now since we've got two values of x, that is, 1/3 and 4. We shall check for its feasibility.

Substitute $x = 1/3$ in $AD = (7x - 4)$, we get

$AD = 7 \times (1/3) - 4 = -1.67$, which is not possible since side of a triangle cannot be negative.

Hence, $x = 4$ cm.

Question: 3 A

D and E are point

Solution:

Here, by applying converse of Thale's theorem we can conclude whether or not $DE \parallel BC$.

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Solving for $\frac{AD}{DB}$,

$$\frac{AD}{DB} = \frac{5.7}{9.5} = \frac{57}{95} = 0.6 \dots(i)$$

Solving for $\frac{AE}{EC}$,

$$\frac{AE}{EC} = \frac{4.8}{8} = 0.6 \dots(ii)$$

As equation (i) is equal to equation (ii),

$$\frac{AD}{DB} = \frac{AE}{EC}$$

it satisfies Thale's theorem.

Hence, we can say $DE \parallel BC$.

Question: 3 B

D and E are point

Solution:

Here, by applying converse of Thale's theorem we can conclude whether or not $DE \parallel BC$.

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Solving for $\frac{AD}{DB}$,

We need to find AD from given $AB = 11.7$ cm and $BD = 6.5$ cm.

$$AD = AB - BD$$

$$\Rightarrow AD = 11.7 - 6.5$$

$$\Rightarrow AD = 5.2$$

$$\frac{AD}{DB} = \frac{5.2}{6.5} = \frac{52}{65} = 0.8 \dots (i)$$

Solving for $\frac{AE}{EC}$,

We need to find EC from given AC = 11.2 cm and AE = 4.2 cm.

$$EC = AC - AE$$

$$\Rightarrow EC = 11.2 - 4.2$$

$$\Rightarrow EC = 7$$

$$\frac{AE}{EC} = \frac{4.2}{7} = 0.6 \dots (ii)$$

As equation (i) is not equal to equation (ii),

$$\frac{AD}{DB} \neq \frac{AE}{EC}$$

it doesn't satisfy Thale's theorem.

Hence, we can say DE not parallel to BC.

Question: 3 C

D and E are point

Solution:

Here, by applying converse of Thale's theorem we can conclude whether or not DE || BC.

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Solving for $\frac{AD}{DB}$,

We need to find DB from given AB = 10.8 cm and AD = 6.3 cm.

$$DB = AB - AD$$

$$\Rightarrow DB = 10.8 - 6.3$$

$$\Rightarrow DB = 4.5$$

$$\frac{AD}{DB} = \frac{6.3}{4.5} = \frac{63}{45} = 1.4 \dots (i)$$

Solving for $\frac{AE}{EC}$,

We need to find AE from given AC = 9.6 cm and EC = 4 cm.

$$AE = AC - EC$$

$$\Rightarrow AE = 9.6 - 4$$

$$\Rightarrow AE = 5.6$$

$$\frac{AE}{EC} = \frac{5.6}{4} = 1.4 \dots (ii)$$

As equation (i) is equal to equation (ii),

$$\frac{AD}{DB} = \frac{AE}{EC}$$

it satisfies Thale's theorem.

Hence, we can say $DE \parallel BC$.

Question: 3 D

D and E are point

Solution:

Here, by applying converse of Thale's theorem we can conclude whether or not $DE \parallel BC$.

By Thale's theorem,

$$\frac{AD}{DB} = \frac{AE}{EC}$$

Solving for $\frac{AD}{DB}$,

We need to find DB from given $AB = 12$ cm and $AD = 7.2$ cm.

$$DB = AB - AD$$

$$\Rightarrow DB = 12 - 7.2$$

$$\Rightarrow DB = 4.8$$

$$\frac{AD}{DB} = \frac{7.2}{4.8} = \frac{72}{48} = 1.5 \dots (i)$$

Solving for $\frac{AE}{EC}$,

We need to find EC from given $AC = 10$ cm and $AE = 6.4$ cm.

$$EC = AC - AE$$

$$\Rightarrow EC = 10 - 6.4$$

$$\Rightarrow EC = 3.6$$

$$\frac{AE}{EC} = \frac{6.4}{3.6} = \frac{64}{36} = 1.78 \dots (ii)$$

As equation (i) is not equal to equation (ii),

$$\frac{AD}{DB} \neq \frac{AE}{EC}$$

it doesn't satisfies Thale's theorem.

Hence, we can say DE is not parallel to BC.

Question: 4 A

In a $\triangle ABC$, AD is

Solution:

Given: $AB = 6.4$ cm, $AC = 8$ cm and $BD = 5.6$ cm

Since AD bisects $\angle A$, we can apply angle-bisector theorem in $\triangle ABC$,

$$\frac{BD}{DC} = \frac{AB}{AC}$$

Substituting given values, we get

$$\frac{5.6}{DC} = \frac{6.4}{8}$$

$$\Rightarrow DC = \frac{5.6 \times 8}{6.4}$$

$$\Rightarrow DC = 7$$

Thus, DC is 7 cm.

Question: 4 B

In a $\triangle ABC$, AD is

Solution:

Given: AB = 10 cm, AC = 14 cm and BC = 6 cm

Since AD bisects $\angle A$, we can apply angle-bisector theorem in $\triangle ABC$,

$$\frac{BD}{DC} = \frac{AB}{AC}$$

Substituting given values, we get

$$\frac{BD}{DC} = \frac{10}{14}$$

To find BD and DC,

Let BD = x cm, and it's given that BC = 6 cm, then DC = (6 - x) cm

Then

$$\frac{x}{6-x} = \frac{10}{14}$$

$$\Rightarrow 14x = 10(6 - x)$$

$$\Rightarrow 14x = 60 - 10x$$

$$\Rightarrow 14x + 10x = 60$$

$$\Rightarrow 24x = 60$$

$$\Rightarrow x = 60/24 = 2.5$$

$$\Rightarrow BD = 2.5 \text{ cm}$$

If BD = 2.5 cm and BC = 6 cm, then DC = (6 - x) = (6 - 2.5) = 3.5

Thus, BD is 2.5 cm and DC = 3.5 cm.

Question: 4 C

In a $\triangle ABC$, AD is

Solution:

Given: AB = 5.6 cm, BC = 6 cm and BD = 3.2 cm

Since AD bisects $\angle A$, we can apply angle-bisector theorem in $\triangle ABC$,

$$\frac{BD}{DC} = \frac{AB}{AC}$$

Substituting given values, we get

$$\frac{3.2}{DC} = \frac{5.6}{AC}$$

Here, DC is given by

$$DC = BC - BD$$

$$\Rightarrow DC = 6 - 3.2 = 2.8$$

$$\frac{3.2}{2.8} = \frac{5.6}{AC}$$

$$\Rightarrow AC = \frac{5.6 \times 2.8}{3.2}$$

$$\Rightarrow AC = 4.9$$

Thus, AC is 4.9 cm.

Question: 4 D

In a $\triangle ABC$, AD is

Solution:

Given: $AB = 5.6$ cm, $AC = 4$ cm and $DC = 3$ cm

Since AD bisects $\angle A$, we can apply angle-bisector theorem in $\triangle ABC$,

$$\frac{BD}{DC} = \frac{AB}{AC}$$

Substituting given values, we get

$$\frac{BD}{3} = \frac{5.6}{4}$$

$$\Rightarrow BD = \frac{5.6 \times 3}{4}$$

$$\Rightarrow BD = 4.2$$

Now, $BC = BD + DC$

$$\Rightarrow BC = 4.2 + 3 = 7.2$$

Thus, BC is 7.2 cm.

Question: 5

M is a point on t

Solution:

(i). Given: ABCD is a parallelogram.

To Prove: $\frac{DM}{MN} = \frac{DC}{BN}$

Proof: In $\triangle DMC$ and $\triangle NMB$,

$$\angle DMC = \angle NMB \text{ [}\because \text{ they are vertically opposite angles]}$$

$$\angle DCM = \angle NBM \text{ [}\because \text{ they are alternate angles]}$$

$$\angle CDM = \angle MNB \text{ [}\because \text{ they are alternate angles]}$$

By AAA-similarity, we can say

$$\triangle DMC \sim \triangle NMB$$

So, from similarity of the triangle, we can say

$$\frac{DM}{MN} = \frac{DC}{BN}$$

Hence, proved.

(ii). Given: ABCD is a parallelogram.

To Prove: $\frac{DN}{DM} = \frac{AN}{DC}$

Proof: As we have already derived

$$\frac{DM}{MN} = \frac{DC}{BN}$$

Add 1 on both sides of the equation, we get

$$\frac{DM}{MN} + 1 = \frac{DC}{BN} + 1$$

$$\Rightarrow \frac{DM+MN}{MN} = \frac{DC+BN}{BN}$$

$\Rightarrow \frac{DM+MN}{MN} = \frac{AB+BN}{BN}$ [$\because$ ABCD is a parallelogram and a parallelogram's opposite sides are always equal $\Rightarrow DC = AB$]

$$\Rightarrow \frac{DN}{MN} = \frac{AN}{BN}$$


Hence, proved.

Question: 6

Show that the lin

Solution:

We can draw the trapezium as



Here, let EF be the line segment joining the oblique sides of the trapezium at midpoints E and F (say) correspondingly.

Construction: Extend AD and BC such that it meets at P.

To Prove: $EF \parallel DC$ and $EF \parallel AB$

Proof: Given that, ABCD is trapezium which means $DC \parallel AB$(statement (i))

In $\triangle PAB$,

$DC \parallel AB$ (by statement (i))

So, this means we can apply Thale's theorem in $\triangle PAB$. We get

$$\frac{PD}{DA} = \frac{PC}{CB} \dots (ii)$$

$\because$ E and F are midpoints of AD and BC respectively, we can write

$$DA = DE + EA$$

$$\text{Or } DA = 2DE \dots (iii)$$

$$CB = CF + FB$$

$$\text{Or } CB = 2CF \dots (iv)$$

Substituting equation (iii) and (iv) in equation (ii), we get

$$\frac{PD}{2DE} = \frac{PC}{2CF}$$

$$\Rightarrow \frac{PD}{DE} = \frac{PC}{CF}$$

By applying converse of Thale's theorem, we can write $DC \parallel EF$.

Now if $DC \parallel EF$, and we already know that $DC \parallel AB$.

$\Rightarrow$ EF is also parallel to AB, that is, $EF \parallel AB$.

This means, $DC \parallel EF \parallel AB$.


Hence, proved.

Question: 7

In the adjoining

Solution:

Given: In the adjoining figure, ABCD is a trapezium in which $CD \parallel AB$ and its diagonals intersect at O. If $AO = (5x - 7)$ cm, $OC = (2x + 1)$ cm, $DO = (7x - 5)$ cm and $OB = (7x + 1)$ cm. **To find:** the



value of x. **Solution:**

In the trapezium ABCD, $AB \parallel DC$ and

its diagonals intersect at O. Through O draw $EO \parallel AB$ meeting AD at E. Now In $\triangle ADC$ As $EO \parallel AB \parallel DC$ By thales theorem which states that If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points then the other two sides are divided in the same

ratio. $\therefore \frac{AE}{ED} = \frac{AO}{OC}$ (i) In $\triangle DAB$, $EO \parallel AB$ By thales theorem, $\therefore \frac{DE}{EA} = \frac{DO}{OB}$

$$\Rightarrow \frac{AE}{ED} = \frac{BO}{OD} \quad \text{..... (ii)} \quad \text{From (i) and (ii)} \quad \frac{AO}{OC} = \frac{BO}{OD}$$

Put the given values as:

$$\Rightarrow \frac{5x-7}{2x+1} = \frac{7x-5}{7x+1}$$

$$\Rightarrow (5x - 7)(7x + 1) = (7x - 5)(2x + 1)$$

$$\Rightarrow 35x^2 + 5x - 49x - 7 = 14x^2 - 10x + 7x - 5$$

$$\Rightarrow 35x^2 - 44x - 7 = 14x^2 - 3x - 5$$

$$\Rightarrow 35x^2 - 14x^2 - 44x + 3x - 7 + 5 = 0$$

$$\Rightarrow 21x^2 - 41x - 2 = 0$$

$$\Rightarrow 21x^2 - 42x + x - 2 = 0$$

$$\Rightarrow 21x(x - 2) + (x - 2) = 0$$

$$\Rightarrow (21x + 1)(x - 2) = 0$$

$$\Rightarrow (21x + 1) = 0 \text{ or } (x - 2) = 0$$

$$\Rightarrow x = -1/21 \text{ or } x = 2$$

But $x = -1/21$ doesn't satisfy the length of intersected lines.

So $x \neq -1/21$


And thus, $x = 2$.

Question: 8

In a $\triangle ABC$

Solution:

We have



To show that, $MN \parallel BC$.

Given that, $\angle B = \angle C$ and $BM = CN$.

So, $AB = AC$ [sides opposite to equal angles ($\angle B = \angle C$) are equal]

Subtract BM from both sides, we get

$$AB - BM = AC - BM$$

$$\Rightarrow AB - BM = AC - CN$$

$$\Rightarrow AM = AN$$

$$\Rightarrow \angle AMN = \angle ANM$$

[angles opposite to equal sides (AM = AN) are equal] ... (i)

We know in $\triangle ABC$,

$$\angle A + \angle B + \angle C = 180^\circ [\because \text{sum of angles of a triangle is } 180^\circ] \dots (ii)$$

And in $\triangle AMN$,

$$\angle A + \angle AMN + \angle ANM = 180^\circ [\because \text{sum of angles of a triangle is } 180^\circ] \dots (iii)$$

Comparing equations (ii) and (iii), we get

$$\angle A + \angle B + \angle C = \angle A + \angle AMN + \angle ANM$$

$$\Rightarrow \angle B + \angle C = \angle AMN + \angle ANM$$

$$\Rightarrow 2\angle B = 2\angle AMN [\because \text{from equation (i), and also } \angle B = \angle C]$$

$$\Rightarrow \angle B = \angle AMN$$

Thus, $MN \parallel BC$ since the corresponding angles, $\angle AMN = \angle B$.

Question: 9

$\triangle ABC$ and $\triangle BDC$ lie

Solution:

We can observe two triangles in the figure.

In $\triangle ABC$,

$PQ \parallel AB$

Applying Thale's theorem, we get

$$\frac{CP}{PB} = \frac{CQ}{QA} \dots (i)$$

In $\triangle BDC$,

$PR \parallel BP$

Applying Thale's theorem, we get

$$\frac{CP}{QA} = \frac{CR}{RO} \dots (ii)$$

Comparing equations (i) and (ii),

$$\frac{CQ}{QA} = \frac{CR}{RO}$$

Now, applying converse of Thale's theorem, we get

$QR \parallel AD$


Hence, QR is parallel to the AD.

Question: 10

In the given figu

Solution:

We have the diagram as,



Given: $BD = DC$ & $OD = DX$

To Prove: $\frac{AO}{AX} = \frac{AF}{AB}$ and also, $EF \parallel BC$

Proof: Since, from the diagram we can see that diagonals OX and BC bisect each other in quadrilateral $BOCX$. Thus, $BOCX$ is a parallelogram.

If $BOCX$ is a parallelogram, $BX \parallel OC$, and $BO \parallel CX$.

$\Rightarrow BX \parallel FC$ (as OC extends to FC) and $CX \parallel BE$ (BO extends to BE)

$\Rightarrow BX \parallel OF$ and $CX \parallel OE$

$\therefore BX \parallel OF$, applying Thale's theorem in $\triangle ABX$, we get

$$\frac{AO}{AX} = \frac{AF}{AB} \dots(i)$$

Now since $CX \parallel OE$, applying Thale's theorem in $\triangle ACX$, we get

$$\frac{AO}{AX} = \frac{AE}{AC} \dots(ii)$$

By equations (i) and (ii), we get

$$\frac{AF}{AB} = \frac{AE}{AC}$$

By applying converse of Thale's theorem in the above equation, we can write

$EF \parallel BC$


Hence, proved.

Question: 11

$ABCD$ is a paralle

Solution:

We have the diagram as



Given: $DP = PC$ &

$CQ = \frac{1}{4}AC \dots(i)$

To Prove: $CR = RB$

Proof: Join B to D

As diagonals of a parallelogram bisect each other at S .

$$CS = \frac{1}{2}AC \dots(ii)$$

Dividing equation (i) by (ii), we get

$$\frac{CQ}{CS} = \frac{AC}{4} \times \frac{2}{AC}$$

$$\Rightarrow \frac{CQ}{CS} = \frac{1}{2}$$

$$\Rightarrow CQ = CS/2$$

$\Rightarrow$ Q is the midpoint of CS.

According to midpoint theorem in ΔCSD , we have

$$PQ \parallel DS$$

Similarly, in ΔCSB , we have

$$QR \parallel SB$$

Also, given that $CQ = QS$

We can conclude that, by the converse of midpoint theorem, $CR = RB$.

That is, R is the midpoint of CB.

Hence, proved.

Question: 12

In the adjoining

Solution:

Given: $AD = AE$...(i)

& $AB = AC$...(ii)

Subtracting AD from both sides of equation (ii), we get

$$AB - AD = AC - AD$$

$$\Rightarrow AB - AD = AC - AE \text{ [from equation (i)]}$$

$$\Rightarrow DB = EC \text{ [}\because AB - AD = DB \text{ \& } AC - AE = EC \text{] ... (iii)}$$

Now, divide equation (i) by (iii), we get

$$\frac{AD}{DB} = \frac{AE}{EC}$$

By converse of Thale's theorem, we can conclude by this equation that $DE \parallel BC$.

So, $\angle DEC + \angle ECB = 180^\circ$ [$\because$ sum of interior angles on the same transversal line is 180°]

Or $\angle DEC + \angle DBC = 180^\circ$ [$\because AB = AC \Rightarrow \angle C = \angle B$]

Hence, we can write DEBC is cyclic and points D, E, B and C are concyclic.

Question: 13

In ΔABC , the bise

Solution:

Given: $\angle PBR = \angle QBR$ & $PQ \parallel AC$.

In ΔBQP ,

BR bisects $\angle B$ such that $\angle PBR = \angle QBR$.

Since angle-bisector theorem says that, if two angles are bisected in a triangle then it equates their relative lengths to the relative lengths of the other two sides of the triangles.

So by applying angle-bisector theorem, we get

$$\frac{QR}{PR} = \frac{BQ}{BP}$$

$$\Rightarrow QR \times BP = PR \times BQ$$

Hence, proved.

Exercise : 4B

Question: 1 A

In each of the gi

Solution:

In these triangles ABC and PQR, observe that

$$\angle BAC = \angle PQR = 50^\circ$$

$$\angle ABC = \angle QPR = 60^\circ$$

$$\angle ACB = \angle PRQ = 70^\circ$$

Thus, by angle-angle-angle similarity, i.e., AAA similarity,

$$\Delta ABC \sim \Delta PQR$$

Question: 1 B

In each of the gi

Solution:

In triangles ABC & EFD,

$$\angle ABC \neq \angle EDF$$

$$\frac{AB}{DF} = \frac{3}{6} = \frac{1}{2}$$

$$\frac{BC}{DE} = \frac{4.5}{9} = \frac{1}{2}$$

So, clearly, since no criteria satisfies, ΔABC is not similar to ΔEFD .

Question: 1 C

In each of the gi

Solution:

In triangles ABC & PQR,

$$\angle ACB = \angle PQR$$

$$\frac{CA}{QR} = \frac{8}{6} = \frac{4}{3}$$

$$\frac{BC}{PQ} = \frac{6}{4.5} = \frac{4}{3}$$

By SAS criteria, we can say

$$\Delta ABC \sim \Delta PQR$$

Question: 1 D

In each of the gi

Solution:

In triangles DEF & PQR,

$$\frac{DE}{QR} = \frac{2.5}{5} = \frac{1}{2}$$

$$\frac{EF}{PQ} = \frac{2}{4} = \frac{1}{2}$$

$$\frac{DF}{PR} = \frac{3}{6} = \frac{1}{2}$$

By SSS criteria, we can write

$$\triangle DEF \sim \triangle PQR$$

Question: 1 E

In each of the gi

Solution:

In $\triangle ABC$, we can find $\angle ABC$.

$$\angle ABC + \angle BCA + \angle CAB = 180^\circ [\because \text{sum of all the angles of a triangle is } 180^\circ]$$

$$\Rightarrow \angle ABC + 70^\circ + 80^\circ = 180^\circ$$

$$\Rightarrow \angle ABC + 150^\circ = 180^\circ$$

$$\Rightarrow \angle ABC = 180^\circ - 150^\circ$$

$$\Rightarrow \angle ABC = 30^\circ$$

We can observe from triangles ABC & MNR ,

$$\angle ABC = \angle MNR$$

$$\angle CAB = \angle RMN$$

Hence, by AA similarity we can say, $\triangle ABC \sim \triangle MNR$

Question: 2

In the given figu

Solution:

(i) To find $\angle DOC$, we can observe the straight line DB .

$$\angle DOC + \angle COB = 180^\circ [\because \text{sum of all angles in a straight line is } 180^\circ]$$

$$\Rightarrow \angle DOC + 115^\circ = 180^\circ$$

$$\Rightarrow \angle DOC = 180^\circ - 115^\circ$$

$$\Rightarrow \angle DOC = 65^\circ$$

(ii) In $\triangle DOC$,

And given that, $\angle CDO = 70^\circ$, $\angle DOC = 65^\circ$ (from (i))

$$\angle DOC + \angle DCO + \angle CDO = 180^\circ$$

$$\Rightarrow 65^\circ + \angle DCO + 70^\circ = 180^\circ$$

$$\Rightarrow \angle DCO + 135^\circ = 180^\circ$$

$$\Rightarrow \angle DCO = 180^\circ - 135^\circ$$

$$\Rightarrow \angle DCO = 45^\circ$$

(iii) We have derived $\angle DCO$ from (ii), $\angle DCO = 45^\circ$

Thus, $\angle OAB = 45^\circ$ [$\because \angle OAB = \angle DCO$ as $\triangle ODC \sim \triangle OBA$]

(iv) It's given that, $\angle CDO = 70^\circ$

Thus, $\angle OBA = 70^\circ$ [$\because \angle OBA = \angle CDO$ as $\triangle ODC \sim \triangle OBA$]

Question: 3

In the given figu

Solution:

(i). Given that, AB = 8 cm

BO = 6.4 cm,

OC = 3.5 cm

& CD = 5 cm

$\Delta OAB \sim \Delta OCD$

When two triangles are similar, they can be written in the ratio as

$$\frac{OA}{OC} = \frac{AB}{CD}$$

Substitute gave values in the above equations,

$$\frac{OA}{3.5} = \frac{8}{5}$$

$$\Rightarrow OA = \frac{8 \times 3.5}{5}$$

$$\Rightarrow OA = 5.6$$

Thus, OA = 5.6 cm

(ii). Given that, AB = 8 cm

BO = 6.4 cm,

OC = 3.5 cm

& CD = 5 cm

$\Delta OAB \sim \Delta OCD$

When two triangles are similar, they can be written in the ratio as

$$\frac{BO}{DO} = \frac{AB}{CD}$$

Substitute gave values in the above equations,

$$\frac{6.4}{DO} = \frac{8}{5}$$

$$\Rightarrow DO = \frac{5 \times 6.4}{8}$$

$$\Rightarrow DO = 4$$

Thus, DO = 4 cm

Question: 4

In the given figu

Solution:

Given is that $\angle ADE = \angle B$

From the diagram clearly, $\angle EAD = \angle BAC$ [$\because$ they are common angles]

Now, since two of the angles are correspondingly equal. Then by AA similarity criteria, we can say

$\Delta ADE \sim \Delta ABC$

Further, it's given that

$$AD = 3.8 \text{ cm}$$

$$AE = 3.6 \text{ cm}$$

$$BE = 2.1 \text{ cm}$$

$$BC = 4.2 \text{ cm}$$

$$DE = ?$$

To find AB, we can express it in the form $AB = AE + BE = 3.6 + 2.1$

$$\Rightarrow AB = 5.7$$

So for the condition that $\triangle ADE \sim \triangle ABC$,

$$\frac{DE}{BC} = \frac{AD}{AB}$$

Substituting given values in the above equation,

$$\Rightarrow \frac{DE}{4.2} = \frac{3.8}{5.7}$$

$$\Rightarrow DE = \frac{3.8 \times 4.2}{5.7}$$

$$\Rightarrow DE = 2.8$$

Thus, $DE = 2.8 \text{ cm}$

Question: 5

The perimeters of

Solution:

Given that, $\triangle ABC \sim \triangle PQR$

And perimeter of $\triangle ABC = 32 \text{ cm}$ & perimeter of $\triangle PQR = 24 \text{ cm}$

We can write relationship as,

$$\frac{\text{perimeter of } \triangle ABC}{\text{perimeter of } \triangle PQR} = \frac{AB}{PQ}$$

$$\Rightarrow \frac{32}{24} = \frac{AB}{12}$$

$$\Rightarrow AB = \frac{32 \times 12}{24}$$

$$\Rightarrow AB = 16$$

Thus, $AB = 16 \text{ cm}$.

Question: 6

The corresponding

Solution:

Given that, $\triangle ABC \sim \triangle DEF$

Also, $BC = 9.1 \text{ cm}$ & $EF = 6.5 \text{ cm}$

And perimeter of $\triangle DEF = 25 \text{ cm}$

We need to find perimeter of $\triangle ABC = ?$

We can write relationship as,

$$\frac{\text{the perimeter of } \triangle ABC}{\text{the perimeter of } \triangle DEF} = \frac{BC}{EF}$$

$$\Rightarrow \frac{\text{the perimeter of } \triangle ABC}{25} = \frac{9.1}{6.5}$$

$$\Rightarrow \text{perimeter of } \triangle ABC = \frac{9.1 \times 25}{6.5}$$

$$\Rightarrow \text{perimeter of } \triangle ABC = 35$$

Thus, perimeter of $\triangle ABC = 35 \text{ cm}$

Question: 7

In the given figu

Solution:

Given that, $\angle CAB = 90^\circ$

$$AC = 75 \text{ cm}$$

$$AB = 1 \text{ m}$$

$$BC = 1.25 \text{ m}$$

To show that, $\triangle BDA \sim \triangle BAC$

In the diagram, we can see

$$\angle BDA = \angle BAC = 90^\circ$$

$$\angle DBA = \angle CBA \text{ [They are common angles]}$$

So by AA-similarity theorem,

$$\triangle BDA \sim \triangle BAC$$

Thus, now since $\triangle BDA \sim \triangle BAC$, we can write as

$$\frac{AD}{AC} = \frac{AB}{BC}$$

$$\Rightarrow \frac{AD}{75} = \frac{100}{125} \text{ [}\because AC = 75 \text{ cm, } AB = 1 \text{ m} = 100 \text{ cm \& } BC = 1.25 \text{ m} = 125 \text{ cm]}$$

$$\Rightarrow AD = \frac{100 \times 75}{125}$$

$$\Rightarrow AD = 60 \text{ cm}$$

Hence, $AD = 60 \text{ cm}$ or 0.6 m

Question: 8

In the given figu

Solution:

Given that, $\angle ABC = 90^\circ$

$$AB = 5.7 \text{ cm}$$

$$BD = 3.8 \text{ cm}$$

$$CD = 5.4 \text{ cm}$$

In order to find BC, we need to prove that $\triangle BDC$ and $\triangle ABC$ are similar.

$$\angle BDC = \angle ABC = 90^\circ$$

$$\angle ACB = \angle DCB \text{ [They are common angles]}$$

By this we have proved $\triangle BDC \sim \triangle ABC$, by AA-similarity criteria.

So we can write,

$$\frac{BD}{AB} = \frac{DC}{BC}$$

$$\Rightarrow \frac{3.8}{5.7} = \frac{5.4}{BC}$$

$$\Rightarrow BC = \frac{5.4 \times 5.7}{3.8}$$

$$\Rightarrow BC = 8.1$$

Hence, $BC = 8.1$ cm.

Question: 9

In the given figu

Solution:

Given that, $\angle ABC = 90^\circ$

$AD = 4$ cm

$BD = 8$ cm

In order to find CD, we need to prove that $\triangle BDC$ and $\triangle ABC$ are similar.

$$\angle BDC = \angle ADB = 90^\circ$$

$$\angle DBA = \angle DCB$$

We have proved $\triangle DBA \sim \triangle DCB$, by AA-similarity criteria.

So we can write,

$$\frac{BD}{CD} = \frac{AD}{BD}$$

$$\Rightarrow \frac{8}{CD} = \frac{4}{8}$$

$$\Rightarrow CD = \frac{8 \times 8}{4}$$

$$\Rightarrow CD = 16$$

Hence, $CD = 16$ cm.

Question: 10

P and Q are point

Solution:

There are two triangles here, $\triangle APQ$ and $\triangle ABC$. We shall prove these triangles to be similar.

$$\frac{AP}{AB} = \frac{2}{4+2} = \frac{2}{6} = \frac{1}{3}$$

$$\& \frac{AQ}{AC} = \frac{3}{6+3} = \frac{3}{9} = \frac{1}{3}$$

$$\Rightarrow \frac{AP}{AB} = \frac{AQ}{AC}$$

Also, $\angle A = \angle A$ [common angle]

So by AA-similarity criteria,

$$\triangle APQ \sim \triangle ABC$$

Thus,

$$\frac{PQ}{BC} = \frac{AQ}{AC}$$

$$\text{And we know } \frac{PQ}{BC} = \frac{1}{3}$$

$$\Rightarrow BC = 3 \times PQ$$

Hence, proved.

Question: 11

ABCD is a paralle

Solution:

Given that, $AB \parallel DC$ & $AD \parallel BC$

To Prove: $AF \times FB = EF \times FD$

Proof: In $\triangle DAF$ & $\triangle BEF$

$\angle DAF = \angle BEF$ [$\because$ they are alternate angles]

$\angle AFD = \angle EFB$ [$\because$ they are vertically opposite angles]

This implies that $\triangle DAF \sim \triangle BEF$ by AA-similarity criteria.

$$\Rightarrow \frac{AF}{EF} = \frac{FD}{FB}$$

Now cross-multiply them,

$$AF \times FB = FD \times EF$$

Hence, proved.

Question: 12

In the given figu

Solution:

Observe in $\triangle BED$ & $\triangle ACB$, we have

$$\angle BED = \angle ACB = 90^\circ$$

Now according to what's given, $DB \perp BC$ and $AC \perp BC$ we can write,

$$\angle B + \angle C = 180^\circ$$

This clearly means $BD \parallel CA$

$$\Rightarrow \angle EBD = \angle CAB \text{ [They are alternate angles]}$$

AA Similarity theorem: The postulate states that two triangles are similar if they have two corresponding angles that are congruent or equal in measure.

Thus, by AA-similarity theorem, $\triangle BED \sim \triangle ACB$ Now, by property of similarity of triangles,

$$\text{So, } \frac{BE}{AC} = \frac{DE}{BC}$$

Cross-multiplying, we get,

$$\Rightarrow \frac{BE}{DE} = \frac{AC}{BC}$$


Hence, proved.

Question: 13

A vertical pole o

Solution:

We have



Let the two triangles be ΔABC and ΔPQR .

Given that, $AB = 7.5 \text{ cm}$

$BC = 5 \text{ m} = 500 \text{ cm}$

$QR = 24 \text{ m} = 2400 \text{ cm}$

We have to find $PQ = x$ (say).

We need to prove ΔABC is similar to ΔPQR .

We can observe that,

$$\angle ABC = \angle PQR = 90^\circ$$

$$\angle ACB = \angle PRQ [\because \text{the sum of angles is same at all places at the same time}]$$

Thus, by AA-similarity criteria, we can say

$$\Delta ABC \sim \Delta PQR$$

So,

$$\frac{AB}{PQ} = \frac{BC}{QR}$$

Substitute the given values in this equation,

$$\frac{7.5}{x} = \frac{500}{2400}$$

$$\Rightarrow x = \frac{7.5 \times 2400}{500}$$

$$\Rightarrow x = 36 \text{ cm}$$

Thus, height of the tower is 36 cm.

Question: 14

In an isosceles Δ

Solution:

To prove: $\Delta ACP \sim \Delta BCQ$

Proof:

Given that, ΔABC is an isosceles triangle. $\Rightarrow AC = BC$

Also, if ΔABC is an isosceles triangle,

$$\text{then } \angle CAB = \angle CBA \dots (i)$$

Subtracting it by 180° from both sides, we get

$$180^\circ - \angle CAB = 180^\circ - \angle CBA$$

$$\Rightarrow \angle CAP = \angle CBQ \dots (ii)$$

Also, given that $AP \times BQ = AC \times AC$

$$\text{Or } \frac{AP}{AC} = \frac{AC}{BQ}$$

$$\text{Or } \frac{AP}{AC} = \frac{BC}{BQ} [\because AC = BC] \dots (iii)$$

Recollecting equations (i), (ii) and (iii),

By SAS-similarity criteria, we get

$$\Delta ACP \sim \Delta BCQ$$

Hence, proved.

Question: 15

In the given figu

Solution:

To Prove: $\triangle ACB \sim \triangle DCE$

Proof:

Given that, $\angle 1 = \angle 2$

$$\Rightarrow \angle DBC = \angle DCE$$

Also in $\triangle ABC$ & $\triangle DCE$, we get

$\angle DCE = \angle ACB$ [they are common angles to both triangles]

And $\frac{AC}{BD} = \frac{CB}{CE}$

Or $\frac{AC}{CB} = \frac{BD}{CE}$

Or $\frac{AC}{CB} = \frac{DC}{CE}$ [$\because BD = DC$ as $\angle 1 = \angle 2$]

Thus by SAS-similarity criteria, we get

$\triangle ACB \sim \triangle DCE$

Hence, proved.

Question: 16

ABCD is a quadril

Solution:

Given: $AD = BC$

P, Q, R and S are the midpoints of AB, AC, CD and BD respectively.

So in $\triangle ABC$, if P and Q are midpoints of AB and AC respectively $\Rightarrow PQ \parallel BC$

And $PQ = (1/2)BC$... (i)

Similarly in $\triangle ADC$,

$QR = (1/2)AD$... (ii)

In $\triangle BCD$,

$SR = (1/2)BC$... (iii)

In $\triangle ABD$,

$PS = (1/2)AD = (1/2)BC$ [$\because AD = BC$]

Using equations (i), (ii), (iii) & (iv), we get

$PQ = QR = SR = PS$

All these sides are equal.

$\Rightarrow PQRS$ is a rhombus.

Hence, shown that $PQRS$ is a rhombus.

Question: 17

In a circle, two

Solution:

Given: AB and CD are chords of the circle, intersecting at point P.

(a). To Prove: $\triangle PAC \sim \triangle PDB$

Proof: In ΔPAC and ΔPDB ,

$\angle APC = \angle DPB$ [$\because$ they are vertically opposite angles]

$\angle CAP = \angle PDB$ [$\because$ angles in the same segment are equal]

Thus, by AA-similarity criteria, we can say that,

$\Delta PAC \sim \Delta PDB$

Hence, proved.

(b). To Prove: $PA \times PB = PC \times PD$

Proof: As already proved that $\Delta PAC \sim \Delta PDB$

We can write as,

$$\frac{PA}{PD} = \frac{PC}{PB}$$

By cross-multiplying, we get

$$PA \times PB = PC \times PD$$

Hence, proved.

Question: 18

Two chords AB and

Solution:

Given: AB and CD are chords of a circle intersecting at point P outside the circle.

(a). To Prove: $\Delta PAC \sim \Delta PDB$

Proof: We know

$\angle ABD + \angle ACD = 180^\circ$ [$\because$ opposite angles of cyclic quadrilateral are supplementary] ... (i)

$\angle PCA + \angle ACD = 180^\circ$ [$\because$ they are linear pair angle] ... (ii)

Comparing equations (i) & (ii), we get

$$\angle ABD + \angle ACD = \angle PCA + \angle ACD$$

$$\Rightarrow \angle ABD = \angle PCA$$

Also, $\angle APC = \angle BPD$ [$\because$ they are common angles]

Thus, by AA-similarity criteria, $\Delta PAC \sim \Delta PDB$

Hence, proved.

(b). To Prove: $PA \times PB = PC \times PD$

Proof: We have already proved that, $\Delta PAC \sim \Delta PDB$

Thus the ratios can be written as,

$$\frac{PA}{PD} = \frac{PC}{PB}$$

By cross-multiplication, we get

$$PA \times PB = PC \times PD$$

Hence, proved.

Question: 19

In a right triang

Solution:

By the property that says, if a perpendicular is drawn from the vertex of a right triangle

to the hypotenuse then the triangles on both the sides of the perpendicular are similar to the whole triangle and also to each other.

We can conclude by the property in $\triangle BDC$,

$$\triangle CQD \sim \triangle DQB$$

(a). To Prove: $DQ^2 = DP \times QC$

Proof: As already proved, $\triangle CQD \sim \triangle DQB$

We can write the ratios as,

$$\frac{CQ}{DQ} = \frac{DQ}{QB}$$

By cross-multiplication, we get

$$DQ^2 = QB \times QC \dots(i)$$

Now since, quadrilateral PDQB forms a rectangle as all angles are 90° in PDQB.

$$\Rightarrow DP = QB \text{ \& } PB = DQ$$

And thus replacing QB by DP in equation (i), we get

$$DQ^2 = DP \times QC$$

Hence, proved.

(b). To Prove: $DP^2 = DQ \times AP$

Prof: Similarly using same property, we get

$$\triangle APD \sim \triangle DPB$$

We can write the ratios as,

$$\frac{AP}{DP} = \frac{PD}{PB}$$

By cross-multiplication, we get

$$DP^2 = PB \times AP$$

$$\Rightarrow DP^2 = DQ \times AP [\because PB = DQ]$$


Hence, proved.

Exercise : 4C

Question: 1

$\triangle ABC \sim \triangle DEF$ and t

Solution:



We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{64}{121} = \frac{BC^2}{EF^2} = \frac{BC^2}{(15.4)^2}$$


$$\Rightarrow BC^2 = \frac{64}{121} \times (15.4)^2$$

$$\Rightarrow BC = \sqrt{\frac{64}{121} \times (15.4)^2} = \frac{8}{11} \times 15.4 = 8 \times 1.4 = 11.2 \text{ cm}$$

Question: 2

The areas of two

Solution:



We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{9}{16} = \frac{BC^2}{QR^2} = \frac{(4.5)^2}{QR^2}$$


$$\Rightarrow QR^2 = \frac{16}{9} \times (4.5)^2$$

$$\Rightarrow QR = \sqrt{\frac{16}{9} \times (4.5)^2} = \frac{4}{3} \times 4.5 = 1.5 \times 4 = 6 \text{ cm}$$

Question: 3

$\triangle ABC \sim \triangle PQR$ and a

Solution:



Given that $\frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{4}{1}$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{4}{1} = \frac{BC^2}{QR^2} = \frac{(12)^2}{QR^2}$$


$$\Rightarrow QR^2 = \frac{1}{4} \times (12)^2$$

$$\Rightarrow QR = \sqrt{\frac{1}{4} \times (12)^2} = \frac{1}{2} \times 12 = 6 \text{ cm}$$

Question: 4

The areas of two

Solution:



Let the two triangles be ABC and PQR and their longest sides are BC and QR.

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their longest sides.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{169}{121} = \frac{BC^2}{QR^2} = \frac{(26)^2}{QR^2}$$


$$\Rightarrow QR^2 = \frac{121}{169} \times (26)^2$$

$$\Rightarrow QR = \sqrt{\frac{121}{169} \times (26)^2} = \frac{11}{13} \times 26 = 22 \text{ cm}$$

Question: 5

$\triangle ABC \sim \triangle DEF$ and th

Solution:



Let the two triangles ABC and DEF have their altitudes as AS and DT.

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding altitudes.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{100}{49} = \frac{AS^2}{DT^2} = \frac{5^2}{DT^2}$$


$$\Rightarrow DT^2 = \frac{49}{100} \times (5)^2$$

$$\Rightarrow DT = \sqrt{\frac{49}{100} \times (5)^2} = \frac{7}{10} \times 5 = 3.5 \text{ cm}$$

Question: 6

The corresponding

Solution:



Let the two triangles ABC and DEF have their altitudes as AS and DT.

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding altitudes.


$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{AS^2}{DT^2} = \frac{6^2}{9^2}$$

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{36}{81} = \frac{4}{9}$$

Question: 7

The areas of two

Solution:



Let the two triangles ABC and DEF have their altitudes as AS and DT.

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding altitudes.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{81}{49} = \frac{AS^2}{DT^2} = \frac{(6.3)^2}{DT^2}$$


$$\Rightarrow DT^2 = \frac{49}{81} \times (6.3)^2$$

$$\Rightarrow DT = \sqrt{\frac{49}{81} \times (6.3)^2} = \frac{7}{9} \times 6.3 = 4.9 \text{ cm}$$

Question: 8

The areas of two

Solution:



Let the two triangles ABC and DEF have their medians as AS and DT.

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding medians.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DEF)} = \frac{100}{64} = \frac{AS^2}{DT^2} = \frac{AS^2}{(5.6)^2}$$

$$\Rightarrow AS^2 = \frac{100}{64} \times (5.6)^2$$

$$\Rightarrow DT = \sqrt{\frac{100}{64} \times (5.6)^2} = \frac{10}{8} \times 5.6 = 7 \text{ cm}$$

Question: 9

In the given figu

Solution:

We have

$$\frac{AP}{AB} = \frac{1}{4} \text{ and } \frac{AQ}{AC} = \frac{1.5}{6} = \frac{1}{4}$$

Also $\angle A = \angle A$

So, by SAS similarity criterion $\triangle APQ \sim \triangle ABC$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle APQ)}{\text{ar}(\triangle ABC)} = \frac{AP^2}{AB^2} = \frac{1^2}{4^2} = \frac{1}{16}$$

$$\Rightarrow \text{ar}(\triangle APQ) = \frac{1}{16} \times \text{ar}(\triangle ABC)$$

Hence, proved.

Question: 10

In the given figu

Solution:

It is given that $DE \parallel BC$

$\therefore \angle ADE = \angle ABC$ (Corresponding angles)

$\angle AED = \angle ACB$ (Corresponding angles)

So, by AA similarity criterion $\triangle ADE \sim \triangle ABC$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{DE^2}{BC^2}$$

$$\Rightarrow \text{ar}(\triangle ABC) = \frac{6^2}{3^2} \times \text{ar}(\triangle ADE)$$

$$\Rightarrow \text{ar}(\triangle ABC) = 4 \times 15 = 60\text{cm}^2$$

Hence, proved.

Question: 11

$\triangle ABC$ is right-ang

Solution:

In $\triangle ABC$ and $\triangle ADC$

$\therefore \angle BAC = \angle ADC$ (90° angle)

$\angle ACB = \angle ACD$ (Common)

So, by AA similarity criterion $\triangle ADC \sim \triangle ABC$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle ADC)} = \frac{BC^2}{AC^2}$$

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle ADC)} = \frac{13^2}{5^2} = \frac{169}{25} = 169:25$$

Question: 12

In the given figu

Solution:

It is given that $DE \parallel BC$

$\therefore \angle ADE = \angle ABC$ (Corresponding angles)

$\angle AED = \angle ACB$ (Corresponding angles)

So, by AA similarity criterion $\triangle ADE \sim \triangle ABC$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{DE^2}{BC^2}$$

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{3^2}{5^2} = \frac{9}{25}$$

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(BCED)} = \frac{9}{25 - 9} = \frac{9}{16} = 9:16$$

Hence, proved.

Question: 13

In $\triangle ABC$, D and E

Solution:

In $\triangle ABC$ and $\triangle ADE$

It is given that $AD = DB$ and $AE = EC$

$$\therefore \frac{AD}{AB} = \frac{1}{2} \text{ and } \frac{AE}{AC} = \frac{1}{2}$$

Also $\angle A = \angle A$

So, by SAS similarity criterion $\triangle ADE \sim \triangle ABC$

We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding sides.

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{AE^2}{AC^2}$$

$$\Rightarrow \frac{\text{ar}(\triangle ADE)}{\text{ar}(\triangle ABC)} = \frac{1^2}{2^2} = \frac{1}{4} = 1:4$$

Exercise : 4D

Question: 1

The sides of cert

Solution:

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(i) \text{ L.H.S.} = (\text{Hypotenuse})^2 = (18)^2 = 324$$

$$\text{R.H.S.} = (\text{Base})^2 + (\text{Height})^2 = (9)^2 + (16)^2 = 81 + 256 = 337$$

$$\Rightarrow \text{L.H.S.} \neq \text{R.H.S.}$$

$\therefore$ It is not a right triangle.

$$(ii) \text{ L.H.S.} = (\text{Hypotenuse})^2 = (27)^2 = 729$$

$$\text{R.H.S.} = (\text{Base})^2 + (\text{Height})^2 = (7)^2 + (25)^2 = 49 + 625 = 674$$

$$\Rightarrow \text{L.H.S.} \neq \text{R.H.S.}$$

$\therefore$ It is not a right triangle.

$$\text{(iii) L.H.S.} = (\text{Hypotenuse})^2 = (5)^2 = 25$$

$$\text{R.H.S.} = (\text{Base})^2 + (\text{Height})^2 = (1.4)^2 + (4.8)^2 = 1.96 + 23.04 = 25$$

$$\Rightarrow \text{L.H.S.} = \text{R.H.S.}$$

$\therefore$ It is a right triangle.

$$\text{(iv) L.H.S.} = (\text{Hypotenuse})^2 = (4)^2 = 16$$

$$\text{R.H.S.} = (\text{Base})^2 + (\text{Height})^2 = (1.6)^2 + (3.8)^2 = 2.56 + 14.44 = 17$$

$$\Rightarrow \text{L.H.S.} \neq \text{R.H.S.}$$

$\therefore$ It is not a right triangle.

$$\text{(v) L.H.S.} = (\text{Hypotenuse})^2 = (a + 1)^2$$

$$\text{R.H.S.} = (\text{Base})^2 + (\text{Height})^2 = (a-1)^2 + (2\sqrt{a})^2 = a^2 + 1 - 2a + 4a = a^2 + 1 + 2a = (a + 1)^2$$


$$\Rightarrow \text{L.H.S.} = \text{R.H.S.}$$

$\therefore$ It is a right triangle.

Question: 2

A man goes 80 m d

Solution:



The starting point of the man is A and the last point is B so we need to find AB. From the figure, $\triangle ABC$ is a right triangle.

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(\text{AB})^2 = (\text{AC})^2 + (\text{BC})^2$$


$$\Rightarrow \text{AB}^2 = (80)^2 + (150)^2 = 6400 + 22500 = 28900$$

$$\Rightarrow \text{AB} = 170 \text{ m}$$

Question: 3

A man goes 10 m d

Solution:



The starting point of the man is B and the last point is A so we need to find AB. From the figure, $\triangle ABC$ is a right triangle.

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(AB)^2 = (AC)^2 + (BC)^2$$


$$\Rightarrow AB^2 = (24)^2 + (10)^2 = 576 + 100 = 676$$

$$\Rightarrow AB = 26 \text{ m}$$

Question: 4

A 13-m-long ladder

Solution:



Ladder $AB = 13 \text{ m}$ and distance from the window $BC = 12 \text{ m}$.

AC is the distance of the ladder from the building.

From the figure, $\triangle ABC$ is a right triangle.

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(AB)^2 = (AC)^2 + (BC)^2$$

$$\Rightarrow 13^2 = (AC)^2 + (12)^2$$


$$\Rightarrow AC^2 = 169 - 144 = 25$$

$$\Rightarrow AC = 5 \text{ m}$$

Question: 5

A ladder is place

Solution:



Ladder AB and distance from the window BC = 20 m.

AC is the distance of the ladder from the building = 15 m.

From the figure, $\triangle ABC$ is a right triangle.

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(AB)^2 = (AC)^2 + (BC)^2$$

$$\Rightarrow AB^2 = (20)^2 + (15)^2$$


$$\Rightarrow AB^2 = 400 + 225 = 625$$

$$\Rightarrow AB = 25 \text{ m}$$

Question: 6

Two vertical pole

Solution:



AE (height of the first building) = 14 m , CD (height of the second building) = 9 m ,
ED (distance between their feet) = BC = 12 m

$$AE - AB = 14 \text{ m} - 9 \text{ m} = 5 \text{ m}$$

From the figure, $\triangle ABC$ is a right triangle.

In a right angled triangle

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Height})^2$$

where hypotenuse is the longest side.

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$\Rightarrow AC^2 = (5)^2 + (12)^2$$


$$\Rightarrow AB^2 = 25 + 144 = 169$$

$$\Rightarrow AB = 13 \text{ m}$$

Question: 7

A guy wire attach

Solution:



Pole AB = 18 m and distance from the window BC.

AC is the length of the wire = 24 m.

From the figure, ΔABC is a right triangle.

In a right angled triangle

$$\text{(Hypotenuse)}^2 = \text{(Base)}^2 + \text{(Height)}^2$$

where hypotenuse is the longest side.

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$\Rightarrow 24^2 = (18)^2 + (BC)^2$$

$$\Rightarrow BC^2 = 576 - 324 = 252$$

$$\Rightarrow BC = 6\sqrt{7} \text{ m}$$

Question: 8

In the given figu

Solution:

ΔPOR is a right triangle because $\angle O = 90^\circ$.

In a right angled triangle

$$\text{(Hypotenuse)}^2 = \text{(Base)}^2 + \text{(Height)}^2$$

where hypotenuse is the longest side.

$$(PR)^2 = (OP)^2 + (OR)^2$$

$$\Rightarrow PR^2 = (6)^2 + (8)^2$$

$$\Rightarrow PR^2 = 36 + 64 = 100$$

$$\Rightarrow PR = 10 \text{ m}$$

$$\text{Now, } PR^2 + PQ^2 = 10^2 + 24^2 = 100 + 576 = 676$$

$$\text{Also, } QR^2 = 26^2 = 676$$

$$\Rightarrow PR^2 + PQ^2 = QR^2$$


which satisfies Pythagoras theorem.

Hence, ΔPQR is right angled triangle.

Question: 9

ΔABC is an isosce

Solution:



ΔABC is an isosceles triangle.

Also, $AB = AC = 13 \text{ cm}$

Suppose the altitude from A on BC meets BC at D. Therefore, D is the midpoint of BC.

$AD = 5 \text{ cm}$

ΔADB and ΔADC are right-angled triangles.

Applying Pythagoras theorem,

$$AB^2 = BD^2 + AD^2$$

$$\Rightarrow BD^2 = 13^2 - 5^2$$

$$\Rightarrow BD^2 = 169 - 25 = 144$$


$$\Rightarrow BD = 12 \text{ cm}$$

$$\text{So, } BC = 2 \times 12 = 24 \text{ cm}$$

Question: 10

Find the length o

Solution:



ΔABC is an isosceles triangle.

Also, $AB = AC = 2a$

The AD is the altitude. Therefore, D is the midpoint of BC.

$$BD = \frac{a}{2}$$

ΔADB and ΔADC are right-angled triangles.

Applying Pythagoras theorem,

$$AB^2 = BD^2 + AD^2$$

$$\Rightarrow (2a)^2 = \frac{a^2}{4} + AD^2$$


$$\Rightarrow AD^2 = \frac{16a^2 - a^2}{4} = \frac{15a^2}{4}$$

$$\Rightarrow AD = \frac{a\sqrt{15}}{2}$$

Question: 11

ΔABC is an equila

Solution:



ΔABC is an equilateral triangle.

Also, $BC = AB = AC = 2a$

The AD, CE, and BF are the altitude at BC, AB and AC respectively. Therefore, D, E, and F are the midpoint of BC, AB and AC respectively.

Now, ΔADB and ΔADC are right-angled triangles.

Applying Pythagoras theorem,

$$AB^2 = BD^2 + AD^2$$

$$\Rightarrow (2a)^2 = a^2 + AD^2$$

$$\Rightarrow AD^2 = 3a^2$$

$$\Rightarrow AD = a\sqrt{3} \text{ units}$$

Similarly ΔACE and ΔBEC are right-angled triangles.

Applying Pythagoras theorem,

$$CE = a\sqrt{3} \text{ units}$$

And ΔABF and ΔBFC are right-angled triangles.


Applying Pythagoras theorem,

$$BF = a\sqrt{3} \text{ units}$$

Question: 12

Find the height o

Solution:



ΔABC is an equilateral triangle.

Also, $BC = AB = AC = 12 \text{ cm}$

The AD is the altitude at BC. Therefore, D is the midpoint of BC.

Now, ΔADB and ΔADC are right-angled triangles.

Applying Pythagoras theorem,

$$AB^2 = BD^2 + AD^2$$

$$\Rightarrow (12)^2 = 6^2 + AD^2$$


$$\Rightarrow AD^2 = 144 - 36 = 108$$

$$\Rightarrow AD = 6\sqrt{3} \text{ cm}$$

Question: 13

Find the length o

Solution:



Given that $AB = 30\text{cm}$ and $AD = 16\text{ cm}$

$\therefore \angle A = 90^\circ$

$\therefore \triangle ADB$ is a right-angled triangle.

Applying Pythagoras theorem,

$$BD^2 = BA^2 + AD^2$$

$$\Rightarrow BD^2 = 30^2 + 16^2$$

$$\Rightarrow BD^2 = 900 + 256 = 1156$$

$$\Rightarrow BD = 34\text{ cm}$$


$\therefore$ Diagonals of a rectangle are equal

$\therefore AC = 34\text{ cm}$

Question: 14

Find the length o

Solution:



ABCD is a rhombus where $AC = 24\text{ cm}$ and $BD = 10\text{ cm}$.

We know that diagonals of a rhombus bisect each other at 90° .

$$\Rightarrow \angle AOB = 90^\circ, OA = 12\text{ cm and } OB = 5\text{ cm}$$

$\therefore \triangle AOB$ is a right-angled triangle.

Applying Pythagoras theorem,

$$BA^2 = BO^2 + AO^2$$

$$\Rightarrow BA^2 = 5^2 + 12^2$$

$$\Rightarrow BA^2 = 25 + 144 = 169$$

$$\Rightarrow BA = AD = CD = BC = 13\text{ cm}$$


$\therefore$ Sides of a rhombus are equal.

Question: 15

In $\triangle ABC$, D is the

Solution:

In right-angled triangle AED, applying Pythagoras theorem,



$$AB^2 = AE^2 + BE^2$$

$$\Rightarrow AE^2 = AB^2 - BE^2 \dots(i)$$

In right-angled triangle AED, applying Pythagoras theorem,

$$AD^2 = AE^2 + ED^2$$

$$\Rightarrow AE^2 = AD^2 - ED^2 \dots(ii)$$

Therefore,

$$AB^2 - BE^2 = AD^2 - ED^2$$

$$AB^2 = AD^2 - ED^2 + \left(\frac{1}{2}BC - DE\right)^2$$

$$\Rightarrow AB^2 = AD^2 - ED^2 + \frac{1}{4}BC^2 + DE^2 - BC \times DE$$

$$\Rightarrow AB^2 = AD^2 + \frac{1}{4}BC^2 - BC \times DE$$

Question: 16

In the given figu

Solution:

In $\triangle ACB$ and $\triangle CDB$,

$$\angle ABC = \angle CBD \text{ (Common)}$$

$$\angle ACB = \angle CDB (90^\circ)$$

So, by AA similarity criterion $\triangle ACB \sim \triangle CDB$

Similarly, In $\triangle ACB$ and $\triangle ADC$,

$$\angle ABC = \angle ADC \text{ (Common)}$$

$$\angle ACB = \angle ADC (90^\circ)$$

So, by AA similarity criterion $\triangle ACB \sim \triangle ADC$

We know that if two triangles are similar then the ratio of their corresponding sides is equal.

$$\Rightarrow \frac{BC}{BD} = \frac{AB}{BC} \text{ and } \frac{AC}{AD} = \frac{AB}{AC}$$

$$\Rightarrow BC^2 = AB \times BD \dots(i)$$

$$\text{And } AC^2 = AB \times AD \dots(ii)$$

Dividing (i) and (ii), we get

$$\frac{BC^2}{AC^2} = \frac{AB \times BD}{AB \times AD} = \frac{BD}{AD}$$

Hence, proved.

Question: 17

In the given figure

Solution:

(i) $\triangle AEC$ and $\triangle AED$ are right triangles.

Applying Pythagoras theorem we get,

$$AC^2 = EC^2 + AE^2$$

$$\text{And } AD^2 = ED^2 + AE^2$$

$$\Rightarrow b^2 = h^2 + \left(\frac{a}{2} + x\right)^2$$

$$\Rightarrow b^2 = h^2 + \left(\frac{a}{2}\right)^2 + x^2 + xa \dots (i)$$

$$\text{And } p^2 = h^2 + x^2 \dots (ii)$$

Putting (ii) in (i),

$$\Rightarrow b^2 = p^2 + \left(\frac{a}{2}\right)^2 + xa$$

$$\Rightarrow b^2 = p^2 + \frac{a^2}{4} + xa \dots (iii)$$

Hence, proved.

(ii) $\triangle AEB$ is a right triangle.

Applying Pythagoras theorem we get,

$$AB^2 = EB^2 + AE^2$$

$$\Rightarrow c^2 = h^2 + \left(a - \frac{a}{2} - x\right)^2$$

$$\Rightarrow c^2 = h^2 + \left(\frac{a}{2} - x\right)^2$$

$$\Rightarrow c^2 = h^2 + \left(\frac{a}{2}\right)^2 + x^2 - xa \dots (iv)$$

Putting (ii) in (iv),

$$\Rightarrow c^2 = p^2 + \left(\frac{a}{2}\right)^2 - xa$$

$$\Rightarrow c^2 = p^2 + \frac{a^2}{4} - xa \dots (v)$$

Hence, proved.

(iii) Adding (iii) and (v),

$$c^2 + b^2 = p^2 + \frac{a^2}{4} + xa + p^2 + \frac{a^2}{4} - xa$$

$$\Rightarrow c^2 + b^2 = 2p^2 + \frac{2a^2}{4}$$

$$\Rightarrow c^2 + b^2 = 2p^2 + \frac{a^2}{2}$$

Hence, proved.

(iv) Subtracting (iii) and (v),

$$b^2 - c^2 = p^2 + \frac{a^2}{4} + xa - p^2 - \frac{a^2}{4} + xa$$


$$\Rightarrow b^2 - c^2 = 2xa$$

Hence, proved.

Question: 18

In $\triangle ABC$, $AB = AC$.

Solution:



Draw $AE \perp BC$. Applying Pythagoras theorem in right-angled triangle AED,

Since, ABC is an isosceles triangle and AE is the altitude and we know that the altitude is also the median of the isosceles triangle.

So, $BE = CE$

And $DE + CE = DE + BE = BD$

$$AD^2 = AE^2 + ED^2$$

$$\Rightarrow AE^2 = AD^2 - ED^2 \dots(i)$$

In $\triangle ACE$,

$$AC^2 = AE^2 + EC^2$$

$$\Rightarrow AE^2 = AC^2 - EC^2 \dots(ii)$$

Using (i) and (ii),

$$\Rightarrow AD^2 - ED^2 = AC^2 - EC^2$$

$$\Rightarrow AD^2 - AC^2 = ED^2 - EC^2$$

$$\Rightarrow AD^2 - AC^2 = (DE + CE)(DE - CE)$$

$$\Rightarrow AD^2 - AC^2 = (DE + BE) CD$$

$$\Rightarrow AD^2 - AC^2 = BD \cdot CD$$

Question: 19

ABC is an isoscel

Solution:

$\triangle ABC$ is right triangle.

Applying Pythagoras theorem we get,

$$AC^2 = AB^2 + BC^2 \{ \because AB = BC \}$$

$$\Rightarrow AC^2 = 2AB^2$$

Given that the two triangles $\triangle ACD$ and $\triangle ABE$ are similar.


We know that if two triangles are similar then the ratio of their areas is equal to the ratio of the squares of their corresponding altitudes.

$$\Rightarrow \frac{\text{ar}(\triangle ACD)}{\text{ar}(\triangle ABE)} = \frac{AC^2}{AB^2} = \frac{AB^2}{2AB^2} = \frac{1}{2}$$

Question: 20

An aeroplane leav

Solution:



Let A be the first aeroplane flired due north at a speed of 1000 km/hr and B be the second aeroplane flired due west at a speed of 1200 km/hr

Distance covered by plane A in 1.5 hrs = $1000 \times 32 = 1500\text{km}$

Distance covered by plane B in 1.5 hrs = $1200 \times 32 = 1800\text{km}$

Now, in right triangle ABC

By using Pythagoras theorem, we have

$$AB^2 = BC^2 + AC^2$$

$$\Rightarrow AB^2 = (1800)^2 + (1500)^2$$

$$\Rightarrow AB^2 = 3240000 + 2250000$$

$$\Rightarrow AB^2 = 5490000$$

$$\Rightarrow AB = 300\sqrt{61} \text{ km}$$

Question: 21

In a ΔABC , AD is

Solution:

(a) In right triangle ALC

Using Pythagoras theorem, we have

$$AC^2 = AL^2 + LC^2$$

$$\Rightarrow AC^2 = AD^2 - DL^2 + (DL + DC)^2$$

$$\Rightarrow AC^2 = AD^2 - DL^2 + \left(DL + \frac{BC}{2}\right)^2$$

$$\Rightarrow AC^2 = AD^2 - DL^2 + DL^2 + \frac{BC^2}{4} + DL \times BC$$

$$\Rightarrow AC^2 = AD^2 + \frac{BC^2}{4} + DL \times BC \dots(1)$$

(b) In right triangle ALD

Using Pythagoras theorem, we have

$$AL^2 = AD^2 - LD^2$$

Again, in ΔABL

Using Pythagoras theorem, we have

$$AB^2 = AL^2 + LB^2$$

$$\Rightarrow AB^2 = AD^2 - DL^2 + LB^2$$

$$\Rightarrow AB^2 = AD^2 - DL^2 + (BD - DL)^2$$

$$\Rightarrow AB^2 = AD^2 - DL^2 + \left(\frac{BC}{2} - DL\right)^2$$

$$\Rightarrow AB^2 = AD^2 - DL^2 + DL^2 + \frac{BC^2}{4} - DL \times BC$$

$$\Rightarrow AB^2 = AD^2 + \frac{BC^2}{4} - DL \times BC \dots\dots(2)$$

(c) Adding (1) and (2)

$$AC^2 + AB^2 = AD^2 + \frac{BC^2}{4} - DL \times BC + AD^2 + \frac{BC^2}{4} + DL \times BC$$

$$\Rightarrow AC^2 + AB^2 = 2AD^2 + \frac{BC^2}{2}$$

Question: 22

Naman is doing fl

Solution:

Naman pulls in the string at the rate of 5 cm per second.

Hence, after 12 seconds the length of the string he will pull is given by

$$12 \times 5 = 60 \text{ cm or } 0.6 \text{ m}$$

Now, in ΔBMC

By using Pythagoras theorem, we have

$$BC^2 = CM^2 + MB^2$$

$$\Rightarrow BC^2 = (2.4)^2 + (1.8)^2 = 9$$

$$\therefore BC = 3 \text{ m}$$

$$\text{Now, } BC' = BC - 0.6 = 3 - 0.6 = 2.4 \text{ m}$$

Now, in $\Delta BC'M$

By using Pythagoras theorem, we have

$$C'M^2 = BC'^2 - MB^2$$

$$\Rightarrow C'M^2 = (2.4)^2 - (1.8)^2 = 2.52$$

$$\therefore C'M = 1.6 \text{ m}$$

The horizontal distance of the fly from him after 12 seconds is given by

$$C'A = C'M + MA = 1.6 + 1.2 = 2.8 \text{ m}$$

Exercise : 4E

Question: 1

State the two pro

Solution:

Two triangles are similar, if

(i) their corresponding angles are equal and


(ii) their corresponding sides are in the same ratio (or proportion).

Question: 2

State the basic p

Solution:

Basic Proportionality Theorem: If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio.




According to the theorem: $\frac{AD}{BD} = \frac{AE}{CE}$

Question: 3

State the convers

Solution:

Converse of Thales' Theorem: If a line divides any two sides of a triangle in the same ratio, then the line is parallel to the third side.



According to figure above, $DE \parallel BC$.

Question: 4

State the midpoin

Solution:

Midpoint Theorem: The line joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

Question: 5

State the AAA-sim

Solution:

If in two triangles, corresponding angles are equal, then their corresponding sides are in the same ratio (or proportion) and hence the two triangles are similar. This criterion is referred to as the AAA (Angle-Angle-Angle) criterion of similarity of two triangles.

Question: 6

State the AA-simi

Solution:

If two angles of one triangle are respectively equal to two angles of another triangle, then the two triangles are similar. This is referred to as the AA-similarity criterion for two triangles.

Question: 7

State the SSS-cri

Solution:

If in two triangles, sides of one triangle are proportional to (i.e., in the same ratio of) the sides of the other triangle, then their corresponding angles are equal and hence the two triangles are similar. This criterion is referred to as the SSS (Side-Side-Side)-similarity criterion for two triangles.

Question: 8

State the SAS-sim

Solution:


If one angle of a triangle is equal to one angle of the other triangle and the sides including these angles are proportional, then the two triangles are similar. This criterion is referred to as the SAS (Side-Angle-Side) similarity criterion for two triangles.

Question: 9

State Pythagoras'

Solution:

In a right angled trianglePythagoras' Theorem: In a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.



As shown in a right angled triangle ABC above,

$$(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Perpendicular})^2 \quad (AC)^2 = (BC)^2 + (AB)^2$$

Question: 10

State the convers


Solution:

Converse of Pythagoras' Theorem: In a triangle, if square of one side is equal to the sum of the squares of the other two sides, then the angle opposite the first side is a right angle.

Question: 11

If D, E and F are

Solution:



The figure is shown below:

We know that the midpoint theorem

states that the line joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

Since D, E and F are respectively the midpoints of sides AB, BC and CA of ΔABC ,

$$DE = AB/2; EF = BC/2; DF = AC/2$$

$$\Rightarrow DE/AB = 1/2; EF/BC = 1/2; DF/AC = 1/2$$

$$\Rightarrow DE/AB = EF/BC = DF/AC = 1/2$$

We know that if in two triangles, sides of one triangle are proportional to (i.e., in the same ratio of) the sides of the other triangle, then their corresponding angles are equal and hence the two triangles are similar (SSS criteria).

So $\Delta ABC \sim \Delta DEF$.

We know that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\therefore \text{ar}(\Delta ABC)/\text{ar}(\Delta DEF) = (AB/DE)^2$$

$$\Rightarrow \text{ar}(\Delta ABC)/\text{ar}(\Delta DEF) = (2DE/DE)^2$$

$$\Rightarrow \text{ar}(\Delta ABC)/\text{ar}(\Delta DEF) = (2/1)^2$$

$$\Rightarrow \text{ar}(\Delta ABC)/\text{ar}(\Delta DEF) = (4/1)$$

But we need to find the ratio of the areas of ΔDEF and ΔABC .

$$\therefore \text{ar}(\Delta DEF)/\text{ar}(\Delta ABC) = (1/4)$$

$$\therefore \text{ar}(\Delta ABC):\text{ar}(\Delta DEF) = 1:4$$

1: 4

Question: 12

Two triangles ABC

Solution:

We know that if one angle of a triangle is equal to one angle of the other triangle and the sides including these angles are proportional, then the two triangles are similar (SAS criteria).

Here in the given triangles, $\angle A = \angle P = 70^\circ$.

And $AB/PQ = AC/PR$

$$\text{i.e. } 6/4.5 = 6/9$$

$$\Rightarrow 2/3 = 2/3$$

Hence $\Delta ABC \sim \Delta PQR$.

SAS-similarity

Question: 13

If $\Delta ABC \sim \Delta DEF$ such

Solution:

Given: $\Delta ABC \sim \Delta DEF$ such that $2AB = DE$ and $BC = 6 \text{ cm}$.

From SSS-similarity criterion,

We get

$$\underline{AB/DE = BC/EF}$$

Substituting the given values,

$$AB/2AB = 6\text{cm}/EF$$

$$1/2 = 6\text{cm}/EF$$

$$EF = 2 \times 6\text{cm}$$

$$EF = 12\text{cm}$$

12cm

Question: 14

In the given figu

Solution:

We know that the basic proportionality theorem states that

"If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio."

So if $DE \parallel BC$,

Then $AD/DB = AE/EC$

By substituting the given values,

$$\Rightarrow x \text{ cm}/(3x + 4) \text{ cm} = (x + 3) \text{ cm}/(3x + 19) \text{ cm}$$

Cross multiplying, we get

$$\Rightarrow 3x^2 + 19x = 3x^2 + 9x + 4x + 12$$

$$\Rightarrow 3x^2 + 19x - 3x^2 - 9x - 4x = 12$$

$$\Rightarrow 6x = 12$$


$$\Rightarrow x = 2$$

$$\underline{x = 2}$$

Question: 15

A ladder 10 m long

Solution:



Let AB be the ladder and CA be the wall with the window at A.

Let the distance of foot of ladder from base of wall BC be x.

Also, $AB = 10\text{m}$ and $CA = 8\text{m}$

From Pythagoras Theorem,

$$\underline{\text{we have: } AB^2 = BC^2 + CA^2}$$

$$\Rightarrow (10)^2 = x^2 + 8^2$$

$$\Rightarrow x^2 = 100 - 64$$

$$\Rightarrow x^2 = 36$$

$$\Rightarrow x = 6\text{m}$$


So, $BC = 6\text{m}$.

Length of the ladder is 6m.

Question: 16

Find the length o

Solution:



Let $\triangle ABC$ be the equilateral triangle whose side is $2a$ cm.

Let us draw altitude AD such that $AD \perp BC$.

We know that altitude bisects the opposite side.

So, $BD = DC = a$ cm.

In $\triangle ADC$, $\angle ADC = 90^\circ$.

We know that the Pythagoras Theorem states that in a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

So, by applying Pythagoras Theorem,

$$AC^2 = AD^2 + DC^2$$

$$(2a \text{ cm})^2 = AD^2 + (a \text{ cm})^2$$

$$4a^2 \text{ cm}^2 = AD^2 + a^2 \text{ cm}^2$$

$$AD^2 = 3a^2 \text{ cm}^2$$

$$AD = \sqrt{3} a \text{ cm}$$

The length of altitude is $\sqrt{3} a$ cm.

Question: 17

$$\triangle ABC \sim \triangle DEF$$

Solution:

We know that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\text{i.e. } \text{ar}(\triangle ABC)/\text{ar}(\triangle DEF) = (BC/EF)^2$$

Substituting the given values, we get

$$\Rightarrow 64\text{cm}^2/169\text{cm}^2 = (4\text{cm}/EF \text{ cm})^2$$

$$\Rightarrow 64/169 = 16/EF^2$$

$$\Rightarrow EF^2 = 42.25$$


$$\Rightarrow EF = 6.5\text{cm}$$

6.5 cm

Question: 18

In a trapezium AB

Solution:



Let us consider $\triangle AOB$ and $\triangle COD$.

$$\angle AOB = \angle COD (\because \text{vertically opposite angles})$$

$$\angle OBA = \angle ODC (\because \text{alternate interior angles})$$

$$\angle OAB = \angle OCD (\because \text{alternate interior angles})$$

We know that if in two triangles, corresponding angles are equal,

then their corresponding sides are in the same ratio (or proportion) and hence the two triangles are similar (AAA criteria).

So, $\triangle AOB \cong \triangle COD$.

Given, $AB = 2CD$ and $\text{ar}(\triangle AOB) = 84 \text{ cm}^2$

We know that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\therefore \text{ar}(\triangle AOB)/\text{ar}(\triangle COD) = (AB/CD)^2$$

$$\Rightarrow 84\text{cm}^2/\text{ar}(\triangle COD) = (2CD/CD)^2$$

$$\Rightarrow 84\text{cm}^2/\text{ar}(\triangle COD) = 4$$

$$\Rightarrow \text{ar}(\triangle COD) = 84\text{cm}^2/4$$

$$\Rightarrow \text{ar}(\triangle COD) = 21\text{cm}^2$$

$$\underline{\text{ar}(\triangle COD) = 21\text{cm}^2}$$

Question: 19

The corresponding

Solution:

Let the smaller triangle be $\triangle ABC$ and larger triangle be $\triangle DEF$.

We know that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\text{i.e. } \text{ar}(\triangle ABC)/\text{ar}(\triangle DEF) = (AB/DE)^2$$

Substituting the given values, we get

$$\Rightarrow 48\text{cm}^2/\text{ar}(\triangle DEF) = (2/3)^2$$

$$\Rightarrow 48\text{cm}^2/\text{ar}(\triangle DEF) = 4/9$$

$$\Rightarrow \text{ar}(\triangle DEF) = (48 \times 9)/4 \text{ cm}^2$$


$$\Rightarrow \text{ar}(\triangle DEF) = 108\text{cm}^2$$

$$\underline{108\text{cm}^2}$$

Question: 20

In an equilateral

Solution:



Let $\triangle ABC$ be the equilateral triangle whose side is a cm.

Let us draw altitude AD(h) such that $AD \perp BC$.

We know that altitude bisects the opposite side.

So, $BD = DC = a \text{ cm}$.

In $\triangle ADC$, $\angle ADC = 90^\circ$.

We know that the Pythagoras Theorem states that in a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

So, by applying Pythagoras Theorem,

$$AC^2 = AD^2 + DC^2$$

$$(a \text{ cm})^2 = AD^2 + (a/2 \text{ cm})^2$$

$$a^2 \text{ cm}^2 = AD^2 + a^2/4 \text{ cm}^2$$

$$AD^2 = 3a^2/4 \text{ cm}^2$$

$$AD = \sqrt{3} a/2 \text{ cm} = h$$

We know that area of a triangle = $1/2 \times \text{base} \times \text{height}$

$$\text{Ar}(\triangle ABC) = 1/2 \times a \text{ cm} \times \sqrt{3} a/2 \text{ cm}$$

$$\Rightarrow \text{ar}(\triangle ABC) = \sqrt{3} a^2/4 \text{ cm}^2$$


Hence proved.

$$\underline{\text{ar}(\triangle ABC) = \sqrt{3} a^2/4 \text{ cm}^2}$$

Question: 21

Find the length o

Solution:



The diagonals of a rhombus bisect each other at right angles.

Let the intersecting point be O.

So, we get right angled triangles.

We know that the Pythagoras Theorem states that in a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

Let us consider $\triangle AOB$.

By Pythagoras Theorem,

$$AB^2 = AO^2 + OB^2$$

$$AB^2 = 12^2 + 5^2$$

$$AB^2 = 144 + 25$$

$$AB^2 = 169$$

$$AB = 13 \text{ cm}$$


The length of side of the rhombus is 13cm.

Question: 22

Two triangles DEF

Solution:

Given that $\triangle DEF \cong \triangle GHK$.



We know that if in two triangles, corresponding angles are equal, then their corresponding sides are in the same ratio (or proportion) and hence the two triangles are similar (AAA criteria).

$$\therefore \angle D = 48^\circ = \angle G$$

$$\angle H = 57^\circ = \angle E$$

$$\angle F = \angle K = x^\circ$$

We know that the sum of angles in a triangle = 180° .

So, in $\triangle DEF$,

$$\Rightarrow 48^\circ + 57^\circ + x^\circ = 180^\circ$$

$$\Rightarrow 105^\circ + x^\circ = 180^\circ$$

$$\Rightarrow x^\circ = 180^\circ - 105^\circ$$

$$\Rightarrow x^\circ = 75^\circ = \angle F$$

Ans. $\angle F = 75^\circ$

Question: 23

In the given figu

Solution:

We have $MN \parallel BC$,

So, $\angle AMN = \angle B$ and $\angle ANM = \angle C$ (Corresponding angles)

We know that if two angles of one triangle are respectively equal to two angles of another triangle, then the two triangles are similar (AA criteria).

$$\therefore \triangle AMN \sim \triangle ABC.$$

We know that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\text{i.e. } \frac{\text{ar}(\triangle AMN)}{\text{ar}(\triangle ABC)} = \left(\frac{AM}{AB}\right)^2$$

Given that $AM:MB = 1:2$.

Since $AB = AM + MB$,

$$AB = 1 + 2 = 3.$$

$$\Rightarrow \frac{\text{ar}(\triangle AMN)}{\text{ar}(\triangle ABC)} = \left(\frac{1}{3}\right)^2$$

$$\Rightarrow \frac{\text{ar}(\triangle AMN)}{\text{ar}(\triangle ABC)} = \frac{1}{9}$$

$$\underline{\text{area}(\triangle AMN) / \text{area}(\triangle ABC) = 1/9}$$

Question: 24**In triangles BMP****Solution:****Given: PB = 5 cm,****MP = 6 cm,****BM = 9 cm and,****NR = 9 cm****Now, it is also given that: $\triangle BMP \sim \triangle CNR$** **When two triangles are similar, then the ratios of the lengths of their corresponding sides are proportional.**

$$\Rightarrow \frac{BM}{CN} = \frac{BP}{CR} = \frac{MP}{NR} \dots (i)$$

$$\Rightarrow \frac{BM}{CN} = \frac{MP}{NR}$$

$$\Rightarrow CN = \frac{BM \times NR}{MP}$$

$$\Rightarrow CN = \frac{9 \text{ cm} \times 9 \text{ cm}}{6 \text{ cm}}$$

$$\Rightarrow CN = 54/6 = 13.5 \text{ cm.}$$

Similarly,


$$\Rightarrow \frac{BM}{CN} = \frac{BP}{CR}$$

$$\Rightarrow CR = \frac{BP \times CN}{BM}$$

$$\Rightarrow CR = \frac{5 \text{ cm} \times 13.5 \text{ cm}}{9 \text{ cm}}$$

$$\Rightarrow CR = 7.5 \text{ cm}$$

$$\therefore \text{Perimeter of } \triangle CNR = CN + NR + CR = 13.5 + 9 + 7.5 = 30 \text{ cm}$$

Question: 25**Each of the equal****Solution:****Let $\triangle ABC$ be the isosceles triangle whose sides are $AB = AC = 25\text{cm}$, $BC = 14\text{cm}$.****Let us draw altitude AD such that $AD \perp BC$.****We know that altitude bisects the opposite side.****So, $BD = DC = 7\text{cm}$.****In $\triangle ADC$, $\angle ADC = 90^\circ$.****We know that the Pythagoras Theorem states that in a right triangle, the square of the**

hypotenuse is equal to the sum of the squares of the other two sides.

So, by applying Pythagoras Theorem,

$$AC^2 = AD^2 + DC^2$$

$$(25 \text{ cm})^2 = AD^2 + (7 \text{ cm})^2$$

$$625 \text{ cm}^2 = AD^2 + 49 \text{ cm}^2$$

$$AD^2 = 576 \text{ cm}^2$$


$$AD = 24 \text{ cm}$$

The length of altitude is 24 cm.

Question: 26

A man goes 12 m d

Solution:



From $\triangle ABC$, we note that

A is the starting point.

$$AB = 12\text{m}, BC = 35\text{m}$$

$$CA = \text{distance from starting point} = x \text{ m}$$

We know that the Pythagoras Theorem states that in a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

By Pythagoras Theorem,

$$CA^2 = AB^2 + BC^2$$

$$CA^2 = 12^2 + 35^2$$

$$CA^2 = 144 + 1225$$

$$CA^2 = 1369$$


$$CA = 37\text{m}$$

The man is 37 m far from the starting point.

Question: 27

If the lengths of

Solution:



Given that $\triangle ABC$ is the triangle whose sides are $AB = c$, $AC = b$, $BC = a$

And AD is the bisector of $\angle A$.

We know that altitude bisects the opposite side.

So, let $BD = DC = x$.

Since AD bisects $\angle A$,

$$AC/AB = CD/DB$$

Substituting the given values,

$$b/c = CD/(a-CD)$$

Cross multiplying,

$$\Rightarrow b(a - CD) = c(CD)$$

$$\Rightarrow ba - b(CD) = c(CD)$$

$$\Rightarrow ba = CD(b + c)$$

$$\Rightarrow CD = ba/(b + c)$$

Since $CD = BD$,

$$BD = ba/(b + c)$$

$$\underline{BD = ba/(b + c) \text{ and } DC = ba/(b + c)}$$

Question: 28

In the given figu

Solution:

In $\triangle AMN$ and $\triangle ABC$

$$\angle AMN = \angle ABC = 76^\circ \text{ (Given)}$$

$$\angle A = \angle A \text{ (common)}$$

By AA Similarity criterion, $\triangle AMN \sim \triangle ABC$

If two triangles are similar, then the ratio of their corresponding sides are proportional

$$\therefore \frac{AM}{AB} = \frac{MN}{BC}$$

$$\Rightarrow \frac{AM}{AM + MB} = \frac{MN}{BC}$$


$$\Rightarrow \frac{a}{a + b} = \frac{MN}{c}$$

$$\Rightarrow MN = \frac{ac}{a + b}$$

Question: 29

The lengths of th

Solution:



The diagonals of a rhombus bisect each other at right angles.

Let the intersecting point be O.

So, we get right angled triangles.

We know that the Pythagoras Theorem states that in a right triangle, the square of the hypotenuse is equal to the sum of the squares of the other two sides.

Let us consider $\triangle AOB$.

By Pythagoras Theorem,

$$AB^2 = AO^2 + OB^2$$

$$AB^2 = 21^2 + 20^2$$

$$AB^2 = 441 + 400$$

$$AB^2 = 841$$

$$AB = 29\text{cm}$$

The length of each side of the rhombus is 29cm.

Question: 30

For each of the f

Solution:

(i) T

Two similar figures have the same shape but not necessarily the same size. Therefore, all circles are similar.

(ii) F

Two polygons of the same number of sides are similar, if (i) their corresponding angles are equal and (ii) their corresponding sides are in the same ratio (or proportion).

Consider an example,

Let a rectangle have sides 2cm and 3cm and another rectangle have sides 2cm and 5cm.

Here, the corresponding angles are equal but the corresponding sides are not in the same ratio.

(iii) F

Two triangles are similar, if

(i) their corresponding angles are equal and

(ii) their corresponding sides are in the same ratio (or proportion).

(iv) T

Midpoint Theorem states that the line joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

(v) F

Two triangles are similar, if

(i) their corresponding angles are equal and

(ii) their corresponding sides are in the same ratio (or proportion).

But here, the corresponding sides are

$$AB/DE = 6/12 = 1/2 \text{ and } AC/DF = 8/9$$

$$AB/DE \neq AC/DF$$

(vi) F

The polygon formed by joining the midpoints of sides of any quadrilateral is a parallelogram.


(vii) T

The ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

(viii) T

The perimeters of the two triangles are in the same ratio as the sides. The corresponding medians also will be in this same ratio.

(ix) T



Let us construct perpendiculars OP, OQ, OR and OS from point O.

$$\text{Let us take LHS} = OA^2 + OC^2$$

From Pythagoras theorem,

$$= (AS^2 + OS^2) + (OQ^2 + QC^2)$$

As also $AS = BQ$, $QC = DS$ and $OQ = BP = OS$,


$$= (BQ^2 + OQ^2) + (OS^2 + DC^2)$$

Again by Pythagoras theorem,

$$= OB^2 + OD^2 = \text{RHS}$$

As $\text{LHS} = \text{RHS}$, hence proved.

(x) T



In rhombus ABCD, $AB = BC = CD = DA$. We know that diagonals of a rhombus bisect each other perpendicularly, i.e. $AC \perp BD$, $\angle AOB = \angle BOC = \angle COD = \angle AOD = 90^\circ$ and $OA = OC = AC/2$, $OB = OD = BD/2$. Let us consider right angled triangle AOB.

$$\text{By Pythagoras theorem, } AB^2 = OA^2 + OB^2$$

$$\Rightarrow AB^2 = (AC/2)^2 + (BD/2)^2$$


$$\Rightarrow AB^2 = AC^2/4 + BD^2/4 \Rightarrow 4AB^2 = AC^2 + BD^2 \Rightarrow AB^2 + AB^2 + AB^2 + AB^2 = AC^2 + BD^2 \therefore AB^2 + BC^2 + CD^2 + DA^2 = AC^2 + BD^2$$

Exercise : MULTIPLE CHOICE QUESTIONS (MCQ)

Question: 1

A man goes 24 m d

Solution:



Since the man goes C to A = 24 m west and then A to B = 10 m north, he is forming a right angle triangle with respect to starting point C.

His distance from the starting point can be calculated by using Pythagoras theorem

$$(AC)^2 = (AB)^2 + (BC)^2$$

$$\Rightarrow (AC)^2 = (24)^2 + (10)^2$$

$$\Rightarrow (AC)^2 = 576 + 100$$


$$\Rightarrow (AC)^2 = 676$$

$$\Rightarrow AC = 26$$

Question: 2

Two poles of heig

Solution:



Let AB and CE be the two poles of the height 13 cm and 7 cm each which are perpendicular to the ground. The distance between them is 8 cm.

Now since CE and AB are $\perp$ ground AE

$BD \perp$ to CE and $BD = 8$ cm

Top of pole AB is B and top of pole CE is C

Now $\triangle BDC$ is right angled at D and BC , the hypotenuse is the distance between the top of the poles and $CD = 13 - 7 = 6$

$$(BC)^2 = (BD)^2 + (CD)^2$$

$$\Rightarrow (BC)^2 = 64 + 36$$

$$\Rightarrow (BC)^2 = 100$$

$$\Rightarrow (BC) = 10 \text{ cm}$$

The distance between the top of the poles is 10 cm

Question: 3


A vertical stick

Solution:

Let DF be the stick of 1.8 m height and AB be the pole of 6 m height.

AC and FE are the shadows of the pole and stick respectively.

$$FE = 45\text{cm} = .45 \text{ m}$$



Since the shadows are formed at the same time, the two $\triangle$ s are similar by AA similarity criterion

$$\text{So } \frac{AB}{DF} = \frac{AC}{FE} = \frac{BC}{DE}$$


$$\Rightarrow \frac{6}{1.8} = \frac{x}{45}$$

$$\Rightarrow x = 1.5 \text{ m}$$

Question: 4

A vertical pole 6

Solution:



Let DE be the pole of 6 m length casting shadow of 3.6 m . Let AB be the tower x meter height casting shadow of 18m at the same time.

Since pole and tower stands vertical to the ground, they form right angled triangle with ground.

ΔABC and ΔEDF are similar by AA similarity criterion

$$\therefore x/6 = 18 / 3.6$$


$$\Rightarrow x = 30$$

The height of the tower is 30 m

Question: 5

The shadow of a 5

Solution:



SINCE BOTH the tree and the stick are forming shadows at the same time the sides of the triangles so formed, would be in same ratio $\therefore$ of AA similarity criterion

$$12.5 / 5 = x / 2$$


$$\Rightarrow x = 5$$

Shadow of the tree would be 5 m long.

Question: 6

A ladder 25 m lon

Solution:



Let BC be the ladder placed against the wall AB. The distance of the ladder from the wall is the base of the right angled triangle as building stands vertically straight to the ground.

By Pythagoras theorem

$$(BC)^2 = (AB)^2 + (AC)^2$$

$$(25)^2 = (24)^2 + (x)^2$$

$$x = 7$$

the distance of ladder from the wall is 7m

Question: 7

In the given figu

Solution:

The ΔMOP is right angled at O so MP is hypotenuse

$$(MP)^2 = (OM)^2 + (OP)^2$$

$$(MP)^2 = (16)^2 + (12)^2$$

$$(MP)^2 = 400$$

$$MP = 20 \text{ cm}$$

ΔNMP is right angled at M so NP is the hypotenuse so


$$(NP)^2 = (21)^2 + (20)^2$$

$$NP = 29$$

Question: 8

The hypotenuse of

Solution:



Given $(BC) = 25 \text{ cm}$

By Pythagoras theorem

$$(BC)^2 = (AB)^2 + (AC)^2$$

$$(25)^2 = (x + 5)^2 + x^2$$

$$625 = x^2 + 25 + 10x + x^2 \because (a + b)^2 = a^2 + b^2 + 2ab$$

$$x^2 + 5x - 300 = 0$$


$$x(x + 15) - 10(x + 15) = 0$$

Since $x = -15$ is not possible so side of the triangle is 15 cm and 20 cm

Question: 9

The height of an

Solution:



Since ΔABC is an equilateral triangle so the altitude (Height = h) from the C is the median for AB dividing AB into two equal halves of 6 cm each

Now there are two right angled Δ s

$$h^2 = a^2 - \frac{1}{2} (AB)^2$$


$$h^2 = (12)^2 - 6^2$$

$$h = 6\sqrt{3}$$

Question: 10

ΔABC is an isosceles

Solution:



The given triangle is isosceles so the altitude from the one of the vertex is median for the side opposite to it.

$$AB = AC = 13 \text{ cm}$$

$$h = 5 \text{ cm (altitude)}$$

ΔABX is a right angled triangle, right angled at X

$$(AB)^2 = h^2 + (BX)^2 \quad (BX = \frac{1}{2} BC)$$

$$169 = 25 + (BX)^2$$

$$BX = 12$$

$$\Rightarrow BC = 24$$

Question: 11

In a ΔABC it is g

Solution:

By internal angle bisector theorem, the bisector of vertical angle of a triangle divides the base in the ratio of the other two sides.

Hence in ΔABC , we have

$$\frac{AB}{AC} = \frac{BD}{DC}$$

Here $AB = 6 \text{ cm}$, $AC = 8 \text{ cm}$

$$\text{So } \frac{AB}{AC} = \frac{6}{8} = \frac{3}{4} = \frac{BD}{DC}$$

Question: 12

In a $\triangle ABC$ i

Solution:

By internal angle bisector theorem, the bisector of vertical angle of a triangle divides the base in the ratio of the other two sides”Hence in $\triangle ABC$, we have

$$\frac{AB}{AC} = \frac{BD}{DC}$$

$$\frac{6}{x} = \frac{4}{5}$$

$$\Rightarrow x = 7.5\text{cm}$$

Question: 13

In a $\triangle ABC$, it is

Solution:

By internal angle bisector theorem, the bisector of vertical angle of a triangle divides the base in the ratio of the other two sides”Hence in $\triangle ABC$

$$\frac{AB}{AC} = \frac{BD}{DC}$$

$$\frac{10}{14} = \frac{(6-x)}{x}$$


$$\Rightarrow 10x - 84 + 14x = 0$$

$$\Rightarrow x = CD = 3.5 \text{ cm}$$

Question: 14

In a triangle, th

Solution:



In $\triangle ABC$, AD bisects $\angle A$ and meets BC in D such that $BD = DC$

Extend AD to E and join C to E such that CE is $\parallel$ to AB

$$\angle BAD = \angle CAD$$

Now $AB \parallel CE$ and AE is transversal

$$\angle BAD = \angle CED \text{ (alternate interior } \angle\text{s)}$$

$$\text{But } \angle BAD = \angle CED = \angle CAD$$

In $\triangle AEC$

$$\angle CEA = \angle CAE$$

$$\therefore AC = CE \dots\dots\dots 1$$

In ΔABD and ΔDCE

$\angle BAD = \angle CED$ (alternate interior $\angle$ s)

$\angle ADB = \angle CDE$ (vertically opposite $\angle$ s)

$BD = BC$ (given)

$\Delta ABD \cong \Delta DCE$

$AB = EC$ (CPCT)

$AC = EC$ (from 1)

$\Rightarrow AB = AC$

$\Rightarrow ABC$ is an isosceles Δ with $AB = AC$

Question: 15

In an equilateral

Solution:

ΔABC is an equilateral triangle

By Pythagoras theorem in triangle ABD

$$AB^2 = AD^2 + BD^2$$

but $BD = \frac{1}{2} BC$ ($\because$ In a triangle, the perpendicular from the vertex to the base bisects the base)

$$\text{thus } AB^2 = AD^2 + \left\{\frac{1}{2} BC\right\}^2$$

$$AB^2 = AD^2 + \frac{1}{4} BC^2$$

$$4 AB^2 = 4AD^2 + BC^2$$

$$4 AB^2 - BC^2 = 4 AD^2$$


(as $AB = BC$ we can subtract them)

$$\text{Thus } 3AB^2 = 4AD^2$$

Question: 16

In a rhombus of s

Solution:



Since the diagonals of the rhombus bisect each other at 90°

$$\therefore DO = OB = 6 \text{ cm}$$

$$\angle AOD = \angle DOC = \angle COB = \angle BOA = 90^\circ$$

ΔAOD is right angled Δ with

$$AD = 10 \text{ cm (given)}$$

$$OD = 6 \text{ cm}$$

$$\angle AOD = 90^\circ$$

So $DA = 10 = \text{hypotenuse}$

$$(DA)^2 = (DO)^2 + (AO)^2$$

$$100 - 36 = (AO)^2$$


$$8 = AO$$

$$\therefore AC = 16$$

Question: 17

The lengths of th

Solution:



In a rhombus the diagonals bisect each other at 90°

$$AC = 24\text{cm (given)}$$

$$\therefore AD = 12\text{cm}$$

$$BD = 10\text{cm (given)}$$

$$\therefore BO = 5\text{cm}$$

In right angled ΔAOB

$$AB^2 = AO^2 + OB^2$$

$$AB^2 = (12)^2 + 5^2$$

$$AB^2 = 144 + 25$$

$$AB^2 = 169$$


$$AB = 13\text{cm}$$

Hence the length of the sides of the rhombus is 13 cm

Question: 18

If the diagonals

Solution:



Given that ABCD is a quadrilateral and diagonals AC and BD intersect at O such that

$$\frac{AO}{OC} = \frac{OB}{OD}$$

IN ΔAOD and ΔBOC

$$\frac{AO}{OC} = \frac{BO}{OD}$$

$$\angle AOD = \angle COB$$

Thus $\Delta AOC \sim \Delta BOC$ (SAS similarity criterion)

$$\Rightarrow \angle OAD = \angle OCB \dots\dots\dots 1$$

Now transversal AC intersect AD and BC such the $\angle CAD = \angle ACB$

(from1) (alternate opposite angles)

So AD $\parallel$ BC

Hence ABCD is a trapezium

Question: 19

In the given figu

Solution:

ABCD is a Trapezium with AC and BD as diagonals and AB $\parallel$ DC

In ΔAOB and ΔDOC

$\angle AOB = \angle DOC$ (vertically opposite angles)

$\angle CDO = \angle OBA$ (alternate interior angles) (AB $\parallel$ DC and BD is transversal)

$\Delta AOB \sim \Delta DOC$ (AA similarity criterion)

$$\frac{AO}{OC} = \frac{OB}{OD}$$

$$\frac{3x - 1}{5x - 3} = \frac{2x + 1}{6x - 5}$$

$$(3x - 1)(6x - 5) = (5x - 3)(2x + 1)$$

$$18x^2 - 21x + 5 = 10x^2 - x - 3$$

$$8x^2 - 20x + 8 = 0$$

$$2x^2 - 5x + 2 = 0$$

$$2x^2 - 4x - x + 2 = 0$$

$$2x(x - 2) - 1(x - 2) = 0$$

$$(2x - 1)(x - 2) = 0$$


$$x = 1/2, 2$$

x = 1/2 is not possible so x = 2 cm

Question: 20

The line segments

Solution:



In the given quadrilateral ABCD

P, Q, R, S are the midpoints of the sides AB, BC, CD and AD respectively.

Construction: - Join AC

In $\triangle ABC$ and $\triangle ADC$

P and Q are midpoints of AB and CB

S and R are midpoints of AD and DC

So by Mid Point Theorem

$PQ \parallel AC$ and $PQ = \frac{1}{2} AC$1

And $SR \parallel AC$ and $SR = \frac{1}{2} AC$2

From 1 and 2

$PQ \parallel SR$ and $PQ = SR$


Since a pair of opposite side is equal ($=$) and parallel ($\parallel$)

PQRS is a parallelogram

Question: 21

If the bisector o

Solution:



Given in $\triangle ABC$, AD bisects the $\angle A$ meeting BC at D

$BD = DC$ and $\angle BAD = \angle CAD$ 1

Construction:- Extend BA to E and join C to E such $CE \parallel AD$ 4

$\angle BAD = \angle AEC$ (corresponding $\angle$ s)..... 2

$\angle CAD = \angle ACE$ (alternate interior $\angle$ s)..... 3

From 1 , 2 and 3

$\angle ACE = \angle AEC$

In $\triangle AEC$

$\angle ACE = \angle AEC$

$\therefore AC = AE$ (sides opposite to equal angles are equal)..... 5

In $\triangle BEC$

$AD \parallel CE$ (From4)

And D is midpoint of BC (given)

By converse of midpoint theorem

A line drawn from the midpoint of a side, parallel to the opposite side of the triangle meets the third side in its middle and is half of it

$\therefore A$ is midpoint of BE

$BA = AE$ 6

From 5 and 6

$$AB = BC$$

$\Rightarrow \triangle ABC$ is an isosceles triangle

Question: 22

In $\triangle ABC$ it is given

Solution:

It is given that in $\triangle ABC$,

$$\frac{AB}{AC} = \frac{BD}{DC}$$

$$\angle B = 70^\circ \text{ and } \angle C = 50^\circ$$

$$\angle A = 180^\circ - (70^\circ + 50^\circ) \text{ (Angle sum property of triangle)}$$

$$= 180^\circ - 120^\circ$$

$$= 60^\circ$$

$$\text{Since, } \frac{AB}{AC} = \frac{BD}{DC}$$

$\therefore AD$ is the bisector of $\angle A$

$$\text{Hence, } \angle BAD = 60^\circ / 2 = 30^\circ$$

Question: 23

In $\triangle ABC$, $DE \parallel BC$

Solution:

By Basic *Proportionality Theorem*

If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio. ..

In $\triangle ABC$, $DE \parallel BC$

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\frac{2.4}{DB} = \frac{3.2}{4.8}$$

$$DB = 3.6 \text{ cm}$$

$$AB = AD + DB$$

$$AB = 6 \text{ cm}$$

Question: 24

In $\triangle ABC$, if DE

Solution:

By Basic *Proportionality Theorem*:

If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio.

$$\frac{AD}{BD} = \frac{AE}{EC}$$

$$\text{Or } \frac{BD}{AD} = \frac{EC}{AE}$$

Adding 1 to both sides

$$\frac{BD}{AD} + 1 = \frac{EC}{AE} + 1$$

$$\frac{BD + AD}{AD} = \frac{EC + AE}{AE}$$

$$\frac{AB}{AD} = \frac{AC}{AE}$$

$$7.2 / 4.5 = 6.4 / AE$$

$$AE = 4 \text{ cm}$$

Question: 25

In $\triangle ABC$, $DE \parallel BC$

Solution:

By Basic Proportionality Theorem:

If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio.

$$\frac{AD}{BD} = \frac{AE}{EC}$$

$$(7x - 4) / (3x + 4) = (5x - 2) / 3x$$

$$3x(7x - 4) = (5x - 2)(3x + 4)$$

$$21x^2 - 12x = 15x^2 + 14x - 8$$

$$6x^2 - 26x + 8 = 0$$

$$3x^2 - 13x + 4 = 0$$

$$3x^2 - 12x - x + 4 = 0$$

$$3x(x - 4) - 1(x - 4) = 0$$

$$X = 1/3, 4$$

Since x cannot be $1/3$ so $x = 4$

Question: 26

In $\triangle ABC$, $DE \parallel BC$

Solution:

BY Basic Proportionality Theorem:

If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, the other two sides are divided in the same ratio.

$$\frac{AD}{BD} = \frac{AE}{EC}$$

$$\frac{3}{5} = \frac{AE}{AC - AE}$$

$$\frac{3}{5} = \frac{AE}{5.6 - AE}$$

$$3(5.6 - AE) = 5AE$$

$$16.8 = 8AE$$

$$AE = 2.1 \text{ cm}$$

Question: 27

$\triangle ABC \sim \triangle DEF$ and t

Solution:

Since the $\triangle ABC \sim \triangle DEF$

Their sides will be same ratios. Let the ratio be K

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = K \dots\dots\dots 1$$

$$AB + BC + AC = K (DE + EF + DF)$$

$$\frac{30}{18} = K$$

$$1.67 = k$$

From..... 1

$$\frac{BC}{EF} = 1.67$$

$$EF = 5.4 \text{ cm}$$

Question: 28

$\triangle ABC \sim \triangle DEF$ such

Solution:

Since the $\triangle ABC \sim \triangle DEF$

So the sides of the triangles are in the same ratio be k

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = K \dots\dots\dots 1$$

$$\frac{AB}{DE} = \frac{9.1}{6.5} = K$$

$$K = 1.4$$

$$\frac{\text{Perimeter of } \triangle ABC}{\text{Perimeter of } \triangle DEF} = K$$

$$\frac{\text{Perimeter of } \triangle ABC}{25} = 1.4$$

$$\text{Perimeter of } \triangle ABC = 1.4 \times 25$$

$$\text{Perimeter of } \triangle ABC = 35 \text{ cm}$$

Question: 29

In $\triangle ABC$, it is gi

Solution:

$$\text{Perimeter of } \triangle ABC = AB + BC + CA$$

$$= 9 + 6 + 7.5$$

$$= 22.5 \text{ cm}$$

Since the $\triangle ABC \sim \triangle DEF$

$$\frac{\text{Perimeter of } \triangle ABC}{\text{Perimeter of } \triangle DEF} = \frac{BC}{EF} \text{ (ratio of perimeter of triangles is equal to the ratio of the sides of the triangle)}$$

$$\frac{22.5}{\text{Perimeter of } \triangle DEF} = \frac{6}{8}$$


$$\text{Perimeter of } \triangle DEF = \frac{22.5 \times 8}{6}$$

$$\text{Perimeter of } \triangle DEF = 30 \text{ cm}$$

Question: 30

ABC and BDE are t

Solution:



ΔABC and ΔBDE are two equilateral triangles

Let a be the side of ΔABC

Since D is midpoint of BC

So the side of equilateral $\Delta BDE = \frac{a}{2}$

Area of equilateral $\Delta = \frac{\sqrt{3}}{4} (\text{side})^2$

Area of $\Delta ABC = \frac{\sqrt{3}}{4} a^2 \dots\dots\dots 1$

Area of $\Delta BDE = \frac{\sqrt{3}}{4} \left(\frac{a}{2}\right)^2$

$= \frac{\sqrt{3}}{4} \frac{a^2}{4}$

Putting value of $\frac{\sqrt{3}}{4} \times a^2 \dots\dots\dots$ from 1

Area of $\Delta BDE = \frac{1}{4}$ Area of ΔABC

$\frac{\text{Area of } \Delta ABC}{\text{Area of } \Delta BDE} = \frac{4}{1}$

Question: 31

It is given that

Solution:

In ΔABC , $\angle A + \angle B + \angle C = 180^\circ$

$30^\circ + \angle B + 50^\circ = 180^\circ$

$\angle B = 100^\circ$ Given that $\Delta ABC \sim \Delta DEF$ $\angle D = \angle A = 30^\circ$

$\angle E = \angle B = 100^\circ$ $\angle F = \angle C = 50^\circ$ Also, $\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$ $\frac{AB}{DE} = \frac{AC}{DF}$ $\frac{5}{DE} = \frac{8}{7.5}$ $DE = 4.6875$ And as

neither BC nor EF is given we can not find either of them. So, none of the given options is correct. Now, $\Delta ABC \sim \Delta DFE$, Then, $\angle D = \angle A = 30^\circ$ $\angle F = \angle B = 100^\circ$ $\angle E = \angle C =$

50° Also, $\frac{AB}{DF} = \frac{BC}{FE} = \frac{AC}{DE}$ $\frac{AB}{DF} = \frac{AC}{DE}$ $\frac{5}{DF} = \frac{8}{7.5}$ $DF = 12\text{cm}$ Therefore, the correct option is (b).

Question: 32

In the given figure

Solution:

$\angle ABD + \angle BAD = 90^\circ$

$\angle ABD + (90 - \angle CAD) = 90^\circ$ $\angle ABD = \angle CAD$ In ΔBDA and ΔADC , $\angle ABD = \angle CAD$ $\angle BDA = \angle ADC = 90^\circ$ Therefore, ΔBDA and ΔADC are similar by AAA. $\frac{BD}{AD} = \frac{AD}{CD}$ $BD \cdot CD =$

AD^2 Therefore the correct option is (c).

Question: 33

In $\triangle ABC$, $AB = 6\sqrt{3}$

Solution:

$$AB = 6\sqrt{3}\text{cm. In } \triangle ABC, AB^2 + BC^2 = AC^2 \quad (6\sqrt{3})^2 + (6)^2 = 12^2$$

Since the square of the longest side is equal to the sum of the squares of the remaining two sides of $\triangle ABC$. Therefore $\triangle ABC$ is right angled at B.

$$\angle B = 90^\circ$$

Question: 34

In $\triangle ABC$ and $\triangle DEF$

Solution:

$$\frac{AB}{DE} = \frac{BC}{FD} = \frac{AC}{EF} \quad \text{With the ratio given, we can observe that } \triangle ABC \sim \triangle EDF, \angle A = \angle E, \angle B = \angle D, \angle C = \angle F$$

Question: 35

In $\triangle DEF$ and $\triangle PQR$,

Solution:

Given $\angle D = \angle Q$ and $\angle E = \angle R$ By AA similarity, $\triangle DEF \sim \triangle QRP$ $\frac{DE}{QR} = \frac{EF}{RP} = \frac{DF}{QP}$ We have to find the option which is not true.

Question: 36

If $\triangle ABC \sim \triangle EDF$ and

Solution:

$$\triangle ABC \sim \triangle EDF, \text{ Therefore, } \frac{AB}{DE} = \frac{BC}{DF} = \frac{AC}{EF}$$

We have to find the option which is not true. $\therefore$ The correct option is (c) .

Question: 37

In $\triangle ABC$ and $\triangle DEF$,

Solution:

$\triangle ABC \sim \triangle DEF$ By AA similarity, the triangles are similar

For triangles to be congruent, $AB = DE$, but given that $AB = 3DE$.

Question: 38

If in $\triangle ABC$ and $\triangle P$

Solution:

$$\text{Given } \frac{AB}{QR} = \frac{BC}{PR} = \frac{CA}{PQ} \quad \text{Therefore } \triangle ABC \sim \triangle QRP \text{ or } \triangle PQR \sim \triangle CAB.$$

Question: 39

In the given figure

Solution:

$$\text{In } \triangle APB \text{ and } \triangle DPC, \angle APB = \angle DPC = 50^\circ \quad \frac{AP}{PB} = \frac{6}{3} = 2$$

$$\frac{PD}{PC} = \frac{5}{2.5} = 2 \quad \text{By SAS property, } \triangle APB \sim \triangle DPC \quad \angle PBA = \angle DPC \quad \text{In } \triangle DPC, \angle D + \angle P + \angle C = 180^\circ$$

$$\angle C = 100^\circ \therefore \angle PBA = \angle DPC = 100^\circ$$

Question: 40

Corresponding sides

Solution:

If two triangles are similar, the ratio of the area of triangle is equal to the square of the ratio of the sides. Ratio of Area = (Ratio of Side)² = $\left(\frac{4}{9}\right)^2 = 16:81$. $\therefore$ The correct option is (d).

Question: 41

It is given that

Solution:

If two triangles are similar, the ratio of the area of triangle is equal to the square of the ratio of the sides. $\frac{\text{Area}(\triangle PQR)}{\text{Area}(\triangle ABC)} = \left(\frac{QR}{BC}\right)^2 = \left(\frac{3}{2}\right)^2 = \frac{9}{4}$

Question: 42

In an equilateral

Solution:

Given that D and E are of AB and AC respectively, Therefore, $\frac{AD}{AB} = \frac{AE}{AC} = \frac{DE}{BC} = \frac{1}{2}$

$\triangle ABC \sim \triangle ADE$ If two triangles are similar, the ratio of the area of triangle is equal to the square of the ratio of the sides. $\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle ADE)} = \left(\frac{AB}{AD}\right)^2 = \left(\frac{2}{1}\right)^2 = \frac{4}{1}$. $\therefore$ The correct option is (b).

Question: 43

In $\triangle ABC$ and $\triangle DEF$,

Solution:

$\frac{AD}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$ Therefore $\triangle ABC \sim \triangle DEF$ If two triangles are similar, the ratio of the area of triangle is equal to the square of the ratio of the sides.

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = \left(\frac{AB}{DE}\right)^2 = \left(\frac{5}{7}\right)^2 = \frac{25}{49}$$

Question: 44

$\triangle ABC \sim \triangle DEF$ such

Solution:

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = \frac{36}{49} = \left(\frac{AB}{DE}\right)^2 \frac{AB}{DE} = \sqrt{\frac{36}{49}} = \frac{6}{7}$$

Question: 45

Two isosceles triangles

Solution:


It is given that the corresponding angles are equal, that implies that the triangles are similar. $\frac{\text{Area}(\triangle 1)}{\text{Area}(\triangle 2)} = \frac{25}{36} = \left(\frac{h_1}{h_2}\right)^2$

$$\frac{h_1}{h_2} = \sqrt{\frac{25}{36}} = \frac{5}{6}$$

Question: 46

The line segments

Solution:



In this figure, As given that the inner triangle is formed by joining the

midpoints of the sides. Therefore the outer three triangles are similar to bigger triangle. By Basic Proportionality Theorem, The inner triangle is also similar to the bigger triangle.

Question: 47

If $\Delta ABC \sim \Delta QRP$ &

Solution:

$$\Delta ABC \sim \Delta QRP \quad \frac{\text{Area}(\Delta ABC)}{\text{Area}(\Delta QRP)} = \frac{9}{4} = \left(\frac{AC}{PR}\right)^2 \frac{BC}{PR} = \sqrt{\frac{9}{4}} = \frac{3}{2}$$

$$PR = \frac{2}{3} BC \quad PR = \frac{2 \times 15}{3} = 10 \text{ cm}$$

Question: 48

In the given figure

Solution:

In ΔDOB and ΔAOC , $\angle DOB = \angle AOC = 45^\circ$ (vertically opposite angle) $\angle OAC = \angle ODB$ (angles in the same segment)

$\angle OCA = \angle OBD$ (angles in the same segment) Therefore, $\Delta DOB \sim \Delta AOC$ by AA similarity,

$$\frac{OD}{OA} = \frac{OB}{OC} = \frac{OB}{OD} = 1 \text{ Therefore, } OC = OA.$$

Question: 49

In an isosceles Δ

Solution:

$$AB^2 = 2AC^2$$

$$AB^2 = AC^2 + AC^2$$

$$AB^2 = AC^2 + BC^2$$

Therefore, it is an isosceles triangle right angled at C. $\angle C = 90^\circ$

Question: 50

In ΔABC , if $AB =$

Solution:

$$AB^2 + BC^2 = 16^2 + 12^2 = 256 + 144 = 400 = 20^2 = AC^2$$

Therefore, ABC is a right angled triangle.

Question: 51

Which of the foll

Solution:

If two triangles ABC and PQR are similar,

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{CA}{RP}$$

That is their corresponding sides are proportional.

Question: 52

Which of the foll


Solution:

The ratio of the areas of two similar triangles is equal to the ratio of squares of their corresponding sides.

Question: 53

Match the followi

Solution:



Given that $DE \parallel BC$, by B.P.T., $\frac{AD}{DB} = \frac{AE}{EC}$

Let $AE = x$

Then, from the figure, $EC = 5.6 - x$

$$\frac{AD}{DB} = \frac{x}{5.6 - x} = \frac{3}{5}$$

$$5x = 3(5.6 - x)$$

$$5x = 16.8 - 3x$$

$$8x = 16.8$$

$$x = 2.1 \text{ cm}$$

Therefore, (A)-(s)

B) As $\triangle ABC \sim \triangle DEF$, $\frac{AB}{DE} = \frac{BC}{EF} = \frac{3}{2}$

$$3EF = 2BC$$

$$3EF = 2 \times 6$$

$$EF = 4 \text{ cm}$$

Therefore, (B)-(q)


C)

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle PQR)} = \frac{9}{16} = \left(\frac{BC}{QR}\right)^2 \frac{BC}{QR} = \sqrt{\frac{9}{16}} = \frac{3}{4}$$

$$QR = \frac{4}{3}BC = \frac{4 \times 4.5}{3} = 6 \text{ cm}$$

Therefore (C)-(p)

D)



$$\frac{OA}{OB} = \frac{OC}{OD} \text{ (BPT)}$$

$$\frac{2x + 4}{9x - 21} = \frac{2x - 1}{3}$$

$$\Rightarrow 3(2x + 4) = (2x - 1)(9x - 21)$$

$$\Rightarrow 6x + 12 = 18x^2 - 42x - 9x + 21$$

$$\Rightarrow 18x^2 - 57x + 9 = 0$$

$$\Rightarrow 18x^2 - 54x - 3x + 9 = 0$$

$$\Rightarrow 18x(x - 3) - 3(x - 3) = 0$$

$$\Rightarrow (18x - 3)(x - 3) = 0$$


$$\text{So, } x = 3 \text{ or } x = \frac{1}{6}$$

But for $x = \frac{1}{6}$, $2x - 1 < 0$ which is not possible. Therefore, (D)-(r)

Question: 54

Match the followi

Solution:



The man starts from A, goes east 10m to B. From B, he goes 20m to C.

$$AC^2 = AB^2 + BC^2$$


$$AC^2 = 10^2 + 20^2$$

$$AC^2 = 100 + 400 = 500$$

$$AC = \sqrt{500} = 10\sqrt{5}$$

Therefore, (A)-(R)

B)



In $\triangle ABD$,

$$AB^2 = AD^2 + BD^2$$

$$10^2 = AD^2 + 5^2$$

$$AD^2 = 100 - 25 = 75$$

$$AD = \sqrt{75} = 5\sqrt{3}$$


Therefore, (B)-(Q)

C)

$$\text{Area of an equilateral triangle} = \frac{\sqrt{3}}{4} \times (\text{Side})^2 = \frac{\sqrt{3}}{4} \times 10^2 = 25\sqrt{3} \text{ cm}$$

Therefore, (C)-(P)

D)



In $\triangle ABC$,

$$AC^2 = AB^2 + BC^2$$

$$AC^2 = 6^2 + 8^2$$

$$AD^2 = 36 + 64 = 100$$

$$AD = \sqrt{100} = 10$$

Therefore, (D)-(S)

Exercise : FORMATIVE ASSESSMENT (UNIT TEST)

Question: 1

$\triangle ABC \sim \triangle DEF$ and t

Solution:

Given: $\triangle ABC \sim \triangle DEF$

Perimeter of $\triangle ABC = 32 \text{ cm}$

Perimeter of $\triangle DEF = 24 \text{ cm}$

$AB = 10 \text{ cm}$

To find: DE

$\therefore \triangle ABC \sim \triangle DEF$

$\therefore$ The ratio of the corresponding sides of $\triangle ABC$ and $\triangle DEF$ are equal to the ratio of the perimeter of the corresponding triangles.

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = \frac{32}{24}$$

$$\Rightarrow \frac{AB}{DE} = \frac{32}{24} \Rightarrow \frac{10}{DE} = \frac{4}{3} \Rightarrow DE = 10 \times \frac{3}{4} = \frac{30}{4} = 7.5 \text{ cm}$$

Question: 2

In the given figu

Solution:

Given: $DE \parallel BC$

$$DE = 5 \text{ cm}$$

$$BC = 8 \text{ cm}$$

$$AD = 3.5 \text{ cm}$$

To find: AB

$$\therefore DE \parallel BC$$

$\therefore$ By Basic proportionality theorem, we have

$$\frac{AD}{AB} = \frac{AE}{AC} \dots\dots\dots(i)$$

Now, in ΔADE and ΔABC , we have

$$\frac{AD}{AB} = \frac{AE}{AC} \text{ [By (i)]}$$

$$\angle DAE = \angle BAC \text{ [Common angle]}$$

$\therefore$ By SAS criterion,

$$\Delta ADE \sim \Delta ABC$$

$$\Rightarrow \frac{AD}{AB} = \frac{AE}{AC} = \frac{DE}{BC}$$

$$\Rightarrow \frac{AD}{AB} = \frac{DE}{BC}$$

$$\Rightarrow \frac{3.5}{AB} = \frac{5}{8} \Rightarrow AB = 8 \times \frac{3.5}{5} = 8 \times 0.7 = 5.6 \text{ cm}$$

Question: 3

Two poles of height


Solution:

Given: Height of pole 1 = 6 m

Height of pole 2 = 11 m

Distance between the feet of pole 1 and pole 2 = 12 m

To find: Distance between the tops of both the poles



Clearly, In ΔADE ,

$$DE = 5 \text{ m}$$

$$AD = 12 \text{ m}$$

Also, $\angle ADE = 90^\circ$ [$\because$ Both the poles stand vertically upright]

$\therefore$ By applying Pythagoras theorem, we have

$$AE^2 = AD^2 + DE^2$$

$$\Rightarrow AE^2 = (12)^2 + (5)^2 = 144 + 25 = 169$$

$$\Rightarrow AE = \sqrt{169} = 13 \text{ m}$$

Question: 4

The areas of two

Solution:

Given: Area of triangle 1 = 25 cm^2

Area of triangle 2 = 36 cm^2

Altitude of triangle 1 = 3.5 cm

To find: Altitude of triangle 2

Let the altitude of triangle 2 be x .

$\therefore$ The areas of two similar triangles are in the ratio of the squares of the corresponding altitudes.

$\therefore$ We have,

$$\frac{\text{Area of triangle 1}}{\text{Area of triangle 2}} = \frac{(\text{Altitude of triangle 1})^2}{(\text{Altitude of triangle 2})^2}$$

$$\Rightarrow \frac{25}{36} = \frac{(3.5)^2}{x^2} \Rightarrow x^2 = 12.25 \times \frac{36}{25} = 17.64$$

$$\Rightarrow x = \sqrt{17.64} = 4.2 \text{ cm}$$

Question: 5

If $\triangle ABC \sim \triangle DEF$ su

Solution:

Given: $\triangle ABC \sim \triangle DEF$

$2AB = DE$ (i)

$BC = 6 \text{ cm}$

To find: EF

$\therefore \triangle ABC \sim \triangle DEF$

$\therefore$ Ratio of all the corresponding sides of $\triangle ABC$ and $\triangle DEF$ are equal.

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} \text{(ii)}$$

Also, from (i), we have

$$2AB = DE$$

$$\Rightarrow \frac{AB}{DE} = \frac{1}{2} \text{(iii)}$$

$$\Rightarrow \frac{BC}{EF} = \frac{1}{2} \text{ [By (ii) and (iii)]}$$

$$\Rightarrow \frac{6}{EF} = \frac{1}{2} \Rightarrow EF = 6 \times 2 = 12 \text{ cm}$$

Question: 6

In the given figu

Solution:

Given: $DE \parallel BC$

$$AD = x \text{ cm}$$

$$DB = (3x + 4) \text{ cm}$$

$$AE = (x + 3) \text{ cm}$$

$$EC = (3x + 19) \text{ cm}$$

To find: x

$$\because DE \parallel BC$$

$\therefore$ By Basic Proportionality theorem, we have

$$\frac{AD}{DB} = \frac{AE}{EC} \Rightarrow \frac{x}{3x+4} = \frac{x+3}{3x+19}$$

$$\Rightarrow x(3x+19) = (x+3)(3x+4)$$

$$\Rightarrow 3x^2 + 19x = 3x^2 + 13x + 12$$

$$\Rightarrow 19x - 13x = 3x^2 + 12 - 3x^2$$

$$\Rightarrow 6x = 12 \text{ or } x = 2$$

Question: 7

A ladder 10 m long


Solution:

Given: Height of the window from the ground = 8 m

Length of the ladder = 10 m

To find: Distance of the foot of the ladder from the base of the wall.

Consider the following diagram corresponding to the question.



Here, AB = Height of the window from the ground = 8 m

AC = Length of the ladder = 10 m

BC = Distance of the foot of the ladder from the base of the wall

Now, in ΔABC ,

By Pythagoras theorem, we have

$$AC^2 = AB^2 + BC^2$$

$$\Rightarrow BC^2 = AC^2 - AB^2$$

$$\Rightarrow BC^2 = (10)^2 - (8)^2 = 100 - 64 = 36$$

$$\Rightarrow BC = \sqrt{36} = 6 \text{ m}$$


Question: 8

Find the length o

Solution:

Given: Side of equilateral triangle = $2a$ cm

To find: Length of altitude



Let ΔABC be an equilateral triangle with side $2a$ cm.

Let AD be the altitude of ΔABC .

Here, $BD = DC = a$

In ΔABD ,

Using Pythagoras theorem, we have

$$AB^2 = AD^2 + BD^2 \Rightarrow AD^2 = AB^2 - BD^2 \Rightarrow AD^2 = (2a)^2 - (a)^2 = 4a^2 - a^2 = 3a^2$$

$$\Rightarrow AD^2 = 3a^2$$

$$\Rightarrow AD = \sqrt{3a^2} = \sqrt{3} a \text{ cm}$$

Question: 9

$\Delta ABC \sim \Delta DEF$ such

Solution:

Given: $\Delta ABC \sim \Delta DEF$

$$\text{ar}(\Delta ABC) = 64 \text{ cm}^2, \text{ar}(\Delta DEF) = 169 \text{ cm}^2$$

$$BC = 4 \text{ cm}$$

To find: EF

$\therefore$ The ratios of the areas of two similar triangles are equal to the ratio of squares of any two corresponding sides.

$\therefore$ We have

$$\frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{BC^2}{EF^2}$$

$$\Rightarrow \frac{64}{169} = \frac{4^2}{EF^2} \Rightarrow EF^2 = 16 \times \frac{169}{64} = \frac{169}{4} \Rightarrow EF = \sqrt{\frac{169}{4}} = \frac{13}{2} = 6.5 \text{ cm}$$

Question: 10

In a trapezium AB


Solution:

Given: $AB \parallel CD$

$$AB = 2CD \dots\dots\dots(i)$$

$$\text{ar}(\Delta AOB) = 84 \text{ cm}^2$$

To find: $\text{ar}(\Delta COD)$



In ΔAOB and ΔCOD ,

$\angle AOB = \angle COD$ [Vertically Opposite angles]

$\angle OAB = \angle OCD$ [Alternate interior angles ($AB \parallel CD$)]

$\angle OBA = \angle ODC$ [Alternate interior angles ($AB \parallel CD$)]

$\Rightarrow \Delta AOB \sim \Delta COD$ [By AAA criterion]

Now,

$\therefore$ The ratios of the areas of two similar triangles are equal to the ratio of squares of any two corresponding sides.

$\therefore$ We have

$$\frac{\text{ar}(\Delta AOB)}{\text{ar}(\Delta COD)} = \frac{AB^2}{CD^2} = \left(\frac{AB}{CD}\right)^2 \Rightarrow \frac{84}{\text{ar}(\Delta COD)} = \left(\frac{AB}{CD}\right)^2$$

Also, from (i), we have

$$\frac{AB}{CD} = 2$$

$$\Rightarrow \frac{84}{\text{ar}(\Delta COD)} = 2^2 = 4 \Rightarrow \text{ar}(\Delta COD) = \frac{84}{4} = 21 \text{ cm}^2$$

Question: 11

The corresponding

Solution:

Given: Let the smaller triangle be ΔABC and the larger triangle be ΔDEF .

The ratio of AB and $DE = 2 : 3$

$$\Rightarrow \frac{AB}{DE} = \frac{2}{3} \dots\dots\dots(i)$$

$$\text{ar}(\Delta ABC) = 48 \text{ cm}^2$$

To find: $\text{ar}(\Delta DEF)$

$\therefore$ The ratios of the areas of two similar triangles are equal to the ratio of squares of any two corresponding sides.

$\therefore$ We have

$$\frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DEF)} = \frac{AB^2}{DE^2} \Rightarrow \frac{48}{\text{ar}(\Delta DEF)} = \frac{2^2}{3^2} = \frac{4}{9} \Rightarrow \text{ar}(\Delta DEF) = 48 \times \frac{9}{4} = 12 \times 9 = 108 \text{ cm}^2$$

Question: 12

In the given figu

Solution:

Given: $LM \parallel CB$ and $LN \parallel CD$

To prove: $\frac{AM}{AB} = \frac{AN}{AD}$

In ΔAML , $LM \parallel CB$

$\therefore$ By Basic proportionality theorem, we have

$$\frac{AM}{AB} = \frac{AL}{AC} \dots\dots\dots(i)$$

In ΔALN , $LN \parallel CD$

$\therefore$ By Basic proportionality theorem, we have

$$\frac{AL}{AC} = \frac{AN}{AD} \dots\dots\dots(ii)$$

By (i) and (ii), we have

$$\frac{AM}{AB} = \frac{AL}{AC} = \frac{AN}{AD} \Rightarrow \frac{AM}{AB} = \frac{AN}{AD}$$

Question: 13


Prove that the in

Solution:

Given: ΔABC with the internal bisector AD of $\angle A$ which intersects BC at D .

To prove: $\frac{BD}{DC} = \frac{AB}{AC}$

First, we construct a line $EC \parallel AD$ which meets BA produced in E .



Now, we have

$CE \parallel DA \Rightarrow \angle 2 = \angle 3$ [Alternate interior angles are equal (transversal AC)]

Also, $\angle 1 = \angle 4$ [Corresponding angles are equal (transversal AE)]

We know that AD bisects $\angle A \Rightarrow \angle 1 = \angle 2$

$$\Rightarrow \angle 4 = \angle 1 = \angle 2 = \angle 3$$

$$\Rightarrow \angle 3 = \angle 4$$

$$\Rightarrow AE = AC \text{ [Sides opposite to equal angles are equal] } \dots\dots\dots(i)$$

Now, consider ΔBCE ,

$AD \parallel EC$

$$\Rightarrow \frac{BD}{DC} = \frac{BA}{AE} \text{ [By Basic Proportionality theorem]}$$

$$\Rightarrow \frac{BD}{DC} = \frac{AB}{AC} \text{ [}\because BA = AB \text{ and } AE = AC \text{ (From (i))]}$$


Question: 14

In an equilateral

Solution:

Let ΔABC be an equilateral triangle with side a .

To prove: Area of $\Delta ABC = \frac{\sqrt{3}}{4} a^2$



In ΔABC , AD bisects BC

$$\Rightarrow BD = DC = \frac{a}{2}$$

Now, in ΔACD

Using Pythagoras theorem,

$$AC^2 = AD^2 + DC^2$$

$$\Rightarrow AD^2 = AC^2 - DC^2$$

$$\Rightarrow AD^2 = a^2 - \left(\frac{a}{2}\right)^2 = a^2 - \frac{a^2}{4} = \frac{4a^2 - a^2}{4} = \frac{3a^2}{4}$$

$$\Rightarrow AD = \sqrt{\frac{3a^2}{4}} = \frac{\sqrt{3}a}{2}$$

Now, in ΔABC

$$\text{Area of } \Delta ABC = \frac{1}{2} \times \text{base} \times \text{height} = \frac{1}{2} \times BC \times AD = \frac{1}{2} \times a \times \frac{\sqrt{3}a}{2} = \frac{\sqrt{3}a^2}{4}$$

Question: 15


Find the length o

Solution:

Given: Length of one of the diagonals = 24 cm

Length of the other diagonal = 10 cm

To find: Length of the side of the rhombus



$\therefore$ The length of all sides of rhombus is equal.

∴ Let side of rhombus ABCD be x cm.

Also, we know that the diagonals of a rhombus are perpendicular bisectors of each other.

$$\Rightarrow AO = OC = 12 \text{ cm and } BO = OD = 5 \text{ cm}$$

$$\text{Also, } \angle AOB = \angle BOC = \angle COD = \angle AOD = 90^\circ$$

Now, consider ΔAOD

$$AO = 12 \text{ cm and } OD = 5 \text{ cm}$$

$$\angle AOD = 90^\circ$$

So, using Pythagoras theorem, we have

$$AD^2 = AO^2 + OD^2 = 12^2 + 5^2 = 144 + 25 = 169$$

$$\Rightarrow AD = \sqrt{169} = 13 \text{ cm}$$

Question: 16

Prove that the ra

Solution:

Let ΔABC and ΔDEF be two similar triangles, i.e., $\Delta ABC \sim \Delta DEF$.

$\Rightarrow$ Ratio of all the corresponding sides of ΔABC and ΔDEF are equal.

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

Let these ratios be equal to some number α .

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = \alpha$$

$$\Rightarrow AB = \alpha DE, BC = \alpha EF, AC = \alpha DF \dots\dots\dots(i)$$

$$\text{Now, perimeter of } \Delta ABC = AB + BC + AC$$

$$= \alpha DE + \alpha EF + \alpha DF \text{ [From (i)]}$$

$$= \alpha (DE + EF + DF)$$

$$= \alpha (\text{perimeter of } \Delta DEF)$$

$$\Rightarrow \frac{\text{Perimeter of } \Delta ABC}{\text{Perimeter of } \Delta DEF} = \alpha$$

$$\Rightarrow \frac{\text{Perimeter of } \Delta ABC}{\text{Perimeter of } \Delta DEF} = \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

Question: 17

In the given figu


Solution:

Given: ΔABC and ΔDBC have the same base BC.

AD and BC intersect at O.

$$\text{To show: } \frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DBC)} = \frac{AO}{DO}$$

First, we construct the altitudes, AE and DF, of ΔABC and ΔDBC , respectively.



Consider, ΔAOE and ΔDOF ,

$\angle DFO = \angle AEO$ [Right angles]

$\angle DOF = \angle AOE$ [Vertically Opposite angles]

So, by AA criterion,

$\Delta AOE \sim \Delta DOF$

$\Rightarrow$ Ratio of all the corresponding sides of ΔAOE and ΔDOF are equal.

$$\Rightarrow \frac{AO}{DO} = \frac{AE}{DF} \dots\dots\dots(i)$$

Now, we know that

$$\text{Area of triangle} = \frac{1}{2} \times \text{base} \times \text{height}$$

$$\Rightarrow \text{Area of } \Delta ABC = \text{ar}(\Delta ABC) = \frac{1}{2} \times BC \times AE \dots\dots\dots(ii)$$

$$\text{Similarly, Area of } \Delta DBC = \text{ar}(\Delta DBC) = \frac{1}{2} \times BC \times DF \dots\dots\dots(iii)$$

Dividing (ii) by (iii),

$$\frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DBC)} = \frac{\frac{1}{2} \times BC \times AE}{\frac{1}{2} \times BC \times DF} = \frac{AE}{DF}$$

$$\Rightarrow \frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DBC)} = \frac{AE}{DF}$$

$$\Rightarrow \frac{\text{ar}(\Delta ABC)}{\text{ar}(\Delta DBC)} = \frac{AO}{DO} \text{ [From (i)]}$$

Question: 18

In the given figu

Solution:

Given: $XY \parallel AC$

$$\text{ar}(\Delta XBY) = \text{ar}(\Delta XCY) \dots\dots\dots(i)$$

$$\text{To show: } \frac{AX}{AB} = \frac{2-\sqrt{2}}{2}$$

Consider ΔABC , $XY \parallel AC$

So, Using Basic Proportionality theorem, we have

$$\frac{XB}{AB} = \frac{YC}{CB} \dots\dots\dots(ii)$$

Now, in ΔXBY and ΔABC ,

$$\angle XBY = \angle ABC \text{ [common angle]}$$

$$\frac{XB}{AB} = \frac{YB}{CB} \text{ [Using (ii)]}$$

$$\Rightarrow \Delta XBY \sim \Delta ABC \text{ [By SAS criterion]}$$

Now, we know that the ratios of the areas of two similar triangles are equal to the ratio of squares of any two corresponding sides.

$$\Rightarrow \frac{\text{ar}(\Delta XBY)}{\text{ar}(\Delta ABC)} = \frac{XB^2}{AB^2}$$

From (i), we have

$$\text{ar}(\Delta XBY) = \text{ar}(\Delta XCY)$$

$$\text{Let ar}(\Delta XBY) = x = \text{ar}(\Delta XCY) \Rightarrow \text{ar}(\Delta ABC) = \text{ar}(\Delta XBY) + \text{ar}(\Delta XCY) = x + x = 2x$$

$$\Rightarrow \frac{\text{ar}(\Delta XBY)}{\text{ar}(\Delta ABC)} = \frac{x}{2x} = \frac{1}{2}$$

$$\Rightarrow \frac{\text{ar}(\Delta XBY)}{\text{ar}(\Delta ABC)} = \frac{XB^2}{AB^2}$$

$$\Rightarrow \frac{XB^2}{AB^2} = \frac{1}{2} \Rightarrow \frac{XB}{AB} = \sqrt{\frac{1}{2}} \Rightarrow \frac{XB}{AB} = \frac{1}{\sqrt{2}}$$

Now, we know that

$$XB = AB - AX$$

$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{XB}{AB} = \frac{AB - AX}{AB} \Rightarrow \frac{AB - AX}{AB} = \frac{1}{\sqrt{2}} \Rightarrow \frac{AB}{AB} - \frac{AX}{AB} = \frac{1}{\sqrt{2}} \Rightarrow 1 - \frac{AX}{AB} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \frac{AX}{AB} = 1 - \frac{1}{\sqrt{2}} = \frac{\sqrt{2} - 1}{\sqrt{2}}$$

Rationalizing the denominator, we have

$$\frac{AX}{AB} = \frac{\sqrt{2} - 1}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{2 - \sqrt{2}}{2}$$

$$\Rightarrow \frac{AX}{AB} = \frac{2 - \sqrt{2}}{2}$$

Question: 19

In the given figure

Solution:

Given: $AD \perp CB$ (produced)

To prove: $AC^2 = AB^2 + BC^2 + 2BC \cdot BD$

In ΔADC , $DC = DB + BC$ (i)

First, in ΔADB ,

Using Pythagoras theorem, we have

$$AB^2 = AD^2 + DB^2 \Rightarrow AD^2 = AB^2 - DB^2 \text{(ii)}$$

Now, applying Pythagoras theorem in ΔADC , we have

$$AC^2 = AD^2 + DC^2$$

$$= (AB^2 - DB^2) + DC^2 \text{ [Using (ii)]}$$

$$= AB^2 - DB^2 + (DB + BC)^2 \text{ [Using (i)]}$$

Now, $\because (a + b)^2 = a^2 + b^2 + 2ab$

$\therefore AC^2 = AB^2 - DB^2 + DB^2 + BC^2 + 2DB \cdot BC$

$\Rightarrow AC^2 = AB^2 + BC^2 + 2BC \cdot BD$

Question: 20

In the given figure

Solution:

Given: $PA \perp AC$, $QB \perp AC$ and $RC \perp AC$

$AP = x$, $QB = z$, $RC = y$, $AB = a$ and $BC = b$

To show: $\frac{1}{x} + \frac{1}{y} = \frac{1}{z}$

In ΔPAC , we have

$QB \parallel PA$

So, by Basic Proportionality theorem, we have

$$\frac{PC}{QC} = \frac{AC}{BC} \dots\dots\dots(i)$$

In ΔARC , we have

$QB \parallel RC$

So, by Basic Proportionality theorem, we have

$$\frac{AR}{AQ} = \frac{AC}{AB} \dots\dots\dots(ii)$$

Now, Consider ΔPAC and ΔQBC ,

$\angle PCA = \angle QCB$ [Common angle]

$$\frac{PC}{QC} = \frac{AC}{BC} \text{ [By (i)]}$$

So, by SAS criterion,

$\Delta PAC \sim \Delta QBC$

$\Rightarrow$ Ratio of all the corresponding sides of ΔABC and ΔDEF are equal.

$$\Rightarrow \frac{QB}{PA} = \frac{BC}{AC}$$

$$\Rightarrow \frac{z}{x} = \frac{b}{a+b} \dots\dots\dots(iii)$$

Now, consider ΔARC and ΔAQB ,

$\angle RAC = \angle QAB$ [Common angle]

$$\frac{AR}{AQ} = \frac{AC}{AB} \text{ [By (ii)]}$$

So, by SAS criterion,

$\Delta ARC \sim \Delta AQB$

$\Rightarrow$ Ratio of all the corresponding sides of ΔARC and ΔAQB are equal.

$$\Rightarrow \frac{QB}{RC} = \frac{AB}{AC}$$

$$\Rightarrow \frac{z}{y} = \frac{a}{a+b} \dots\dots\dots(iv)$$

Now, adding (iii) and (iv), we get

$$\frac{z}{x} + \frac{z}{y} = \frac{b}{a+b} + \frac{a}{a+b}$$

$$\Rightarrow z \left(\frac{1}{x} + \frac{1}{y} \right) = \frac{b+a}{a+b} = 1$$

$$\Rightarrow \frac{1}{x} + \frac{1}{y} = \frac{1}{z}$$