

Topics : Method of Differentiation, Continuity & Derivability, Application of Derivatives, Sequence & Series, Straight Line

Type of Questions	M.M., Min.
Single choice Objective (no negative marking) Q.1,2,3,4,5,6 (3 marks, 3 min.)	[18, 18]
Multiple choice objective (no negative marking) Q.7 (5 marks, 4 min.)	[5, 4]
Subjective Questions (no negative marking) Q.8 (4 marks, 5 min.)	[4, 5]

- If $f(x) = \max \{\sin x, \sin^{-1}(\cos x)\}$, then

(A) f is differentiable everywhere (B) f is continuous but not differentiable everywhere

(C) f is discontinuous at $x = \frac{n\pi}{2}, n \in \mathbb{I}$ (D) none of these
- The radius of a right circular cylinder increases at a constant rate. Its altitude is a linear function of the radius and increases three times as fast as radius. When the radius is 1 cm the altitude is 6 cm. When the radius is 6 cm, the volume is increasing at the rate of 1 Cu cm/sec. When the radius is 36 cm, the volume is increasing at a rate of n cu. cm/sec. The value of ' n ' is equal to :

(A) 12 (B) 22 (C) 30 (D) 33
- If $y = (A + Bx)e^{mx} + (m-1)^{-2} \cdot e^x$, then $\frac{d^2y}{dx^2} - 2m \frac{dy}{dx} + m^2y$ is equal to

(A) e^{mx} (B) e^{-mx} (C) $e^{(1-m)x}$ (D) e^x
- If $\tan y = \frac{2^x}{1+2^{2x+1}}$, then $\frac{dy}{dx}$ at $x = 0$ is

(A) $-\frac{3}{10}$ (B) $-\frac{3}{10} \ln 2$ (C) $-\frac{1}{10}$ (D) $-\frac{1}{10} \ln 2$
- If $y = \sin x$, then $\frac{d^2(\cos^7 x)}{dx^2}$ is equal to

(A) $35 \cos^3 x - 42 \cos^5 x$ (B) $35 \cos^3 x + 42 \cos^5 x$

(C) $42 \cos^3 x - 35 \cos^5 x$ (D) $-35 \cos^3 x - 42 \cos^5 x$
- If $2a + 3b + c = 3$; $a > 0, b > 0, c > 0$, then the greatest value of $a^2 b^5 c^2$

(A) $\frac{5^5 \cdot 2^2}{3^{23}}$ (B) $\frac{5^5 \cdot 2^2}{3^{14}}$ (C) $\frac{4 \cdot 5^5}{9^9}$ (D) $\frac{5^6 \cdot 2^2}{3^4 \cdot 9^{10}}$
- The function $f(x) = (\tan^{-1}x)^3 - (\cot^{-1}x)^2 + \tan^{-1}x + 2$ is

(A) decreasing $\forall x \in \mathbb{R}$. (B) Increasing $\forall x \in \mathbb{R}$.

(C) Bounded (D) Many one function.
- The tangent to $y = ax^2 + bx + \frac{7}{2}$ at $(1, 2)$ is parallel to the normal at the point $(-2, 2)$ on the curve $y = x^2 + 6x + 10$. Find the value of a and b .

Answers Key

1. (B) 2. (D) 3. (D) 4. (D)

5. (A) 6. (B) 7. (B)(C) 8. $a = 1, b = -\frac{5}{2}$