

AREA OF PARALLELOGRAM AND TRIANGLES

4

CHAPTER

▶ IMPORTANT POINTS

- ◆ Parallelograms on the same base and between the same parallels are equal in area.
- ◆ Area of a parallelogram is the product of its any side and the corresponding altitude.
- ◆ Parallelogram on the same base and having equal areas lie between the same parallels.
- ◆ If a parallelogram and a triangle are on the same base and between the same parallels, then area of the triangle, is half the area of the parallelogram
- ◆ Two congruent figures having same area.

❖ EXAMPLES ❖

Ex.1 ABCD is a quadrilateral and BD is one of its diagonals as shown in fig. Show that ABCD is a parallelogram and find its area.

Sol. Since diagonal BD intersects transversals AB and DC at B and D respectively such that

$$\angle ABD = \angle CDB \quad [\text{Each equal to } 90^\circ]$$

i.e., alternate interior angles are equal.


$$\therefore AB \parallel DC$$

Also, $AB = DC$ [Each equal to 5.2 cms (Given)]

Thus, one pair of opposite sides AB and DC of quadrilateral ABCD are equal and parallel.

Hence, ABCD is a parallelogram.

Now



$$\begin{aligned} \text{ar} (\parallel^{\text{gm}} ABCD) &= \text{Base} \times \text{Corresponding altitude} \\ &= AB \times BD = 5.2 \times 4 \text{ sq.cm} \end{aligned}$$


Ex.2 In parallelogram ABCD, $AB = 10$ cm. The altitudes corresponding to the sides AB and AD are respectively 7 cm and 8 cm. Find AD.

Sol. We have,

$$\text{Area of a } \parallel^{\text{gm}} = \text{Base} \times \text{Height.}$$

$$\begin{aligned} \therefore \text{ar} (\parallel^{\text{gm}} ABCD) &= AB \times DM \\ &= (10 \times 7) \text{ cm}^2 \quad \dots (i) \end{aligned}$$

$$\begin{aligned} \text{Also, ar} (\parallel^{\text{gm}} ABCD) &= AD \times BN \\ &= (AD \times 8) \text{ cm}^2 \quad \dots (ii) \end{aligned}$$




From (i) and (ii), we get

$$10 \times 7 = AD \times 8$$

$$\Rightarrow AD = \frac{10 \times 7}{8} \text{ cm} = 8.75 \text{ cm.}$$

Ex.3 In the adjoining figure, ABCD is a $\parallel^{\text{gm}}$ whose diagonals AC and BD intersect at O. A line segment through O meets AB at P and DC at Q. Prove that $\text{ar} (\triangle APQD) = \frac{1}{2} \text{ar} (\parallel^{\text{gm}} ABCD)$.



Sol. Diagonal AC of ||gm ABCD divides it into two triangles of equal area.

$$\therefore \text{ar}(\triangle ACD) = \frac{1}{2} \text{ar}(\text{||gm ABCD}) \dots (i)$$

In $\triangle OAP$ and $\triangle OCQ$, we have

$$OA = OC$$

[diagonals of a ||gm bisect each other]

$$\angle AOP = \angle COQ \text{ [vert. opp. } \angle]$$

$$\angle PAO = \angle QCO \text{ [alt. int. } \angle]$$

$$\therefore \triangle OAP \cong \triangle OCQ$$

$$\therefore \text{ar}(\triangle OAP) = \text{ar}(\triangle OCQ)$$

$$\begin{aligned} \Rightarrow \text{ar}(\triangle OAP) + \text{ar}(\text{quad. AOQD}) \\ = \text{ar}(\triangle OCQ) + \text{ar}(\text{quad. AOQD}) \end{aligned}$$

$$\begin{aligned} \Rightarrow \text{ar}(\text{quad. APQD}) &= \text{ar}(\triangle ACD) \\ &= \frac{1}{2} \text{ar}(\text{||gm ABCD}) \text{ [using (i)]} \end{aligned}$$

$$\therefore \text{ar}(\triangle APQD) = \frac{1}{2} \text{ar}(\text{||gm ABCD})$$

- ◆ Triangles on the same base and between the same parallels are equal in area.
- ◆ The area of a triangle is half the product of any of its sides and the corresponding altitude.
- ◆ If a triangle and a parallelogram are on the same base and between the same parallels, the area of the triangle is equal to half of the parallelogram.
- ◆ The area of a trapezium is half the product of its height and the sum of parallel sides.
- ◆ Triangles having equal areas and having one side of one of the triangles equal to one side of the other, have their corresponding altitudes equal.


Ex.4 Show that a median of a triangle divides it into two triangles of equal area.

Sol. Given : A $\triangle ABC$ in which AD is the median.

To Prove $\text{ar}(\triangle ABD) = \text{ar}(\triangle ADC)$

Construction : Draw $AL \perp BC$.

Proof Since AD is the median of $\triangle ABC$. Therefore,



D is the mid-point of BC.

$$\Rightarrow BD = DC$$

$$\Rightarrow BD \times AL = DC \times AL$$

[Multiplying both sides by AL]


$$\Rightarrow \frac{1}{2} (BD \times AL) = \frac{1}{2} (DC \times AL)$$

$$\Rightarrow \text{ar}(\triangle ABD) = \text{ar}(\triangle ADC)$$

ALITER Since $\triangle s$ ABD and ADC have equal bases and the same altitude AL. Therefore, $\text{ar}(\triangle ABD) = \text{ar}(\triangle ADC)$.

Ex.5 In figure, AD is a median of $\triangle ABC$ and DE is a median of $\triangle DAC$. Show that

$$\text{ar}(\triangle AED) = \frac{1}{4} \text{ar}(\triangle ABC)$$



Sol. AD is a median of $\triangle ABC$

$$\Rightarrow \text{ar}(\triangle ABD) = \text{ar}(\triangle ADC)$$

$$\Rightarrow \text{ar}(\triangle ADC) = \frac{1}{2} \text{ar}(\triangle ABC) \dots (1)$$

DE is a median of $\triangle DAC$

$$\Rightarrow \text{ar}(\triangle AED) = \frac{1}{2} \text{ar}(\triangle ADC) \dots (2)$$

From (1) and (2),


$$\text{ar}(\triangle AED) = \frac{1}{2} \left\{ \frac{1}{2} \text{ar}(\triangle ABC) \right\} = \frac{1}{4} \text{ar}(\triangle ABC)$$

Ex.6 The diagonals of ABCD, AC and BD intersect in O. Prove that if $BO = OD$, the triangles ABC and ADC are equal in area.

Sol. Given : A quadrilateral ABCD in which its diagonals AC and BD intersect at O such that $BO = OD$.

To Prove : $\text{ar}(\triangle ABC) = \text{ar}(\triangle ADC)$

Proof : In $\triangle ABD$, we have $BO = OD$.
[Given]



$\Rightarrow$ O is the mid-point of BD

$\Rightarrow$ AO is the median

$\Rightarrow \text{ar}(\triangle AOB) = \text{ar}(\triangle AOD) \quad \dots(i)$

⊙ Median divides a $\triangle$ into two $\triangle$ s of equal area

In $\triangle CBD$, O is the mid-point of BD.

$\therefore$ CO is a median


$\Rightarrow \text{ar}(\triangle COB) = \text{ar}(\triangle COD) \quad \dots(ii)$

Adding (i) and (ii), we get

$$\begin{aligned} \text{ar}(\triangle AOB) + \text{ar}(\triangle COB) &= \text{ar}(\triangle AOD) \\ &+ \text{ar}(\triangle COD) \end{aligned}$$

$\Rightarrow \text{ar}(\triangle ABC) = \text{ar}(\triangle ADC)$.

Ex.7 Let P, Q, R, S be respectively the midpoints of the sides AB, BC, CD and DA of quad. ABCD. Show that PQRS is a parallelogram such that $\text{ar}(\text{||gm PQRS}) = \frac{1}{2} \text{ar}(\text{quad. ABCD})$.



Sol. Join AC and AR.

In $\triangle ABC$, P and Q are midpoints of AB and BC respectively.

$\therefore PQ \parallel AC$ and $PQ = \frac{1}{2} AC$

In $\triangle DAC$, S and R are midpoints of AD and DC respectively.

$\therefore SR \parallel AC$ and $SR = \frac{1}{2} AC$

Thus, $PQ \parallel SR$ and $PQ = SR$.

$\therefore$ PQRS is a ||gm.

Now, median AR divides $\triangle ACD$ into two $\triangle$ of equal area.

$\therefore \text{ar}(\triangle ARD) = \frac{1}{2} \text{ar}(\triangle ACD) \quad \dots(1)$

Median RS divides $\triangle ARD$ into two $\triangle$ of equal area.

$\therefore \text{ar}(\triangle DSR) = \frac{1}{2} \text{ar}(\triangle ARD) \quad \dots(2)$

From (1) and (2), we get

$$\text{ar}(\triangle DSR) = \frac{1}{4} \text{ar}(\triangle ACD)$$

Similarly, $\text{ar}(\triangle BQP) = \frac{1}{4} \text{ar}(\triangle ABC)$

$\Rightarrow \text{ar}(\triangle DSR) + \text{ar}(\triangle BQP)$

$$= \frac{1}{4} [\text{ar}(\triangle ACD) + \text{ar}(\triangle ABC)]$$

$\Rightarrow \text{ar}(\triangle DSR) + \text{ar}(\triangle BQP)$

$$= \frac{1}{4} \text{ar}(\text{quad. ABCD}) \quad \dots(3)$$

Similarly, $\text{ar}(\triangle CRQ) + \text{ar}(\triangle ASP)$

$$= \frac{1}{4} \text{ar}(\text{quad. ABCD}) \quad \dots(4)$$

Adding (3) and (4), we get

$$\begin{aligned} \text{ar}(\triangle DSR) + \text{ar}(\triangle BQP) + \text{ar}(\triangle CRQ) \\ + \text{ar}(\triangle ASP) \end{aligned}$$

$$= \frac{1}{2} \text{ar}(\text{quad. ABCD}) \quad \dots(5)$$

But, $\text{ar}(\triangle DSR) + \text{ar}(\triangle BQP) + \text{ar}(\triangle CRQ)$

$$+ \text{ar}(\triangle ASP) + \text{ar}(\text{||gm PQRS})$$


$$= \text{ar}(\text{quad. ABCD}) \quad \dots(6)$$

Subtracting (5) from (6), we get

$$\text{ar}(\text{||gm PQRS}) = \frac{1}{2} \text{ar}(\text{quad. ABCD})$$

Ex.8 The medians BE and CF of a triangle ABC intersect at G. Prove that area of ΔGBC = area of quadrilateral AFGE.

Sol. Join EF. Since the line segment Joining the mid-points of two sides of a triangle is parallel to the third side. So, $EF \parallel BC$.



Clearly, Δs BEF and CEF are on the same base EF and between the same parallel lines. So,

$$\text{ar}(\Delta BEF) = \text{ar}(\Delta CEF)$$

$$\Rightarrow \text{ar}(\Delta BEF) - \text{ar}(\Delta GEF)$$

$$= \text{ar}(\Delta CEF) - \text{ar}(\Delta GEF)$$

$$\Rightarrow \text{ar}(\Delta BFG) = \text{ar}(\Delta CEG) \quad \dots (i)$$

We know that a median of a triangle divides it into two triangles of equal area.

Therefore,

$$\text{ar}(\Delta BEC) = \text{ar}(\Delta ABE)$$

$$\Rightarrow \text{ar}(\Delta BGC) + \text{ar}(\Delta CEG) = \text{ar}(\text{quad. AFGE}) + \text{ar}(\Delta BFG)$$

$$\Rightarrow \text{ar}(\Delta BGC) + \text{ar}(\Delta BFG) = \text{ar}(\text{quad. AFGE}) + \text{ar}(\Delta BFG) \quad [\text{Using (i)}]$$

$$\Rightarrow \text{ar}(\Delta BGC) = \text{ar}(\text{quad. AFGE})$$


Ex.9 E, F, G, H are respectively, the mid-points of the sides AB, BC, CD and DA of parallelogram ABCD. Show that the area of quadrilateral EFGH is half the area of the parallelogram ABCD.

Sol. Given: A quadrilateral ABCD in which E, F, G, H are respectively the mid-points of the sides AB, BC, CD and DA.

To Prove :

$$(i) \text{ar}(\text{||}^{\text{gm}} \text{EFGH}) = \frac{1}{2} \text{ar}(\text{||}^{\text{gm}} \text{ABCD})$$

Construction : Join AC and HF



Since ΔHGF and $\text{||}^{\text{gm}} \text{HDCF}$ are on the same base HF and between the same parallel lines.

$$\therefore \text{ar}(\Delta HGF) = \frac{1}{2} \text{ar}(\text{||}^{\text{gm}} \text{HDCF}) \quad \dots (i)$$

Similarly, ΔHEF and $\text{||}^{\text{gm}} \text{HABF}$ are on the same base HF and between the same parallels.

$$\therefore \text{ar}(\Delta HEF) = \frac{1}{2} \text{ar}(\text{||}^{\text{gm}} \text{HABF}) \quad \dots (ii)$$

Adding (iii) and (iv), we get

$$\begin{aligned} \text{ar}(\Delta HGF) + \text{ar}(\Delta HEF) &= \frac{1}{2} [\text{ar}(\text{||}^{\text{gm}} \text{HDCF}) \\ &\quad + \text{ar}(\text{||}^{\text{gm}} \text{HABF})] \end{aligned}$$


$$\Rightarrow \text{ar}(\text{||}^{\text{gm}} \text{EFGH}) = \frac{1}{2} \text{ar}(\text{||}^{\text{gm}} \text{ABCD}).$$

Ex.10 Two segments AC and BD bisect each other at O. Prove that ABCD is a parallelogram.

Sol. Given: AC and BD are two segments bisecting each other at O.

To Prove: ABCD is a parallelogram.

Construction : Join AB, BC, CD and DA.



Proof : In Δs AOB and COD, we have

$$AO = CO \quad [\text{Given}]$$

$$BO = DO \quad [\text{Given}]$$

$$\text{and, } \angle AOB = \angle COD \quad [\text{Vertically opp. } \angle s]$$

So, by SAS criterion of congruence

$$\Delta AOB \cong \Delta COD$$

$$\Rightarrow AB = CD \quad [\because \text{Corresponding parts of congruent triangles are equal}]$$

$$\text{and, } \angle 1 = \angle 2.$$

Thus, AB and DC intersect AC at A and C respectively such that $\angle 1 = \angle 2$ i.e. alternate interior angles are equal.

$$\therefore AB \parallel DC$$

Thus, in quadrilateral ABCD, we have

$$AB = DC \text{ and } AB \parallel DC$$

i.e. a pair of opposite sides are equal and parallel. Hence, ABCD is a parallelogram.

Hence, ABCD is a parallelogram.

Ex.11 ABCD is a parallelogram. L and M are points on AB and DC respectively and AL = CM. Prove that LM and BD bisect each other.

Sol. We have, AL = CM


$$\Rightarrow AB - BL = CD - DM$$

$$\Rightarrow -BL = -DM \quad [\ominus \text{ ABCD is a } \parallel^{\text{gm}}]$$

$$\therefore AB = DC]$$

$$\Rightarrow BL = DM \quad \dots (i)$$

Now, AB $\parallel$ DC and transversals BD and LM intersect them.



$$\therefore \angle 3 = \angle 4 \text{ and } \angle 1 = \angle 2 \quad \dots (ii)$$

Thus, in Δ s OBL and ODM, we have

$$\angle 1 = \angle 2 \quad [\text{From (ii)}]$$

$$BL = MD \quad [\text{From (i)}]$$

$$\angle 3 = \angle 4 \quad [\text{From (ii)}]$$

So, by ASA criterion of congruence

$$\Delta OBL \cong \Delta ODM$$

$$\Rightarrow OB = OD \text{ and } OL = OM$$

$$\left[\begin{array}{l} \ominus \text{ Corresponding parts of} \\ \text{congruent triangles are equal} \end{array} \right]$$

$$\Rightarrow O \text{ is the mid-point of BD and LM both.}$$

$$\Rightarrow BD \text{ and LM bisect each other.}$$

Ex.12 A point O inside a rectangle ABCD is joined to the vertices. Prove that the sum of the areas of a pair of opposite triangles so formed is equal to the sum of the other pair of triangles.

Sol. Given : A rectangle ABCD and O is a point inside it, OA, OB, OC and OD have been joined.

$$\text{To Prove : } \ar(\Delta AOD) + \ar(\Delta BOC)$$

$$= \ar(\Delta AOB) + \ar(\Delta COD).$$

Construction : Draw EOF $\parallel$ AB and LOM $\parallel$ AD.


Proof : We have,

$$\ar(\Delta AOD) + \ar(\Delta BOC)$$

$$= \frac{1}{2} (AD \times OE) + \frac{1}{2} (BC \times OF)$$

$$= \frac{1}{2} (AD \times OE) + \frac{1}{2} (AD \times OF) \quad [\because AD = BC]$$

$$= \frac{1}{2} (AD \times (OE + OF))$$



$$= \frac{1}{2} (AD \times EF)$$

$$= \frac{1}{2} (AD \times AB) \quad [\because EF = AB]$$

$$= \frac{1}{2} \ar(\text{rect ABCD}) \quad \text{and,}$$

$$\ar(\Delta AOB) + \ar(\Delta COD)$$

$$= \frac{1}{2} (AB \times OL) + \frac{1}{2} (CD \times OM) \quad \frac{1}{2}$$

$$= (AB \times OL) + \frac{1}{2} (AB \times OM) \quad [\ominus AB = CD]$$

$$= \frac{1}{2} AB \times (OL + OM)$$

$$= \frac{1}{2} (AB \times LM) \quad [\ominus LM = AD]$$

$$\frac{1}{2} = (AB \times AD) = \frac{1}{2} \ar(\text{rect ABCD})$$

$$\therefore \ar(\Delta AOD) + \ar(\Delta BOC)$$


$$= \ar(\Delta AOB) + \ar(\Delta COD)$$

Ex.13 #ABCD is a rhombus and P, Q, R, S are the mid-points of AB, BC, CD, DA respectively. Prove that #PQRS is a rectangle.

Sol. Given: A rhombus #ABCD in which P, Q, R, S are the mid-points of sides AB, BC, CD and DA respectively. PQ, QR, RS and SP are joined.

To Prove: #PQRS is rectangle.

Construction : Join AC.



Proof : In order to prove that PQRS is a rectangle, it is sufficient to show that it is a parallelogram whose one angle is a right angle. First we shall prove that PQRS is parallelogram.

In $\triangle ABC$, P and Q are the mid-points of AB and BC respectively.

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC \quad \dots(i)$$

In $\triangle ADC$, R and S are the mid-points of CD and AD respectively.

$$\therefore RS \parallel AC \text{ and } RS = \frac{1}{2} AC \quad \dots(ii)$$

From (i) and (ii), we have

$$PQ \parallel RS \text{ and } PQ = RS$$

Thus, PQRS is a quadrilateral such that one pair of opposite sides PQ and SR is equal and parallel.

So, PQRS is a parallelogram.

Now, we shall prove that one angle of parallelogram PQRS is a right angle.

ABCD is a rhombus

$$\Rightarrow AB = BC$$

[Θ All sides of a rhombus are equal]

$$\Rightarrow \frac{1}{2} AB = \frac{1}{2} BC$$

$$\Rightarrow PB = BQ \quad \dots(i)$$

$$\left[\begin{array}{l} \Theta \text{ P and Q are the mid - points} \\ \text{of AB and BC respectively} \end{array} \right]$$

Now, in $\triangle PBQ$ we have,

$$PB = BQ$$

$$\Rightarrow \angle 1 = \angle 2 \quad \dots(ii)$$

[Angles opposite to
equal sides are equal]

Again, ABCD is a rhombus

$$\Rightarrow AB = BC = CD = AD$$

$$\Rightarrow AB = BC, CD = AD$$

$$\Rightarrow \frac{1}{2} AB = \frac{1}{2} BC, \frac{1}{2} CD = \frac{1}{2} AD$$

$$\Rightarrow AP = CQ, CR = AS \quad \dots(iii)$$

Now, in $\triangle APS$ and $\triangle CQR$, we have

$$AP = CQ \quad [\text{From (iii)}]$$

$$AS = CR \quad [\text{From (iii)}]$$

and $PS = QR$ [Θ PQRS is $\parallel^{\text{gm}}$ $\therefore PS = QR$]

So, by SSS criterion of congruence

$$\triangle APS \cong \triangle CQR$$

$$\Rightarrow \angle 3 = \angle 4 \quad \dots(iv)$$

[Θ Corresponding parts of
congruent triangles are equal]

$$\text{Now, } \angle 3 + \angle SPQ + \angle 2 = 180^\circ$$

$$\text{and, } \angle 1 + \angle PQR + \angle 4 = 180^\circ$$

$$\therefore \angle 3 + \angle SPQ + \angle 2 = \angle 1 + \angle PQR + \angle 4$$

$$\Rightarrow \angle SPQ = \angle PQR \quad \dots(v)$$

[$\Theta \angle 1 = \angle 2$, from (ii)
and $\angle 3 = \angle 4$, from (iv)]

Now, transversal PQ cuts parallel lines SP and RQ at P and Q respectively.

$$\therefore \angle SPQ + \angle PQR = 180^\circ$$

$$\Rightarrow \angle SPQ + \angle SPQ = 180^\circ \quad [\text{Using (v)}]$$

$$\Rightarrow \angle SPQ = 90^\circ$$

Thus, PQRS is a parallelogram such that $\angle SPQ = 90^\circ$.

Hence, PQRS is a rectangle.

Ex.14 Triangles ABC and DBC are on the same base BC with A, D on opposite sides of line BC, such that $\text{ar}(\triangle ABC) = \text{ar}(\triangle DBC)$. Show that BC bisects AD.

Sol. Since $\triangle ABC$ and $\triangle DBC$ are equal in area and have a common side BC. Therefore the altitudes corresponding to BC are equal i.e.


$$AE = DF.$$

Now, in $\triangle AEO$ and $\triangle DFO$, we have

$$\angle 1 = \angle 2 \quad [\text{Vertically opp. angles}]$$

$$\angle AEO = \angle DFO \quad [\text{Each equal to } 90^\circ]$$

and, $AE = DF$



So, by AAS criterion of congruence,

$$\triangle AEO \cong \triangle DFO$$

$$\Rightarrow AO = DO$$

$$\Rightarrow BC \text{ bisects } AD.$$

Ex.15 ABCD is a parallelogram and O is any point in its interior. Prove that :

$$(i) \ar(\triangle AOB) + \ar(\triangle COD) = \frac{1}{2} \ar(\text{gm ABCD})$$

$$(ii) \ar(\triangle AOB) + \ar(\triangle COD) = \ar(\triangle BOC) + \ar(\triangle AOD)$$

Sol. Given: A parallelogram ABCD and O is a point in its interior.

- (i) Since $\triangle AOB$ and parallelogram ABFE are on the same base AB and between the same parallel lines AB and EF.

$$\therefore \ar(\triangle AOB) = \frac{1}{2} \ar(\text{gm ABFE}) \quad \dots(i)$$

Similarly,

$$\ar(\triangle COD) = \frac{1}{2} \ar(\text{gm DEFC}) \quad \dots(ii)$$

Adding (i) and (ii), we get


$$\ar(\triangle AOB) + \ar(\triangle COD)$$

$$= \frac{1}{2} \ar(\text{gm ABCD})$$

- (ii) To Prove: $\ar(\triangle AOB) + \ar(\triangle COD)$

$$= \ar(\triangle BOC) + \ar(\triangle AOD).$$

Construction: Draw $EOF \parallel AB$ and $GOH \parallel AD$.



Proof: Since $GH \parallel DE$ and $EF \parallel DC$

$$\therefore OG \parallel DE \text{ and } OE \parallel GD$$

$\Rightarrow$ EOGD is a parallelogram

Similarly, EAHO, HBFO and FOGC are parallelograms.

Now, OD is a diagonal of parallelogram EOGD

$$\Rightarrow \ar(\triangle EOD) = \ar(\triangle DOG) \quad \dots (iii)$$

OA is a diagonal of parallelogram EAHO

$$\Rightarrow \ar(\triangle EOA) = \ar(\triangle AOH) \quad \dots(iv)$$

OB is a diagonal of parallelogram HBFO

$$\Rightarrow \ar(\triangle BOF) = \ar(\triangle BOH) \quad \dots(v)$$

OC is a diagonal of parallelogram FOGC

$$\Rightarrow \ar(\triangle FOC) = \ar(\triangle COG)$$

Adding (iii), (iv) and (v), we get

$$\ar(\triangle EOD) + \ar(\triangle EOA) + \ar(\triangle BOF) + \ar(\triangle FOC)$$

$$= \ar(\triangle DOG) + \ar(\triangle AOH) + \ar(\triangle BOH)$$

$$+ \ar(\triangle COG)$$

$$\Rightarrow \ar(\triangle AOD) + \ar(\triangle BOC)$$

$$= \ar(\triangle AOB) + \ar(\triangle COD)$$

Ex.16 A quadrilateral ABCD is such that diagonal BD divides its area in two equal parts. Prove that BD bisects AC.

Sol. Given: A quadrilateral ABCD in which diagonal BD bisects it, i.e.

$$\ar(\triangle ABD) = \ar(\triangle BDC)$$

Construction: Join AC.


Suppose AC and BD intersect at O. Draw $AL \perp BD$ and $CM \perp BD$.

To Prove : $AO = OC$.

Proof: We have, $\ar(\triangle ABD) = \ar(\triangle BDC)$

Thus, $\triangle ABD$ and $\triangle BDC$ are on the same base AB and have equal area. Therefore, their corresponding altitudes are equal.

i.e., $AL = CM$



Now, in Δ s ALO and CMO, we have

$$\angle 1 = \angle 2 \quad [\text{Vertically opposite angles}]$$

$$\angle ALO = \angle CMO \quad [\text{Each equal to } 90^\circ]$$


and, $AL = CM$ [Proved above]

So, by AAS criterion of congruence

$$\Delta ALO \cong \Delta CMO$$

$$\Rightarrow AO = OC \Rightarrow BD \text{ bisects } AC.$$

Ex.17 In Fig. ABCD is a trapezium in which side AB is parallel to side DC and E is the mid-point of side AD. If F is a point on the side BC such that the segment EF is parallel to side DC.



Prove that $EF = \frac{1}{2} (AB + DC)$.

Sol. Given : A trapezium ABCD in which $AB \parallel DC$, E is the mid-point of AD and F is a point on BC such that $EF \parallel DC$.

To Prove: $EF = \frac{1}{2} (AB + DC)$

Proof: In ΔADC , E is the mid-point of AD and $EG \parallel DC$ (Given)

$\therefore$ G is the mid-point of AC

Since segment joining the mid-points of two sides of a triangle is half of the third side.

$$\therefore EG = \frac{1}{2} DC \quad \dots(i)$$

Now, ABCD is a trapezium in which $AB \parallel DC$.

But, $EF \parallel DC$

$\therefore EF \parallel AB$

$\Rightarrow GF \parallel AB$

In ΔABC , G is the mid-point of AC (proved above) and $EF \parallel AB$.

$\therefore$ F is the mid-point of BC

$$\Rightarrow GF = \frac{1}{2} AB \quad \dots(ii)$$

[$\therefore$ Segment joining the mid-points of two sides of a Δ is half of the third sides]

From (i) and (ii), we have


$$EG + GF = \frac{1}{2} (DC) + \frac{1}{2} (AB)$$

$$\Rightarrow EF = \frac{1}{2} (AB + DC)$$

Ex.18 In ΔABC , D is the mid-point of AB. P is any point of BC. $CQ \parallel PD$ meets AB in Q. Show that $\text{ar}(\Delta BPQ) = \frac{1}{2} \text{ar}(\Delta ABC)$.

Sol. To Prove: $\text{ar}(\Delta BPQ) = \frac{1}{2} \text{ar}(\Delta ABC)$

Construction: Join CD.



Proof: Since D is the mid-point of AB. So, in ΔABC , CD is the median.

$$\text{ar}(\Delta BCD) = \frac{1}{2} \text{ar}(\Delta ABC) \quad \dots(i)$$

Since Δ s PDQ and PDC are on the same base PD and between the same parallel lines PD and QC.

$$\therefore \text{ar}(\Delta PDQ) = \text{ar}(\Delta PDC) \quad \dots(ii)$$

Now, from (i)

$$\text{ar}(\Delta BCD) = \frac{1}{2} \text{ar}(\Delta ABC)$$

$$\Rightarrow \text{ar}(\Delta BPD) + \text{ar}(\Delta PDC) = \frac{1}{2} \text{ar}(\Delta ABC)$$

$$\Rightarrow \text{ar}(\Delta BPD) + \text{ar}(\Delta PDQ) = \frac{1}{2} \text{ar}(\Delta ABC)$$

[Using (ii)]

$$\Rightarrow \text{ar}(\Delta BPQ) = \frac{1}{2} \text{ar}(\Delta ABC)$$

Ex.19 If the medians of a ΔABC intersect at G, show that $\text{ar}(\Delta AGB) = \text{ar}(\Delta AGC)$
 $= \text{ar}(\Delta BGC) = \frac{1}{3} \text{ar}(\Delta ABC)$.

Sol. Given: A ΔABC such that its medians AD, BE and CF intersect at G.

To Prove : $\text{ar}(\Delta AGB) = \text{ar}(\Delta BGC)$
 $= \text{ar}(\Delta CGA) = \frac{1}{3} \text{ar}(\Delta ABC)$

Proof : We know that the median of a triangle divides it into two triangles of equal area.

In ΔABC , AD is the median

$$\Rightarrow \text{ar}(\Delta ABD) = \text{ar}(\Delta ACD) \quad \dots (i)$$

In ΔGBC , GD is the median


$$\Rightarrow \text{ar}(\Delta GBD) = \text{ar}(\Delta GCD) \quad \dots (ii)$$

Subtracting (ii) from (i), we get

$$\text{ar}(\Delta ABD) - \text{ar}(\Delta GBD) = \text{ar}(\Delta ACD) - \text{ar}(\Delta GCD)$$

$$\Rightarrow \text{ar}(\Delta AGB) = \text{ar}(\Delta AGC) \quad \dots (iii)$$

Similarly,



$$\text{ar}(\Delta AGB) = \text{ar}(\Delta BGC) \quad \dots (iv)$$

From (iii) and (iv), we get

$$\text{ar}(\Delta AGB) = \text{ar}(\Delta BGC) = \text{ar}(\Delta AGC) \quad \dots (v)$$

$$\text{But, } \text{ar}(\Delta AGB) + \text{ar}(\Delta BGC) + \text{ar}(\Delta AGC) = \text{ar}(\Delta ABC)$$


$$\therefore 3 \text{ar}(\Delta AGB) = \text{ar}(\Delta ABC)$$

$$\Rightarrow \text{ar}(\Delta AGB) = \frac{1}{3} \text{ar}(\Delta ABC)$$

$$\text{Hence, } \text{ar}(\Delta AGB) = \text{ar}(\Delta AGC) = \text{ar}(\Delta BGC) = \frac{1}{3} \text{ar}(\Delta ABC).$$

Ex.20 In a parallelogram ABCD, E, F are any two point on the sides AB and BC respectively. Show that $\text{ar}(\Delta ADF) = \text{ar}(\Delta DCE)$.

Sol. Construction: Draw EG $\parallel$ AD and FH $\parallel$ AB.



Proof: Since FH $\parallel$ AB (by construction). Therefore, ABFH is a parallelogram.

Now, AF is a diagonal of $\parallel^{\text{gm}}$ ABFH

$$\therefore \text{ar}(\Delta AFH) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{ABFH}) \quad \dots (i)$$

In $\parallel^{\text{gm}}$ DCFH, DF is a diagonal.

$$\therefore \text{ar}(\Delta DFH) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{DCFH}) \quad \dots (ii)$$

From (i) and (ii), we have

$$\begin{aligned} \text{ar}(\Delta AFH) + \text{ar}(\Delta DFH) &= \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{ABFH}) + \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{DCFH}) \\ \Rightarrow \text{ar}(\Delta AFH) + \text{ar}(\Delta DFH) &= \frac{1}{2} [\text{ar}(\parallel^{\text{gm}} \text{ABFH}) + \text{ar}(\parallel^{\text{gm}} \text{DCFH})] \end{aligned}$$

$$\Rightarrow \text{ar}(\Delta AFH) + \text{ar}(\Delta DFH) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{ABCD})$$

$$\Rightarrow \text{ar}(\Delta ADF) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{ABCD}) \quad \dots (iii)$$

In $\parallel^{\text{gm}}$ AEGD, DE is a diagonal.

$$\therefore \text{ar}(\Delta DEG) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{AEGD}) \quad \dots (iv)$$

In $\parallel^{\text{gm}}$ CBEG, CE is a diagonal.

$$\therefore \text{ar}(\Delta CEG) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{CBEG}) \quad \dots (v)$$

From (iv) and (v), we have

$$\begin{aligned} \text{ar}(\Delta DEG) + \text{ar}(\Delta CEG) &= \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{AEGD}) + \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{CBEG}) \\ \Rightarrow \text{ar}(\Delta DEG) + \text{ar}(\Delta CEG) &= \frac{1}{2} [\text{ar}(\parallel^{\text{gm}} \text{AEGD}) + \text{ar}(\parallel^{\text{gm}} \text{CBEG})] \\ \Rightarrow \text{ar}(\Delta DEG) + \text{ar}(\Delta CEG) &= \frac{1}{2} \text{ar}(\parallel^{\text{gm}} \text{ABCD}) \end{aligned}$$

$$\Rightarrow \text{ar}(\Delta DCE) = \frac{1}{2} \text{ar}(\text{||gm } ABCD) \quad \dots(\text{vi})$$

From (iii) and (vi), we get

$$\text{ar}(\Delta ADF) = \text{ar}(\Delta DCE).$$


Ex.21 In Fig. PQRS is a parallelogram, PQ and QO are respectively, the angle bisectors of $\angle P$ and $\angle Q$. Line LOM is drawn parallel to PQ. Prove that :

$$(i) \text{ PL} = \text{QM} \quad (ii) \text{ LO} = \text{OM}.$$

Sol. Since PQRS is a parallelogram.

$$\therefore \text{PS} \parallel \text{QR} \Rightarrow \text{PL} \parallel \text{QM}$$

Thus, we have



$$\text{PL} \parallel \text{QM} \text{ and } \text{LM} \parallel \text{PQ} \quad [\text{Given}]$$

$$\Rightarrow \text{PQML is parallelogram.}$$

$$\Rightarrow \text{PL} = \text{QM} \quad [\Theta \text{ Opp. sides of a || gm are equal}]$$

This proves (i).

Now, OP is the bisector of $\angle P$

$$\therefore \angle 1 = \angle 2 \quad \dots(\text{i})$$

Now, PQ $\parallel$ LM and transversal OP intersects them

$$\therefore \angle 1 = \angle 3 \quad \dots(\text{ii})$$

$$\text{From (i) and (ii), we get} \\ \angle 2 = \angle 3$$

Thus, in ΔOPL , we have

$$\angle 2 = \angle 3$$

$$\Rightarrow \text{OL} = \text{PL} \quad \dots(\text{iii})$$

$$\left[\begin{array}{l} \Theta \text{ Opposite sides of equal} \\ \text{angles in triangle are equal} \end{array} \right]$$

Since OQ is the bisector of $\angle Q$

$$\therefore \angle 4 = \angle 5 \quad \dots(\text{iv})$$

Also, PQ $\parallel$ LM and transversal OQ intersects them

$$\angle 4 = \angle 6 \quad \dots(\text{v})$$

From (iv) and (v), we get

$$\angle 5 = \angle 6$$

Thus, in ΔOQM , we have

$$\angle 5 = \angle 6$$

$$\Rightarrow \text{OM} = \text{QM} \quad \dots(\text{vi})$$

[Θ Opp. sides of equal angles are equal]

$$\text{But, PL} = \text{QM} \quad \dots(\text{vii}) \quad [\text{As proved above}]$$


So, from (iii), (vi) and (vii), we get

$$\text{OL} = \text{OM}.$$

Ex.22 The diagonals of a quadrilateral ABCD are perpendicular. Show that the quadrilateral, formed by joining the mid-points of its sides, is a rectangle.

Sol. Given: A quadrilateral whose diagonals AC and BD are perpendicular to each other, P, Q, R, S are the mid-points of sides AB, BC, CD and DA respectively. PQ, QR, RS and SP are joined.

To Prove: PQRS is a rectangle.



Proof: In ΔABC , P and Q are the mid-points of AB and BC respectively.

$$\therefore \text{PQ} \parallel \text{AC} \text{ and } \text{PQ} = \frac{1}{2} \text{AC} \quad \dots(\text{i})$$

In ΔADC , R and S are the mid-points of CD and AD respectively.

$$\therefore \text{RS} \parallel \text{AC} \text{ and } \text{RS} = \frac{1}{2} \text{AC} \quad \dots(\text{ii})$$

From (i) and (ii), we have

$$\text{PQ} \parallel \text{RS} \text{ and } \text{PQ} = \text{RS}$$

Thus, in quadrilateral PQRS, a pair of opposite sides are equal and parallel. So, PQRS is a parallelogram.

Suppose the diagonals AC and BD of quadrilateral ABCD intersect at O.

Now in ΔABD , P is the mid-point of AB and S is the mid-point of AD.

$$\therefore PS \parallel BD \Rightarrow PN \parallel MO$$

Also, from (i), $PQ \parallel AC$

$$\Rightarrow PM \parallel NO$$

Thus, in quadrilateral PMON, we have

$$PN \parallel MO \text{ and } PM \parallel NO$$

$\Rightarrow$ PMON is a parallelogram.

$$\Rightarrow \angle MPN = \angle MON$$

[Θ Opposite angles of a $\parallel^{\text{gm}}$ are equal]

$$\Rightarrow \angle MPN = \angle BOA \quad [\Theta \angle BOA = \angle MON]$$

$$\Rightarrow \angle MPN = 90^\circ \quad [\Theta AC \perp BD \therefore \angle BOA = 90^\circ]$$

$$\Rightarrow \angle QPS = 90^\circ \quad [\Theta \angle MPN = \angle QPS]$$

Thus, PQRS is a parallelogram whose one angle $\angle QPS = 90^\circ$ Hence PQRS is a rectangle.

Ex.23 In a parallelogram ABCD diagonals AC and BD intersect at O and $AC = 6.8\text{cm}$ and $BD = 13.6\text{ cm}$. Find the measures of OC and CD.

Sol. Since the diagonals of a parallelogram bisect each other. Therefore, O is the mid-point of AC and BD.

$$\therefore OC = \frac{1}{2} AC = \frac{1}{2} \times 6.8\text{ cm} = 3.4\text{ cm}$$


$$\text{and, } OD = \frac{1}{2} BD = \frac{1}{2} \times 5.6\text{ cm} = 2.8\text{ cm}$$

Ex.24 Prove that the figure formed by joining the mid-points of the pairs of consecutive sides of a quadrilateral is a parallelogram.

Sol. Given: ABCD is a quadrilateral in which P, Q, R and S are the mid-points of sides AB, BC, CD and DA respectively.

To Prove: PQRS is a parallelogram.

Construction: Join A and C.



Proof: In $\triangle ABC$, P and Q are the mid-points of sides AB and BC respectively.

$$\therefore PQ \parallel AC \text{ and } PQ = \frac{1}{2} AC \quad \dots (i)$$

In $\triangle ADC$, R and S are the mid-points of CD and AD respectively.

$$\therefore RS \parallel AC \text{ and } RS = \frac{1}{2} AC \quad \dots (ii)$$

From (i) and (ii), we have

$$PQ = RS \text{ and } PQ \parallel RS$$


Thus, in quadrilateral PQRS one pair of opposite sides are equal and parallel.

Hence, PQRS is a parallelogram.

Ex.25 The side AB of a parallelogram ABCD is produced to any point P. A line through A parallel to CF meets CB produced in Q and the parallelogram PBQR completed.

Show that $\text{ar}(\parallel^{\text{gm}} ABCD) = \text{ar}(\parallel^{\text{gm}} BPRQ)$.

Sol. Construction: Join AC and PQ.



To prove: $\text{ar}(\parallel^{\text{gm}} ABCD) = \text{ar}(\parallel^{\text{gm}} BPRQ)$

Proof: Since AC and PQ are diagonals of parallelograms ABCD and BPQR respectively.

$$\therefore \text{ar}(\triangle ABC) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} ABCD) \quad \dots (i)$$

$$\text{and, } \text{ar}(\triangle PBQ) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} BPRQ) \quad \dots (ii)$$

Now, $\triangle ACQ$ and $\triangle AQP$ are on the same base AQ and between the same parallels AQ and CP

$$\therefore \text{ar}(\triangle ACQ) = \text{ar}(\triangle AQP)$$

$$\Rightarrow \text{ar}(\triangle ACQ) - \text{ar}(\triangle ABQ)$$

$$= \text{ar}(\triangle AQP) - \text{ar}(\triangle ABQ)$$

[Subtracting $\text{ar}(\triangle ABQ)$ from both sides]

$$\Rightarrow \text{ar}(\triangle ABC) = \text{ar}(\triangle BPQ)$$

$$\Rightarrow \frac{1}{2} \text{ar}(\parallel^{\text{gm}} ABCD) = \frac{1}{2} \text{ar}(\parallel^{\text{gm}} BPRQ)$$

[Using (i) and (ii)]

$$\Rightarrow \text{ar}(\parallel^{\text{gm}} ABCD) = \text{ar}(\parallel^{\text{gm}} BPRQ).$$

Ex.26 In a parallelogram ABCD, the bisector of $\angle A$ also bisects BC at X. Prove that $AD = 2AB$.

Sol. Since AX is the bisector of $\angle A$.


$$\therefore \angle 1 = \frac{1}{2} \angle A \quad \dots (i)$$

Since ABCD is a parallelogram.

Therefore, $AD \parallel BC$ and AB intersects them.

$$\Rightarrow \angle A + \angle B = 180^\circ$$

[Θ Sum of interior angles is 180°]



$$\Rightarrow \angle B = 180^\circ - \angle A$$

In $\triangle ABX$, we have

$$\angle 1 + \angle 2 + \angle B = 180^\circ$$

$$\Rightarrow \frac{1}{2} \angle A + \angle 2 + 180^\circ - \angle A = 180^\circ$$

$$\Rightarrow \angle 2 - \frac{1}{2} \angle A = 0$$

$$\Rightarrow \angle 2 = \frac{1}{2} \angle A \quad \dots (ii)$$

From (i) and (ii), we have

$$\angle 1 = \angle 2.$$

Thus, in $\triangle ABX$, we have

$$\angle 1 = \angle 2.$$

$\Rightarrow BX = AB$ [Θ Sides opposite to equal angles in a $\triangle$ are equal]

$$\Rightarrow 2BX = 2AB \text{ [Multiplying both sides by 2]}$$

$$\Rightarrow BC = 2AB \text{ [}\Theta \text{ X is the mid-point of BC]}$$

$$\therefore AD = BC$$

$$\Rightarrow AD = 2AB$$

[Θ ABCD is a $\parallel^{\text{gm}}$ $\therefore AD = BC$]


Ex.27 In Fig. $BC \parallel XY$, $BX \parallel CA$ and $AB \parallel YC$. Prove that:

$$\text{ar}(\triangle ABX) = \text{ar}(\triangle ACY).$$

Sol. Join XC and BY.

Since $\triangle$ s BXC and BCY are on the same base BC and between the same parallels BC and XY

$$\therefore \text{ar}(\triangle BXC) = \text{ar}(\triangle BCY) \quad \dots (i)$$



Also, $\triangle$ s BXC and ABX are on the same base BX and between the same parallels BX and AC.

$$\therefore \text{ar}(\triangle BXC) = \text{ar}(\triangle ABX) \quad \dots (ii)$$


Clearly, $\triangle$ s BCY and ACY are on the same base CY and between the same parallels AB and CY.

$$\therefore \text{ar}(\triangle BCY) = \text{ar}(\triangle ACY) \quad \dots (iii)$$

From (i), (ii) and (iii), we get

$$\text{ar}(\triangle ABX) = \text{ar}(\triangle ACY).$$

Ex.28 In Fig. AD and BE are medians of $\triangle ABC$ and $BE \parallel DF$.



Prove that $CF = \frac{1}{4} AC$

Sol. In $\triangle BEC$, DF is a line through the mid-point D of BC and parallel to BE intersecting CE at F. Therefore, F is the mid-point of CE. Because the line drawn through the mid-point of one side of a triangle and parallel to another side bisects the third side.

Now, F is the mid-point of CE

$$\Rightarrow CF = \frac{1}{2} CE$$


$$\Rightarrow CF = \frac{1}{2} \left(\frac{1}{2} AC \right) \left[\Theta \text{ E is the mid-point of AC } \therefore EC = \frac{1}{2} AC \right]$$

$$\Rightarrow CF = \frac{1}{4} AC$$

Ex.29 D, E, F are the mid-points of the sides BC, CA and AB respectively of $\triangle ABC$, prove that BDEF is a parallelogram whose area is half that of $\triangle ABC$. Also, show that

$$\text{ar}(\triangle DEF) = \frac{1}{4} \text{ar}(\triangle ABC).$$

Sol. Since D and E are the mid-points of sides BC and AC respectively.



Therefore, $DE \parallel BA \Rightarrow DE \parallel BF$

Similarly, $FE \parallel BD$. So, BDEF is a parallelogram. Similarly, DCEF and AFDE are parallelograms.

Now, DF is a diagonal of $\parallel^{\text{gm}}$ BDEF.

$$\therefore \text{ar}(\triangle BDF) = \text{ar}(\triangle DEF) \quad \dots (i)$$

DE is a diagonal of $\parallel^{\text{gm}}$ DCEF

$$\therefore \text{ar}(\triangle DCE) = \text{ar}(\triangle DEF) \quad \dots (ii)$$

FE is a diagonal of $\parallel^{\text{gm}}$ AFDE

$$\therefore \text{ar}(\triangle AFE) = \text{ar}(\triangle DEF) \quad \dots (iii)$$

From (i), (ii) and (iii), we have

$$\text{ar}(\triangle BDF) = \text{ar}(\triangle DCE) = \text{ar}(\triangle AFE) = \text{ar}(\triangle DEF)$$

$$\begin{aligned} \text{But, } \text{ar}(\triangle BDF) + \text{ar}(\triangle DCE) + \text{ar}(\triangle AFE) \\ + \text{ar}(\triangle DEF) = \text{ar}(\triangle ABC) \end{aligned}$$

$$\therefore 4 \text{ar}(\triangle DEF) = \text{ar}(\triangle ABC)$$

$$\Rightarrow \text{ar}(\triangle DEF) = \frac{1}{4} \text{ar}(\triangle ABC).$$


Now, $\text{ar}(\parallel^{\text{gm}} \text{BDEF}) = 2 \text{ar}(\triangle DEF)$

$$\begin{aligned} \Rightarrow \text{ar}(\parallel^{\text{gm}} \text{BDEF}) &= 2 \times \frac{1}{4} \text{ar}(\triangle ABC) \\ &= \frac{1}{2} \text{ar}(\triangle ABC) \end{aligned}$$

Ex.30 $\triangle ABC$ and $\triangle DEF$ are two triangles such that AB, BC are respectively equal and parallel to DE, EF; show that AC is equal and parallel to DF.

Sol. Given: Two triangles ABC and DEF such that $AB = DE$ and $AB \parallel DE$. Also $BC = EF$ and $BC \parallel EF$

To Prove: $AC = DF$ and $AC \parallel DF$



Proof : Consider the quadrilateral ABED.

We have, $AB = DE$ and $AB \parallel DE$

$\Rightarrow$ One pair of opposite sides are equal and parallel

$\Rightarrow$ ABED is a parallelogram.

$$\Rightarrow AD = BE \text{ and } AD \parallel BE \quad \dots (i)$$

Now, consider quadrilateral DCEF.

We have, $BC = EF$ and $BC \parallel EF$

$\Rightarrow$ One pair of opposite sides are equal and parallel

$\Rightarrow$ DCEF is a parallelogram.

$$\Rightarrow CF = BE \text{ and } CF \parallel BE \quad \dots (ii)$$

From (i) and (ii), we have

$$AD = CF \text{ and } AD \parallel CF$$

$\Rightarrow$ ACFD is a parallelogram


$$AC = DF \text{ and } AC \parallel DF$$

Ex.31 Parallelogram ABCD & rectangle ABEF have the same base AB and also have equal areas. Show that perimeter of the parallelogram is greater than that of the rectangle.

Sol. Given: A $\parallel^{\text{gm}}$ ABCD and a rectangle ABEF with the same base AB and equal areas.

To Prove: Perimeter of $\parallel^{\text{gm}}$ ABCD $>$ Perimeter of rectangle ABEF

$$\text{i.e. } AB + BC + CD + AD > AB + BE + EF + AF.$$



Proof: Since opposite sides of a parallelogram and a rectangle are equal.

$$\therefore AB = DC \quad [\ominus ABCD \text{ is a } \parallel^{\text{gm}}]$$

$$\text{and, } AB = EF \quad [\ominus ABEF \text{ is a rectangle}]$$

$$\therefore DC = EF \quad \dots (i)$$

$$\Rightarrow AB + DC = AB + EF \quad \dots(ii)$$

Since, of all the segments that can be drawn to a given line from a point not lying on it, the perpendicular segment is the shortest.

$$\therefore BE < BC \text{ and } AF < AD \Rightarrow BC > BE \text{ and } AD > AF$$

$$\Rightarrow BC + AD > BE + AF \quad \dots(iii)$$


Adding (ii) and (iii), we get

$$AB + DC + BC + AD > AB + EF + BE + AF$$

$$\Rightarrow AB + BC + CD + DA > AB + BE + EF + FA.$$

Ex.32 In $\triangle ABC$, AD is the median through A and E is the mid-point of AD. BE produced meets AC in F. Prove that $AF = \frac{1}{3} AC$.

Sol. Through D, draw $DK \parallel BF$. In $\triangle ADK$, E is the mid-point of AD and $EF \parallel DK$.



$$\therefore F \text{ is the mid-point of } AK$$

$$\Rightarrow AF = FK \quad \dots (i)$$

In $\triangle BCF$, D is the mid-point of BC and $DK \parallel BF$

$$\therefore K \text{ is the mid-point of } FC$$

$$\therefore FK = KC \quad \dots(ii)$$

From (i) and (ii), we have

$$AF = FK = KC \quad \dots(iii)$$

$$\text{Now, } AC = AF + FK + KC$$

$$\Rightarrow AC = AF + AF + AF \text{ [Using (iii)]}$$

$$\Rightarrow AC = 3 (AF)$$

$$\Rightarrow AF = \frac{1}{3} AC$$

IMPORTANT POINTS TO BE REMEMBERED

1. Two figures are said to be on the same base and between the same parallels, if they have a common side (base) and the vertices (or the vertex) opposite to the common base of each figure lie on a line parallel to the base.
2. Two congruent figures have equal areas but the converse need not be true.
3. A diagonal of a parallelogram divides it into two triangles of equal area.
4. Parallelograms on the same base and between the same parallels are equal in area.
5. The area of a parallelogram is the product of its base and the corresponding altitude.
6. Parallelograms on equal bases and between the same parallels are equal in area.
7. Triangles on the same bases and between the same parallels are equal in area.
8. The area of a triangle is half the product of any of its sides and the corresponding altitude.
9. If a triangle and a parallelogram are on the same base and between the same parallels, then the area of the triangle is equal to the half of the parallelogram.
10. The area of a trapezium is half the product of its height and the sum of parallel sides.
11. Triangles having equal areas and having one side of one of the triangles, equal to one side of the other, have their corresponding altitudes equal.
12. If each diagonal of a quadrilateral separates it into two triangles of equal area, then the quadrilateral is a parallelogram.
13. The area of a rhombus is half the product of the lengths of its diagonals.
14. Diagonals of a parallelogram divide it into four triangles of equal area.
15. A median of a triangle divides it into two triangles of equal area.

EXERCISE

Q.1 Two opposite angles of a parallelogram are $(3x - 2)^\circ$ and $(50 - x)^\circ$. Find the measure of each angle of the parallelogram.

Q.2 If an angle of a parallelogram is two-third of its adjacent angle, find the angles of the parallelogram.


Q.3 Find the measure of all the angles of a parallelogram, if one angle is 24° less than twice the smallest angle.

Q.4 The perimeter of a parallelogram is 22 cm. If the longer side measures 6.5 cm what is the measure of the shorter side?

Q.5 In a parallelogram ABCD, $\angle D = 135^\circ$, determine the measures of $\angle A$ and $\angle B$.


Q.6 ABC is a triangle. D is a point on AB such that $AD = \frac{1}{4} AB$ and E is a point on AC such that $AE = \frac{1}{4} AC$. Prove that $DE = \frac{1}{4} BC$.

Q.7 In fig. ABCD is a parallelogram in which P is the mid-point of DC and Q is a point on AC such that $CQ = \frac{1}{4} AC$. If PQ produced meets BC at R, prove that R is a mid-point of BC.



Q.8 ABCD is a parallelogram, E and F are the mid-points of AB and CD respectively. GH is any line intersecting AD, EF and BC at G, P and H respectively. Prove that $GP = PH$.


Q.9 In the figure, ABCD is a parallelogram and a line through A cuts DC at P and BC produced at Q. Prove that $\text{arc}(\triangle BPC) = \text{arc}(\triangle DPQ)$



Q.10 If one angle of a parallelogram is 24° less than twice the smallest angle, then find the largest angle of the parallelogram.

Q.11 If an angle of a parallelogram is two-third of its adjacent angle, then find smallest angle of the parallelogram.

Q.12 In the given figure, ABCD is a parallelogram in which $\angle DAB = 75^\circ$ and $\angle DBC = 60^\circ$. Then find $\angle BDC$.



Q.13 Two parallelograms stand on equal bases and between the same parallels. Then find ratio of their areas.

Q.14 If a rectangle and a parallelogram are equal in area and have the same base and are situated on the same side, then the quotient:


$$\frac{\text{Perimeter of rectangle}}{\text{Perimeter of } \parallel \text{ gm}}$$

Q.15 ABCD is a parallelogram, E, F are the mid points of BC and AD respectively and G is any point on EF. Then find area of $\triangle GAB$.


Q.16 P and Q are any two points lying on the sides DC and AD respectively of a parallelogram ABCD. Show that $\text{ar}(\text{APB}) = \text{ar}(\text{BQC})$.

Q.17 The altitude of a parallelogram is twice the length of the base and its area is 1250 cm^2 . Then find lengths of the base and the altitude.

Q.18 In the figure if area of parallelogram ABCD is 30 cm^2 , then find $\text{ar}(\text{ADE}) + \text{ar}(\text{BCE})$.




Q.19 In the figure, ABCD is a parallelogram, if area of $\triangle AEB$ is 16 cm^2 , then find area of $\triangle BFC$.




Q.20 If the ratio of the altitude and the area of the parallelogram is $2 : 11$, then find the length of the base of the parallelogram.

Q.21 Two adjacent sides of a parallelogram are 24 cm and 18 cm . If the distance between the longer sides is 12 cm , then find the distance between the shorter sides.

Q.22 ABCD is a trapezium with parallel sides $AB = a \text{ cm}$ and $DC = b \text{ cm}$. E and F are the mid-points of the non-parallel sides. Then find ratio of $\text{ar}(\text{ABFE})$ and $\text{ar}(\text{EFCD})$.



Q.23 In figure, diagonals AC and BD of quadrilateral ABCD, intersect at O such that $OB = OD$. If $AB = CD$, then show that :



(i) $\text{ar}(\text{DOC}) = \text{ar}(\text{AOB})$


(ii) $\text{ar}(\text{DCB}) = \text{ar}(\text{ACB})$

(iii) $DA \parallel CB$ or ABCD is a parallelogram

Q.24 The area of a triangle is equal to the area of a rectangle whose length and breadth are 18 cm and 12 cm respectively. If the base of the triangle is 24 cm , then find its altitude.

Q.25 In figure, ABC is a triangle, AD is a median and E is the mid-point of AD. BE is joined and produced to intersect AC in a point F. Prove that :


$$AF = \left(\frac{1}{3}\right) AC$$



Q.26 ABC is triangle right-angled at B and P is the mid-point of AC. Prove that :


$$PB = PA = \frac{1}{2} AC$$

Q.27 The diagonals of a parallelogram ABCD intersect at a point O. Through O, a line is drawn to intersect AD at P and BC at Q. Show that PQ divides the parallelogram into two parts of equal area. (See figure)




Q.28 ABCD is a quadrilateral. A line through D, parallel to AC meets BC produced in P. Prove that area of $\triangle ABP$ = area of quad. ABCD.

Q.29 In figure, ABCD is a parallelogram, P is any point on AB produced, AQ is drawn parallel to CP to intersect CB produced at Q and parallelogram BQRP is completed. Show that area of ||gm ABCD = area BQRP.




➤ True/False Type Questions

- Q.30** In a parallelogram, the diagonals are equal.
- Q.31** In a parallelogram, the diagonals bisect each other.
- Q.32** In a parallelogram, the diagonals intersect each other at right angles.
- Q.33** If ABCD is a parallelogram with two adjacent angles A and B equal to each other, then the parallelogram is a rectangle.
- Q.34** If ABCD is a parallelogram and E, F are the centroids of $\triangle$ s ABD and BCD respectively, then $EF = AE$.
- Q.35** ABCD is a parallelogram one of whose diagonals is AC. Then $\text{ar}(\triangle ADC) = \text{ar}(\triangle CBA)$




➤ Fill in the blanks Types Questions


- Q.36** Fill in the blanks :
- The triangle formed by joining the mid-points of the sides of an isosceles triangle is
 - The triangle formed by joining the mid-points of the sides of a right triangle is
 - The figure formed by joining the mid-points of the consecutive sides of a quadrilateral is
- Q.37** In figure $CD \parallel AE$ and $CY \parallel BA$.
- Name a triangle equal in area of $\triangle CBX$
 - Prove that $\text{ar}(\triangle ZDE) = \text{ar}(\triangle CZA)$
 - Prove that $\text{ar}(\triangle BCZY) = \text{ar}(\triangle EDZ)$



- Q.38** In figure PSDA is a parallelogram in which $PQ = QR = RS$ and $AP \parallel BQ \parallel CR$. Prove that $\text{ar}(\triangle PQE) = \text{ar}(\triangle CQF)$.



- Q.39** In figure ABCD is a trapezium in which $AB \parallel DC$ and $DC = 40$ cm and $AB = 60$ cm. If X and Y are, respectively, the mid-point of AD and BC, prove that :



(i) $XY = 50$ cm

(ii) DCYX is a trapezium

(iii) $\text{ar}(\text{trap. DCYX}) = \frac{9}{11} \text{ar}(\text{trap. (XYBA)})$

Q.40 D is the mid-point of side BC of $\triangle ABC$ and E is the mid-point of BD. If O is the mid-point of AE, prove that $\text{ar}(\triangle BOE) = \frac{1}{8} \text{ar}(\triangle ABC)$.

ANSWER KEY

- | | | |
|--|---|---|
| 1. $37^\circ, 143^\circ, 37^\circ, 143^\circ$ | 2. $108^\circ, 72^\circ, 108^\circ, 72^\circ$ | 3. $68^\circ, 112^\circ, 68^\circ, 112^\circ$ 4.45 cm |
| 4. $\angle A = 45^\circ, \angle B = 135^\circ$ | 10. 112° | 11. 72° |
| 12. 45° | 13. 1 : 1 | 14. Less than 1 |
| 15. $\frac{1}{4}$ (gm ABCD) | 17. 25 cm, 50 cm | 18. 15 cm^2 |
| 19. 16 cm^2 | 20. 5.5 units | 21. 16 cm |
| 22. $(3a + b) : (a + 3b)$ | 24. 18 cm | 30. Flase |
| 31. True | 32. Flase | 33. True |
| 34. True | 35. True | |
| 36. (i) Also isosceles Δ | (ii) Also right angle Δ | (iii) Parallelograme |