

Topics : Fundamentals of Mathematics, Quadratic Equation, Complex Number

Type of Questions

		M.M.,	Min.
Comprehension (no negative marking) Q.1 to Q.3	(3 marks, 3 min.)	[9,	9]
Single choice Objective (no negative marking) Q.4, 5, 6, 7	(3 marks, 3 min.)	[12,	12]
Subjective Questions (no negative marking) Q.8,9	(4 marks, 5 min.)	[8,	10]

COMPREHENSION (Q. No. 1 to 3)

Consider the equation $|2x - 1| - 2|x - 2| = \lambda$

- If the above equation has only one solution, then λ belongs to
 (A) $\{-3, 3\}$ (B) $[-3, 3]$ (C) $(-3, 3)$ (D) ϕ
- If the above equation has more than one solutions then λ belongs to
 (A) $\{-3, 3\}$ (B) $[-3, 3]$ (C) $(-3, 3)$ (D) ϕ
- If $\lambda = 6$, then the above equation has
 (A) only one solution (B) only two solutions. (C) no solution. (D) more than two solutions.
- If the roots of the equation $x^2 + 2cx + ab = 0$ are real and unequal, then the roots of the equation $x^2 - 2(a + b)x + (a^2 + b^2 + 2c^2) = 0$ are :
 (A) real and unequal (B) real and equal
 (C) imaginary (D) rational
- If $-3 + 5i$ is a root of the equation $x^2 + px + q = 0$, then the ordered pair (p, q) is $(p, q \in \mathbb{R})$
 (A) $(-6, 34)$ (B) $(6, 34)$ (C) $(34, -6)$ (D) $(34, 6)$
- If the quadratic equation $ax^2 + bx + a^2 + b^2 + c^2 - ab - bc - ca = 0$, where a, b, c are distinct reals, has imaginary roots then :
 (A) $2(a - b) + (a - b)^2 + (b - c)^2 + (c - a)^2 > 0$
 (B) $2(a - b) + (a - b)^2 + (b - c)^2 + (c - a)^2 < 0$
 (C) $2(a - b) + (a - b)^2 + (b - c)^2 + (c - a)^2 = 0$
 (D) none
- If the quadratic equations $ax^2 + 2cx + b = 0$ & $ax^2 + 2bx + c = 0$ ($b \neq c$) have a common root, then $a + 4b + 4c$ is equal to :
 (A) -2 (B) -2 (C) 0 (D) 1
- Solve the equation : $|x+1| - |x| + 3|x-1| - 2|x-2| = x+2$
- Solve the equation : $\left| \frac{x+1}{x} \right| + |x+1| = \frac{(x+1)^2}{x}$

Answers Key

1. (C) 2. (A) 3. (C) 4. (C) 5. (B)
6. (A) 7. (C) 8. $x \in [2, \infty) \cup \{-2\}$
9. $x \in \{-1\} \cup (0, \infty)$