

# MIND MAP : LEARNING MADE SIMPLE CHAPTER - 4

## Determinants

Minor of an element  $a_{ij}$  in a determinant of matrix  $A$  is the determinant obtained by deleting  $i^{\text{th}}$  row and  $j^{\text{th}}$  column and is denoted by  $M_{ij}$ . If  $M_{ij}$  is the minor of  $a_{ij}$  and cofactor of  $a_{ij}$  is  $A_{ij}$  given by  $A_{ij} = (-1)^{i+j} M_{ij}$ .

- If  $A_{3 \times 3}$  is a matrix, then  $|A| = a_{11} \cdot A_{11} + a_{12} \cdot A_{12} + a_{13} \cdot A_{13}$ .
  - If elements of one row (or column) are multiplied with cofactors of elements of any other row (or column), then their sum is zero. For e.g.,  $a_{11} A_{21} + a_{12} A_{22} + a_{13} A_{23} = 0$ .
- e.g., if  $A = \begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix}$ , then  $M_{11} = 4$  and  $A_{11} = (-1)^{1+1} 4 = 4$ .

if  $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ , then  $\text{adj. } A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix}$ , where  $A_{ij}$  is the cofactor of  $a_{ij}$ .

- $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$ ,  $A$  - square matrix of order 'n'
- if  $|A| = 0$ , then  $A$  is singular. Otherwise,  $A$  is non-singular.
- if  $AB = BA = I$ , where  $B$  is a square matrix, then  $B$  is called the inverse of  $A$ ,  $A^{-1} = B$  or  $B^{-1} = A$ ,  $(A^{-1})^{-1} = A$ .

Inverse of a square matrix exists if  $A$  is non-singular i.e.  $|A| \neq 0$ , and is given by  $A^{-1} = \frac{1}{|A|} (\text{adj. } A)$

- if  $a_1x + b_1y + c_1z = d_1$ ,  $a_2x + b_2y + c_2z = d_2$ ,  $a_3x + b_3y + c_3z = d_3$  then we can write  $AX = B$ ,

where  $A = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$ ,  $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$  and  $B = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix}$

- Unique solution of  $AX = B$  is  $X = A^{-1}B$ ,  $|A| \neq 0$ .
- $AX = B$  is consistent or inconsistent according as the solution exists or not.
- For a square matrix  $A$  in  $AX = B$ , if
  - $|A| \neq 0$  then there exists unique solution.
  - $|A| = 0$  and  $(\text{adj. } A)B \neq 0$ , then no solution.
  - if  $|A| = 0$  and  $(\text{adj. } A)B = 0$  then system may or may not be consistent.

Minors and cofactors of a matrix

Adjoint and inverse of a matrix

Applications of determinants & matrices

Determinant of a square matrix 'A',  $|A|$  is given by

Properties of  $|A|$

Area of a triangle

- (i) if  $A = [a_{11}]_{1 \times 1}$ , then  $|A| = a_{11}$
  - (ii) if  $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}_{2 \times 2}$ , then  $|A| = a_{11} a_{22} - a_{12} a_{21}$
  - (iii) if  $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3}$ , then  $|A| = a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$
- For eg. if  $A = \begin{bmatrix} 2 & 3 \\ 2 & 4 \end{bmatrix}$ , then  $|A| = 2 \times 4 - 3 \times 2 = 2$

- (i)  $|A|$  remains unchanged, if the rows and columns of  $A$  are interchanged i.e.,  $|A| = |A'|$
- (ii) if any two rows (or columns) of  $A$  are interchanged, then the sign of  $|A|$  changes.
- (iii) if any two rows (or columns) of  $A$  are identical, then  $|A| = 0$
- (iv) if each element of a row (or a column) of  $A$  is multiplied by  $B$  (const.), then  $|A|$  gets multiplied by  $B$ .
- (v) if  $A = [a_{ij}]_{3 \times 3}$  then  $|kA| = k^3|A|$ .
- (vi) if elements of a row or a column in a determinant  $|A|$  can be expressed as sum of two or more elements, then  $|A|$  can be expressed as  $|B| + |C|$ .
- (vii) if  $R_i \rightarrow R_i + kR_j$  or  $C_i = C_i + kC_j$  in  $|A|$ , then the value of  $|A|$  remains same

if  $(x_1, y_1)$ ,  $(x_2, y_2)$  and  $(x_3, y_3)$   $\Delta = \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$

For eg: if  $(1, 2)$ ,  $(3, 4)$  and  $(-2, 5)$  are the vertices, then area of the triangle is

$$\Delta = \begin{vmatrix} 1 & 2 & 1 \\ 3 & 4 & 1 \\ -2 & 5 & 1 \end{vmatrix} = 1(4 - 5) - 2(3 + 2) + 1(15 + 8) = 12 \text{ sq. units.}$$

we take positive value of the determinant.