



Assignment

Types of Matrices

Basic Level

- If $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & b & 1 \end{bmatrix}$ be a diagonal matrix, then $b =$
 - 2
 - 0
 - 1
 - 3
- Which of the following is a diagonal matrix
 - $\begin{bmatrix} 2 & 0 & 2 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$
 - $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$
 - $\begin{bmatrix} 2 & 2 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$
 - None of these
- If I is a unit matrix, then $3I$ will be
 - A unit matrix
 - A triangular matrix
 - A scalar matrix
 - None of these
- If $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 4 & 7 \\ 5 & 1 & 6 \end{bmatrix}$, then the value of X where $A+X$ is a unit matrix, is
 - $\begin{bmatrix} 0 & -2 & 1 \\ -3 & -3 & -7 \\ -5 & -1 & -6 \end{bmatrix}$
 - $\begin{bmatrix} 0 & -3 & 5 \\ -2 & -3 & 1 \\ -1 & -7 & 6 \end{bmatrix}$
 - $\begin{bmatrix} 0 & -1 & -2 \\ 3 & 3 & 7 \\ 5 & 1 & 6 \end{bmatrix}$
 - None of these
- If A is diagonal matrix of order 2×2 , then wrong statement is
 - $AB = BA$, where B is a diagonal matrix of order 2×2
 - AB is a diagonal matrix
 - $A^T = A$
 - A is a scalar matrix

Algebra of Matrices

Basic Level

- If $M = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$ and $M^2 - \lambda M - I_2 = O$, then $\lambda =$ [MP PET 1990, 2001]
 - 2
 - 2
 - 4
 - 4
- If $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ and $B = \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix}$, then the correct relation is
 - $A^2 = B^2$
 - $A + B = B - A$
 - $AB = BA$
 - None of these
- If $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, then
 - 2
 - 2
 - 4
 - 4

- (a) $AB = BA$ (b) $AB = BA = O$ (c) $AB = O, BA \neq O$ (d) $AB \neq BA = O$
9. If $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, then $A^n =$ [Rajasthan PET 1995]
 (a) $\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} n & n \\ 0 & n \end{bmatrix}$ (c) $\begin{bmatrix} n & 1 \\ 0 & n \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 1 \\ 0 & n \end{bmatrix}$
10. If $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$, then $A^2 =$
 (a) A (b) $2A$ (c) $-A$ (d) $-2A$
11. If $2A + \begin{bmatrix} 2 & -1 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} -3 & 7 \\ 4 & 3 \end{bmatrix}$, then $A =$
 (a) $\begin{bmatrix} -5 & 8 \\ 2 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} -5/2 & 4 \\ 1 & 3/2 \end{bmatrix}$ (c) $\begin{bmatrix} -5 & 6 \\ 2 & 3 \end{bmatrix}$ (d) None of these
12. If $[m \ n] \begin{bmatrix} m \\ n \end{bmatrix} = [25]$ and $m < n$, then $(m, n) =$
 (a) $(2, 3)$ (b) $(3, 4)$ (c) $(4, 3)$ (d) None of these
13. If $A = \begin{bmatrix} \cos^2 \theta & \sin \theta \cos \theta \\ \sin \theta \cos \theta & \sin^2 \theta \end{bmatrix}$, $B = \begin{bmatrix} \cos^2 \phi & \sin \phi \cos \phi \\ \sin \phi \cos \phi & \sin^2 \phi \end{bmatrix}$ and θ and ϕ differs by $\frac{\pi}{2}$, then $AB =$
 (a) I (b) O (c) $-I$ (d) None of these
14. If $A = \begin{bmatrix} 0 & r & -q \\ -r & 0 & p \\ q & -p & 0 \end{bmatrix}$ and $B = \begin{bmatrix} p^2 & pq & pr \\ pq & q^2 & qr \\ pr & qr & r^2 \end{bmatrix}$, then $AB =$
 (a) $\begin{bmatrix} p & 0 & 0 \\ 0 & q & 0 \\ 0 & 0 & r \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$
15. If $A = \begin{bmatrix} 4 & 1 \\ 3 & 2 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then $A^2 - 6A =$ [MP PET 1987]
 (a) $3I$ (b) $5I$ (c) $-5I$ (d) None of these
16. If $A = \begin{bmatrix} \lambda & 1 \\ -1 & -\lambda \end{bmatrix}$, then for what value of λ , $A^2 = O$ [MP PET 1992]
 (a) 0 (b) ± 1 (c) -1 (d) 1
17. If $A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$ and $A^n = O$, then the minimum value of n is
 (a) 2 (b) 3 (c) 4 (d) 5
18. If $A = \begin{bmatrix} 1/3 & 2 \\ 0 & 2x-3 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 6 \\ 0 & -1 \end{bmatrix}$ and $AB = I$, then $x =$ [MP PET 1987]
 (a) -1 (b) 1 (c) 0 (d) 2
19. If $2A + B = \begin{bmatrix} 6 & 4 \\ 6 & -11 \end{bmatrix}$ and $A - B = \begin{bmatrix} 0 & 2 \\ 6 & 2 \end{bmatrix}$, then $A =$

374 Matrices

(a) $\begin{bmatrix} 2 & 2 \\ 4 & -3 \end{bmatrix}$

(b) $\begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix}$

(c) $\begin{bmatrix} 2 & 2 \\ 4 & 3 \end{bmatrix}$

(d) None of these

20. If $A = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} -5 & 4 & 0 \\ 0 & 2 & -1 \\ 1 & -3 & 2 \end{bmatrix}$, then $AB =$

[MP PET 1988]

(a) $\begin{bmatrix} -5 & 4 & 0 \\ 0 & 4 & -2 \\ 3 & -9 & 6 \end{bmatrix}$

(b) $\begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$

(c) $\begin{bmatrix} -2 & -1 & 4 \end{bmatrix}$

(d) $\begin{bmatrix} -5 & 8 & 0 \\ 0 & 4 & -3 \\ 1 & -6 & 6 \end{bmatrix}$

21. If $A = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix}$, then AB is

(a) Diagonal matrix

(b) Null matrix

(c) Unit matrix

(d) None of these

22. If $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$, then $A^5 =$

(a) $5A$

(b) $10A$

(c) $16A$

(d) $32A$

23. If $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $AB = O$, then $B =$

[MP PET 1989]

(a) $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

(d) $\begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix}$

24. If $R(t) = \begin{bmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{bmatrix}$, then $R(s)R(t) =$

[Roorkee 1981]

(a) $R(s) + R(t)$

(b) $R(st)$

(c) $R(s+t)$

(d) None of these

25. If $A = \begin{bmatrix} 0 & 1 & 2 \\ 2 & 3 & 4 \\ 4 & 5 & 6 \end{bmatrix}$ and $3A - 4B = \begin{bmatrix} -4 & 3 & 6 \\ 6 & 5 & 12 \\ 12 & 15 & 14 \end{bmatrix}$, then $B =$

(a) $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

(d) None of these

26. If $A = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$, then A^4 is equal to

[MP PET 1993]

(a) $\begin{bmatrix} 1 & a^4 \\ 0 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 4 & 4a \\ 0 & 4 \end{bmatrix}$

(c) $\begin{bmatrix} 4 & a^4 \\ 0 & 4 \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 4a \\ 0 & 1 \end{bmatrix}$

27. If $\begin{bmatrix} 3 & 1 \\ 4 & 1 \end{bmatrix} X = \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$, then $X =$

(a) $\begin{bmatrix} -3 & 4 \\ 14 & -13 \end{bmatrix}$

(b) $\begin{bmatrix} 3 & -4 \\ -14 & 13 \end{bmatrix}$

(c) $\begin{bmatrix} 3 & 4 \\ 14 & 13 \end{bmatrix}$

(d) $\begin{bmatrix} -3 & 4 \\ -14 & 13 \end{bmatrix}$

28. If $A = \begin{bmatrix} 5 & -3 \\ 2 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 6 & -4 \\ 3 & 6 \end{bmatrix}$, then $A - B =$

[Rajasthan PET 1995]

(a) $\begin{bmatrix} 11 & -7 \\ 5 & 10 \end{bmatrix}$

(b) $\begin{bmatrix} -1 & 1 \\ -1 & -2 \end{bmatrix}$

(c) $\begin{bmatrix} 11 & 7 \\ 5 & -10 \end{bmatrix}$

(d) $\begin{bmatrix} 12 & -7 \\ 5 & -10 \end{bmatrix}$

29. If $3X + 2Y = I$ and $2X - Y = O$, where I and O are unit and null matrices of order 3 respectively, then [MP PET 1995]
- (a) $X = \frac{1}{7}, Y = \frac{2}{7}$ (b) $X = \frac{2}{7}, Y = \frac{1}{7}$ (c) $X = \frac{1}{7}I, Y = \frac{2}{7}I$ (d) $X = \frac{2}{7}I, Y = \frac{1}{7}I$
30. If $A = \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$ and I is the identity matrix of order 2, then $(A - 2I)(A - 3I) =$ [Rajasthan PET 2002]
- (a) I (b) O (c) $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$
31. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, then $A^4 =$ [EAMCET 1994]
- (a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
32. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, then $A^2 =$ [Karnataka CET 1994]
- (a) $\begin{bmatrix} 8 & -5 \\ -5 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 8 & -5 \\ 5 & 3 \end{bmatrix}$ (c) $\begin{bmatrix} 8 & -5 \\ -5 & -3 \end{bmatrix}$ (d) $\begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix}$
33. If $X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, then the value of X^n is [EAMCET 1991]
- (a) $\begin{bmatrix} 3n & -4n \\ n & -n \end{bmatrix}$ (b) $\begin{bmatrix} 2+n & 5-n \\ n & -n \end{bmatrix}$ (c) $\begin{bmatrix} 3^n & (-4)^n \\ 1^n & (-1)^n \end{bmatrix}$ (d) None of these
34. If $A = \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix}$, then $A^2 =$ [EAMCET 1983]
- (a) $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$
35. If $A + B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ and $A - 2B = \begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}$, then $A =$ [Karnataka CET 1994]
- (a) $\begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{bmatrix}$ (c) $\begin{bmatrix} 1/3 & 1/3 \\ 2/3 & 1/3 \end{bmatrix}$ (d) None of these
36. If $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, then $AB =$ [EAMCET 1984]
- (a) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$
37. If $A = \begin{bmatrix} 1 & 3 & 0 \\ -1 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 2 & 3 \\ -1 & 1 & 2 \end{bmatrix}$, then $AB =$ [EAMCET 1987]
- (a) $\begin{bmatrix} 5 & 9 & 13 \\ -1 & 2 & 4 \\ -1 & 2 & 4 \end{bmatrix}$ (b) $\begin{bmatrix} 5 & 9 & 13 \\ -1 & 2 & 4 \\ -2 & 2 & 4 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 2 & 4 \\ -1 & 2 & 4 \\ -2 & 2 & 4 \end{bmatrix}$ (d) None of these
38. If $\begin{bmatrix} x & 0 \\ 1 & y \end{bmatrix} + \begin{bmatrix} -2 & 1 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 5 \\ 6 & 3 \end{bmatrix} - \begin{bmatrix} 2 & 4 \\ 2 & 1 \end{bmatrix}$, then [Rajasthan PET 1994]
- (a) $x = -3, y = -2$ (b) $x = 3, y = -2$ (c) $x = 3, y = 2$ (d) $x = -3, y = 2$

376 Matrices

39. If $A = \begin{bmatrix} 1 & -2 \\ 3 & 0 \end{bmatrix}$, $B = \begin{bmatrix} -1 & 4 \\ 2 & 3 \end{bmatrix}$, $C = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then $5A - 3B - 2C =$
- (a) $\begin{bmatrix} 8 & 20 \\ 7 & 9 \end{bmatrix}$ (b) $\begin{bmatrix} 8 & -20 \\ 7 & -9 \end{bmatrix}$ (c) $\begin{bmatrix} -8 & 20 \\ -7 & 9 \end{bmatrix}$ (d) $\begin{bmatrix} 8 & 7 \\ -20 & -9 \end{bmatrix}$
40. If $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, I is the unit matrix of order 2 and a, b are arbitrary constants, then $(aI + bA)^2$ is equal to
- (a) $a^2I + abA$ (b) $a^2I + 2abA$ (c) $a^2I + b^2A$ (d) None of these
41. If $U = \begin{bmatrix} 2 & -3 & 4 \end{bmatrix}$, $X = \begin{bmatrix} 0 & 2 & 3 \end{bmatrix}$, $V = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ and $Y = \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix}$, then $UV + XY =$ [MP PET 1997]
- (a) 20 (b) [-20] (c) -20 (d) [20]
42. Which one of the following is not true [Kurukshetra CEE 1998]
- (a) Matrix addition is commutative (b) Matrix addition is associative
(c) Matrix multiplication is commutative (d) Matrix multiplication is associative
43. If $A = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 5 \\ 0 & 2 \end{bmatrix}$, then which of the following is defined [Rajasthan PET 1996]
- (a) AB (b) BA (c) $(AB).C$ (d) $(AC).B$
44. If $A = \begin{bmatrix} 1 & 2 & 3 \\ -2 & 3 & -1 \\ 3 & 1 & 2 \end{bmatrix}$ and I is a unit matrix of 3rd order, then $(A^2 + 9I)$ equals [Rajasthan PET 1999]
- (a) $2A$ (b) $4A$ (c) $6A$ (d) None of these
45. If $A = \begin{pmatrix} i & 1 \\ 0 & i \end{pmatrix}$, then A^4 equals [AMU 1999]
- (a) $\begin{pmatrix} 1 & -4i \\ 0 & 1 \end{pmatrix}$ (b) $\begin{pmatrix} -1 & 4i \\ 0 & -1 \end{pmatrix}$ (c) $\begin{pmatrix} -i & 4 \\ 0 & i \end{pmatrix}$ (d) $\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$
46. $\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} [2 \ 1 \ -1] =$ [MP PET 2000]
- (a) [-1] (b) $\begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 1 & -1 \\ -2 & -1 & 1 \\ 4 & 2 & -2 \end{bmatrix}$ (d) Not defined
47. If $2X - \begin{bmatrix} 1 & 2 \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 0 & -2 \end{bmatrix}$, then X is equal to [Rajasthan PET 2001]
- (a) $\begin{bmatrix} 2 & 2 \\ 7 & 4 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 2 \\ 7/2 & 2 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 2 \\ 7/2 & 1 \end{bmatrix}$ (d) None of these
48. If $A = \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix}$ and $kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$, then the values of k, a, b are respectively [EAMCET 2001]
- (a) -6, -12, -18 (b) -6, 4, 9 (c) -6, -4, -9 (d) -6, 12, 18
49. If $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, then $A^n =$ [Kerala (Engg.) 2001]

(a) $\begin{bmatrix} 1 & 2n \\ 0 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 2 & n \\ 0 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 2n \\ 0 & -1 \end{bmatrix}$

(d) $\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$

50. If matrix $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then $A^{16} =$

[Karnataka CET 2002]

(a) $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

(b) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

51. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$, then $A^2 - 5A =$

[Rajasthan PET 2002]

(a) I

(b) $14I$

(c) O

(d) None of these

52. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then $A^{100} =$

[UPSEAT 2002]

(a) $2^{100} A$

(b) $2^{99} A$

(c) $2^{101} A$

(d) None of these

53. Which is true about matrix multiplication

[UPSEAT 2002]

(a) It is commutative

(b) It is associative

(c) Both (a) and (b)

(d) None of these

54. If $P = \begin{pmatrix} i & 0 & -i \\ 0 & -i & i \\ -i & i & 0 \end{pmatrix}$ and $Q = \begin{pmatrix} -i & i \\ 0 & 0 \\ i & -i \end{pmatrix}$, then PQ is equal to

[Kerala (Engg.) 2002]

(a) $\begin{pmatrix} -2 & 2 \\ 1 & -1 \\ 1 & -1 \end{pmatrix}$

(b) $\begin{pmatrix} 2 & -2 \\ -1 & 1 \\ -1 & 1 \end{pmatrix}$

(c) $\begin{pmatrix} 2 & -2 \\ -1 & 1 \end{pmatrix}$

(d) $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

55. $\begin{bmatrix} 7 & 1 & 2 \\ 9 & 2 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ 2 \end{bmatrix}$ is equal to

(a) $\begin{bmatrix} 43 \\ 44 \end{bmatrix}$

(b) $\begin{bmatrix} 43 \\ 45 \end{bmatrix}$

(c) $\begin{bmatrix} 45 \\ 44 \end{bmatrix}$

(d) $\begin{bmatrix} 44 \\ 45 \end{bmatrix}$

56. If $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$, then AB is

[MP PET 2003]

(a) $\begin{bmatrix} 5 & 1 & -3 \\ 3 & 2 & 6 \\ 14 & 5 & 0 \end{bmatrix}$

(b) $\begin{bmatrix} 11 & 4 & 3 \\ 1 & 2 & 3 \\ 0 & 3 & 3 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 8 & 4 \\ 2 & 9 & 6 \\ 0 & 2 & 0 \end{bmatrix}$

(d) $\begin{bmatrix} 0 & 1 & 2 \\ 5 & 4 & 3 \\ 1 & 8 & 2 \end{bmatrix}$

57. For 2×2 matrices A , B and I , if $A + B = I$ and $2A - 2B = I$, then A equals

(a) $\begin{bmatrix} \frac{1}{4} & 0 \\ 0 & \frac{1}{4} \end{bmatrix}$

(b) $\begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$

(c) $\begin{bmatrix} \frac{3}{4} & 0 \\ 0 & \frac{3}{4} \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

58. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $A^2 + 2A$ equals

[AMU 1988]

(a) A

(b) $2A$

(c) $3A$

(d) $4A$

59. If $A = \begin{bmatrix} 1 & -6 & 2 \\ 0 & -1 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$, then AB equals [AMU 1987]
- (a) $\begin{bmatrix} -8 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} -8 \\ 3 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & -12 & 2 \\ 0 & -2 & 5 \end{bmatrix}$ (d) $\begin{bmatrix} 2 & 12 & 4 \\ 0 & -2 & -10 \end{bmatrix}$
60. Let $A = \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 0 \\ 1 & 12 \end{bmatrix}$, then [DCE 1999]
- (a) $AB = O, BA = O$ (b) $AB = O, BA \neq O$ (c) $AB \neq O, BA = O$ (d) $AB \neq O, BA \neq O$
61. If A, B are square matrices of order $n \times n$, then $(A - B)^2$ is equal to
- (a) $A^2 - B^2$ (b) $A^2 - 2BA + B^2$ (c) $A^2 - AB - BA + B^2$ (d) $A^2 - 2AB + B^2$
62. If $A = \begin{bmatrix} 3 & 5 \\ -2 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -3 \\ 2 & -7 \end{bmatrix}$, then $2A - 3B$ is equal to [Rajasthan PET 1989, 90]
- (a) $\begin{bmatrix} 3 & -19 \\ 10 & 29 \end{bmatrix}$ (b) $\begin{bmatrix} 3 & 19 \\ -10 & 29 \end{bmatrix}$ (c) $\begin{bmatrix} -3 & 19 \\ 10 & 29 \end{bmatrix}$ (d) None of these
63. If $A = \begin{bmatrix} 2 & 4 \\ 0 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ 0 & 5 \end{bmatrix}$, then $4A - 3B$ is equal to [Rajasthan PET 1993]
- (a) $\begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$ (b) $\begin{bmatrix} 7 & 14 \\ 0 & 7 \end{bmatrix}$ (c) $\begin{bmatrix} 5 & 10 \\ 0 & -3 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & -2 \\ 0 & -12 \end{bmatrix}$
64. If $A = \begin{pmatrix} 1 & -2 \\ 3 & 4 \end{pmatrix}$, $B = \begin{pmatrix} 2 & 1 \\ -4 & 3 \end{pmatrix}$ and $C = \begin{pmatrix} 2 & 12 \\ -17 & 9 \end{pmatrix}$, then $5A - 3B + C$ equals [Rajasthan PET 1993]
- (a) $\begin{pmatrix} 1 & 10 \\ -1 & 20 \end{pmatrix}$ (b) $\begin{pmatrix} 1 & -1 \\ 10 & 20 \end{pmatrix}$ (c) $\begin{pmatrix} 1 & 1 \\ -10 & 20 \end{pmatrix}$ (d) $\begin{pmatrix} -1 & 1 \\ 1 & 20 \end{pmatrix}$
65. If $A = \begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then $A + B - C$ equals
- (a) $\begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 5 & 2 \\ 5 & 5 \end{bmatrix}$ (c) $\begin{bmatrix} 5 & 2 \\ 3 & 5 \end{bmatrix}$ (d) $\begin{bmatrix} 3 & 2 \\ 5 & 5 \end{bmatrix}$
66. If $\begin{bmatrix} x & y \\ -2 & 2 \end{bmatrix} + \begin{bmatrix} x & 1 \\ 4 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ y & 2 \end{bmatrix}$, then x and y are [Rajasthan PET 1994]
- (a) 1, 1 (b) 1, 2 (c) 2, 2 (d) 2, 1
67. If $X = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$ and $3X - \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$, then the value of a is
- (a) -2 (b) 0 (c) 2 (d) 1
68. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$, then [Rajasthan PET 1985]
- (a) $AB = BA$ (b) $AB = B^2$ (c) $AB = -BA$ (d) None of these
69. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 3 & 4 \end{bmatrix}$, then AB is equal to [Rajasthan PET 1989, 90, 98]

- (a) $\begin{bmatrix} 8 & 15 & 16 \\ 5 & 9 & 10 \end{bmatrix}$ (b) $\begin{bmatrix} 8 & 5 \\ 15 & 9 \\ 16 & 10 \end{bmatrix}$ (c) $\begin{bmatrix} 8 & 5 \\ 15 & 9 \end{bmatrix}$ (d) None of these

70. If $A = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & -2 \\ -2 & 5 \end{pmatrix}$, then AB equals [Rajasthan PET 1991]

- (a) $\begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$ (b) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ (c) $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ (d) $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$

71. $A \begin{bmatrix} -1 & 2 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} -4 & 1 \\ 7 & 7 \end{bmatrix}$, then A equals [EAMCET 1996]

- (a) $\begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 1 \\ -2 & 3 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & 1 \\ 2 & 3 \end{bmatrix}$

72. If $A = \begin{bmatrix} 4 & 6 & -1 \\ 3 & 0 & 2 \\ 1 & -2 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 4 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & 4 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}$, then which of the following is not defined

- (a) AB (b) $B'C$ (c) CC' (d) $A^2 + 2B - 2A$

73. If a matrix B is obtained by multiplying each element of a matrix A of order 2×2 by 3, then relation between A and B is

- (a) $A = 3B$ (b) $3A = B$ (c) $9A = B$ (d) $A = 9B$ [Rajasthan PET 1986]

Advance Level

74. For each real number x such that $-1 < x < 1$, let $A(x)$ be the matrix $(1-x)^{-1} \begin{bmatrix} 1 & -x \\ -x & 1 \end{bmatrix}$ and $z = \frac{x+y}{1+xy}$. Then

- (a) $A(z) = A(x) + A(y)$ (b) $A(z) = A(x)[A(y)]^{-1}$ (c) $A(z) = A(x)A(y)$ (d) $A(z) = A(x) - A(y)$

75. If $A = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}$, then the value of A^{40} is [Rajasthan PET 1999]

- (a) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}$

76. If $A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 3 & -1 & 1 \end{bmatrix}$, then [Kurukshetra CEE 2002]

- (a) $A^3 + 3A^2 + A - 9I_3 = 0$ (b) $A^3 - 3A^2 + A + 9I_3 = 0$ (c) $A^3 + 3A^2 - A + 9I_3 = 0$ (d) $A^3 - 3A^2 - A + 9I_3 = 0$

77. If $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ and I is the unit matrix of order 2, then A^2 equals

- (a) $4A - 3I$ (b) $3A - 4I$ (c) $A - I$ (d) $A + I$

78. If $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$, then $A^n =$

380 Matrices

- (a) $\begin{bmatrix} na & 0 & 0 \\ 0 & nb & 0 \\ 0 & 0 & nc \end{bmatrix}$ (b) $\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$ (c) $\begin{bmatrix} a^n & 0 & 0 \\ 0 & b^n & 0 \\ 0 & 0 & c^n \end{bmatrix}$ (d) None of these

79. If $A_\alpha = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, then which of following statement is true

- (a) $A_\alpha \cdot A_\beta = A_{\alpha\beta}$ and $(A_\alpha)^n = \begin{bmatrix} \cos^n \alpha & \sin^n \alpha \\ -\sin^n \alpha & \cos^n \alpha \end{bmatrix}$ (b) $A_\alpha \cdot A_\beta = A_{\alpha\beta}$ and $(A_\alpha)^n = \begin{bmatrix} \cos n\alpha & \sin n\alpha \\ -\sin n\alpha & \cos n\alpha \end{bmatrix}$
(c) $A_\alpha \cdot A_\beta = A_{\alpha+\beta}$ and $(A_\alpha)^n = \begin{bmatrix} \cos^n \alpha & \sin^n \alpha \\ -\sin^n \alpha & \cos^n \alpha \end{bmatrix}$ (d) $A_\alpha \cdot A_\beta = A_{\alpha+\beta}$ and $(A_\alpha)^n = \begin{bmatrix} \cos n\alpha & \sin n\alpha \\ -\sin n\alpha & \cos n\alpha \end{bmatrix}$

Properties of Matrices

Basic Level

80. $AB = 0$, if and only if [MNR 1981; Karnataka CET 1993]
(a) $A \neq 0, B \neq 0$ (b) $A = 0, B \neq 0$ (c) $A = 0$ or $B = 0$ (d) None of these
81. If $A \times \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 2 \end{bmatrix}$, then the order of A is
(a) 1×1 (b) 2×1 (c) 1×2 (d) 2×2
82. If $AB = C$, then matrices A, B, C are [MP PET 1991]
(a) $A_{2 \times 3}, B_{3 \times 2}, C_{2 \times 3}$ (b) $A_{3 \times 2}, B_{2 \times 3}, C_{3 \times 2}$ (c) $A_{3 \times 3}, B_{2 \times 3}, C_{3 \times 3}$ (d) $A_{3 \times 2}, B_{2 \times 3}, C_{3 \times 3}$
83. $A = [a_{ij}]_{m \times n}$ is a square matrix, if
(a) $m < n$ (b) $m > n$ (c) $m = n$ (d) None of these
84. If $A = \begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 3 \\ -2 & 2 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 4 & 5 \\ 5 & -4 & 0 \end{bmatrix}$, then the element of 3rd row and third column in AB will be
(a) -18 (b) 4 (c) -12 (d) None of these
85. If A and B be symmetric matrices of the same order, then $AB - BA$ will be a
(a) Symmetric matrix (b) Skew-symmetric matrix (c) Null matrix (d) None of these
86. If A and B are square matrices of order 2, then $(A + B)^2 =$
(a) $A^2 + 2AB + B^2$ (b) $A^2 + AB + BA + B^2$ (c) $A^2 + 2BA + B^2$ (d) None of these
87. If the order of the matrices A and B be 2×3 and 3×2 respectively, then the order of $A + B$ will be
(a) 2×2 (b) 3×3 (c) 2×3 (d) None of these
88. In a lower triangular matrix element $a_{ij} = 0$, if
(a) $i \leq j$ (b) $i \geq j$ (c) $i > j$ (d) $i < j$
89. If A is a square matrix of order n and $A = kB$, where k is a scalar, then $|A| =$ [Karnataka CET 1992]
(a) $|B|$ (b) $k|B|$ (c) $k^n|B|$ (d) $n|B|$
90. Let $A = \begin{bmatrix} 4 & 6 & -1 \\ 3 & 0 & 2 \\ 1 & -2 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 4 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} 3 & 1 & 2 \end{bmatrix}$. The expression which is not defined is

- (a) $B'B$ (b) CAB (c) $A+B'$ (d) $A^2 + A$
91. If $A = \begin{bmatrix} a & b \end{bmatrix}$, $B = \begin{bmatrix} -b & -a \end{bmatrix}$ and $C = \begin{bmatrix} a \\ -a \end{bmatrix}$, then the correct statement is [AMU 1987]
- (a) $A = -B$ (b) $A+B = A-B$ (c) $AC = BC$ (d) $CA = CB$
92. If A and B are two matrices and $(A+B)(A-B) = A^2 - B^2$, then
- (a) $AB = BA$ (b) $A^2 + B^2 = A^2 - B^2$ (c) $A'B' = AB$ (d) None of these
93. If A and B are square matrices of same order, then [Roorkee 1995]
- (a) $A+B = B+A$ (b) $A+B = A-B$ (c) $A-B = B-A$ (d) $AB = BA$
94. Which of the following is incorrect
- (a) $A^2 - B^2 = (A+B)(A-B)$ (b) $(A^T)^T = A$
- (c) $(AB)^n = A^n B^n$, where A, B commute (d) $(A-I)(I+A) = 0 \Leftrightarrow A^2 = I$
95. Which of the following is/are incorrect
- (i) Adjoint of a symmetric matrix is symmetric,
(ii) Adjoint of unit matrix is a unit matrix,
(iii) $A(\text{adj } A) = (\text{adj } A)A \neq A|I$ and
(iv) Adjoint of a diagonal matrix is a diagonal matrix
- (a) (i) (b) (ii) (c) (iii) and (iv) (d) None of these
96. Let $A = [a_{ij}]_{n \times n}$ be a square matrix and let c_{ij} be cofactor of a_{ij} in A . If $C = [c_{ij}]$, then
- (a) $|C| = |A|$ (b) $|C| = |A|^{n-1}$ (c) $|C| = |A|^{n-2}$ (d) None of these
97. A, B are n -rowed square matrices such that $AB = 0$ and B is non-singular. Then
- (a) $A \neq 0$ (b) $A = 0$ (c) $A = I$ (d) None of these
98. If A and B are two matrices such that $AB = B$ and $BA = A$, then $A^2 + B^2 =$ [EAMCET 1994]
- (a) $2AB$ (b) $2BA$ (c) $A+B$ (d) AB
99. If A and B are two matrices such that $A+B$ and AB are both defined, then [Pb. CET 1990]
- (a) A and B are two matrices not necessarily of same order
(b) A and B are square matrices of same order
(c) Number of columns of A = number of rows of B
(d) None of these
100. If $A = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}$ and A^2 is the identity matrix, then $x =$ [EAMCET 1993]
- (a) 1 (b) 2 (c) 3 (d) 0
101. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$, then $(A+B)^2$ equals [Rajasthan PET 1994]
- (a) $A^2 + B^2$ (b) $A^2 + B^2 + 2AB$ (c) $A^2 + B^2 + AB - BA$ (d) None of these
102. If $A = \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix}$, then $(A+B)(A-B)$ is equal to [Rajasthan PET 1994]
- (a) $A^2 - B^2$ (b) $A^2 + B^2$ (c) $A^2 - B^2 + BA + AB$ (d) None of these
103. If A is 3×4 matrix and B is a matrix such that $A'B$ and BA' are both defined. Then B is of the type [Himachal Pradesh]
- (a) 3×4 (b) 3×3 (c) 4×4 (d) 4×3
104. Which of the following is not true
- (a) Every skew-symmetric matrix of odd order is non-singular

382 Matrices

- (b) If determinant of a square matrix is non-zero, then it is non-singular
 (c) Adjoint of a symmetric matrix is symmetric
 (d) Adjoint of a diagonal matrix is diagonal
- 105.** Which one of the following statements is true [MP PET 1996]
 (a) Non-singular square matrix does not have a unique inverse (b) Determinant of a non-singular matrix is zero
 (c) If $A' = A$, then A is a square matrix (d) If $|A| \neq 0$, then $|A \cdot \text{adj } A| = |A|^{(n-1)}$, where $A = (a_{ij})_{n \times n}$
- 106.** If matrix $A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$, then
 (a) $A' = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ (b) $A^{-1} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$
 (c) $A \cdot \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} = 2I$ (d) $\lambda A = \begin{bmatrix} \lambda & -\lambda \\ 1 & 1 \end{bmatrix}$, where λ is a non-zero scalar
- 107.** If $A = \begin{bmatrix} 1 & 2 \\ -3 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 \\ 2 & 3 \end{bmatrix}$, then [MP PET 1996]
 (a) $A^2 = A$ (b) $B^2 = B$ (c) $AB \neq BA$ (d) $AB = BA$
- 108.** Which one of the following is correct [Kurukshetra CEE 1998]
 (a) Skew-symmetric matrix of odd order is non-singular. (b) Skew-symmetric matrix of odd order is singular
 (c) Skew-symmetric matrix of even order is always singular (d) None of these
- 109.** Choose the correct answer
 (a) Every identity matrix is a scalar matrix
 (b) Every scalar matrix is an identity matrix
 (c) Every diagonal matrix is an identity matrix
 (d) A square matrix whose each element is 1 is an identity matrix.
- 110.** If A and B are two square matrices such that $B = -A^{-1}BA$, then $(A + B)^2 =$ [EAMCET 2000]
 (a) 0 (b) $A^2 + B^2$ (c) $A^2 + 2AB + B^2$ (d) $A + B$
- 111.** For a matrix A , $AI = A$ and $AA^T = I$ is true for [Rajasthan PET 2000]
 (a) If A is a square matrix (b) If A is a non singular matrix (c) If A is a symmetric matrix
- 112.** If two matrices A and B are of order $p \times q$ and $r \times s$ respectively, can be subtracted only, if
 (a) $p = q$ (b) $p = q, r = s$ (c) $p = r, q = s$ (d) None of these
- 113.** The set of all 2×2 matrices over the real numbers is not a group under matrix multiplication because
 (a) Identity element does not exist (b) Closure property is not satisfied
 (c) Association property is not satisfied (d) Inverse axiom may not be satisfied
- 114.** If the matrix $AB = O$, then [Pb. CET 2000; Kurukshetra CEE 1998; Rajasthan PET 2001]
 (a) $A = O$ or $B = O$ (b) $A = O$ and $B = O$
 (c) It is not necessary that either $A = O$ or $B = O$ (d) $A \neq O, B \neq O$
- 115.** If $a_{ij} = \frac{1}{2}(3i - 2j)$ and $A = [a_{ij}]_{2 \times 2}$, then A is equal to [Rajasthan PET 2001]

- (a) $\begin{bmatrix} 1/2 & 2 \\ -1/2 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 1/2 & -1/2 \\ 2 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 1 \\ 1/2 & -1/2 \end{bmatrix}$ (d) None of these

116. Assuming that the sums and products given below are defined, which of the following is not true for matrices [Karnat

- (a) $A + B = B + A$ (b) $AB = AC$ does not imply $B = C$
 (c) $AB = O$ implies $A = O$ or $B = O$ (d) $(AB)' = B' A'$

117. Which of the following is true for matrix AB

[Rajasthan PET 2003]

- (a) $(AB)^{-1} = A^{-1} B^{-1}$ (b) $(AB)^{-1} = B^{-1} A^{-1}$ (c) $AB = BA$ (d) All of these

118. If A and B are 3×3 matrices such that $AB = A$ and $BA = B$, then

- (a) $A^2 = A$ and $B^2 \neq B$ (b) $A^2 \neq A$ and $B^2 = B$ (c) $A^2 = A$ and $B^2 = B$ (d) $A^2 \neq A$ and $B^2 \neq B$

119. If A and B are symmetric matrices of order n ($A \neq B$), then

- (a) $A + B$ is skew symmetric (b) $A + B$ is symmetric
 (c) $A + B$ is a diagonal matrix (d) $A - B$ is a zero matrix

120. The possible number of different order which a matrix can have when it has 24 elements is

- (a) 6 (b) 8 (c) 4 (d) 10

121. If $A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$ and $A^n = 0$, then minimum value of n is

- (a) 2 (b) 4 (c) 5 (d) 3

122. If A, B, C are square matrices of the same order, then which of the following is true

[JMIIEE 1997]

- (a) $AB = AC$ (b) $(AB)^2 = A^2 B^2$ (c) $AB = 0 \Rightarrow A = 0$ or $B = 0$ (d) $AB = I \Rightarrow AB = BA$

123. If a matrix has 13 elements, then the possible dimensions (order) it can have are

[MNR 1985]

- (a) $1 \times 13, 13 \times 1$ (b) $1 \times 26, 26 \times 1$ (c) $2 \times 13, 13 \times 2$ (d) None of these

Transpose of Matrices

Basic Level

124. If A, B, C are three $n \times n$ matrices, then $(ABC)' =$

[MP PET 1988]

- (a) $A' B' C'$ (b) $C' B' A'$ (c) $B' C' A'$ (d) $B' A' C'$

125. If $A = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$, then AA' =

[MP PET 1992]

- (a) 14 (b) $\begin{bmatrix} 1 \\ 4 \\ 3 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}$ (d) None of these

126. If $A = \begin{bmatrix} 1 & -2 \\ 5 & 3 \end{bmatrix}$, then $A + A^T$ equals

[Rajasthan PET 1994]

- (a) $\begin{bmatrix} 2 & 3 \\ 3 & 6 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & -4 \\ 10 & 6 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 4 \\ -10 & 6 \end{bmatrix}$ (d) None of these

127. If $A = \begin{bmatrix} 1 & -2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 1 \end{bmatrix}$, then $(AB)^T =$

[Rajasthan PET 1996, 2001]

384 Matrices

(a) $\begin{bmatrix} -3 & -2 \\ 10 & 7 \end{bmatrix}$

(b) $\begin{bmatrix} -3 & 10 \\ -2 & 7 \end{bmatrix}$

(c) $\begin{bmatrix} -3 & 10 \\ 7 & -2 \end{bmatrix}$

(d) $\begin{bmatrix} 3 & 10 \\ 2 & 7 \end{bmatrix}$

128. If $A = \begin{pmatrix} 1 & -2 & 1 \\ 2 & 1 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 1 \end{pmatrix}$, then $(AB)^T$ is equal to

[Rajasthan PET 2001]

(a) $\begin{pmatrix} -3 & -2 \\ 10 & 7 \end{pmatrix}$

(b) $\begin{pmatrix} -3 & 10 \\ -2 & 7 \end{pmatrix}$

(c) $\begin{pmatrix} -3 & 7 \\ 10 & 2 \end{pmatrix}$

(d) None of these

129. If $A = \begin{bmatrix} 0 & 1 & -2 \\ -1 & 0 & 5 \\ 2 & 5 & 0 \end{bmatrix}$, then

[MNR 1982]

(a) $A' = A$

(b) $A' = -A$

(c) $A' = 2A$

(d) None of these

130. Transpose of a row matrix is a

(a) Row matrix

(b) Column matrix

(c) A square matrix

(d) A scalar matrix

131. If $A = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 2 \\ -1 & 1 \end{bmatrix}$, then correct statement is

[Rajasthan PET 1987]

(a) $AB = BA$

(b) $AA^T = A^2$

(c) $AB = B^2$

(d) None of these

132. If matrix A is of order $m \times n$ and B is of order $n \times p$, then order of $(AB)^T$ is equal to

(a) Order of AB

(b) Order of BA

(c) Order of $A^T B^T$

(d) Order of $B^T A^T$

133. If $A = \begin{pmatrix} 4 & 2 & 7 \\ 6 & 0 & 8 \end{pmatrix}$, then AA^T is

[Rajasthan PET 1991]

(a) $\begin{pmatrix} 69 & 80 \\ 80 & 100 \end{pmatrix}$

(b) $\begin{pmatrix} 69 & 80 \\ 100 & 69 \end{pmatrix}$

(c) $\begin{pmatrix} 69 & 80 \\ 80 & 69 \end{pmatrix}$

(d) $\begin{pmatrix} 69 & 100 \\ 100 & 69 \end{pmatrix}$

134. Let A is a skew-symmetric matrix and C is a column matrix, then $C^T AC$ is

[Rajasthan PET 1995]

(a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(b) $[0]$

(c) $[1]$

(d) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

135. If A and B are matrices of suitable order and k is any number, then correct statement is

(a) $(AB)^T = A^T B^T$

(b) $(A+B)^T = A^T + B^T$

(c) $(AB)^{-1} = A^{-1} B^{-1}$

(d) $(kA)^T \neq kA^T$

136. If A and B are matrices of suitable order, then wrong statement is

(a) $(AB)^T = A^T B^T$

(b) $(A^T)^T = A$

(c) $(A-B)^T = A^T - B^T$

(d) $(A^T)^{-1} = (A^{-1})^T$

137. If A is a square matrix such that $|A| = 2$, then $|A'|$, where A' is transpose of A , is equal to

(a) 0

(b) -2

(c) 1/2

(d) 2

Special types of Matrices

Basic Level

138. An orthogonal matrix is

(a) $\begin{bmatrix} \cos \alpha & 2 \sin \alpha \\ -2 \sin \alpha & \cos \alpha \end{bmatrix}$

(b) $\begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$

(c) $\begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

139. Matrix $\begin{bmatrix} 0 & -4 & 1 \\ 4 & 0 & -5 \\ -1 & 5 & 0 \end{bmatrix}$ is
- (a) Orthogonal (b) Idempotent (c) Skew-symmetric (d) Symmetric
140. The inverse of a symmetric matrix is
- (a) Symmetric (b) Skew-symmetric (c) Diagonal matrix (d) None of these
141. If A is a symmetric matrix and $n \in \mathbb{N}$, then A^n is
- (a) Symmetric (b) Skew-symmetric (c) A diagonal matrix (d) None of these
142. If A is a skew-symmetric matrix and n is a positive integer, then A^n is
- (a) A symmetric matrix (b) Skew-symmetric matrix (c) Diagonal matrix (d) None of these
143. If $A = \begin{bmatrix} 4 & x+2 \\ 2x-3 & x+1 \end{bmatrix}$ is symmetric, then $x =$ [Karnataka CET 1994]
- (a) 3 (b) 5 (c) 2 (d) 4
144. If A is a square matrix, then $A + A^T$ is [Rajasthan PET 2001]
- (a) Non-singular matrix (b) Symmetric matrix (c) Skew-symmetric matrix (d) Unit matrix
145. For any square matrix A , AA^T is a [Rajasthan PET 2000]
- (a) Unit matrix (b) Symmetric matrix (c) Skew-symmetric matrix (d) Diagonal matrix
146. If A is a square matrix for which $a_{ij} = i^2 - j^2$, then A is [Rajasthan PET 1999]
- (a) Zero matrix (b) Unit matrix (c) Symmetric matrix (d) Skew-symmetric matrix
147. If A is a square matrix and $A + A^T$ is symmetric matrix, then $A - A^T =$
- (a) Unit matrix (b) Symmetric matrix (c) Skew-symmetric matrix (d) Zero matrix
148. The value of a for which the matrix $A = \begin{pmatrix} a & 2 \\ 2 & 4 \end{pmatrix}$ is singular if
- (a) $a \neq 1$ (b) $a = 1$ (c) $a = 0$ (d) $a = -1$
149. The matrix $A = \begin{bmatrix} i & 1-2i \\ -1-2i & 0 \end{bmatrix}$ is which of the following [Kurukshetra CEE 2002]
- (a) Symmetric (b) Skew-symmetric (c) Hermitian (d) Skew-hermitian
150. The matrix, $A = \begin{bmatrix} 1 & -3 & -4 \\ -1 & 3 & 4 \\ 1 & -3 & -4 \end{bmatrix}$ is nilpotent of index [Kurukshetra CEE 2002]
- (a) 2 (b) 3 (c) 4 (d) 6
151. If $\begin{bmatrix} x & y \\ u & v \end{bmatrix}$ is symmetric matrix, then
- (a) $x + v = 0$ (b) $x - v = 0$ (c) $y + u = 0$ (d) $y - u = 0$
152. The matrix $\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$ is a
- (a) Non-singular (b) Idempotent (c) Nilpotent (d) Orthogonal
153. For any square matrix A , which statement is wrong
- (a) $(adj A)^{-1} = adj(A^{-1})$ (b) $(A^T)^{-1} = (A^{-1})^T$ (c) $(A^3)^{-1} = (A^{-1})^3$ (d) None of these

386 Matrices

154. If $A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 3 & 0 & 0 \\ 4 & 5 & 6 & 0 \\ 7 & 8 & 9 & 10 \end{bmatrix}$, then A is
- (a) An upper triangular matrix (b) A null matrix
(c) A lower triangular matrix (d) None of these
155. If A is a square matrix, then A will be non-singular if
- (a) $|A| = 0$ (b) $|A| > 0$ (c) $|A| < 0$ (d) $|A| \neq 0$
156. The matrix $\begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$ is
- (a) Symmetric (b) Skew-symmetric (c) Scalar (d) None of these
157. If $A = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$, then A^2 is [MNR 1980]
- (a) Null matrix (b) Unit matrix (c) A (d) $2A$
158. If A is a symmetric matrix, then matrix $M'AM$ is [MP PET 1990]
- (a) Symmetric (b) Skew-symmetric (c) Hermitian (d) Skew-Hermitian
159. If A is a square matrix, then which of the following matrices is not symmetric
- (a) $A + A'$ (b) AA' (c) $A'A$ (d) $A - A'$
160. Square matrix $[a_{ij}]_{n \times n}$ will be an upper triangular matrix, if
- (a) $a_{ij} \neq 0$ for $i > j$ (b) $a_{ij} = 0$ for $i > j$ (c) $a_{ij} = 0$ for $i < j$ (d) None of these
161. If the matrix $\begin{bmatrix} 0 & 1 & -2 \\ -1 & 0 & 3 \\ \lambda & -3 & 0 \end{bmatrix}$ is singular, then $\lambda =$ [MP PET 1989]
- (a) -2 (b) -1 (c) 1 (d) 2
162. In order that the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & \lambda & 5 \end{bmatrix}$ be non-singular, λ should not be equal to [Kurukshetra CEE 1998]
- (a) 1 (b) 2 (c) 3 (d) 4
163. If A is involutory matrix and I is unit matrix of same order, then $(I - A)(I + A)$ is
- (a) Zero matrix (b) A (c) I (d) $2A$
164. If $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$, then A is
- (a) Symmetric (b) Skew-symmetric (c) Non-singular (d) Singular
165. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$, then $A^2 =$ [MNR 1980; Pb. CET 1990]
- (a) Unit matrix (b) Null matrix (c) A (d) $-A$

166. If the matrix $\begin{bmatrix} 1 & 3 & \lambda + 2 \\ 2 & 4 & 8 \\ 3 & 5 & 10 \end{bmatrix}$ is singular, then $\lambda =$ [MP PET 1990; Pb. CET 2000]
 (a) -2 (b) 4 (c) 2 (d) -4
167. Out of the following a skew-symmetric matrix is [MP PET 1992]
 (a) $\begin{bmatrix} 0 & 4 & 5 \\ -4 & 0 & -6 \\ -5 & 6 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 4 & 5 \\ -4 & 1 & -6 \\ -5 & 6 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 4 & 5 \\ -4 & 2 & -6 \\ -5 & 6 & 3 \end{bmatrix}$ (d) $\begin{bmatrix} i+1 & 4 & 5 \\ -4 & i & -6 \\ -5 & 6 & i \end{bmatrix}$
168. If $A = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 4 & 4 \\ 3 & 5 & 6 \end{bmatrix}$, then A is
 (a) Singular (b) Non-singular (c) Unitary (d) Symmetric
169. If A, B, C are three square matrices such that $AB = AC$ implies $B = C$, then the matrix A is always [MP PET 1989; Karnataka CET 1992]
 (a) A singular matrix (b) A Non-singular matrix (c) An orthogonal matrix (d) A diagonal matrix
170. The matrix $A = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}$ is [MP PET 1988]
 (a) Unitary (b) Orthogonal (c) Nilpotent (d) Involutary
171. If a matrix A is symmetric as well as skew symmetric, then
 (a) A is a diagonal matrix (b) A is a null matrix (c) A is a unit matrix (d)
172. A and B are any two square matrices. Which one of the following is a skew symmetric matrix
 (a) $\frac{A+A'}{2}$ (b) $\frac{A+B}{2}$ (c) $\frac{A'-A}{2}$ (d) None of the above.
173. Choose the correct answer
 (a) Every scalar matrix is an identity matrix
 (b) Every identity matrix is a scalar matrix
 (c) Every diagonal matrix is an identity matrix
 (d) A Square matrix whose each element is 1 is an identity matrix
174. For a square matrix A , it is given that $AA' = I$, then A is a [DCE 1998]
 (a) Orthogonal matrix (b) Diagonal matrix (c) Symmetric matrix (d) None of these
175. A square matrix can always be expressed as a [DCE1998]
 (a) Sum of a symmetric matrix and a skew-symmetric matrix (b) Sum of a diagonal matrix and a symmetric matrix
 (c) Skew matrix (d) Skew-symmetric matrix
176. If A is a skew-symmetric matrix and n is odd positive integer, then A^n is
 (a) A symmetric matrix (b) A skew-symmetric matrix (c) A diagonal matrix (d) None of these
177. If A, B symmetric matrices of the same order then $AB - BA$ is
 (a) Symmetric matrix (b) Skew-symmetric matrix (c) Null matrix (d) Unit matrix

178. If k is a scalar and I is a unit matrix of order 3, then $\text{adj}(kI) =$

- (a) $k^3 I$ (b) $k^2 I$ (c) $-k^3 I$ (d) $-k^2 I$

179. If $A = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $\text{adj } A =$

- (a) A (b) I (c) O (d) A^2

180. If A is a $n \times n$ matrix, then $\text{adj}(\text{adj } A) =$

- (a) $|A|^{n-1} A$ (b) $|A|^{n-2} A$ (c) $|A|^n A$ (d) None of these

181. Adjoint of the matrix $N = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix}$ is

[MP PET 1989]

- (a) N (b) $2N$ (c) $-N$ (d) None of these

182. If A is a non-singular matrix, then $A(\text{adj } A) =$

- (a) A (b) I (c) $|A| I$ (d) $|A|^2 I$

183. If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ and $A \text{adj } A = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}$, then k is equal to

- (a) 0 (b) 1 (c) $\sin \alpha \cos \alpha$ (d) $\cos 2\alpha$

184. Let $A = \begin{bmatrix} 1 & 0 & 0 \\ 5 & 2 & 0 \\ -1 & 6 & 1 \end{bmatrix}$, then the adjoint of A is

[MNR 1982]

- (a) $\begin{bmatrix} 2 & -5 & 32 \\ 0 & 1 & -6 \\ 0 & 0 & 2 \end{bmatrix}$ (b) $\begin{bmatrix} -1 & 0 & 0 \\ -5 & -2 & 0 \\ 1 & -6 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} -1 & 0 & 0 \\ -5 & -2 & 0 \\ 1 & -6 & -1 \end{bmatrix}$ (d) None of these

185. If $A = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix}$, then $A(\text{adj } A) =$

[MP PET 1995; Rajasthan PET

1997]

- (a) $\begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$ (b) $\begin{bmatrix} 0 & 10 \\ 10 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 10 & 1 \\ 1 & 10 \end{bmatrix}$ (d) None of these

186. If A is a singular matrix, then $\text{adj } A$ is

[Karnataka CET 1993]

- (a) Singular (b) Non-singular (c) Symmetric (d) Not defined

187. The adjoint of $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & -3 \\ 2 & -1 & 3 \end{bmatrix}$ is

- (a) $\begin{bmatrix} 3 & -9 & -5 \\ -4 & 1 & 3 \\ -5 & 4 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 3 & -4 & -5 \\ -9 & 1 & 4 \\ -5 & 3 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} -3 & 4 & 5 \\ 9 & -1 & -4 \\ 5 & -3 & -1 \end{bmatrix}$ (d) None of these

188. $\text{Adj} \cdot (AB) - (\text{Adj} \cdot B)(\text{Adj} \cdot A) =$

[MP PET 1997]

- (a) $\text{Adj} \cdot A - \text{Adj} \cdot B$ (b) I (c) O (d) None of these

- 189.** If $A = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 2 & 0 & 1 \end{pmatrix}$, then $\text{adj } A =$ [Rajasthan PET 1996]
- (a) $\begin{pmatrix} 1 & 4 & -2 \\ -2 & 1 & 4 \\ 4 & -2 & 1 \end{pmatrix}$ (b) $\begin{pmatrix} 1 & -2 & 4 \\ 4 & 1 & -2 \\ -2 & 4 & 1 \end{pmatrix}$ (c) $\begin{pmatrix} 1 & 2 & 4 \\ -4 & 1 & 2 \\ -4 & -2 & 1 \end{pmatrix}$ (d) None of these
- 190.** If $A = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 4 & 9 \\ 1 & 8 & 27 \end{pmatrix}$, then the value of $|\text{adj } A|$ is [Rajasthan PET 1999]
- (a) 36 (b) 72 (c) 144 (d) None of these
- 191.** If A is a matrix of order 3 and $|A| = 8$, then $|\text{adj } A| =$ [DCE 1999; Karnataka CET 2002]
- (a) 1 (b) 2 (c) 2^3 (d) 2^6
- 192.** If A and B are non-singular square matrices of same order, then $\text{adj}(AB)$ is equal to [AMU 1999]
- (a) $(\text{adj } A)(\text{adj } B)$ (b) $(\text{adj } B)(\text{adj } A)$ (c) $(\text{adj } B^{-1})(\text{adj } A^{-1})$ (d) $(\text{adj } A^{-1})(\text{adj } B^{-1})$
- 193.** If d is the determinant of a square matrix A of order n , then the determinant of its adjoint is
- (a) d^n (b) d^{n-1} (c) d^{n+1} (d) d
- 194.** If $A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$, then $\text{adj } A$ is equal to [Rajasthan PET 2001]
- (a) $\begin{bmatrix} -3 & -1 \\ 2 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} 3 & 1 \\ -2 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 3 & -2 \\ 1 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix}$
- 195.** If $A = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}$, then $A(\text{adj } A) =$ [Rajasthan PET 2002]
- (a) I (b) $|A|$ (c) $|A|I$ (d) None of these
- 196.** If $A = \begin{bmatrix} -2 & 6 \\ -5 & 7 \end{bmatrix}$, then $\text{adj}(A)$ is
- (a) $\begin{bmatrix} 7 & -6 \\ 5 & -2 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & -6 \\ 5 & -7 \end{bmatrix}$ (c) $\begin{bmatrix} 7 & -5 \\ 6 & -2 \end{bmatrix}$ (d) None of these
- 197.** The adjoint matrix of $\begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$ is [MP PET 2003]
- (a) $\begin{bmatrix} 4 & 8 & 3 \\ 2 & 1 & 6 \\ 0 & 2 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix}$ (c) $\begin{bmatrix} 11 & 9 & 3 \\ 1 & 2 & 8 \\ 6 & 9 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & -2 & 1 \\ -1 & 3 & 3 \\ -2 & 3 & -3 \end{bmatrix}$
- 198.** If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$, then $A(\text{adj}(A)) =$ [Rajasthan PET 2003]
- (a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}$
- 199.** If $A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$, then the value of $|A| \cdot |\text{Adj } A|$ is [AMU 1987]
- (a) a^3 (b) a^6 (c) a^9 (d) a^{27}

390 Matrices

200. If $A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{bmatrix}$, then the value of $|adj A|$ is [AMU 1989]

- (a) a^3 (b) a^6 (c) a^9 (d) a^{27}

201. If $A = \begin{bmatrix} 1 & 2 & -1 \\ -1 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}$, then determinant $(adj(adj A))$ is

- (a) $(14)^1$ (b) $(14)^2$ (c) $(14)^3$ (d) $(14)^4$

202. If A is a square matrix, then $adj(A') - (adj A)'$ is equal to

- (a) $2|A|$ (b) $2|A|I$ (c) Null matrix (d) Unit matrix

203. If $A = \begin{bmatrix} 1 & -2 & 3 \\ 4 & 0 & -1 \\ -3 & 1 & 5 \end{bmatrix}$, then $(adj A)_{23}$ is equal to [Rajasthan PET 1984]

- (a) 13 (b) -13 (c) 5 (d) -5

204. For a third order non-singular matrix A , $|A(adj A)|$ equals

- (a) $|A|$ (b) $|A|^2$ (c) $|A|^3$ (d) None of these

Advance Level

205. If A be a square matrix of order n and if $|A| = D$ and $|adj A| = D'$, then [Rajasthan PET 2000]

- (a) $DD' = D^2$ (b) $DD' = D^{n-1}$ (c) $DD' = D^n$ (d) None of these

Inverse of a Matrix

Basic Level

206. Inverse of the matrix $\begin{bmatrix} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{bmatrix}$ is [MP PET 1990]

- (a) $\begin{bmatrix} 1 & 2 & 3 \\ 3 & 3 & 7 \\ -2 & -4 & -5 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -3 & 5 \\ 7 & 4 & 6 \\ 4 & 2 & 7 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 2 & -4 \\ 8 & -4 & -5 \\ 3 & 5 & 2 \end{bmatrix}$

207. If A and B are non-singular matrices, then [MP PET 1991; Kurukshetra CEE 1998]

- (a) $(AB)^{-1} = A^{-1}B^{-1}$ (b) $AB = BA$ (c) $(AB)' = A'B'$ (d) $(AB)^{-1} = B^{-1}A^{-1}$

208. If $A = \begin{bmatrix} i & 0 \\ 0 & i/2 \end{bmatrix}$, ($i = \sqrt{-1}$), then $A^{-1} =$ [MP PET 1992]

- (a) $\begin{bmatrix} i & 0 \\ 0 & i/2 \end{bmatrix}$ (b) $\begin{bmatrix} -i & 0 \\ 0 & -2i \end{bmatrix}$ (c) $\begin{bmatrix} i & 0 \\ 0 & 2i \end{bmatrix}$ (d) $\begin{bmatrix} 0 & i \\ 2i & 0 \end{bmatrix}$

209. If $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$, then $A^{-1} =$

- (a) $\begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ (b) $\begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ (c) $\begin{bmatrix} -\cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{bmatrix}$ (d) None of these

210. If $A = \begin{bmatrix} a & c \\ d & b \end{bmatrix}$, then $A^{-1} =$ [MP PET 1988]

- (a) $\frac{1}{ab-cd} \begin{bmatrix} b & -c \\ -d & a \end{bmatrix}$ (b) $\frac{1}{ad-bc} \begin{bmatrix} b & -c \\ -d & a \end{bmatrix}$ (c) $\frac{1}{ab-cd} \begin{bmatrix} b & d \\ c & a \end{bmatrix}$ (d) None of these

211. The element of second row and third column in the inverse of $\begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$ is [MP PET 1992]

- (a) -2 (b) -1 (c) 1 (d) 2

212. The inverse of the matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is [MP PET 1989; Pb. CET 1989, 93]

- (a) $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$

213. The inverse of $\begin{bmatrix} 2 & -3 \\ -4 & 2 \end{bmatrix}$ is

- (a) $\frac{-1}{8} \begin{bmatrix} 2 & 3 \\ 4 & 2 \end{bmatrix}$ (b) $\frac{-1}{8} \begin{bmatrix} 3 & 2 \\ 2 & 4 \end{bmatrix}$ (c) $\frac{1}{8} \begin{bmatrix} 2 & 3 \\ 4 & 2 \end{bmatrix}$ (d) $\frac{1}{8} \begin{bmatrix} 3 & 2 \\ 2 & 4 \end{bmatrix}$

214. The inverse of the matrix $\begin{bmatrix} 3 & -2 \\ 1 & 4 \end{bmatrix}$ is [MP PET 1994]

- (a) $\begin{bmatrix} \frac{4}{14} & \frac{2}{14} \\ \frac{-1}{14} & \frac{3}{14} \end{bmatrix}$ (b) $\begin{bmatrix} \frac{3}{14} & \frac{-2}{14} \\ \frac{1}{14} & \frac{4}{14} \end{bmatrix}$ (c) $\begin{bmatrix} \frac{4}{14} & \frac{-2}{14} \\ \frac{1}{14} & \frac{3}{14} \end{bmatrix}$ (d) $\begin{bmatrix} \frac{3}{14} & \frac{2}{14} \\ \frac{1}{14} & \frac{4}{14} \end{bmatrix}$

215. If a matrix A is such that $3A^3 + 2A^2 + 5A + I = 0$, then its inverse is

- (a) $-(3A^2 + 2A + 5I)$ (b) $3A^2 + 2A + 5I$ (c) $3A^2 - 2A - 5I$ (d) None of these

216. If $F(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $G(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}$, then $[F(\alpha)G(\beta)]^{-1} =$

- (a) $F(\alpha) - G(\beta)$ (b) $-F(\alpha) - G(\beta)$ (c) $[F(\alpha)]^{-1}[G(\beta)]^{-1}$ (d) $[G(\beta)]^{-1}[F(\alpha)]^{-1}$

217. If $A = \begin{bmatrix} 1 & \tan \theta/2 \\ -\tan \theta/2 & 1 \end{bmatrix}$ and $AB = I$, then $B =$ [MP PET 1995, 98]

- (a) $\cos^2 \frac{\theta}{2} \cdot A$ (b) $\cos^2 \frac{\theta}{2} \cdot A^T$ (c) $\cos^2 \frac{\theta}{2} \cdot I$ (d) None of these

218. If $\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, then the matrix $A =$

- (a) $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$

392 Matrices

- 219.** If A is an invertible matrix, then which of the following is correct
 (a) A^{-1} is multivalued (b) A^{-1} is singular (c) $(A^{-1})^T \neq (A^T)^{-1}$ (d) $|A| \neq 0$
- 220.** If $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$, then $A^{-1} =$ [AMU 1988]
 (a) $\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$ (b) $\begin{bmatrix} -a & 0 & 0 \\ 0 & -b & 0 \\ 0 & 0 & -c \end{bmatrix}$ (c) $\begin{bmatrix} 1/a & 0 & 0 \\ 0 & 1/b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$ (d) None of these
- 221.** $\begin{bmatrix} 1 & 3 \\ 3 & 10 \end{bmatrix}^{-1} =$ [EAMCET 1994; DCE 1999]
 (a) $\begin{bmatrix} 10 & 3 \\ 3 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 10 & -3 \\ -3 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 3 \\ 3 & 10 \end{bmatrix}$ (d) $\begin{bmatrix} -1 & -3 \\ -3 & -10 \end{bmatrix}$
- 222.** If $A = \begin{bmatrix} 5 & 2 \\ 3 & 1 \end{bmatrix}$, then $A^{-1} =$ [EAMCET 1988]
 (a) $\begin{bmatrix} 1 & -2 \\ -3 & 5 \end{bmatrix}$ (b) $\begin{bmatrix} -1 & 2 \\ 3 & -5 \end{bmatrix}$ (c) $\begin{bmatrix} -1 & -2 \\ -3 & -5 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 2 \\ 3 & 5 \end{bmatrix}$
- 223.** $\begin{bmatrix} -6 & 5 \\ -7 & 6 \end{bmatrix}^{-1} =$ [Karnataka CET 1994]
 (a) $\begin{bmatrix} -6 & 5 \\ -7 & 6 \end{bmatrix}$ (b) $\begin{bmatrix} 6 & -5 \\ -7 & 6 \end{bmatrix}$ (c) $\begin{bmatrix} 6 & 5 \\ 7 & 6 \end{bmatrix}$ (d) $\begin{bmatrix} 6 & -5 \\ 7 & -6 \end{bmatrix}$
- 224.** The inverse of matrix $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is [Karnataka CET 1993]
 (a) A (b) A^T (c) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$
- 225.** The inverse of $\begin{bmatrix} 3 & 5 & 7 \\ 2 & -3 & 1 \\ 1 & 1 & 2 \end{bmatrix}$ is [EAMCET 1989]
 (a) $\begin{bmatrix} 7 & 3 & -26 \\ 3 & 1 & -11 \\ -5 & -2 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 7 & 3 & -26 \\ 3 & 1 & 11 \\ -5 & -2 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 3 & 1 & 11 \\ 7 & 3 & -26 \\ -5 & 2 & 1 \end{bmatrix}$ (d) None of these
- 226.** The inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$ is
 (a) $\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$ (d) None of these
- 227.** The matrix $\begin{bmatrix} \lambda & -1 & 4 \\ -3 & 0 & 1 \\ -1 & 1 & 2 \end{bmatrix}$ is invertible, if [Kurukshetra CEE 1996]
 (a) $\lambda \neq -15$ (b) $\lambda \neq -17$ (c) $\lambda \neq -16$ (d) $\lambda \neq -18$
- 228.** If $A = \begin{pmatrix} 3 & 2 \\ 0 & 1 \end{pmatrix}$, then $(A^{-1})^3$ is equal to [MP PET 1997]

(a) $\frac{1}{27} \begin{pmatrix} 1 & -26 \\ 0 & 27 \end{pmatrix}$ (b) $\frac{1}{27} \begin{pmatrix} -1 & 26 \\ 0 & 27 \end{pmatrix}$ (c) $\frac{1}{27} \begin{pmatrix} 1 & -26 \\ 0 & -27 \end{pmatrix}$ (d) $\frac{1}{27} \begin{pmatrix} -1 & -26 \\ 0 & -27 \end{pmatrix}$

229. The matrix $\begin{bmatrix} 1 & a & 2 \\ 1 & 2 & 5 \\ 2 & 1 & 1 \end{bmatrix}$ is not invertible, if 'a' has the value

- (a) 2 (b) 1 (c) 0 (d) -1

230. Inverse matrix of $\begin{bmatrix} 4 & 7 \\ 1 & 2 \end{bmatrix}$

[Rajasthan PET 1996,

2001]

(a) $\begin{bmatrix} 2 & -7 \\ -1 & 4 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & -1 \\ -7 & 4 \end{bmatrix}$ (c) $\begin{bmatrix} -2 & 7 \\ 1 & -4 \end{bmatrix}$ (d) $\begin{bmatrix} -2 & 1 \\ 7 & -4 \end{bmatrix}$

231. If the multiplicative group of 2×2 matrices of the form $\begin{pmatrix} a & a \\ a & a \end{pmatrix}$, for $a \neq 0$ and $a \in R$, then the inverse of $\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix}$ is

[Karnataka CET 1999]

(a) $\begin{pmatrix} \frac{1}{8} & \frac{1}{8} \\ \frac{1}{8} & \frac{1}{8} \end{pmatrix}$ (b) $\begin{pmatrix} \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}$ (c) $\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$ (d) Does not exist

232. The element in the first row and third column of the inverse of the matrix $\begin{bmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$ is

- (a) -2 (b) 0 (c) 1 (d) 7

233. If I_3 is the identity matrix of order 3, then I_3^{-1} is

[Pb. CET 2000]

- (a) 0 (b) $3I_3$ (c) I_3 (d) Does not exist

234. If a matrix A is such that $4A^3 + 2A^2 + 7A + I = O$, then A^{-1} equals

- (a) $(4A^2 + 2A + 7I)$ (b) $-(4A^2 + 2A + 7I)$ (c) $-(4A^2 - 2A + 7I)$ (d) $(4A^2 + 2A - 7I)$

235. If $A = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then $(B^{-1}A^{-1})^{-1} =$

[EAMCET 2001]

(a) $\begin{bmatrix} 2 & -2 \\ 2 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 3 & -2 \\ 2 & 2 \end{bmatrix}$ (c) $\frac{1}{10} \begin{bmatrix} 2 & 2 \\ -2 & 3 \end{bmatrix}$ (d) $\frac{1}{10} \begin{bmatrix} 3 & 2 \\ -2 & 2 \end{bmatrix}$

236. If $A^2 - A + I = 0$, then $A^{-1} =$

[Kerala (Engg.) 2001]

- (a) A^{-2} (b) $A + I$ (c) $I - A$ (d) $A - I$

237. If $A = \begin{bmatrix} 2 & 3 \\ 4 & 6 \end{bmatrix}$, then $A^{-1} =$

[Karnataka CET 2001]

(a) $\begin{bmatrix} 1 & 2 \\ -3/2 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & -3 \\ 4 & 6 \end{bmatrix}$ (c) $\begin{bmatrix} -2 & 4 \\ -3 & 6 \end{bmatrix}$ (d) Does not exist

238. If for the matrix A , $A^3 = I$, then $A^{-1} =$

[Rajasthan PET 2002]

- (a) A^2 (b) A^3 (c) A (d) None of these

239. For two invertible matrices A and B of suitable orders, the value of $(AB)^{-1}$ is

[Rajasthan PET 2000, 02;

Karnataka CET 2001]

- (a) $(BA)^{-1}$ (b) $B^{-1}A^{-1}$ (c) $A^{-1}B^{-1}$ (d) $(AB')^{-1}$

240. If $A = \begin{bmatrix} -1 & 2 \\ 2 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$, $AX = B$, then $X =$

[MP PET 2002]

(a) $\begin{bmatrix} 5 & 7 \end{bmatrix}$ (b) $\frac{1}{3} \begin{bmatrix} 5 \\ 7 \end{bmatrix}$ (c) $\frac{1}{3} \begin{bmatrix} 5 & 7 \end{bmatrix}$ (d) $\begin{bmatrix} 5 \\ 7 \end{bmatrix}$

394 Matrices

241. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$, then $A^{-1} =$

- (a) $\begin{bmatrix} -5 & -2 \\ -3 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (c) $\begin{bmatrix} -\frac{5}{11} & -\frac{2}{11} \\ -\frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (d) $\begin{bmatrix} 5 & 2 \\ 3 & -1 \end{bmatrix}$

242. If $A = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$ and $A^{-1} = \lambda(\text{adj}(A))$, then $\lambda =$

[UPSEAT 2002]

- (a) $-\frac{1}{6}$ (b) $\frac{1}{3}$ (c) $-\frac{1}{3}$ (d) $\frac{1}{6}$

243. The multiplicative inverse of matrix $\begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix}$ is

[DCE 2002]

- (a) $\begin{bmatrix} 4 & -1 \\ -7 & -2 \end{bmatrix}$ (b) $\begin{bmatrix} -4 & -1 \\ 7 & -2 \end{bmatrix}$ (c) $\begin{bmatrix} 4 & -7 \\ 7 & 2 \end{bmatrix}$ (d) $\begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$

244. The inverse matrix of $\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$ is

[MP PET 2003]

- (a) $\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ -4 & 3 & -1 \\ \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{bmatrix}$ (b) $\begin{bmatrix} \frac{1}{2} & -4 & \frac{5}{2} \\ 1 & -6 & 3 \\ 1 & 2 & -1 \end{bmatrix}$ (c) $\frac{1}{2} \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 4 & 2 & 3 \end{bmatrix}$ (d) $\frac{1}{2} \begin{bmatrix} 1 & -1 & -1 \\ -8 & 6 & -2 \\ 5 & -3 & 1 \end{bmatrix}$

245. Inverse of the matrix $\begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix}$ is

- (a) $\frac{1}{10} \begin{bmatrix} 4 & 2 \\ -3 & 1 \end{bmatrix}$ (b) $\frac{1}{10} \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix}$ (c) $\frac{1}{10} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}$ (d) $\begin{bmatrix} 4 & 2 \\ -3 & 1 \end{bmatrix}$

246. If A is an orthogonal matrix, then A^{-1} is equal to

- (a) A (b) A' (c) A^2 (d) None of these

247. The multiplicative inverse of the matrix $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$ is

- (a) $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 0 & 1 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$ (c) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$

248. Let A be an invertible matrix. Which of the following is not true

- (a) $A^{-1} \neq |A|^{-1}$ (b) $(A^2)^{-1} = (A^{-1})^2$ (c) $(A')^{-1} = (A^{-1})'$ (d) None of these

249. Inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 6 \end{bmatrix}$ is

[Kurukshetra CEE 1995]

- (a) $\begin{bmatrix} 2 & 0 & -1 \\ 0 & -3 & 2 \\ -1 & 2 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 6 \end{bmatrix}$ (c) $\begin{bmatrix} -2 & 0 & 1 \\ 0 & 3 & -2 \\ 1 & -2 & 1 \end{bmatrix}$ (d) None of these

250. If $A = \begin{bmatrix} 1 & 2 \\ -4 & -1 \end{bmatrix}$, then $A^{-1} =$

[Karnataka CET 1997]

- (a) $\frac{1}{7} \begin{bmatrix} 1 & 2 \\ -4 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} -1 & -2 \\ 4 & 1 \end{bmatrix}$ (c) $\frac{1}{9} \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix}$ (d) $\frac{1}{7} \begin{bmatrix} -1 & -2 \\ 4 & 1 \end{bmatrix}$

251. If $A = \begin{bmatrix} 0 & -1 & 2 \\ 2 & -2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$ and $M = AB$, then M^{-1} is equal to

- (a) $\begin{bmatrix} 2 & -2 \\ 2 & 1 \end{bmatrix}$ (b) $\begin{bmatrix} 1/3 & 1/3 \\ -1/3 & 1/6 \end{bmatrix}$ (c) $\begin{bmatrix} 1/3 & -1/3 \\ 1/3 & 1/6 \end{bmatrix}$ (d) $\begin{bmatrix} 1/3 & -1/3 \\ -1/3 & 1/6 \end{bmatrix}$

252. If for a square matrix A , $AA^{-1} = I$, then A is

[DCE 1998]

- (a) Orthogonal matrix (b) Symmetric matrix (c) Diagonal matrix (d) Invertible matrix

253. If matrix $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & \lambda & 5 \end{bmatrix}$ is invertible, then

[Kurukshetra CEE 1998]

- (a) $\lambda \neq 4$ (b) $\lambda \neq 3$ (c) $\lambda \neq 2$ (d) $\lambda \neq 0$

254. If $\begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, then

- (a) $a = 1, b = 1$ (b) $a = \cos 2\theta, b = \sin 2\theta$ (c) $a = \sin 2\theta, b = \cos 2\theta$ (d) None of these

Advance Level

255. If $A = \begin{bmatrix} 1 & -2 & 3 \\ 0 & -1 & 4 \\ -2 & 2 & 1 \end{bmatrix}$, then $(A')^{-1} =$

- (a) $\begin{bmatrix} -9 & -8 & -2 \\ 8 & 7 & 2 \\ -5 & -4 & -1 \end{bmatrix}$ (b) $\begin{bmatrix} 1 & 0 & -2 \\ -2 & -1 & 2 \\ 3 & 4 & 1 \end{bmatrix}$ (c) $\begin{bmatrix} -9 & 8 & -5 \\ -8 & 7 & -4 \\ 2 & 2 & -1 \end{bmatrix}$ (d) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

256. If ω is a cube root of unity and $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{bmatrix}$, then $A^{-1} =$

- (a) $\begin{bmatrix} 1 & \omega & \omega^2 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{bmatrix}$ (b) $\frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{bmatrix}$ (c) $\begin{bmatrix} 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \\ 1 & 1 & 1 \end{bmatrix}$ (d) $\frac{1}{2} \begin{bmatrix} 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \\ 1 & 1 & 1 \end{bmatrix}$

257. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then $A^{-1} =$

[DCE 1999]

- (a) A (b) A^2 (c) A^3 (d) A^4

258. If $D = \text{diag}(d_1, d_2, d_3, \dots, d_n)$, where $d_i \neq 0$ for all $i = 1, 2, \dots, n$, then D^{-1} is equal to

- (a) D (b) $\text{diag}(d_1^{-1}, d_2^{-1}, \dots, d_n^{-1})$ (c) I (d) None of these

259. If $A = \text{diag}(d_1, d_2, d_3, \dots, d_n)$, then A^n is equal to

- (a) $\text{diag}(d_1^{n-1}, d_2^{n-1}, d_3^{n-1}, \dots, d_n^{n-1})$ (b) $\text{diag}(d_1^n, d_2^n, d_3^n, \dots, d_n^n)$
(c) A (d) None of these

Relation between Determinants and Matrices

Basic Level

- 260.** If A is a square matrix of order 3, then true statement is (where I is unit matrix) [MP PET 1992]
 (a) $\det(-A) = -\det A$ (b) $\det A = 0$ (c) $\det(A + I) = 1 + \det A$ (d) $\det 2A = 2 \det A$
- 261.** If $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 3 \\ 0 & 0 & 2 \end{bmatrix}$, then $|AB|$ is equal to [Rajasthan PET 1995]
 (a) 4 (b) 8 (c) 16 (d) 32
- 262.** If A and B are square matrices of order 3 such that $|A| = -1$, $|B| = 3$, then $|3AB| =$ [IIT 1988; MP PET 1995, 99]
 (a) -9 (b) -81 (c) -27 (d) 81
- 263.** Which of the following is correct
 (a) Determinant is a square matrix (b) Determinant is a number associated to a matrix
 (c) Determinant is a number associated to a square matrix (d) None of these
- 264.** Let A be a skew-symmetric matrix of odd order, then $|A|$ is equal to
 (a) 0 (b) 1 (c) -1 (d) None of these
- 265.** Let A be a skew-symmetric matrix of even order, then $|A|$
 (a) Is a perfect square (b) Is not a perfect square (c) Is always zero (d) None of these
- 266.** For any 2×2 matrix A , if $A(\text{adj.}A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$, then $|A| =$ [MP PET 1999]
 (a) 0 (b) 10 (c) 20 (d) 100
- 267.** If $A = \begin{bmatrix} 1 & 3 \\ 2 & 1 \end{bmatrix}$, then determinant of $A^2 - 2A$ is [EAMCET 2000]
 (a) 5 (b) 25 (c) -5 (d) -25
- 268.** If $\begin{bmatrix} 2+x & 3 & 4 \\ 1 & -1 & 2 \\ x & 1 & -5 \end{bmatrix}$ is a singular matrix, then x is [Kerala (Engg.) 2001]
 (a) $\frac{13}{25}$ (b) $-\frac{25}{13}$ (c) $\frac{5}{13}$ (d) $\frac{25}{13}$
- 269.** The product of a matrix and its transpose is an identity matrix. The value of determinant of this matrix is
 (a) -1 (b) 0 (c) ± 1 (d) 1
- 270.** If $A = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix}$, then $\det A =$
 (a) 2 (b) 3 (c) 4 (d) 5
- 271.** If $A \neq O$ and $B \neq O$ are $n \times n$ matrix such that $AB = O$, then
 (a) $\text{Det}(A) = 0$ or $\text{Det}(B) = 0$ (b) $\text{Det}(A) = 0$ and $\text{Det}(B) = 0$
 (c) $\text{Det}(A) = \text{Det}(B) \neq 0$ (d) $A^{-1} = B^{-1}$
- 272.** If A is a square matrix such that $A^2 = A$, then $\det(A)$ equals
 (a) 0 or 1 (b) -2 or 2 (c) -3 or 3 (d) None of these
- 273.** If A is a square matrix such that $|A| = 2$, then for any +ve integer n , $|A^n|$ is equal to
 (a) 0 (b) $2n$ (c) 2^n (d) n^2
- 274.** If A is a square matrix of order 3 and entries of A are positive integers, then $|A|$ is
 (a) Different from zero (b) 0 (c) Positive (d) An arbitrary integer.
- 275.** If A and B are any 2×2 matrix, then $\det(A+B) = 0$ implies
 (a) $\text{Det}A + \text{Det}B = 0$ (b) $\text{Det}A = 0$ or $\text{Det}B = 0$ (c) $\text{Det}A = 0$ and $\text{Det}B = 0$ (d) None of these

276. If $\begin{bmatrix} x+y+z \\ x+y \\ y+z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix}$, then $(x, y, z) =$

- (a) (4, 3, 2) (b) (3, 2, 4) (c) (2, 3, 4) (d) None of these

277. The solution of the equation $\begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ is $(x, y, z) =$

- (a) (1, 1, 1) (b) (0, -1, 2) (c) (-1, 2, 2) (d) (-1, 0, 2)

278. If $AX = B$, $B = \begin{bmatrix} 9 \\ 52 \\ 0 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} 3 & -\frac{1}{2} & -\frac{1}{2} \\ -4 & \frac{3}{4} & \frac{5}{4} \\ 2 & -\frac{1}{4} & -\frac{3}{4} \end{bmatrix}$, then X is equal to

- (a) $\begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}$ (b) $\begin{bmatrix} -\frac{1}{2} \\ -\frac{1}{2} \\ 2 \end{bmatrix}$ (c) $\begin{bmatrix} -4 \\ 2 \\ 3 \end{bmatrix}$ (d) $\begin{bmatrix} 3 \\ \frac{3}{4} \\ \frac{3}{4} \\ -\frac{3}{4} \end{bmatrix}$

Rank of a Matrix

Basic Level

279. If A is a non-zero column matrix of order $m \times 1$ and B is a non-zero row matrix of order $1 \times n$, then rank of AB is equal to

- (a) m (b) n (c) 1 (d) None of these

280. If $A = \begin{bmatrix} 1 & 2 & -1 \\ 1 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}$, then

- (a) $\rho(A) = 2$ (b) $\rho(A) = 1$ (c) $\rho(A) = 3$ (d) None of these

281. If I_n is the identity matrix of order n , then rank of I_n is

- (a) 1 (b) n (c) 0 (d) None of these

282. The rank of a null matrix is

- (a) 0 (b) 1 (c) Does not exist (d) None of these

283. If A is a non-singular square matrix of order n , then the rank of A is

- (a) Equal to n (b) Less than n (c) Greater than n (d) None of these

284. If A and B are two matrices such that rank of $A = m$ and rank of $B = n$, then

- (a) rank $(AB) = mn$ (b) rank $(AB) \geq \text{rank}(A)$
(c) rank $(AB) \geq \text{rank}(B)$ (d) rank $(AB) \leq \min(\text{rank } A, \text{rank } B)$

285. If A is an invertible matrix and B is a matrix, then

- (a) rank $(AB) = \text{rank}(A)$ (b) rank $(AB) = \text{rank}(B)$ (c) rank $(AB) > \text{rank}(A)$ (d) rank $(AB) > \text{rank}(B)$

Advance Level


398 Matrices

286. If the points (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are collinear, then the rank of the matrix $\begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$ will always be less than
- [Orissa JEE 2003]
- (a) 3 (b) 2 (c) 1 (d) None of these
287. If A is a matrix such that there exists a square submatrix of order r which is non-singular and every square submatrix of order $r+1$ or more is singular, then
- (a) $\text{rank}(A) = r+1$ (b) $\text{rank}(A) = r$ (c) $\text{rank}(A) > r$ (d) $\text{rank}(A) \geq r+1$
288. The rank of the matrix $\begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 6 & 8 & 7 & 5 \end{bmatrix}$ is
- (a) 1 (b) 2 (c) 3 (d) 4
289. The system of equations $AX = B$ of n equations in n unknown has infinitely many solutions if
- (a) $\det A \neq 0$ (b) $\det A \neq 0, (\text{adj } A)B = 0$ (c) $\det A = 0, (\text{adj } A)B \neq 0$ (d) $\det A = 0, (\text{adj } A)B = 0$
290. The trace of skew symmetric matrix of order $n \times n$ is
- (a) 0 (b) 1 (c) n (d) n^2

Miscellaneous Problems

Basic Level

291. If $A = \begin{bmatrix} -2 & 1 \\ 0 & 3 \end{bmatrix}$ and $f(x) = 2x^2 - 3x$, then $f(A)$ equals
- (a) $\begin{bmatrix} 14 & 1 \\ 0 & -9 \end{bmatrix}$ (b) $\begin{bmatrix} -14 & 1 \\ 0 & 9 \end{bmatrix}$ (c) $\begin{bmatrix} 14 & -1 \\ 0 & 9 \end{bmatrix}$ (d) $\begin{bmatrix} -14 & -1 \\ 0 & -9 \end{bmatrix}$
292. The construction of 3×4 matrix A whose element a_{ij} is given by $\frac{(i+j)^2}{2}$ is
- [IIT 1988]
- (a) $\begin{bmatrix} 2 & 9/2 & 8 & 25 \\ 9 & 4 & 5 & 18 \\ 8 & 25 & 18 & 49 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & 9/2 & 25/2 \\ 9/2 & 5/2 & 5 \\ 25 & 18 & 25 \end{bmatrix}$
- (c) $\begin{bmatrix} 2 & 9/2 & 8 & 25/2 \\ 9/2 & 8 & 25/2 & 18 \\ 8 & 25/2 & 18 & 49/2 \end{bmatrix}$ (d) None of these
293. If A is a square matrix of order n such that its elements are polynomial in x and its r -rows become identical for $x = k$, then
- (a) $(x-k)^r$ is a factor of $|A|$ (b) $(x-k)^{r-1}$ is a factor of $|A|$
- (c) $(x-k)^{r+1}$ is a factor of $|A|$ (d) $(x-k)^r$ is a factor of A
294. If $A = [a_{ij}]$ is a scalar matrix of order $n \times n$ such that $a_{ij} = k$ for all i , then trace of A is equal to
- (a) nk (b) $n+k$ (c) n/k (d) None of these



Matrices

Assignment (Basic and Advance Level)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
b	b	c	d	d	d	c	c	a	b	b	b	b	c	c	b	a	b	a	c
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
b	c	d	c	b	d	a	b	c	b	a	d	d	b	c	b	b	b	b	b
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
d	c	a,b	d	a	c	c	c	a	d	b	b	b	b	a	a	c	c	b	b
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
c	b	c	b	a	b	c	c	a	c	c	d	b	c	b	d	a	c	d	d
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
c	d	c	b	b	b	d	d	c	c	c	a	a	a	d	b	b	c	b	d
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
a	a	a	a	c	c	c	b	a	b	a	c	d	c	b	c	b	c	b	b
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
a	d	a	b	c	a	b	b	b	b	d	d	a	b	b	a	d	b	c	a
141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
a	d	b	b	b	c,d	c	b	d	a	d	b	d	c	d	a	b	a	d	b
161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
d	d	a	c	a	b	a	a	b	c	b	c	b	a	a	b	b	b	a	b
181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
a	c	b	d	a	a	b	c	b	c	d	b	b	b	c	a	b	a	c	b
201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220
d	c	a	c	c	c	d	b	a	a	b	b	a	a	a	d	b	a	d	c
221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
b	b	a	a	d	b	b	a	b	a	d	d	c	b	a	c	d	a	b	b
241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260
b	a	d	a	a	b	d	a	c	d	c	d	a	b	a	b	c	b	b	a
261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280
c	b	c	a	a	b	b	b	c	a	a	a	c	d	d	c	d	a	c	c
281	282	283	284	285	286	287	288	289	290	291	292	293	294						
b	c	a	d	b	b	b	c	c	a	c	c	a	a						