### DAY THIRTY SEVEN

# **Unit Test 6**

## (Organic Chemistry-II)

- 1 Identify the correct statement regarding enzymes.
  - (a) Enzymes are specific biological catalysts that can normally function at very high temperature (*T*~1000 K)
  - (b) Enzymes are normally heterogeneous catalysts that are very specific in their actions
  - (c) Enzymes are specific biological catalysts that cannot be poisoned
  - (d) Enzymes are specific biological catalysts that possess well defined active sites
- 2 Which of the following is not optically active?
  - (a) Glycine
- (b) Alanine
- (c) Tyrosine
- (d) Lysine
- 3 Isoelectric point is a
  - (a) specific temperature
  - (b) suitable concentration of amino acid
  - (c) hydrogen ion concentration that does not allow migration of amino acid under electric field
  - (d) melting point of an amino acid under the influence of electric field
- 4 From the following statements, which one is incorrect?
  - (a) Albumin is a simple protein
  - (b) Amino acid lysine contains a basic side chain
  - (c) Insulin is a hormone
  - (d) Muscles contain the protein keratin
- 5 Which of the following statements is not true about glucose? → NCERT Exemplar
  - (a) It is an aldohexose
  - (b) On heating with HI, it forms *n*-hexane
  - (c) It is present in furanose form
  - (d) It does not give 2,4-DNP test
- **6** When glucose reacts with bromine water, the main product is
  - (a) acetic acid
- (b) saccharic acid
- (c) glyceraldehyde
- (d) gluconic acid

7 Match the following and choose the correct option.

	Column I		Column II
Α.	Analgesics	1.	Treatment of stress
В.	Antihistamines	2.	Pain-killing effect
C.	Tranquilizers	3.	Applied to inanimate objects
D.	Disinfectants	4.	Prevents the interaction of histamine with its receptor


#### Codes

	Α	В	С	D		Α	В	С	D
a)	4	3	1	2	(b)	2	4	1	3
$\sim$	1	2	4	3	(d)	3	4	2	1

- 8 Amylopectin is a polymer of
  - (a)  $\alpha$ -D-glucose (b)  $\alpha$ -D-fructose (c) lactose (d) amylose

**Direction** (Q. Nos. 9-10) In the following questions, Assertion followed by Reason is given. Choose the correct answer out of the following choices.

- (a) Both Assertion and Reason are true and Reason is the correct explanation of Assertion
- (b) Both Assertion and Reason are true but Reason is not the correct explanation of Assertion
- (c) Assertion is true but Reason is false
- (d) Both Assertion and Reason are false
- **9** Assertion (A)  $\beta$  -glycosidic linkage is present in maltose.



**Reason** (R) Maltose is composed of two glucose units in which C-1 of one glucose unit is linked to C-2 of another glucose unit.

10 Assertion (A) Polytetrafluoroethene is used in making non-stick cookwares.

Reason (R) Fluorine has highest electronegativity.

#### → NCERT Exemplar

- 11 Food preservatives prevent spoilage of food due to microbial growth. The most commonly used preservatives are
  - (a) table salt, sugar
  - (b) vegetable oils and sodium benzoate.
  - (c) C<sub>6</sub>H<sub>5</sub>COONa
  - (d) All of the above
- 12 Which is not the correct matching of medicine with its disease/activity?
  - (a) Antihistamines - anti-allergic
  - (b) Barbiturates — hypnotic
  - (c) Chloramphenicol typhoid
  - (d) 1% phenol - antiseptic
- 13 Dettol is a mixture of
  - (a) chloroxylenol and terpeneol
  - (b) phenol and chlorophenol
  - (c) phenol and chloroxylenol
  - (d) chlorophenol and chloroxylenol
- 14 Sodium benzoate is used as food preservative. It is
  - (a) metabolised by conversion to hippuric acid and is excreted in the urine
  - (b) metabolised to benzoic acid and deposited in the bond
  - (c) decomposed by gastric juice and escapes as CO<sub>2</sub>
  - (d) decomposed by heat of the digestion process and escapes as CO2
- 15 Aspartame, an artificial sweetener, is

Functional groups, which are not present in aspartame, are

- (a) ester, peptide, amino, carboxyl
- (b) hydroxyl, keto, methoxy
- (c) Both (a) and (b)
- (d) None of the above
- 16 Glyptal polymer is obtained by the reaction of phthalic acid with
  - (a) glycerol
- (b) ethylene glycol
- (c) acetic acid
- (d) malonic acid
- 17 Structurally, the cellulose is a linear polymer of
  - (a) sucrose molecules
- (b) β-D-glucose molecules
- (c)  $\alpha$ -D-glucose molecules
- (d) fructose molecules
- 18 The baby feeding bottles are made up of
  - (a) polyester
- (b) polyurethane
- (c) polystyrene
- (d) polyamide

**19** The monomer of the polymer

$$\begin{array}{cccc} & \text{CH}_3 & \text{CH}_3 \\ & | & | \\ -\text{(CH}_2 - \text{C} - \text{CH}_2 - \text{C} & | \\ & | & | \\ & \text{CH}_3 & \text{CH}_3 \end{array}$$

- (a)  $CH_2 = C(CH_3)_2$
- (b)  $(CH_3)_2C = C(CH_3)_2$
- (c)  $CH_3CH = CHCH_3$
- (d)  $CH_3CH = CH_2$

for

- **20** The monomers used nylon-2-nylon-6 is/are
- the preparation → [NCERT Exemplar]
- (a) caprolactum
- (b) alanine and amino caproic acid
- (c) glycine and amino caproic acid
- (d) hexamethylenediamine and adipic acid
- 21 Which of the following is a monomer of teflon?
  - (a) Difluoro ethane
- (b) Trifluoro ethane
- (c) Tetrafluoro ethene
- (d) None of these
- 22 A salt is heated first with dil. H<sub>2</sub>SO<sub>4</sub> and then with conc. H<sub>2</sub>SO<sub>4</sub>, no reaction takes place. It may be
- (b) sulphide (c) oxalate
- (d) sulphate
- 23 When Cl<sub>2</sub> water is added to a salt solution containing chloroform, chloroform layer turns violet. The salt contains  $(c) NO_2^-$ (d)  $S^{2}$

- 24 Which will give borax bead test with blue bead? (d) Cd<sup>2+</sup> (a) Cr<sup>3+</sup> (b) Co<sup>3+</sup> (c) Ni<sup>2+</sup>
- 25 Which of the following leaves no residue on heating? (a)  $Pb(NO_3)_2$  (b)  $NH_4NO_3$ (c)  $Cu(NO_3)_2$  (d)  $NaNO_3$
- 26 Mark the compound which turns black with NH<sub>4</sub>OH.
  - (a) Lead chloride
- (b) Mercurous chloride
- (c) Mercuric chloride
- (d) Silver chloride
- 27 There is foul smell in presence of moisture with (d) FeSO<sub>4</sub>
- (a) AICI<sub>3</sub> (b)  $Al_2(SO_4)_3$  (c) FeS 28 A colourless salt changes to yellow on heating. Salt is
  - (a) FeO
- (b) PbO
- also soluble in NaOH as well as in dil. HCl. Salt can be (c) ZnO
  - (d) CdO
- 29 A glycoside is the carbohydrate form of a/an
  - (a) ether
- (b) acetal
- - (c) glycone (d) alcohol
- **30** Reduction of hexose A (mol. formula,  $C_6H_{12}O_6$ ) with  $NaBH_4$  gives compounds B and C. Compound B is optically inactive and compound C is optically active. Which of the following is compound A?
  - (a) D-fructose (b) D-glucose (c) D-mannose (d) D-psicose
- **31** Which of the following is true about teflon?
  - (a) It is linear, unbranched polymer of tetrafluoroethylene
  - (b) It has very high thermal stability
  - (c) The polymer molecules are associated by strong dipole-dipole attraction
  - (d) All of the above
- 32 Cellulose has very high degree of hydrophilicity because of
  - (a) its amorphous nature
  - (b) crystalline nature

- (c) presence of excessive voids in solid state
- (d) presence of many hydroxyl groups on the polymer
- 33 Which one among the following is not an analgesic?
  - (a) Ibuprofen
- (b) Naproxen
- (c) Valium
- (d) Aspirin
- 34 Nucleoside involves the combination of
  - (a) sugar + base +  $H_3PO_4$
- (b) sugar + base
- (c) sugar + acid
- (d) sugar + H<sub>2</sub>PO<sub>4</sub>
- 35 On hydrolysis of caprolactum, a compound (B) is obtained. On polymerisation of (B), product (C) is formed. (C) is
  - (a) saran
- (b) nylon-6
- (c) terylene
- (d) bakelite
- **36** The artificial sweetener aspartame (A) is converted to ...... on storage for extended period of time in aqueous solution.

(a) 
$$H_3 \stackrel{\oplus}{N}$$
 —  $CH$  —  $CH_2$  —  $NH$  —  $CH$  —  $C$  —  $OCH_3$  —  $CH_2$  —  $CH_2$  —  $CH_2$  —  $CH_2$  —  $COOH$ 

- (d) No change, remains as A
- **37** A carbohydrate is treatd with  $\alpha$  –naphthol and conc. H<sub>2</sub>SO<sub>4</sub>. What colour will be formed at the junction of two liquids?
  - (a) Blood red
- (b) Violet
- (c) Brown
- (d) Orange
- 38 (A) is a ternary salt with divalent cation. (A) gives yellow ppt. with K2CrO4 as well as with AgNO3. (A) is precipitated by H2S neither in acidic nor ammoniacal medium, but addition of (NH<sub>4</sub>)<sub>2</sub>CO<sub>3</sub> in NH<sub>4</sub>OH gave white ppt. (B). 0.297 g of (A) gave 0.197 g of (B). Identify (A). (c) FeSO<sub>4</sub> (a) BaBr<sub>2</sub> (b) Cu<sub>2</sub>S (d) ZnCl<sub>2</sub>

39 A shallow eutrophic water lake located in a region where the bedrock and sediments contain limestone, has a pH 7.2 and equilibrium constant  $K_a = 4.7 \times 10^{-11}$  for the reaction.

$$H_2O + HCO_3^- \longrightarrow H_3O^+ + CO_3^{2-}$$

Concentrations of HCO<sub>3</sub> and H<sub>3</sub>O<sup>+</sup> are respectively  $1.06 \times 10^{-3}$  mol L<sup>-1</sup> and  $6.3 \times 10^{-8}$  mol L<sup>-1</sup>. If the concentration of  $Ca^{2+}$  is  $1.5 \times 10^{-3} \text{mol}^{-1}$  and  $K_{sp}$  of  $CaCO_3 = 5 \times 10^{-9}$ . Would the precipitation of CaCO<sub>3</sub> take place?

- (a) Precipitation will take place
- (b) Precipitation will not take place
- (c) It may or may not take place
- (d) Cannot be predicted
- **40** To an aqueous solution containing anions from a few drops of acidified KMnO<sub>4</sub> are added. Which one of the following anions, if present will not decolourise the KMnO<sub>4</sub> solution?
  - (a) I<sup>-</sup>

(b)  $CO_3^{2-}$ 

(c) S<sup>2-</sup>

- $(d) NO_2$
- 41 In the brown ring test, the brown colour of the ring is due to
  - (a) ferrous nitrate
- (b) ferric nitrate
- (c) a mixture of NO and NO<sub>2</sub> (d) ferrous nitrososulphate
- 42 What colour is imparted into the flame when lithium is
  - (a) Golden yellow
- (b) Brick red
- (c) Crimson red
- (d) Grassy green
- 43 Give the pOH range for the isoelectric point of the amphoteric ion of an amino acid?
  - (a) 5.5 to 6.3
- (b) 2.5 to 5.0
- (c) 7.7 to 8.5
- (d) 9.0 to 10.7
- 44 Which of the following describes the overall three dimensional folding of a polypeptide?
  - (a) Primary structure
- (b) Secondary structure
- (c) Tertiary structure
- (d) Quaternary structure

**Direction** (Q. Nos. 45-46) Each of these questions contains two statements: Statement I (Assertion) and Statement II (Reason). Each of these questions also has four alternative choices, only one of which is the correct answer. You have to select one of the codes (a), (b), (c) and (d) given below:

- (a) Statement I is true, Statement II is true; Statement II is the correct explanation for Statement I
- (b) Statement I is true, Statement II is true; Statement II is not the correct explanation for Statement I
- (c) Statement I is true; Statement II is false
- (d) Statement I is false; Statement II is true
- **45.** Statement | Fructose because of the presence of keto group does not reduce Tollen's reagent.
  - Statement II Fructose is also called fruit sugar.
- **46.** Statement I MnS is pink in colour which on dissolution in dil. HCl and then heating with NaOH and Br<sub>2</sub> gives pink colour solution changing to brown on heating.
  - Statement II Mn<sup>2+</sup> is oxidised to Mn<sup>4+</sup>.

### **ANSWERS**

<b>1.</b> (d)	<b>2.</b> (a)	<b>3.</b> (c)	<b>4.</b> (d)	<b>5.</b> (c)	<b>6.</b> (d)	<b>7.</b> (b)	<b>8.</b> (a)	<b>9.</b> (d)	<b>10.</b> (c)
<b>11.</b> (d)	<b>12.</b> (d)	<b>13.</b> (a)	<b>14.</b> (a)	<b>15.</b> (b)	<b>16.</b> (b)	<b>17.</b> (b)	<b>18.</b> (c)	<b>19.</b> (a)	<b>20.</b> (c)
<b>21.</b> (c)	<b>22.</b> (d)	<b>23.</b> (b)	<b>24.</b> (b)	<b>25.</b> (b)	<b>26.</b> (b)	<b>27.</b> (c)	<b>28.</b> (c)	<b>29.</b> (b)	<b>30.</b> (d)
<b>31.</b> (d)	<b>32.</b> (d)	<b>33.</b> (c)	<b>34.</b> (b)	<b>35.</b> (b)	<b>36.</b> (c)	<b>37.</b> (b)	<b>38.</b> (a)	<b>39.</b> (b)	<b>40.</b> (b)
<b>41.</b> (d)	<b>42.</b> (c)	<b>43.</b> (c)	<b>44.</b> (c)	<b>45.</b> (d)	<b>46.</b> (a)				

## **Hints and Explanations**

- 1. Normal optimum temperature of enzymes is between 25°C to 40°C, hence (a) is false, enzymes have well defined active sites and hence, their actions are specific in nature.
- 2. Glycine (an amino acid) is optically inactive compound.
- 3. Isoelectric point is a pH at which Zwitter ion does not migrate towards any of the electrodes. Since, amino acids are also Zwitter ions, hence, they do not migrate under the influence of electric field at isoelectric point.
- 4. Muscles contain myosin protein. Keratin is present in hair, wool and silk.
- 5. Glucose is never present in furanose form.
- **6.** When glucose reacts with  $\mathrm{Br}_2$  water, gluconic acid is obtained as main product.

$$\begin{array}{c|c} \text{CHO} & \text{COOH} \\ | & & | \\ \text{(CHOH)}_4 & + \text{[O]} \xrightarrow{\text{Br}_2/\text{H}_2\text{O}} & \text{(CHOH)}_4 \\ | & & | \\ \text{CH}_2\text{OH} & & \text{CH}_2\text{OH} \\ \text{Gluconic acid} & & \text{Gluconic acid} \\ \end{array}$$

7.	Α.	Analgesics	Pain killing effect				
	В.	Antihistamines	Prevents the interaction of histamine with its receptor				
	C.	Tranquilizers	Treatment of stress				
	D.	Disinfectants	Applied to inanimate objects				

- **8.** Amylopectin is a highly branched polymer of  $\alpha$  D glucose units which are joined together through α-glycosidic linkages involving  $C_1$  of one glucose unit with  $C_4$  of the other.
- **9.** Correct Assertion  $\alpha$ -glycosidic linkage is present in maltose. Correct reason maltose is a composed of two  $\alpha$ -D-glucose units in which C-1 of one glucose unit (I) is linked to C-4 of another glucose unit (II).
- 10. Correct Explanation Because of great chemical inertness and high thermal stability, teflon (polytetrafluoroethene) is used in making non-stick cookwares.
- **11.** The chemicals which are used to protect food from microbes action are known as food preservatives. Table salt, vegetable oil, sugar, vinegar, sodium benzoate (C<sub>6</sub>H<sub>5</sub>COO<sup>-</sup>Na<sup>+</sup>) sodium metabisulphite (Na 2S2O5), vitamin E etc. are food preservatives
- 12. 1% phenol is a disinfectant while 0.2% phenol is an antiseptic

- 13. Dettol is a mixture of chloroxylenol and terpeneol.
- 14. Sodium benzoate is used as food preservative. It is metabolised to hippuric acid and excreted in urine. It is used in soft drinks and acidic foods.
- **15.** Hydroxyl, keto, methoxy groups are absent in aspartame while ester, peptide amino an carboxyl groups are present.
- **16.** Glyptal is obtained by the reaction of phthalic acid with ethylene

Structure of glyptals:

- 17. Cellulose is a linear condensation polymer of  $\beta$ -D- glucose in which C<sub>1</sub> of one glucose unit is connected to C<sub>4</sub> of the other through  $\beta$  -D-glycosidic linkage.
- 18. Polystyrene is used for the production of baby feeding bottles.
- **19.**  $CH_2 = C(CH_3)_2$  is the repeating unit of given polymer.

**20.** 
$$nH_2NCH_2COOH+ nH_2N(CH_2)_5COOH \longrightarrow$$
Glycine Amino caproic acid
$$- \left\{ NH - CH_2 - CONH - (CH_2)_5 - C - \right\}$$
Nylon-2-nylon-6

21. Teflon is an addition polymer of tetrafluoroethylene.

$$\begin{array}{ccc} n \mathbf{F_2} \mathbf{C} = \mathbf{C} \mathbf{F_2} & \xrightarrow{\quad \mathbf{Heat} \quad } & \mathbf{\left[ \mathbf{F_2} \mathbf{C} - \mathbf{C} \mathbf{F_2} \right]_n} \\ \mathbf{Tetrafluoroethylene} & & \mathbf{Teflon} \end{array}$$

- 22.  $SO_4^{2-}$  does not react with either dilute or conc.  $H_2SO_4$ .
- **23.**  $2I^- + CI_2 \longrightarrow I_2 + 2CI^$ l<sub>2</sub> gives violet colour to the chloroform layer.
- 24. Co<sup>3+</sup> will give blue bead in borax bead test.
- **25.**  $NH_4NO_3 \xrightarrow{\Delta} N_2O \uparrow + 2H_2O \uparrow$
- 26. Mercurous chloride (Hg<sub>2</sub>Cl<sub>2</sub>) compound turns black with NH<sub>4</sub>OH  $\begin{aligned} & \text{Hg}_2\text{Cl}_2 + 2\text{NH}_4\text{OH} \longrightarrow \text{Hg} \downarrow + \text{Hg} \cdot \text{NH}_2\text{Cl} + 2\text{H}_2\text{O} + \text{NH}_4\text{Cl} \\ & \text{Black} \end{aligned}$

27. 
$$FeS + 2H_2O \longrightarrow Fe(OH)_2 + H_2S$$
Foul sme

28. ZnO is colourless and changes to yellow on heating.

$$\label{eq:ZnO+2HCl} \begin{split} &\text{ZnO} + 2\text{HCl} \longrightarrow \text{ZnCl}_2 + \text{H}_2\text{O} \\ &\text{ZnO} + 2\text{NaOH} \longrightarrow \text{Na}_2\text{ZnO}_2 + \text{H}_2\text{O} \end{split}$$

29. Acetal form of carbohydrates are known as glycoside.

30. 
$$H \xrightarrow{CH_2OH} CH_2OH CH_2OH CH_2OH$$
 $C=O$ 
 $H \xrightarrow{OH} OH$ 
 $H \xrightarrow{OH} OH$ 

- 31. Due to symmetry of F<sub>2</sub>C=CF<sub>2</sub> (teflon) and strength of C—F bond, branching doesn't takes place and a linear polymer is formed. High bond energy of C—F bond gives high thermal stability to polymer. Large electronegativity difference of C and F makes C—F bonds highly polar.
- **32.** Cellulose fibres are hydrophilic due to the presence of OH groups at their surfaces.
- **33.** Valium is white all other gives option are the examples of analgesics.
- **34.** A combination of sugar and base is called nucleoside and a combination of sugar, base and phosphate is called nucleotide.

35. 

Caprolactum 
$$(CH_2)_5 - COO^{-1}$$
 $(B)$ 
 $(B)$ 

- **37.** This is Molisch's test for carbohydrates. In this experiment, violet ring is formed at the junction of two liquids.
- **38.** (A) gives yellow ppt. with  $K_2CrO_4$  as well as with  $AgNO_3$ . Therefore, (A) can have  $Pb^{2+}$  or  $Ba^{2+}$  (both divalent cations) and  $Br^-$ . (A) is not precipitated by  $H_2S$  in acidic and ammoniacal medium. Hence, (A) does not have  $Pb^{2+}$  (precipitated as PbS in acidic medium by  $H_2S$  gas).
  - (A) contains Ba<sup>2+</sup> (apple green flame).

$$\begin{split} \therefore \text{(A) is BaBr}_2. \\ & \text{BaBr}_2 + 2 \text{AgNO}_3 \longrightarrow 2 \text{AgBr} \downarrow + \text{Ba(NO}_3)_2 \\ & \text{BaBr}_2 + \text{K}_2 \text{CrO}_4 \longrightarrow \text{BaCrO}_4 \downarrow + 2 \text{KBr} \\ & \text{BaBr}_2 + (\text{NH}_4)_2 \text{CO}_3 \longrightarrow \text{BaCO}_3 + 2 \text{NH}_4 \text{Br} \end{split}$$

39. Equilibrium constant for reaction:

$$K_{a} = \frac{[H_{3}O^{+}] [CO_{3}^{2-}]}{[HCO_{3}^{-}] [H_{2}O]}$$
$$[CO_{3}^{2-}] = \frac{4.7 \times 10^{-11} \times 1.06 \times 10^{-3}}{6.3 \times 10^{-8}}$$
$$= 7.9 \times 10^{-7} \text{mol L}^{-1}$$

The reaction quotient for the reaction

Ca<sup>2+</sup> + CO<sub>3</sub><sup>2-</sup> 
$$\longrightarrow$$
 CaCO<sub>3</sub>  
Q = [Ca<sup>2+</sup>] [CO<sub>3</sub><sup>2-</sup>]  
= 1.5 × 10<sup>-3</sup> × 7.9 × 10<sup>-7</sup>  
= 1.2 × 10<sup>-9</sup> < 5 × 10<sup>-9</sup> =  $K_{ep}$ 

Hence, no precipitation will take place.

- **40.** In CO<sub>3</sub><sup>2-</sup>, C is present in its highest oxidation state, i.e +4 state, so its further oxidation is not possible, it only undergoes reduction. Acidified KMnO<sub>4</sub> is a strong oxidising agent but it cannot oxidise CO<sub>3</sub><sup>2-</sup>.
- 41. On adding freshly prepared FeSO<sub>4</sub> solution and then conc. H<sub>2</sub>SO<sub>4</sub> to water extract of salt mixture carefully by the sides of test-tube, a dark brown ring of ferrous nitrososulphate, FeSO<sub>4</sub>NO is formed.

42.	Colour	Cation
	Golden yellow	Na <sup>+</sup>
	Brick red	Ca <sup>2+</sup>
	Crimson red	Li <sup>+</sup>
	Grassy green	Cu <sup>2+</sup> ,BO <sub>3</sub> <sup>3-</sup> ,TI <sup>3+</sup>

**43.** The pH at which a particular amino acid does not migrate under the influence of an electric field is called isoelectric point of that amino acid.

The pH range for the isoelectric point is from 5.5 to 6.3 on the pOH range for the isoelectric point is from 7.7 to 8.5.

- **44.** The tertiary structure of proteins refer to three dimensional folding of polymer chain.
- **45.** Fructose although have keto group but it reduces Tollen's reagent to give silver mirror. It is also called fruit sugar.

46. MnS + dil.HCl
$$\longrightarrow$$
 MnCl<sub>2</sub>

MnCl<sub>2</sub> + 2NaOH $\longrightarrow$  Mn(OH)<sub>2</sub> + 2NaCl

Mn(OH)<sub>2</sub> + [O]  $\xrightarrow{\text{NaOH} + \text{Br}_2}$  MnO<sub>2</sub> + H<sub>2</sub>O