

10

Definite Integrals

10.1 FUNDAMENTAL THEOREM OF INTEGRAL CALCULUS

Let f be a continuous function on the closed interval $[a, b]$ and ϕ be an anti-derivative of f , then

$$\int_a^b f(x) dx = \phi(b) - \phi(a). \quad (\text{We assume it without proof})$$

In words, the above theorem tells us that

$$\int_a^b f(x) dx = (\text{value of an anti-derivative at } b, \text{ the upper limit}) \\ - (\text{value of the same anti-derivative at } a, \text{ the lower limit}).$$

Remarks

1. We often write $\phi(b) - \phi(a)$ as $[\phi(x)]_a^b$.
2. No matter which anti-derivative we take as ϕ , the value of the definite integral comes out to be the same.

10.1.1 Evaluation of Definite Integrals

The fundamental theorem enables us to evaluate the definite integrals by making use of *anti-derivatives*.

ILLUSTRATIVE EXAMPLES

Example 1. Evaluate the following :

$$(i) \int_0^1 (2x^3 + 3)^2 dx \quad (ii) \int_3^5 \frac{dt}{1+3t}.$$

Solution. (i) $\int_0^1 (2x^3 + 3)^2 dx = \int_0^1 (4x^6 + 12x^3 + 9) dx$

$$= \left[4 \cdot \frac{x^7}{7} + 12 \cdot \frac{x^4}{4} + 9x \right]_0^1 = \left[\frac{4}{7}x^7 + 3x^4 + 9x \right]_0^1$$
$$= \left(\frac{4}{7} + 3 + 9 \right) - (0 + 0 + 0) = \frac{88}{7}.$$

$$(ii) \int_3^5 \frac{dt}{1+3t} = \left[\frac{\log |1+3t|}{3} \right]_3^5 = \frac{1}{3} (\log 16 - \log 10)$$
$$= \frac{1}{3} \log \frac{16}{10} = \frac{1}{3} \log \frac{8}{5}.$$

Example 2. Evaluate the following :

$$(i) \int_0^{\pi/2} \cos^2 x \, dx \qquad (ii) \int_0^{\pi/2} \cos^4 x \, dx.$$

Solution. (i) $\int_0^{\pi/2} \cos^2 x \, dx = \int_0^{\pi/2} \frac{1 + \cos 2x}{2} \, dx = \frac{1}{2} \left[x + \frac{\sin 2x}{2} \right]_0^{\pi/2}$

$$= \frac{1}{2} \left[\left(\frac{\pi}{2} + \frac{1}{2} \sin \pi \right) - \left(0 + \frac{1}{2} \sin 0 \right) \right]$$

$$= \frac{1}{2} \left[\left(\frac{\pi}{2} + \frac{1}{2} \cdot 0 \right) - \frac{1}{2} \cdot 0 \right] = \frac{\pi}{4}.$$

(ii) $\int_0^{\pi/2} \cos^4 x \, dx = \int_0^{\pi/2} (\cos^2 x)^2 \, dx = \int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right)^2 \, dx$

$$= \frac{1}{4} \int_0^{\pi/2} (1 + 2 \cos 2x + \cos^2 2x) \, dx = \frac{1}{4} \int_0^{\pi/2} \left(1 + 2 \cos 2x + \frac{1 + \cos 4x}{2} \right) \, dx$$

$$= \frac{1}{8} \int_0^{\pi/2} (3 + 4 \cos 2x + \cos 4x) \, dx = \frac{1}{8} \left[3x + 4 \cdot \frac{\sin 2x}{2} + \frac{\sin 4x}{4} \right]_0^{\pi/2}$$

$$= \frac{1}{8} \left[3 \left(\frac{\pi}{2} - 0 \right) + 2(\sin \pi - \sin 0) + \frac{1}{4}(\sin 2\pi - \sin 0) \right]$$

$$= \frac{1}{8} \left[\frac{3\pi}{2} + 2(0 - 0) + \frac{1}{4}(0 - 0) \right] = \frac{3\pi}{16}.$$

Example 3. Evaluate the following :

$$(i) \int_0^{\pi/2} \sqrt{1 + \cos 2x} \, dx \qquad (ii) \int_0^{\pi/2} \sqrt{1 + \sin 2x} \, dx.$$

Solution. (i) $\int_0^{\pi/2} \sqrt{1 + \cos 2x} \, dx = \int_0^{\pi/2} \sqrt{2 \cos^2 x} \, dx = \sqrt{2} \int_0^{\pi/2} |\cos x| \, dx$

$$= \sqrt{2} \int_0^{\pi/2} \cos x \, dx$$

$$(\text{As } 0 \leq x \leq \frac{\pi}{2} \Rightarrow \cos x \geq 0 \Rightarrow |\cos x| = \cos x)$$

$$= \sqrt{2} \left[\sin x \right]_0^{\pi/2} = \sqrt{2} \left[\sin \frac{\pi}{2} - \sin 0 \right]$$

$$= \sqrt{2} (1 - 0) = \sqrt{2}.$$

(ii) $\int_0^{\pi/2} \sqrt{1 + \sin 2x} \, dx = \int_0^{\pi/2} \sqrt{1 + \cos \left(\frac{\pi}{2} - 2x \right)} \, dx = \int_0^{\pi/2} \sqrt{2 \cos^2 \left(\frac{\pi}{4} - x \right)} \, dx$

$$= \sqrt{2} \int_0^{\pi/2} \left| \cos \left(\frac{\pi}{4} - x \right) \right| \, dx = \sqrt{2} \int_0^{\pi/2} \cos \left(\frac{\pi}{4} - x \right) \, dx$$

$$\left[\text{As } 0 \leq x \leq \frac{\pi}{2} \Rightarrow 0 \geq -x \geq -\frac{\pi}{2} \Rightarrow \frac{\pi}{4} \geq \frac{\pi}{4} - x \geq -\frac{\pi}{4} \right]$$

$$\Rightarrow -\frac{\pi}{4} \leq \frac{\pi}{4} - x \leq \frac{\pi}{4} \Rightarrow \cos \left(\frac{\pi}{4} - x \right) > 0 \Rightarrow \left| \cos \left(\frac{\pi}{4} - x \right) \right| = \cos \left(\frac{\pi}{4} - x \right)$$

$$\begin{aligned}
 &= \sqrt{2} \left[\frac{\sin\left(\frac{\pi}{4} - x\right)}{-1} \right]_0^{\pi/2} = -\sqrt{2} \left[\sin\left(-\frac{\pi}{4}\right) - \sin\frac{\pi}{4} \right] \\
 &= -\sqrt{2} \left[-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right] = \sqrt{2} \cdot \frac{2}{\sqrt{2}} = 2.
 \end{aligned}$$

Example 4. Evaluate the following integrals :

$$(i) \int_0^{\pi/4} \sqrt{1 - \sin 2x} \, dx \quad (ii) \int_{\pi}^{3\pi/2} \sqrt{1 - \cos 2x} \, dx.$$

Solution. (i) $\int_0^{\pi/4} \sqrt{1 - \sin 2x} \, dx = \int_0^{\pi/4} \sqrt{1 - \cos\left(\frac{\pi}{2} - 2x\right)} \, dx = \int_0^{\pi/4} \sqrt{2 \sin^2\left(\frac{\pi}{4} - x\right)} \, dx$

$$= \sqrt{2} \int_0^{\pi/4} \left| \sin\left(\frac{\pi}{4} - x\right) \right| \, dx = \sqrt{2} \int_0^{\pi/4} \sin\left(\frac{\pi}{4} - x\right) \, dx$$

$$\begin{aligned}
 &\left[\text{As } 0 \leq x \leq \frac{\pi}{4} \Rightarrow 0 \geq -x \geq -\frac{\pi}{4} \Rightarrow \frac{\pi}{4} \geq \frac{\pi}{4} - x \geq 0 \right. \\
 &\Rightarrow 0 \leq \frac{\pi}{4} - x \leq \frac{\pi}{4} \Rightarrow \sin\left(\frac{\pi}{4} - x\right) \geq 0 \Rightarrow \left| \sin\left(\frac{\pi}{4} - x\right) \right| = \sin\left(\frac{\pi}{4} - x\right) \left. \right] \\
 &= \sqrt{2} \left[\frac{-\cos\left(\frac{\pi}{4} - x\right)}{-1} \right]_0^{\pi/4} = \sqrt{2} \left[\cos 0 - \cos \frac{\pi}{4} \right] \\
 &= \sqrt{2} \left(1 - \frac{1}{\sqrt{2}} \right) = \sqrt{2} - 1.
 \end{aligned}$$

$$\begin{aligned}
 (ii) \int_{\pi}^{3\pi/2} \sqrt{1 - \cos 2x} \, dx &= \int_{\pi}^{3\pi/2} \sqrt{2 \sin^2 x} \, dx = \sqrt{2} \int_{\pi}^{3\pi/2} |\sin x| \, dx \\
 &= \sqrt{2} \int_{\pi}^{3\pi/2} (-\sin x) \, dx \\
 &\quad \left[\text{As } \pi \leq x \leq \frac{3\pi}{2} \Rightarrow \sin x \leq 0 \Rightarrow |\sin x| = -\sin x \right] \\
 &= -\sqrt{2} \left[-\cos x \right]_{\pi}^{3\pi/2} = \sqrt{2} \left(\cos \frac{3\pi}{2} - \cos \pi \right) \\
 &= \sqrt{2} (0 - (-1)) = \sqrt{2}.
 \end{aligned}$$

Example 5. Evaluate the following integrals :

$$(i) \int_{\pi/4}^{\pi/2} \sqrt{1 - \sin 2x} \, dx \quad (ii) \int_0^{2\pi} \sqrt{1 + \sin \frac{x}{2}} \, dx.$$

Solution. (i) $\int_{\pi/4}^{\pi/2} \sqrt{1 - \sin 2x} \, dx = \int_{\pi/4}^{\pi/2} \sqrt{1 - \cos\left(\frac{\pi}{2} - 2x\right)} \, dx = \int_{\pi/4}^{\pi/2} \sqrt{2 \sin^2\left(\frac{\pi}{4} - x\right)} \, dx$

$$= \sqrt{2} \int_{\pi/4}^{\pi/2} \left| \sin\left(\frac{\pi}{4} - x\right) \right| \, dx = \sqrt{2} \int_{\pi/4}^{\pi/2} \left(-\sin\left(\frac{\pi}{4} - x\right) \right) \, dx$$

$$\begin{aligned}
 &\left[\text{As } \frac{\pi}{4} \leq x \leq \frac{\pi}{2} \Rightarrow -\frac{\pi}{4} \geq -x \geq -\frac{\pi}{2} \Rightarrow 0 \geq \frac{\pi}{4} - x \geq -\frac{\pi}{4} \right. \\
 &\Rightarrow -\frac{\pi}{4} \leq \frac{\pi}{4} - x \leq 0 \Rightarrow \sin\left(\frac{\pi}{4} - x\right) \leq 0 \Rightarrow \left| \sin\left(\frac{\pi}{4} - x\right) \right| = -\sin\left(\frac{\pi}{4} - x\right) \left. \right]
 \end{aligned}$$

$$= -\sqrt{2} \left[\frac{-\cos\left(\frac{\pi}{4} - x\right)}{-1} \right]_{\pi/4}^{\pi/2} = -\sqrt{2} \left(\cos\left(-\frac{\pi}{4}\right) - \cos 0 \right)$$

$$= -\sqrt{2} \left(\frac{1}{\sqrt{2}} - 1 \right) = \sqrt{2} - 1.$$

$$(ii) \quad \int_0^{2\pi} \sqrt{1 + \sin \frac{x}{2}} dx = \int_0^{2\pi} \sqrt{1 + \cos\left(\frac{\pi}{2} - \frac{x}{2}\right)} dx = \int_0^{2\pi} \sqrt{2 \cos^2\left(\frac{\pi}{4} - \frac{x}{4}\right)} dx$$

$$= \sqrt{2} \int_0^{2\pi} \left| \cos\left(\frac{\pi}{4} - \frac{x}{4}\right) \right| dx = \sqrt{2} \int_0^{2\pi} \cos\left(\frac{\pi}{4} - \frac{x}{4}\right) dx$$

$$\left[\text{As } 0 \leq x \leq 2\pi \Rightarrow 0 \leq \frac{x}{4} \leq \frac{\pi}{2} \Rightarrow 0 \geq -\frac{x}{4} \geq -\frac{\pi}{2} \Rightarrow \frac{\pi}{4} \geq \frac{\pi}{4} - \frac{x}{4} \geq -\frac{\pi}{4} \right.$$

$$\Rightarrow -\frac{\pi}{4} \leq \frac{\pi}{4} - \frac{x}{4} \leq \frac{\pi}{4} \Rightarrow \cos\left(\frac{\pi}{4} - \frac{x}{4}\right) > 0 \Rightarrow \left| \cos\left(\frac{\pi}{4} - \frac{x}{4}\right) \right| = \cos\left(\frac{\pi}{4} - \frac{x}{4}\right) \Big]$$

$$= \sqrt{2} \left[\frac{\sin\left(\frac{\pi}{4} - \frac{x}{4}\right)}{-\frac{1}{4}} \right]_0^{2\pi} = -4\sqrt{2} \left(\sin\left(-\frac{\pi}{4}\right) - \sin \frac{\pi}{4} \right)$$

$$= -4\sqrt{2} \left(-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right) = 4\sqrt{2} \cdot \frac{2}{\sqrt{2}} = 8.$$

Example 6. Evaluate the following integrals :

$$(i) \quad \int_0^{\pi/4} \sec x \sqrt{\frac{1 - \sin x}{1 + \sin x}} dx$$

$$(ii) \quad \int_0^{\pi/4} (\tan x + \cot x)^{-2} dx.$$

Solution. (i) $\int_0^{\pi/4} \sec x \sqrt{\frac{1 - \sin x}{1 + \sin x}} dx = \int_0^{\pi/4} \sec x \sqrt{\frac{1 - \sin x}{1 + \sin x} \times \frac{1 - \sin x}{1 - \sin x}} dx$

$$= \int_0^{\pi/4} \sec x \sqrt{\frac{(1 - \sin x)^2}{\cos^2 x}} dx = \int_0^{\pi/4} \sec x \left| \frac{1 - \sin x}{\cos x} \right| dx$$

$$\left[\text{As } 0 \leq x \leq \frac{\pi}{4} \Rightarrow \cos x > 0, 1 - \sin x > 0 \Rightarrow \frac{1 - \sin x}{\cos x} > 0 \right]$$

$$= \int_0^{\pi/4} \sec x \cdot \frac{1 - \sin x}{\cos x} dx = \int_0^{\pi/4} \sec x (\sec x - \tan x) dx$$

$$= \int_0^{\pi/4} (\sec^2 x - \sec x \tan x) dx = \left[\tan x - \sec x \right]_0^{\pi/4}$$

$$= \left(\tan \frac{\pi}{4} - \sec \frac{\pi}{4} \right) - (\tan 0 - \sec 0)$$

$$= (1 - \sqrt{2}) - (0 - 1) = 2 - \sqrt{2}.$$

$$(ii) \quad \int_0^{\pi/4} (\tan x + \cot x)^{-2} dx = \int_0^{\pi/4} \left(\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} \right)^{-2} dx = \int_0^{\pi/4} \left(\frac{\sin^2 x + \cos^2 x}{\sin x \cos x} \right)^{-2} dx$$

$$= \int_0^{\pi/4} \left(\frac{1}{\sin x \cos x} \right)^{-2} dx = \int_0^{\pi/4} (\sin x \cos x)^2 dx$$

$$\begin{aligned}
&= \frac{1}{4} \int_0^{\pi/4} (2 \sin x \cos x)^2 dx = \frac{1}{4} \int_0^{\pi/4} \sin^2 2x dx \\
&= \frac{1}{4} \int_0^{\pi/4} \frac{1 - \cos 4x}{2} dx = \frac{1}{8} \left[x - \frac{\sin 4x}{4} \right]_0^{\pi/4} \\
&= \frac{1}{8} \left[\left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) \right] = \frac{1}{8} \left[\left(\frac{\pi}{4} - 0 \right) - 0 \right] = \frac{\pi}{32}.
\end{aligned}$$

Example 7. Evaluate the following integrals :

$$(i) \int_0^1 \frac{(\sin^{-1} x)^2}{\sqrt{1-x^2}} dx \quad (ii) \int_0^1 \frac{x+1}{(x^2+2x+3)^2} dx.$$

Solution. (i) $\int_0^1 \frac{(\sin^{-1} x)^2}{\sqrt{1-x^2}} dx = \int_0^1 (\sin^{-1} x)^2 \cdot \frac{1}{\sqrt{1-x^2}} dx$ | $\int (f(x))^n f'(x) dx$ form

$$\begin{aligned}
&= \left[\frac{(\sin^{-1} x)^3}{3} \right]_0^1 = \frac{1}{3} [(\sin^{-1} 1)^3 - (\sin^{-1} 0)^3] \\
&= \frac{1}{3} \left[\left(\frac{\pi}{2} \right)^3 - 0^3 \right] = \frac{\pi^3}{24}.
\end{aligned}$$

$$\begin{aligned}
(ii) \int_0^1 \frac{x+1}{(x^2+2x+3)^2} dx &= \frac{1}{2} \int_0^1 (x^2+2x+3)^{-2} (2x+2) dx \\
&= \frac{1}{2} \cdot \left[\frac{(x^2+2x+3)^{-1}}{-1} \right]_0^1 = -\frac{1}{2} \left[\frac{1}{x^2+2x+3} \right]_0^1 \\
&= -\frac{1}{2} \left[\frac{1}{1+2+3} - \frac{1}{0+0+3} \right] = -\frac{1}{2} \left(\frac{1}{6} - \frac{1}{3} \right) = -\frac{1}{2} \left(-\frac{1}{6} \right) = \frac{1}{12}.
\end{aligned}$$

Example 8. Evaluate the following integrals :

$$(i) \int_a^b \frac{\log x}{x} dx \quad (\text{I.S.C. 2000}) \quad (ii) \int_0^{\pi/4} \frac{2 \cos 2x}{1 + \sin 2x} dx. \quad (\text{I.S.C. 2002})$$

Solution. (i) $\int_a^b \frac{\log x}{x} dx = \int_a^b (\log x)^1 \cdot \frac{1}{x} dx$ | $\int (f(x))^n f'(x) dx$ form

$$\begin{aligned}
&= \left[\frac{(\log x)^2}{2} \right]_a^b = \frac{1}{2} [(\log b)^2 - (\log a)^2] \\
&= \frac{1}{2} (\log b + \log a) (\log b - \log a) \\
&= \frac{1}{2} \log ab \log \left(\frac{b}{a} \right).
\end{aligned}$$

(ii) As $\frac{d}{dx} (1 + \sin 2x) = 0 + \cos 2x \cdot 2 = 2 \cos 2x$,

$$\begin{aligned}
\therefore \int_0^{\pi/4} \frac{2 \cos 2x}{1 + \sin 2x} dx &= \left[\log |1 + \sin 2x| \right]_0^{\pi/4} \quad \left| \int \frac{f'(x)}{f(x)} dx \text{ form} \right. \\
&= \log |1 + \sin \frac{\pi}{2}| - \log |1 + \sin 0| \\
&= \log (1 + 1) - \log (1 + 0) = \log 2 - \log 1 \\
&= \log 2 - 0 = \log 2.
\end{aligned}$$

Example 9. Evaluate the following :

$$(i) \int_0^1 \frac{x^9}{5+x^{10}} dx \quad (ii) \int_0^{\pi/2} x \cos 2x dx.$$

Solution. (i) $\int_0^1 \frac{x^9}{5+x^{10}} dx = \frac{1}{10} \int_0^1 \frac{10 \cdot x^9}{5+x^{10}} dx \quad \left| \int \frac{f'(x)}{f(x)} dx \text{ form} \right.$

$$= \frac{1}{10} [\log |5+x^{10}|]_0^1 = \frac{1}{10} [\log 6 - \log 5] = \frac{1}{10} \log \frac{6}{5}.$$

(ii) $\int_0^{\pi/2} x \cos 2x dx = \left[x \cdot \frac{\sin 2x}{2} \right]_0^{\pi/2} - \int_0^{\pi/2} 1 \cdot \frac{\sin 2x}{2} dx \quad (\text{using integration by parts})$

$$= \frac{1}{2} \left(\frac{\pi}{2} \sin \pi - 0 \right) - \frac{1}{2} \int_0^{\pi/2} \sin 2x dx = \frac{1}{2} \left(\frac{\pi}{2} \cdot 0 \right) - \frac{1}{2} \left[-\frac{\cos 2x}{2} \right]_0^{\pi/2}$$

$$= \frac{1}{4} (\cos \pi - \cos 0) = \frac{1}{4} (-1 - 1) = -\frac{1}{2}.$$

Example 10. Evaluate the following :

$$(i) \int_0^{\pi/2} x \sin^2 x dx \quad (ii) \int_0^1 x^2 e^x dx.$$

Solution. (i) $\int_0^{\pi/2} x \sin^2 x dx = \int_0^{\pi/2} x \cdot \frac{1 - \cos 2x}{2} dx = \frac{1}{2} \int_0^{\pi/2} x dx - \frac{1}{2} \int_0^{\pi/2} x \cos 2x dx$

$$= \frac{1}{2} \left[\frac{x^2}{2} \right]_0^{\pi/2} - \frac{1}{2} \left(\left[x \cdot \frac{\sin 2x}{2} \right]_0^{\pi/2} - \int_0^{\pi/2} 1 \cdot \frac{\sin 2x}{2} dx \right)$$

$$= \frac{1}{4} \left(\frac{\pi^2}{4} - 0 \right) - \frac{1}{4} \left(\frac{\pi}{2} \sin \pi - 0 \right) + \frac{1}{4} \int_0^{\pi/2} \sin 2x dx$$

$$= \frac{\pi^2}{16} - \frac{1}{4} \left(\frac{\pi}{2} \cdot 0 \right) + \frac{1}{4} \left[-\frac{\cos 2x}{2} \right]_0^{\pi/2} = \frac{\pi^2}{16} - \frac{1}{8} [\cos 2x]_0^{\pi/2}$$

$$= \frac{\pi^2}{16} - \frac{1}{8} (\cos \pi - \cos 0) = \frac{\pi^2}{16} - \frac{1}{8} (-1 - 1) = \frac{\pi^2}{16} + \frac{1}{4}.$$

(ii) $\int_0^1 x^2 e^x dx = [x^2 \cdot e^x]_0^1 - \int_0^1 2x e^x dx$

$$= (1 \cdot e^1 - 0) - 2 \int_0^1 x e^x dx = e - 2 \left([x e^x]_0^1 - \int_0^1 1 \cdot e^x dx \right)$$

$$= e - 2 (1 \cdot e^1 - 0) + 2 \int_0^1 e^x dx = e - 2e + 2 [e^x]_0^1$$

$$= -e + 2(e^1 - e^0) = -e + 2(e - 1) = e - 2.$$

Example 11. Evaluate the following integrals :

$$(i) \int_{\pi/4}^{\pi/2} \cos 2x \log \sin x dx \quad (ii) \int_0^1 x \tan^{-1} x dx. \quad (\text{I.S.C. 2002})$$

Solution. (i) $\int_{\pi/4}^{\pi/2} \cos 2x \log \sin x dx = \int_{\pi/4}^{\pi/2} \log \sin x \cdot \cos 2x dx \quad (\text{integrate by parts})$

$$= \left[\log \sin x \cdot \frac{\sin 2x}{2} \right]_{\pi/4}^{\pi/2} - \int_{\pi/4}^{\pi/2} \frac{1}{\sin x} \cdot \cos x \cdot \frac{\sin 2x}{2} dx$$

$$\begin{aligned}
&= \frac{1}{2} \left[0 - \log \frac{1}{\sqrt{2}} \right] - \int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin x} \cdot \frac{2 \sin x \cos x}{2} dx \\
&= -\frac{1}{2} (\log 1 - \frac{1}{2} \log 2) - \int_{\pi/4}^{\pi/2} \cos^2 x dx \\
&= \frac{1}{4} \log 2 - \int_{\pi/4}^{\pi/2} \frac{1 + \cos 2x}{2} dx \\
&= \frac{1}{4} \log 2 - \frac{1}{2} \left[x + \frac{\sin 2x}{2} \right]_{\pi/4}^{\pi/2} \\
&= \frac{1}{4} \log 2 - \frac{1}{2} \left[\left(\frac{\pi}{2} - \frac{\pi}{4} \right) + \frac{1}{2} (0 - 1) \right] = \frac{1}{4} \log 2 - \frac{\pi}{8} + \frac{1}{4}.
\end{aligned}$$

$$(ii) \int_0^1 x \tan^{-1} x dx = \int_0^1 \tan^{-1} x \cdot x dx \quad (\text{integrate by parts})$$

$$\begin{aligned}
&= \left[\tan^{-1} x \cdot \frac{x^2}{2} \right]_0^1 - \int_0^1 \frac{1}{1+x^2} \cdot \frac{x^2}{2} dx \\
&= \frac{1}{2} [\tan^{-1} 1 - 0] - \frac{1}{2} \int_0^1 \frac{x^2}{1+x^2} dx \\
&= \frac{1}{2} \cdot \frac{\pi}{4} - \frac{1}{2} \int_0^1 \left(1 - \frac{1}{1+x^2} \right) dx \\
&= \frac{\pi}{8} - \frac{1}{2} [x - \tan^{-1} x]_0^1 = \frac{\pi}{8} - \frac{1}{2} [(1 - \tan^{-1} 1) - (0 - \tan^{-1} 0)] \\
&= \frac{\pi}{8} - \frac{1}{2} \left[\left(1 - \frac{\pi}{4} \right) - 0 \right] = \frac{\pi}{4} - \frac{1}{2}.
\end{aligned}$$

Example 12. Evaluate the following :

$$(i) \int_0^{\pi/2} \frac{\cos x}{1 + \cos x + \sin x} dx \quad (ii) \int_1^5 \frac{\log x}{(x+1)^2} dx.$$

Solution. (i) $\int_0^{\pi/2} \frac{\cos x}{1 + \cos x + \sin x} dx = \int_0^{\pi/2} \frac{\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}}{2 \cos^2 \frac{x}{2} + 2 \sin \frac{x}{2} \cos \frac{x}{2}} dx$

$$\begin{aligned}
&= \int_0^{\pi/2} \frac{\cos \frac{x}{2} - \sin \frac{x}{2}}{2 \cos \frac{x}{2}} dx = \frac{1}{2} \int_0^{\pi/2} \left(1 - \tan \frac{x}{2} \right) dx \\
&= \frac{1}{2} \left[x + \frac{\log \left| \cos \frac{x}{2} \right|}{\frac{1}{2}} \right]_0^{\pi/2} \\
&= \frac{1}{2} \left[\left(\frac{\pi}{2} - 0 \right) + 2 \left(\log \frac{1}{\sqrt{2}} - \log 1 \right) \right] \\
&= \frac{\pi}{4} + \log 1 - \log \sqrt{2} = \frac{\pi}{4} - \frac{1}{2} \log 2.
\end{aligned}$$

$$\begin{aligned}
 (ii) \int_1^5 \frac{\log x}{(x+1)^2} dx &= \int_1^5 \log x \cdot (x+1)^{-2} dx && \text{(integrate by parts)} \\
 &= \left[\log x \cdot \frac{(x+1)^{-1}}{-1} \right]_1^5 - \int_1^5 \frac{1}{x} \cdot \frac{(x+1)^{-1}}{-1} dx \\
 &= - \left[\frac{\log x}{x+1} \right]_1^5 + \int_1^5 \frac{dx}{x(x+1)} \\
 &= - \left(\frac{\log 5}{6} - \frac{\log 1}{2} \right) + \int_1^5 \left(\frac{1}{x} - \frac{1}{x+1} \right) dx && \text{(by partial fractions)} \\
 &= - \left(\frac{\log 5}{6} - 0 \right) + [\log |x| - \log |x+1|]_1^5 \\
 &= - \frac{1}{6} \log 5 + (\log 5 - \log 6) - (\log 1 - \log 2) \\
 &= \frac{5}{6} \log 5 - (\log 6 - \log 2) = \frac{5}{6} \log 5 - \log 3.
 \end{aligned}$$

Example 13. Evaluate the following :

$$(i) \int_0^{\sqrt{2}} \sqrt{2-x^2} dx \qquad (ii) \int_1^2 \frac{2}{4x^2-1} dx.$$

$$\begin{aligned}
 \text{Solution. (i)} \int_0^{\sqrt{2}} \sqrt{2-x^2} dx &= \int_0^{\sqrt{2}} \sqrt{(\sqrt{2})^2 - x^2} dx \\
 &= \left[\frac{x\sqrt{2-x^2}}{2} + \frac{(\sqrt{2})^2}{2} \sin^{-1} \frac{x}{\sqrt{2}} \right]_0^{\sqrt{2}} \\
 &= \left(\frac{\sqrt{2} \cdot 0}{2} + \sin^{-1} 1 \right) - \left(\frac{0 \cdot \sqrt{2}}{2} + \sin^{-1} 0 \right) \\
 &= \left(0 + \frac{\pi}{2} \right) - (0 + 0) = \frac{\pi}{2}.
 \end{aligned}$$

$$\begin{aligned}
 (ii) \int_1^2 \frac{2}{4x^2-1} dx &= 2 \cdot \frac{1}{4} \int_1^2 \frac{dx}{x^2 - \left(\frac{1}{2}\right)^2} = \frac{1}{2} \cdot \frac{1}{2 \cdot \frac{1}{2}} \left[\log \left| \frac{x - \frac{1}{2}}{x + \frac{1}{2}} \right| \right]_1^2 \\
 &= \frac{1}{2} \left(\log \frac{3}{5} - \log \frac{1}{3} \right) = \frac{1}{2} \log \left(\frac{3}{5} \times \frac{3}{1} \right) = \frac{1}{2} \log \frac{9}{5}.
 \end{aligned}$$

Example 14. Evaluate the following integrals :

$$(i) \int_0^1 \frac{x e^x}{(x+1)^2} dx \quad (\text{I.S.C. 2011}) \qquad (ii) \int_1^5 \frac{\log x}{(x+1)^2} dx \qquad (iii) \int_0^1 \sin^{-1} \left(\frac{2x}{1+x^2} \right) dx.$$

$$\begin{aligned}
 \text{Solution. (i)} \int_0^1 \frac{x e^x}{(x+1)^2} dx &= \int_0^1 \frac{(x+1)-1}{(x+1)^2} e^x dx \\
 &= \int_0^1 \frac{1}{x+1} \cdot e^x dx - \int_0^1 \frac{1}{(x+1)^2} e^x dx
 \end{aligned}$$

(evaluate the first integral by parts, taking $\frac{1}{x+1}$ as the first function)

$$\begin{aligned}
&= \left[\frac{1}{x+1} \cdot e^x \right]_0^1 - \int_0^1 (-1)(x+1)^{-2} \cdot e^x dx - \int_0^1 \frac{1}{(x+1)^2} e^x dx \\
&= \left(\frac{1}{2} \cdot e^1 - 1 \cdot e^0 \right) + \int_0^1 \frac{1}{(x+1)^2} e^x dx - \int_0^1 \frac{1}{(x+1)^2} e^x dx = \frac{1}{2} e - 1.
\end{aligned}$$

$$(ii) \quad \int_1^5 \frac{\log x}{(x+1)^2} dx = \int_1^5 \log x \cdot (x+1)^{-2} dx \quad (\text{by parts})$$

$$\begin{aligned}
&= \left[\log x \cdot \frac{(x+1)^{-1}}{-1} \right]_1^5 - \int_1^5 \frac{1}{x} \cdot \frac{(x+1)^{-1}}{-1} dx \\
&= - \left[\frac{\log x}{x+1} \right]_1^5 + \int_1^5 \frac{1}{x(x+1)} dx \\
&= - \left[\frac{\log 5}{6} - \frac{\log 1}{2} \right] + \int_1^5 \left(\frac{1}{x} - \frac{1}{x+1} \right) dx \\
&= - \left(\frac{\log 5}{6} - 0 \right) + \left[\log |x| - \log |x+1| \right]_1^5 \\
&= - \frac{\log 5}{6} + (\log 5 - \log 6) - (\log 1 - \log 2) \\
&= \frac{5}{6} \log 5 - (\log 6 - \log 2) = \frac{5}{6} \log 5 - \log 3.
\end{aligned}$$

$$\begin{aligned}
(iii) \quad \int_0^1 \sin^{-1} \left(\frac{2x}{1+x^2} \right) dx &= \int_0^1 2 \tan^{-1} x dx = 2 \int_0^1 \tan^{-1} x \cdot 1 dx \quad (\text{integrate by parts}) \\
&= 2 \left[\left[\tan^{-1} x \cdot x \right]_0^1 - \int_0^1 \frac{1}{1+x^2} \cdot x dx \right] \\
&= 2 (\tan^{-1} 1 - 0) - \int_0^1 \frac{2x}{1+x^2} dx \\
&= 2 \left(\frac{\pi}{4} - 0 \right) - \left[\log(1+x^2) \right]_0^1 \\
&= \frac{\pi}{2} - (\log 2 - \log 1) = \frac{\pi}{2} - (\log 2 - 0) \\
&= \frac{\pi}{2} - \log 2.
\end{aligned}$$

Example 15. Evaluate the following :

$$(i) \quad \int_{1/4}^{1/2} \frac{dx}{\sqrt{x-x^2}} \quad (ii) \quad \int_2^3 \frac{x^3+1}{x(x-1)} dx.$$

$$\begin{aligned}
\text{Solution. } (i) \quad \int_{1/4}^{1/2} \frac{dx}{\sqrt{x-x^2}} &= \int_{1/4}^{1/2} \frac{dx}{\sqrt{\frac{1}{4} - \left(x^2 - x + \frac{1}{4} \right)}} = \int_{1/4}^{1/2} \frac{dx}{\sqrt{\left(\frac{1}{2} \right)^2 - \left(x - \frac{1}{2} \right)^2}} \\
&= \left[\sin^{-1} \left(\frac{x - \frac{1}{2}}{\frac{1}{2}} \right) \right]_{1/4}^{1/2} = \left[\sin^{-1} (2x - 1) \right]_{1/4}^{1/2} \\
&= \sin^{-1} 0 - \sin^{-1} \left(-\frac{1}{2} \right) = 0 - \left(-\frac{\pi}{6} \right) = \frac{\pi}{6}.
\end{aligned}$$

$$(ii) \int_2^3 \frac{x^3+1}{x(x-1)} dx = \int_2^3 \left(x+1 + \frac{x+1}{x(x-1)} \right) dx \quad (\text{by division})$$

$$\text{Let } \frac{x+1}{x(x-1)} = \frac{A}{x} + \frac{B}{x-1} \Rightarrow x+1 = A(x-1) + Bx.$$

On putting $x = 0$ and $x = 1$, we get

$$1 = -A \text{ and } 2 = B \Rightarrow A = -1 \text{ and } B = 2.$$

$$\begin{aligned} \therefore \int_2^3 \frac{x^3+1}{x(x-1)} dx &= \int_2^3 \left(x+1 - \frac{1}{x} + \frac{2}{x-1} \right) dx \\ &= \left[\frac{x^2}{2} + x - \log|x| + 2 \log|x-1| \right]_2^3 \\ &= \left(\frac{9}{2} + 3 - \log 3 + 2 \log 2 \right) - (2 + 2 - \log 2 - 2 \log 1) \\ &= \frac{15}{2} - \log 3 + 2 \log 2 - 4 + \log 2 - 2.0 = \frac{7}{2} + 3 \log 2 - \log 3. \end{aligned}$$

Example 16. Prove that : $\int_0^{\pi/2} \frac{3 \sin \theta + 4 \cos \theta}{\sin \theta + \cos \theta} d\theta = \frac{7\pi}{4}.$ (I.S.C. 2005)

Solution. Let $3 \sin \theta + 4 \cos \theta = l(\sin \theta + \cos \theta) + m(\cos \theta - \sin \theta).$

Equating coefficients of $\sin \theta$ and $\cos \theta$ on both sides, we get

$$3 = l - m \text{ and } 4 = l + m.$$

Solving these for l and m , we get $l = \frac{7}{2}$ and $m = \frac{1}{2}.$

$$\begin{aligned} \therefore \int_0^{\pi/2} \frac{3 \sin \theta + 4 \cos \theta}{\sin \theta + \cos \theta} d\theta &= \int_0^{\pi/2} \frac{\frac{7}{2}(\sin \theta + \cos \theta) + \frac{1}{2}(\cos \theta - \sin \theta)}{\sin \theta + \cos \theta} d\theta \\ &= \int_0^{\pi/2} \left(\frac{7}{2} + \frac{1}{2} \cdot \frac{\cos \theta - \sin \theta}{\sin \theta + \cos \theta} \right) d\theta \\ &= \frac{7}{2} [\theta]_0^{\pi/2} + \frac{1}{2} [\log|\sin \theta + \cos \theta|]_0^{\pi/2} \\ &= \frac{7}{2} \left[\frac{\pi}{2} - 0 \right] + \frac{1}{2} [\log 1 - \log 1] = \frac{7\pi}{4}. \end{aligned}$$

EXERCISE 10.1

Evaluate the following (1 to 21) definite integrals :

1. (i) $\int_0^8 \left(\sqrt{8x} - \frac{x^2}{8} \right) dx$

(ii) $\int_0^1 \frac{1}{2x-3} dx.$

2. (i) $\int_{-4}^{-1} \frac{1}{x} dx$

(ii) $\int_{\pi/6}^{\pi/4} \operatorname{cosec} x dx.$

3. (i) $\int_0^1 \sqrt{5x+4} dx$

(ii) $\int_0^1 \frac{dx}{\sqrt{1+x} + \sqrt{x}}.$

4. (i) $\int_0^{\pi} \frac{dx}{1 + \sin x}$

(ii) $\int_0^1 \frac{dx}{\sqrt{1-x^2}}.$

$$5. (i) \int_0^1 \frac{1-x}{1+x} dx$$

$$(ii) \int_0^{\pi/2} \frac{\cos x}{5+4\sin x} dx.$$

$$6. (i) \int_1^3 (x^2 + e^x) (x^3 + 3e^x + 4) dx$$

$$(ii) \int_1^2 \frac{3x}{9x^2-1} dx.$$

$$7. (i) \int_0^{\pi/4} \tan^3 x \sec^2 x dx$$

$$(ii) \int_0^{\pi/2} \sin^4 x dx.$$

$$8. (i) \int_0^{\pi/2} \sqrt{1-\cos 2x} dx$$

$$(ii) \int_0^{\pi/4} \sqrt{1+\sin 2x} dx.$$

$$9. (i) \int_0^{\pi/2} \frac{\sin^2 x}{(1+\cos x)^2} dx$$

$$(ii) \int_0^{\pi/2} \frac{\sin \theta}{\sqrt{1+\cos \theta}} d\theta.$$

$$10. (i) \int_0^{\pi/4} \sin 2x \sin 3x dx$$

$$(ii) \int_0^{\pi/2} (a^2 \cos^2 x + b^2 \sin^2 x) dx.$$

$$11. (i) \int_0^{\pi/4} 2 \tan^3 x dx$$

$$(ii) \int_0^{\pi/4} \frac{\tan^3 x}{1+\cos 2x} dx.$$

$$12. (i) \int_0^{\pi/4} \frac{\sin x}{\cos 3x + 3 \cos x} dx$$

$$(ii) \int_{\pi/3}^{\pi/4} (\tan x + \cot x)^2 dx$$

$$(iii) \int_0^{\pi/4} (\tan x + \cot x)^{-1} dx.$$

(I.S.C. 2003)

$$13. (i) \int_0^4 \frac{dx}{\sqrt{x^2+2x+3}}$$

$$(ii) \int_0^a \frac{dx}{\sqrt{ax-x^2}}.$$

$$14. (i) \int_0^1 \frac{\tan^{-1} x}{1+x^2} dx$$

$$(ii) \int_0^1 \frac{x^5}{1+x^6} dx.$$

$$15. (i) \int_0^1 x e^x dx$$

$$(ii) \int_1^2 \frac{x+3}{x(x+2)} dx.$$

$$16. (i) \int_0^{\pi/2} x^2 \cos 2x dx$$

$$(ii) \int_{-a}^a \sqrt{\frac{a-x}{a+x}} dx.$$

$$17. (i) \int_1^3 \frac{dx}{x^2(x+1)}$$

$$(ii) \int_1^2 \frac{dx}{(x+1)(x^2-7x+12)}.$$

$$18. (i) \int_0^2 \frac{1}{\sqrt{3+2x-x^2}} dx$$

$$(ii) \int_0^2 \frac{dx}{4+x-x^2}.$$

$$19. (i) \int_0^1 \sin^{-1} x dx$$

$$(ii) \int_0^1 \tan^{-1} x dx.$$

$$20. (i) \int_1^2 e^x \left(\frac{1}{x} - \frac{1}{x^2} \right) dx$$

$$(ii) \int_0^{\pi/2} \frac{x + \sin x}{1 + \cos x} dx.$$

$$21. (i) \int_{\pi/4}^{\pi/2} e^x (\log(\sin x) + \cot x) dx \quad (ii) \int_{\pi/2}^{\pi} e^x \left(\frac{1 - \sin x}{1 - \cos x} \right) dx.$$

$$22. \text{ If } \int_0^a 3x^2 dx = 8, \text{ find the value of } a.$$

10.1.2 Evaluation of Definite Integrals by Substitution

ILLUSTRATIVE EXAMPLES

Example 1. Evaluate the following :

$$(i) \int_0^1 x^2 e^{x^3} dx \quad (ii) \int_0^1 \frac{x}{1+x^4} dx.$$

Solution. (i) Put $x^3 = t \Rightarrow 3x^2 dx = dt \Rightarrow x^2 dx = \frac{1}{3} dt$.

When $x = 0$, $t = 0$ and when $x = 1$, $t = 1^3 = 1$.

$$\begin{aligned} \therefore \int_0^1 x^2 e^{x^3} dx &= \int_0^1 e^t \frac{1}{3} dt = \frac{1}{3} [e^t]_0^1 \\ &= \frac{1}{3} [e^1 - e^0] = \frac{1}{3} (e - 1). \end{aligned}$$

(ii) Put $x^2 = t \Rightarrow 2x dx = dt \Rightarrow x dx = \frac{1}{2} dt$.

When $x = 0$, $t = 0$ and when $x = 1$, $t = 1$.

$$\begin{aligned} \therefore \int_0^1 \frac{x}{1+x^4} dx &= \int_0^1 \frac{1}{1+t^2} \cdot \frac{1}{2} dt = \frac{1}{2} \cdot \frac{1}{1} \left[\tan^{-1} t \right]_0^1 \\ &= \frac{1}{2} [\tan^{-1} 1 - \tan^{-1} 0] = \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] = \frac{\pi}{8}. \end{aligned}$$

Example 2. Evaluate the following :

$$(i) \int_0^1 \frac{e^x}{1+e^{2x}} dx \quad (ii) \int_0^{\pi/2} \frac{\sin x}{1+\cos^2 x} dx \quad (iii) \int_0^{\pi/4} \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx.$$

Solution. (i) Put $e^x = t \Rightarrow e^x dx = dt$.

When $x = 0$, $t = e^0 = 1$ and when $x = 1$, $t = e^1 = e$.

$$\begin{aligned} \therefore \int_0^1 \frac{e^x}{1+e^{2x}} dx &= \int_1^e \frac{dt}{1+t^2} = \frac{1}{1} \left[\tan^{-1} t \right]_1^e \\ &= \tan^{-1} e - \tan^{-1} 1 = \tan^{-1} e - \frac{\pi}{4}. \end{aligned}$$

(ii) Put $\cos x = t \Rightarrow -\sin x dx = dt \Rightarrow \sin x dx = -dt$.

When $x = 0$, $t = \cos 0 = 1$ and when $x = \frac{\pi}{2}$, $t = \cos \frac{\pi}{2} = 0$.

$$\begin{aligned} \therefore \int_0^{\pi/2} \frac{\sin x}{1+\cos^2 x} dx &= \int_1^0 \frac{-dt}{1+t^2} = -\frac{1}{1} \left[\tan^{-1} t \right]_1^0 \\ &= -[\tan^{-1} 0 - \tan^{-1} 1] = -\left(0 - \frac{\pi}{4}\right) = \frac{\pi}{4}. \end{aligned}$$

15. (i) 1 (ii) $\frac{1}{2} \log 6$. 16. (i) $-\frac{\pi}{4}$ (ii) $a\pi$.
 17. (i) $\frac{2}{3} + \log \frac{2}{3}$ (ii) $\frac{2}{5} \log 2 - \frac{3}{20} \log 3$.
 18. (i) $\frac{\pi}{3}$ (ii) $\frac{1}{\sqrt{17}} \log \left(\frac{21 + 5\sqrt{17}}{4} \right)$.
 19. (i) $\frac{\pi}{2} - 1$ (ii) $\frac{\pi}{4} - \frac{1}{2} \log 2$. 20. (i) $\frac{1}{2} e^2 - e$ (ii) $\frac{\pi}{2}$.
 21. (i) $\frac{1}{2} e^{\pi/4} \log 2$ (ii) $e^{\pi/2}$. 22. 2.

EXERCISE 10.2

1. (i) $\frac{1}{2} (e - 1)$ (ii) $\frac{1}{2} \tan^{-1} e^2 - \frac{\pi}{2}$. 2. (i) $2(\sqrt{2} - 1)$ (ii) $\frac{8}{21}$.
 3. (i) $\tan^{-1} \left(\frac{1}{3} \right)$ (ii) $\tan^{-1} \left(\frac{1}{3} \right)$. 4. (i) $\sin (\log 3)$ (ii) $\frac{\log 2}{1 + \log 2}$.
 5. (i) $\frac{16}{15} (2 + \sqrt{2})$ (ii) $\frac{132}{7} \sqrt[3]{4}$. 6. (i) $1 - \log 2$ (ii) $\frac{\pi}{2} - 1$.
 7. (i) $\log \frac{4}{3}$ (ii) $\log \frac{4}{3}$. 8. (i) $\frac{\pi}{8}$ (ii) $\log \frac{2e - 1}{e}$.
 9. (i) $\frac{3}{16} \pi a^4$ (ii) $\frac{\pi}{2ab}$. 10. $\frac{\pi}{2} - \log 2$ (ii) $\frac{\pi}{2} - \log 2$.
 11. (i) $\frac{\pi^2}{16} - \frac{\pi}{4} + \frac{1}{2} \log 2$ (ii) $e - \frac{2}{\log 2}$.
 12. (i) $\frac{2}{3} \tan^{-1} \frac{1}{3}$ (ii) $\frac{1}{\sqrt{5}} \log \left(\frac{\sqrt{5} + 1}{\sqrt{5} - 1} \right)$.
 13. (i) $\frac{\pi}{3}$ (ii) $\frac{\pi}{\sqrt{35}}$. 14. (i) $\frac{1}{2} \log 3$ (ii) $\frac{1}{24}$.
 15. (i) $\frac{\pi}{4}$ (ii) $\frac{\pi}{2}$.

EXERCISE 10.3

1. (i) $\frac{13}{2}$ (ii) 34 (iii) $\frac{25}{2}$.
 2. (i) $2 - \sqrt{2}$ (ii) 2 (iii) 1.
 3. (i) 4 (ii) $2(e - 1)$ (iii) $2 - \frac{2}{e}$.
 4. (i) 0 (ii) 4. 5. 37. 6. 47.
 7. (i) 0 (ii) 0 (iii) 0.
 8. (i) $\frac{\pi}{2}$ (ii) $\frac{3\pi}{8}$ (iii) 0 (iv) 0.
 9. (i) 0 (ii) 0 (iii) 0 (iv) 0.
 10. (i) $\frac{1}{42}$ (ii) $\frac{4}{63}$ (iii) $\frac{144}{35} \sqrt{3}$ (iv) $\frac{16}{315} a^{9/2}$.
 11. (i) $\frac{\pi}{4}$ (ii) $\frac{\pi}{4}$ (iii) $\frac{\pi}{4}$ (iv) $\frac{\pi}{4}$.
 12. (i) $\frac{\pi}{4}$ (ii) $\frac{\pi}{4}$ (iii) $\frac{\pi}{4}$ (iv) $\frac{\pi}{4}$.
 13. (i) $\frac{\pi^2}{4}$ (ii) $\frac{2\pi}{3}$. 14. (i) $\frac{\pi}{5}$ (ii) $\frac{\pi}{2\sqrt{2}} \log (\sqrt{2} + 1)$.