

Chapter - Continuity & Differentiability



Topic-1: Continuity



1 MCQs with One Correct Answer

- The function $f(x) = [x]^2 - [x^2]$ (where $[y]$ is the greatest integer less than or equal to y), is discontinuous at [1999 - 2 Marks]
 - all integers
 - all integers except 0 and 1
 - all integers except 0
 - all integers except 1
- The function $f(x) = [x] \cos\left(\frac{2x-1}{2}\right)\pi$, $[.]$ denotes the greatest integer function, is discontinuous at [1995S]
 - All x
 - All integer points
 - No x
 - x which is not an integer



4 Fill in the Blanks

- Let $f(x) = [x] \sin\left(\frac{\pi}{[x+1]}\right)$, where $[.]$ denotes the greatest integer function. The domain of f is... and the points of discontinuity of f in the domain are..... [1996 - 2 Marks]
- Let $f(x) = \begin{cases} \frac{(x^3+x^2-16x+20)}{(x-2)^2}, & \text{if } x \neq 2 \\ k, & \text{if } x = 2 \end{cases}$
If $f(x)$ is continuous for all x , then $k = \dots\dots\dots$ [1981 - 2 Marks]



6 MCQs with One or More Than One Correct

- Let $[x]$ be the greatest integer less than or equals to x . Then, at which of the following point(s) the function $f(x) = x \cos(\pi(x + [x]))$ is discontinuous? [Adv. 2017]

- $x = -1$
- $x = 0$
- $x = 1$
- $x = 2$

- For every pair of continuous functions $f, g : [0, 1] \rightarrow \mathbb{R}$ such that $\max \{f(x) : x \in [0, 1]\} = \max \{g(x) : x \in [0, 1]\}$, the correct statement(s) is (are): [Adv. 2014]
 - $(f(c))^2 + 3f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0, 1]$
 - $(f(c))^2 + f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0, 1]$
 - $(f(c))^2 + 3f(c) = (g(c))^2 + g(c)$ for some $c \in [0, 1]$
 - $(f(c))^2 = (g(c))^2$ for some $c \in [0, 1]$

- For every integer n , let a_n and b_n be real numbers. Let function $f : \mathbb{R} \rightarrow \mathbb{R}$ be given by [2012]

$$f(x) = \begin{cases} a_n + \sin \pi x, & \text{for } x \in [2n, 2n+1] \\ b_n + \cos \pi x, & \text{for } x \in (2n-1, 2n) \end{cases}$$

for all integers n . If f is continuous, then which of the following hold(s) for all n ?

- $a_{n-1} - b_{n-1} = 0$
- $a_n - b_n = 1$
- $a_n - b_{n+1} = 1$
- $a_{n-1} - b_n = -1$

- The following functions are continuous on $(0, \pi)$. [1991 - 2 Marks]

- $\tan x$
- $\int_0^x t \sin \frac{1}{t} dt$
- $\begin{cases} 1, & 0 < x \leq \frac{3\pi}{4} \\ 2 \sin \frac{2}{9}x, & \frac{3\pi}{4} < x < \pi \end{cases}$
- $\begin{cases} x \sin x, & 0 < x \leq \frac{\pi}{2} \\ \frac{\pi}{2} \sin(\pi + x), & \frac{\pi}{2} < x < \pi \end{cases}$

9. If $f(x) = \frac{1}{2}x - 1$, then on the interval $[0, \pi]$ [1989 - 2 Marks]

- (a) $\tan [f(x)]$ and $1/f(x)$ are both continuous
 (b) $\tan [f(x)]$ and $1/f(x)$ are both discontinuous
 (c) $\tan [f(x)]$ and $f^{-1}(x)$ are both continuous
 (d) $\tan [f(x)]$ is continuous but $1/f(x)$ is not.



10 Subjective Problems

10. Let $f(x) = \begin{cases} \{1 + |\sin x|\}^{a/|\sin x|} & ; \quad -\frac{\pi}{6} < x < 0 \\ b & ; \quad x = 0 \\ e^{\tan 2x / \tan 3x} & ; \quad 0 < x < \frac{\pi}{6} \end{cases}$

[1994 - 4 Marks]

Determine a and b such that $f(x)$ is continuous at $x = 0$

11. Let $f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2}, & x < 0 \\ a, & x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x}} - 4}, & x > 0 \end{cases}$ [1990 - 4 Marks]

Determine the value of a , if possible, so that the function is continuous at $x = 0$

12. Find the values of a and b so that the function

$$f(x) = \begin{cases} x + a\sqrt{2} \sin x, & 0 \leq x < \pi/4 \\ 2x \cot x + b, & \pi/4 \leq x \leq \pi/2 \\ a \cos 2x - b \sin x, & \pi/2 < x \leq \pi \end{cases}$$

is continuous for $0 \leq x \leq \pi$.

[1989 - 2 Marks]

13. Let $f(x)$ be a continuous and $g(x)$ be a discontinuous function. prove that $f(x) + g(x)$ is a discontinuous function. [1987 - 2 Marks]

14. Let $f(x) = \begin{cases} 1 + x, & 0 \leq x < 2 \\ 3 - x, & 2 \leq x \leq 3 \end{cases}$ [1983 - 2 Marks]

Determine the form of $g(x) = f(f(x))$ and hence find the points of discontinuity of g , if any

15. Let $f(x + y) = f(x) + f(y)$ for all x and y . If the function $f(x)$ is continuous at $x = 0$, then show that $f(x)$ is continuous at all x . [1981 - 2 Marks]



Topic-2: Differentiability



1 MCQs with One Correct Answer

1. Let $f(x)$ be a continuously differentiable function on the interval $(0, \infty)$ such that $f(1) = 2$ and

$$\lim_{t \rightarrow x} \frac{t^{10} f(x) - x^{10} f(t)}{t^9 - x^9} = 1 \text{ for each } x > 0. \text{ Then, for all } x >$$

$0, f(x)$ is equal to

[Adv. 2024]

- (a) $\frac{31}{11x} - \frac{9}{11}x^{10}$ (b) $\frac{9}{11x} + \frac{13}{11}x^{10}$
 (c) $\frac{-9}{11x} + \frac{31}{11}x^{10}$ (d) $\frac{13}{11x} + \frac{9}{11}x^{10}$

2. Let $f(x) = \begin{cases} x^2 \cos \frac{\pi}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, $x \in \mathbb{R}$ then f is [2012]

- (a) differentiable both at $x = 0$ and at $x = 2$
 (b) differentiable at $x = 0$ but not differentiable at $x = 2$
 (c) not differentiable at $x = 0$ but differentiable at $x = 2$
 (d) differentiable neither at $x = 0$ nor at $x = 2$

3. Let $g(x) = \frac{(x-1)^n}{\log \cos^m(x-1)}$; $0 < x < 2$, m and n are integers,

$m \neq 0$, $n > 0$, and let p be the left hand derivative of $|x-1|$

at $x = 1$. If $\lim_{x \rightarrow 1^+} g(x) = p$, then [2008]

- (a) $n = 1, m = 1$ (b) $n = 1, m = -1$
 (c) $n = 2, m = 2$ (d) $n > 2, m = n$

4. Let $f(x)$ be differentiable on the interval $(0, \infty)$ such that

$$f(1) = 1, \text{ and } \lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1 \text{ for each } x > 0. \text{ Then}$$

$f(x)$ is

[2007 - 3 marks]

- (a) $\frac{1}{3x} + \frac{2x^2}{3}$ (b) $\frac{-1}{3x} + \frac{4x^2}{3}$
 (c) $\frac{-1}{x} + \frac{2}{x^2}$ (d) $\frac{1}{x}$

5. If $f(x)$ is continuous and differentiable function and $f(1/n) = 0 \forall n \geq 1$ and $n \in \mathbb{I}$, then [2005S]

(a) $f(x) = 0, x \in (0, 1]$
 (b) $f(0) = 0, f'(0) = 0$
 (c) $f(0) = 0 = f'(0), x \in (0, 1]$
 (d) $f(0) = 0$ and $f'(0)$ need not to be zero

6. The function given by $y = [x - 1]$ is differentiable for all real numbers except the points [2005S]

(a) $\{0, 1, -1\}$ (b) ± 1 (c) 1 (d) -1

7. The domain of the derivative of the function

$$f(x) = \begin{cases} \tan^{-1} x & \text{if } |x| \leq 1 \\ \frac{1}{2}(|x| - 1) & \text{if } |x| > 1 \end{cases} \text{ is [2002S]}$$

(a) $R - \{0\}$ (b) $R - \{1\}$
 (c) $R - \{-1\}$ (d) $R - \{-1, 1\}$

8. Which of the following functions is differentiable at $x = 0$?

(a) $\cos(|x|) + |x|$ (b) $\cos(|x|) - |x|$ [2001S]
 (c) $\sin(|x|) + |x|$ (d) $\sin(|x|) - |x|$

9. Let $f: R \rightarrow R$ be a function defined by $f(x) = \max\{x, x^3\}$. The set of all points where $f(x)$ is NOT differentiable is [2001S]

(a) $\{-1, 1\}$ (b) $\{-1, 0\}$ (c) $\{0, 1\}$ (d) $\{-1, 0, 1\}$

10. The left-hand derivative of $f(x) = [x] \sin(\pi x)$ at $x = k$, k an integer, is [2001S]

(a) $(-1)^k(k-1)\pi$ (b) $(-1)^{k-1}(k-1)\pi$
 (c) $(-1)^k k\pi$ (d) $(-1)^{k-1} k\pi$

11. The function $f(x) = (x^2 - 1)|x^2 - 3x + 2| + \cos(|x|)$ is NOT differentiable at [1999 - 2 Marks]

(a) -1 (b) 0 (c) 1 (d) 2

12. Let $[.]$ denote the greatest integer function and $f(x) = [\tan^2 x]$, then: [1993 - 1 Mark]

(a) $\lim_{x \rightarrow 0} f(x)$ does not exist
 (b) $f(x)$ is continuous at $x = 0$
 (c) $f(x)$ is not differentiable at $x = 0$
 (d) $f'(0) = 1$

13. Let $f: R \rightarrow R$ be a differentiable function and $f(1) = 4$.

Then the value of $\lim_{x \rightarrow 1} \int_4^{f(x)} \frac{2t}{x-1} dt$ is [1990 - 2 Marks]

(a) $8f'(1)$ (b) $4f'(1)$ (c) $2f'(1)$ (d) $f'(1)$

14. If $f(a) = 2, f'(a) = 1, g(a) = -1, g'(a) = 2$, then the

value of $\lim_{x \rightarrow a} \frac{g(x)f(a) - g(a)f(x)}{x - a}$ is [1983 - 1 Mark]

(a) -5 (b) $\frac{1}{5}$
 (c) 5 (d) none of these

15. For a real number y , let $[y]$ denotes the greatest integer less

than or equal to y : Then the function $f(x) = \frac{\tan(\pi[x - \pi])}{1 + [x]^2}$

is [1981 - 2 Marks]

(a) discontinuous at some x
 (b) continuous at all x , but the derivative $f'(x)$ does not exist for some x
 (c) $f'(x)$ exists for all x , but the second derivative $f''(x)$ does not exist for some x
 (d) $f'(x)$ exists for all x



Integer Value Answer Non-Negative Integer

16. Let the functions $f: (-1, 1) \rightarrow R$ and $g: (-1, 1) \rightarrow (-1, 1)$ be defined by

$$f(x) = |2x - 1| + |2x + 1| \text{ and } g(x) = x - [x],$$

where $[x]$ denotes the greatest integer less than or equal to x . Let $f \circ g: (-1, 1) \rightarrow R$ be the composite function

defined by $(f \circ g)(x) = f(g(x))$. Suppose c is the number of points in the interval $(-1, 1)$ at which $f \circ g$ is NOT

continuous, and suppose d is the number of points in the interval $(-1, 1)$ at which $f \circ g$ is NOT differentiable. Then

the value of $c + d$ is [Adv. 2020]

17. Let $f: R \rightarrow R$ and $g: R \rightarrow R$ be respectively given by

$$f(x) = |x| + 1 \text{ and } g(x) = x^2 + 1. \text{ Define } h: R \rightarrow R \text{ by}$$

$$h(x) = \begin{cases} \max\{f(x), g(x)\} & \text{if } x \leq 0, \\ \min\{f(x), g(x)\} & \text{if } x > 0. \end{cases}$$

The number of points at which $h(x)$ is not differentiable is [Adv. 2014]

18. Let $f: [1, \infty) \rightarrow [2, \infty)$ be a differentiable function such

that $f(1) = 2$. If $6 \int_1^x f(t) dt = 3xf(x) - x^3$ for all $x \geq 1$, then

the value of $f(2)$ is [2011]



4 Fill in the Blanks

19. Let $f(x) = x|x|$. The set of points where $f(x)$ is twice differentiable is [1992 - 2 Marks]

$$20. \text{ Let } f(x) = \begin{cases} (x-1)^2 \sin \frac{1}{(x-1)} |x| & \text{if } x \neq 1 \\ -1, & \text{if } x = 1 \end{cases}$$

be a real-valued function. Then the set of points where $f(x)$ is not differentiable is [1981 - 2 Marks]



6 MCQs with One or More Than One Correct

21. Let $S = (0, 1) \cup (1, 2) \cup (3, 4)$ and $T = \{0, 1, 2, 3\}$. Then which of the following statements is(are) true? [Adv. 2023]
- There are infinitely many functions from S to T
 - There are infinitely many strictly increasing functions from S to T
 - The number of continuous functions from S to T is at most 120
 - Every continuous function from S to T is differentiable
22. Let $f: (0, 1) \rightarrow \mathbb{R}$ be the function defined as $f(x) = [4x] \left(x - \frac{1}{4} \right)^2 \left(x - \frac{1}{2} \right)$, where $[x]$ denotes the greatest integer less than or equal to x . Then which of the following statements is(are) true? [Adv. 2023]
- The function f is discontinuous exactly at one point in $(0, 1)$
 - There is exactly one point in $(0, 1)$ at which the function f is continuous but NOT differentiable
 - The function f is NOT differentiable at more than three points in $(0, 1)$
 - The minimum value of the function f is $-\frac{1}{512}$
23. Let the function $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = x^3 - x^2 + (x-1)\sin x$ and let $g: \mathbb{R} \rightarrow \mathbb{R}$ be an arbitrary function. Let $fg: \mathbb{R} \rightarrow \mathbb{R}$ be the product function defined by $(fg)(x) = f(x)g(x)$. Then which of the following statements is/are TRUE? [Adv. 2020]
- If g is continuous at $x = 1$, then fg is differentiable at $x = 1$
 - If fg is differentiable at $x = 1$, then g is continuous at $x = 1$
 - If g is differentiable at $x = 1$, then fg is differentiable at $x = 1$
 - If fg is differentiable at $x = 1$, then g is differentiable at $x = 1$
24. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ given by [Adv. 2019]
- $$f(x) = \begin{cases} x^5 + 5x^4 + 10x^3 + 10x^2 + 3x + 1, & x < 0; \\ x^2 - x + 1, & 0 \leq x < 1; \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3; \\ (x-2)\log_e(x-2) - x + \frac{10}{3}, & x \geq 3 \end{cases}$$
- Then which of the following options is/are correct?
- f' has a local maximum at $x = 1$
 - f is increasing on $(-\infty, 0)$
 - f' is NOT differentiable at $x = 1$
 - f is onto
25. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ be two non-constant differentiable functions. If $f'(x) = (e^{f(x)-g(x)})g'(x)$ for all $x \in \mathbb{R}$, and $f(1) = g(2) = 1$, then which of the following statement (s) is (are) TRUE? [Adv. 2018]
- $f(2) < 1 - \log_e 2$
 - $f(2) > 1 - \log_e 2$
 - $g(1) > 1 - \log_e 2$
 - $g(1) < 1 - \log_e 2$
26. Let $f: \left[-\frac{1}{2}, 2\right] \rightarrow \mathbb{R}$ and $g: \left[-\frac{1}{2}, 2\right] \rightarrow \mathbb{R}$ be functions defined by $f(x) = [x^2 - 3]$ and $g(x) = [x]f(x) + [4x - 7]f(x)$, where $[y]$ denotes the greatest integer less than or equal to y for $y \in \mathbb{R}$. Then [Adv. 2016]
- f is discontinuous exactly at three points in $\left[-\frac{1}{2}, 2\right]$
 - f is discontinuous exactly at four points in $\left[-\frac{1}{2}, 2\right]$
 - g is NOT differentiable exactly at four points in $\left[-\frac{1}{2}, 2\right]$
 - g is NOT differentiable exactly at five points in $\left[-\frac{1}{2}, 2\right]$
27. Let $a, b \in \mathbb{R}$ and $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = a \cos(|x^3 - x|) + b|x|\sin(|x^3 + x|)$. Then f is [Adv. 2016]
- differentiable at $x=0$ if $a=0$ and $b=1$
 - differentiable at $x=1$ if $a=1$ and $b=0$
 - NOT differentiable at $x=0$ if $a=1$ $b=0$
 - NOT differentiable at $x=1$ if $a=1$ and $b=1$
28. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function with $g(0) = 0$, $g'(0) = 0$ and $g'(1) \neq 0$. Let $f(x) = \begin{cases} \frac{x}{|x|} g(x), & x \neq 0 \\ 0, & x = 0 \end{cases}$ and $h(x) = e^{|x|}$ for all $x \in \mathbb{R}$. Let $(f \circ h)(x)$ denote $f(h(x))$ and $(h \circ f)(x)$ denote $h(f(x))$. Then which of the following is (are) true? [Adv. 2015]
- f is differentiable at $x = 0$
 - h is differentiable at $x = 0$
 - $f \circ h$ is differentiable at $x = 0$
 - $h \circ f$ is differentiable at $x = 0$

$$29. \text{ If } f(x) = \begin{cases} -x - \frac{\pi}{2}, & x \leq -\frac{\pi}{2} \\ -\cos x, & -\frac{\pi}{2} < x \leq 0, \text{ then} \\ x-1, & 0 < x \leq 1 \\ \ln x, & x > 1 \end{cases} \quad [2011]$$

- (a) $f(x)$ is continuous at $x = -\frac{\pi}{2}$
 (b) $f(x)$ is not differentiable at $x = 0$
 (c) $f(x)$ is differentiable at $x = 1$
 (d) $f(x)$ is differentiable at $x = -\frac{3}{2}$

30. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(x+y) = f(x) + f(y)$, $\forall x, y \in \mathbb{R}$. If $f(x)$ is differentiable at $x = 0$, then [2011]

- (a) $f(x)$ is differentiable only in a finite interval containing zero
 (b) $f(x)$ is continuous $\forall x \in \mathbb{R}$
 (c) $f'(x)$ is constant $\forall x \in \mathbb{R}$
 (d) $f(x)$ is differentiable except at finitely many points.

31. If $f(x) = \min \{1, x^2, x^3\}$, then [2006 - 5M, -1]

- (a) $f(x)$ is continuous $\forall x \in \mathbb{R}$
 (b) $f(x)$ is continuous and differentiable everywhere.
 (c) $f(x)$ is not differentiable at two points
 (d) $f(x)$ is not differentiable at one point

32. Let $h(x) = \min \{x, x^2\}$, for every real number of x , Then

[1998 - 2 Marks]

- (a) h is continuous for all x
 (b) h is differentiable for all x
 (c) $h'(x) = 1$, for all $x > 1$
 (d) h is not differentiable at two values of x .

33. The function $f(x) = \max \{(1-x), (1+x), 2\}$, $x \in (-\infty, \infty)$ is

- (a) continuous at all points [1995]
 (b) differentiable at all points
 (c) differentiable at all points except at $x = 1$ and $x = -1$
 (d) continuous at all points except at $x = 1$ and $x = -1$, where it is discontinuous

34. Let $g(x) = xf(x)$, where $f(x) = \begin{cases} x \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$. At $x = 0$

- (a) g is differentiable but g' is not continuous [1994]
 (b) g is differentiable while f is not
 (c) both f and g are differentiable
 (d) g is differentiable and g' is continuous

35. Let $f(x) = \begin{cases} 0, & x < 0 \\ x^2, & x \geq 0 \end{cases}$ then for all x [1994]

- (a) f' is differentiable (b) f is differentiable
 (c) f' is continuous (d) f is continuous

36. The function $f(x) = \begin{cases} |x-3|, & x \geq 1 \\ \frac{x^2}{4} - \frac{3x}{2} + \frac{13}{4}, & x < 1 \end{cases}$ is

[1988 - 2 Marks]

- (a) continuous at $x = 1$ (b) differentiable at $x = 1$
 (c) continuous at $x = 3$ (d) differentiable at $x = 3$.

37. The set of all points where the function $f(x) = \frac{x}{(1+|x|)}$ is differentiable, is [1987 - 2 Marks]

- (a) $(-\infty, \infty)$ (b) $[0, \infty)$
 (c) $(-\infty, 0) \cup (0, \infty)$ (d) $(0, \infty)$
 (e) None

38. Let $[x]$ denote the greatest integer less than or equal to x . If $f(x) = [x \sin \pi x]$, then $f(x)$ is [1986 - 2 Marks]

- (a) continuous at $x = 0$ (b) continuous in $(-1, 0)$
 (c) differentiable at $x = 1$ (d) differentiable in $(-1, 1)$
 (e) none of these

39. The function $f(x) = 1 + |\sin x|$ is [1986 - 2 Marks]

- (a) continuous nowhere
 (b) continuous everywhere
 (c) differentiable nowhere
 (d) not differentiable at $x = 0$
 (e) not differentiable at infinite number of points.

40. If $f(x) = x(\sqrt{x} - \sqrt{x+1})$, then [1985 - 2 Marks]

- (a) $f(x)$ is continuous but not differentiable at $x = 0$
 (b) $f(x)$ is differentiable at $x = 0$
 (c) $f(x)$ is not differentiable at $x = 0$
 (d) none of these

41. If $x + |y| = 2y$, then y as a function of x is [1984 - 3 Marks]

- (a) defined for all real x
 (b) continuous at $x = 0$
 (c) differentiable for all x
 (d) such that $\frac{dy}{dx} = \frac{1}{3}$ for $x < 0$



Match the Following

42. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ be functions defined by

$$f(x) = \begin{cases} x|x| \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases} \text{ and}$$

$$g(x) = \begin{cases} 1-2x, & 0 \leq x \leq \frac{1}{2}, \\ 0, & \text{otherwise.} \end{cases}$$

Let $a, b, c, d \in \mathbb{R}$. Define the function $h: \mathbb{R} \rightarrow \mathbb{R}$ by

$$h(x) = a f(x) + b \left(g(x) + g \left(\frac{1}{2} - x \right) \right) + c(x - g(x)) + d g(x), x \in \mathbb{R}$$

Match each entry in List-I to the correct entry in List-II.

List-I

List-II

- (P) If $a = 0, b = 1, c = 0$, and $d = 0$, then
 (Q) If $a = 1, b = 0, c = 0$, and $d = 0$, then

- (1) h is one-one.
 (2) h is onto.

(R) If $a = 0, b = 0, c = 1$, and $d = 0$, then

(S) If $a = 0, b = 0, c = 0$, and $d = 1$ then

(5) the range of h is $\{0, 1\}$.

The correct option is

[Adv. 2024]

- (a) (P) \rightarrow (4) (Q) \rightarrow (3) (R) \rightarrow (1) (S) \rightarrow (2)
 (b) (P) \rightarrow (5) (Q) \rightarrow (2) (R) \rightarrow (4) (S) \rightarrow (3)
 (c) (P) \rightarrow (5) (Q) \rightarrow (3) (R) \rightarrow (2) (S) \rightarrow (4)
 (d) (P) \rightarrow (4) (Q) \rightarrow (2) (R) \rightarrow (1) (S) \rightarrow (3)

43. Let $f_1 : \mathbb{R} \rightarrow \mathbb{R}, f_2 : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}, f_3 : \left(-1, e^{\frac{\pi}{2}} - 2\right) \rightarrow \mathbb{R}$ and $f_4 : \mathbb{R} \rightarrow \mathbb{R}$ be functions defined by [Adv. 2018]

(i) $f_1(x) = \sin \left(\sqrt{1 - e^{-x^2}} \right),$

(ii) $f_2(x) = \begin{cases} \frac{|\sin x|}{\tan^{-1} x} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases},$ where the inverse trigonometric function $\tan^{-1} x$ assumes values in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right),$

(iii) $f_3(x) = [\sin(\log_e(x+2))],$ where, for $t \in \mathbb{R}, [t]$ denotes the greatest integer less than or equal to $t,$

(iv) $f_4(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}.$

LIST-I

- P. The function f_1 is
 Q. The function f_2 is
 R. The function f_3 is
 S. The function f_4 is

The correct option is:

- (a) $P \rightarrow 2; Q \rightarrow 3; R \rightarrow 1; S \rightarrow 4$
 (c) $P \rightarrow 4; Q \rightarrow 2; R \rightarrow 1; S \rightarrow 3$

LIST-II

1. NOT continuous at $x = 0$
 2. continuous at $x = 0$ and NOT differentiable at $x = 0$
 3. differentiable at $x = 0$ and its derivative is NOT continuous at $x = 0$
 4. differentiable at $x = 0$ and its derivative is continuous at $x = 0$

(b) $P \rightarrow 4; Q \rightarrow 1; R \rightarrow 2; S \rightarrow 3$

(d) $P \rightarrow 2; Q \rightarrow 1; R \rightarrow 4; S \rightarrow 3$

44. Let $f_1 : \mathbb{R} \rightarrow \mathbb{R}, f_2 : [0, \infty) \rightarrow \mathbb{R}, f_3 : \mathbb{R} \rightarrow \mathbb{R}$ and $f_4 : \mathbb{R} \rightarrow [0, \infty)$ be defined by $f_1(x) = \begin{cases} |x| & \text{if } x < 0, \\ e^x & \text{if } x \geq 0; \end{cases}$

$f_2(x) = x^2; f_3(x) = \begin{cases} \sin x & \text{if } x < 0, \\ x & \text{if } x \geq 0; \end{cases}$ and $f_4(x) = \begin{cases} f_2(f_1(x)) & \text{if } x < 0, \\ f_2(f_1(x)) - 1 & \text{if } x \geq 0. \end{cases}$

[Adv. 2014]

List-I

- P. f_4 is
 Q. f_3 is
 R. $f_2 \circ f_1$ is
 S. f_2 is
 (a) $P \rightarrow 3; Q \rightarrow 1; R \rightarrow 4; S \rightarrow 2$
 (c) $P \rightarrow 3; Q \rightarrow 1; R \rightarrow 2; S \rightarrow 4$

List-II

1. Onto but not one-one
 2. Neither continuous nor one-one
 3. Differentiable but not one-one
 4. Continuous and one-one
 (b) $P \rightarrow 1; Q \rightarrow 3; R \rightarrow 4; S \rightarrow 2$
 (d) $P \rightarrow 1; Q \rightarrow 3; R \rightarrow 2; S \rightarrow 4$

45. In the following $[x]$ denotes the greatest integer less than or equal to x .

Match the functions in Column I with the properties in Column II and indicate your answer by darkening the appropriate bubbles in the 4×4 matrix given in the ORS. [2007 - 6 marks]

Column I

- (A) $x|x|$
 (B) $\sqrt{|x|}$
 (C) $x + [x]$
 (D) $|x-1| + |x+1|$

Column II

- (p) continuous in $(-1, 1)$
 (q) differentiable in $(-1, 1)$
 (r) strictly increasing in $(-1, 1)$
 (s) not differentiable at least at one point in $(-1, 1)$

46. In this questions there are entries in columns I and II. Each entry in column I is related to exactly one entry in column II. Write the correct letter from column II against the entry number in column I in your answer book.

[1992 - 2 Marks]

Column I

- (A) $\sin(\pi[x])$
 (B) $\sin(\pi(x-[x]))$

Column II

- (p) differentiable everywhere
 (q) nowhere differentiable
 (r) not differentiable at 1 and -1



10 Subjective Problems

47. If $f(x-y) = f(x) \cdot g(y) - f(y) \cdot g(x)$ and

$$g(x-y) = g(x) \cdot g(y) - f(x) \cdot f(y) \text{ for all } x, y \in \mathbb{R}.$$

If right hand derivative at $x=0$ exists for $f(x)$. Find derivative of $g(x)$ at $x=0$ [2005 - 4 Marks]

48. If $|c| \leq \frac{1}{2}$ and $f(x)$ is a differentiable function at $x=0$ given

$$\text{by } f(x) = \begin{cases} b \sin^{-1}\left(\frac{c+x}{2}\right), & -\frac{1}{2} < x < 0 \\ \frac{1}{2}, & x = 0 \\ \frac{e^{ax/2} - 1}{x}, & 0 < x < \frac{1}{2} \end{cases}$$

Find the value of 'a' and prove that $64b^2 = 4 - c^2$

[2004 - 4 Marks]

49. If a function $f: [-2a, 2a] \rightarrow \mathbb{R}$ is an odd function such that $f(x) = f(2a-x)$ for $x \in [a, 2a]$ and the left hand derivative at $x=a$ is 0 then find the left hand derivative at $x=-a$. [2003 - 2 Marks]

50. Let $f(x) = \begin{cases} x+a & \text{if } x < 0 \\ |x-1| & \text{if } x \geq 0, \end{cases}$ and [2002 - 5 Marks]

$$g(x) = \begin{cases} x+1 & \text{if } x < 0 \\ (x-1)^2 + b & \text{if } x \geq 0, \end{cases} \text{ where } a \text{ and } b \text{ are}$$

non-negative real numbers. Determine the composite function $g \circ f$. If $(g \circ f)(x)$ is continuous for all real x , determine the values of a and b . Further, for these values of a and b , is $g \circ f$ differentiable at $x=0$? Justify your answer.

51. Let $\alpha \in \mathbb{R}$. Prove that a function $f: \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at α if and only if there is a function $g: \mathbb{R} \rightarrow \mathbb{R}$ which is continuous at α and satisfies $f(x) - f(\alpha) = g(x)(x - \alpha)$ for all $x \in \mathbb{R}$. [2001 - 5 Marks]

52. Determine the values of x for which the following function fails to be continuous or differentiable: [1997 - 5 Marks]

$$f(x) = \begin{cases} 1-x, & x < 1 \\ (1-x)(2-x), & 1 \leq x \leq 2 \\ 3-x, & x > 2 \end{cases} \text{ Justify your answer.}$$

53. Let $f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}$ for all real x and y . If $f'(0)$ exists and equals -1 and $f(0)=1$, find $f(2)$. [1995 - 5 Marks]

54. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the equation $f(x+y) = f(x)f(y)$ for all x, y in \mathbb{R} and $f(x) \neq 0$ for any x in \mathbb{R} . Let the function be differentiable at $x=0$ and $f'(0)=2$. Show that $f'(x) = 2f(x)$ for all x in \mathbb{R} . Hence, determine $f(x)$. [1990 - 4 Marks]

55. Draw a graph of the function $y = [x] + |1-x|$, $-1 \leq x \leq 3$.

Determine the points, if any, where this function is not differentiable. [1989 - 4 Marks]

56. Let $f(x)$ be a function satisfying the condition $f(-x) = f(x)$ for all real x . If $f'(0)$ exists, find its value. [1987 - 2 Marks]

57. Let $f(x)$ be defined in the interval $[-2, 2]$ such that

$$f(x) = \begin{cases} -1, & -2 \leq x \leq 0 \\ x-1, & 0 < x \leq 2 \end{cases}$$

$$\text{and } g(x) = f(|x|) + |f(x)|$$

Test the differentiability of $g(x)$ in $(-2, 2)$. [1986 - 5 Marks]

58. Let $f(x) = x^3 - x^2 + x + 1$ and

$$\max\{f(t); 0 \leq t \leq x\}, \quad 0 \leq x \leq 1 \quad [1985 - 5 \text{ Marks}]$$

$$g(x) = 3 - x \quad 1 \leq x \leq 2$$

Discuss the continuity and differentiability of the function $g(x)$ in the interval $(0, 2)$.

59. Let $f(x) = \begin{cases} \frac{x^2}{2}, & 0 \leq x < 1 \\ 2x^2 - 3x + \frac{3}{2}, & 1 \leq x \leq 2 \end{cases} \quad [1983 - 2 \text{ Marks}]$

Discuss the continuity of f, f' and f'' on $[0, 2]$.

60. Find the derivative of

$$f(x) = \begin{cases} \frac{x-1}{2x^2-7x+5} & \text{when } x \neq 1 \\ -\frac{1}{3} & \text{when } x = 1 \end{cases}$$

at $x = 1$

[1979]

Topic-3: Chain Rule of Differentiation, Differentiation of Explicit & Implicit Functions, Parametric & Composite Functions, Logarithmic & Exponential Functions, Inverse Functions, Differentiation by Trigonometric Substitution



1 MCQs with One Correct Answer

- If y is a function of x and $\log(x+y) - 2xy = 0$, then the value of $y'(0)$ is equal to [2004S]
(a) 1 (b) -1 (c) 2 (d) 0
- If $y = (\sin x)^{\tan x}$, then $\frac{dy}{dx}$ is equal to [1994]
(a) $(\sin x)^{\tan x} (1 + \sec^2 x \log \sin x)$
(b) $\tan x (\sin x)^{\tan x - 1} \cos x$
(c) $(\sin x)^{\tan x} \sec^2 x \log \sin x$
(d) $\tan x (\sin x)^{\tan x - 1}$
- There exist a function $f(x)$, satisfying $f(0) = 1, f'(0) = -1, f(x) > 0$ for all x , and [1982 - 2 Marks]
(a) $f''(x) > 0$ for all x
(b) $-1 < f''(x) < 0$ for all x
(c) $-2 \leq f''(x) \leq -1$ for all x
(d) $f''(x) < -2$ for all x



2 Integer Value Answer/Non-Negative Integer

4. Let $f(\theta) = \sin\left(\tan^{-1}\left(\frac{\sin \theta}{\sqrt{\cos 2\theta}}\right)\right)$, where $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$.

Then the value of $\frac{d}{d(\tan \theta)}(f(\theta))$ is [2011]

5. If the function $f(x) = x^3 + e^{\frac{x}{2}}$ and $g(x) = f^{-1}(x)$, then the value of $g'(1)$ is [2009]



4 Fill in the Blanks

- If $xe^{xy} = y + \sin^2 x$, then at $x = 0$, $\frac{dy}{dx} = \dots\dots\dots$ [1996 - 1 Mark]
- If $f(x) = |x - 2|$ and $g(x) = f[f(x)]$, then $g'(x) = \dots\dots\dots$ for $x > 0$ [1990 - 2 Marks]
- The derivative of $\sec^{-1}\left(\frac{1}{2x^2 - 1}\right)$ with respect to $\sqrt{1 - x^2}$ at $x = \frac{1}{2}$ is $\dots\dots\dots$ [1986 - 2 Marks]
- If $f(x) = \log_x(\ln x)$, then $f'(x)$ at $x = e$ is $\dots\dots\dots$ [1985 - 2 Marks]
- If $f_r(x), g_r(x), h_r(x), r = 1, 2, 3$ are polynomials in x such that $f_r(a) = g_r(a) = h_r(a), r = 1, 2, 3$

$$\text{and } F(x) = \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix} \text{ then } F'(x) \text{ at } x = a \text{ is}$$

[1985 - 2 Marks]

11. If $y = f\left(\frac{2x-1}{x^2+1}\right)$ and $f'(x) = \sin x^2$, then $\frac{dy}{dx} =$ (a) $g'(2) = \frac{1}{15}$ (b) $h'(1) = 666$

[1982 - 2 Marks]

- (c)
- $h(0) = 16$
- (d)
- $h(g(3)) = 36$



5 True / False

12. The derivative of an even function is always an odd function. [1983 - 1 Mark]



6 MCQs with One or More Than One Correct

13. For any positive integer n , define $f_n : (0, \infty) \rightarrow \mathbb{R}$ as

$$f_n(x) = \sum_{j=1}^n \tan^{-1} \left(\frac{1}{1+(x+j)(x+j-1)} \right) \text{ for all } x \in (0, \infty).$$

(Here, the inverse trigonometric function $\tan^{-1} x$ assumes values in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.)

Then, which of the following statement(s) is (are) TRUE?

[Adv. 2018]

- (a) $\sum_{j=1}^5 \tan^2(f_j(0)) = 55$
 (b) $\sum_{j=1}^{10} (1 + f'_j(0)) \sec^2(f_j(0)) = 10$
 (c) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \tan(f_n(x)) = \frac{1}{n}$
 (d) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \sec^2(f_n(x)) = 1$

14. For every twice differentiable function $f : \mathbb{R} \rightarrow [-2, 2]$ with $(f(0))^2 + (f'(0))^2 = 85$, which of the following statement(s) is (are) TRUE? [Adv. 2018]

- (a) There exist $r, s \in \mathbb{R}$, where $r < s$, such that f is one-one on the open interval (r, s)
 (b) There exists $x_0 \in (-4, 0)$ such that $|f'(x_0)| \leq 1$
 (c) $\lim_{x \rightarrow \infty} f(x) = 1$
 (d) There exists $\alpha \in (-4, 4)$ such that $f'(\alpha) + f''(\alpha) = 0$ and $f'(\alpha) \neq 0$

15. Let $f : \mathbb{R} \rightarrow \mathbb{R}$, $g : \mathbb{R} \rightarrow \mathbb{R}$ and $h : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable functions such that $f(x) = x^3 + 3x + 2$, $g(f(x)) = x$ and $h(g(g(x))) = x$ for all $x \in \mathbb{R}$. Then [Adv. 2016]



9 Assertion and Reason / Statement Type Questions

16. Let $f(x) = 2 + \cos x$ for all real x .

STATEMENT - 1 : For each real t , there exists a point c in $[t, t + \pi]$ such that $f'(c) = 0$ because

STATEMENT - 2 : $f(t) = f(t + 2\pi)$ for each real t .

[2007 - 3 marks]

- (a) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
 (b) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1
 (c) Statement-1 is True, Statement-2 is False
 (d) Statement-1 is False, Statement-2 is True.



10 Subjective Problems

17. If $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$, prove that $\frac{y'}{y} = \frac{1}{x} \left(\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right)$. [1998 - 8 Marks]

18. Find $\frac{dy}{dx}$ at $x = -1$, when

$$(\sin y)^{\sin\left(\frac{\pi}{2}x\right)} + \frac{\sqrt{3}}{2} \sec^{-1}(2x) + 2^x \tan(\ln(x+2)) = 0$$

[1991 - 4 Marks]

19. If $x = \sec \theta - \cos \theta$ and $y = \sec^n \theta - \cos^n \theta$, then show

$$\text{that } (x^2 + 4) \left(\frac{dy}{dx} \right)^2 = n^2 (y^2 + 4) \quad [1989 - 2 \text{ Marks}]$$

20. If α be a repeated root of a quadratic equation $f(x) = 0$ and $A(x)$, $B(x)$ and $C(x)$ be polynomials of degree 3, 4 and 5

$$\text{respectively, then show that } \begin{vmatrix} A(x) & B(x) & C(x) \\ A(\alpha) & B(\alpha) & C(\alpha) \\ A'(\alpha) & B'(\alpha) & C'(\alpha) \end{vmatrix} \text{ is}$$

divisible by $f(x)$, where prime denotes the derivatives.

[1984 - 4 Marks]

21. Let $y = e^{x \sin x^3} + (\tan x)^x$. Find $\frac{dy}{dx}$ [1981 - 2 Marks]

22. Given $y = \frac{5x}{3\sqrt{1-x^2}} + \cos^2(2x+1)$; Find $\frac{dy}{dx}$. [1980]



Topic-4: Differentiation of Infinite Series, Successive Differentiation, nth Derivative of Some Standard Functions, Leibnitz's Theorem, Rolle's Theorem, Lagrange's Mean Value Theorem



1 MCQs with One Correct Answer

1. Let $g(x) = \log f(x)$ where $f(x)$ is twice differentiable positive function on $(0, \infty)$ such that $f(x+1) = xf(x)$. Then, for $N = 1, 2, 3, \dots$ [2008]

$$g''\left(N + \frac{1}{2}\right) - g''\left(\frac{1}{2}\right) =$$

(a) $-4\left\{1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N-1)^2}\right\}$

(b) $4\left\{1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N-1)^2}\right\}$

(c) $-4\left\{1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N+1)^2}\right\}$

(d) $4\left\{1 + \frac{1}{9} + \frac{1}{25} + \dots + \frac{1}{(2N+1)^2}\right\}$

2. $\frac{d^2x}{dy^2}$ equals [2007-3 marks]

(a) $\left(\frac{d^2y}{dx^2}\right)^{-1}$

(b) $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

(c) $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$

(d) $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

3. If $f(x)$ is a twice differentiable function and given that $f(1) = 1; f(2) = 4; f(3) = 9$, then [2005S]

(a) $f''(x) = 2$ for $\forall x \in (1, 3)$

(b) $f''(x) = f'(x) = 5$ for some $x \in (2, 3)$

(c) $f''(x) = 3$ for $\forall x \in (2, 3)$

(d) $f''(x) = 2$ for some $x \in (1, 3)$

4. If $x^2 + y^2 = 1$ then [2000]

(a) $yy'' - 2(y')^2 + 1 = 0$

(b) $yy'' + (y')^2 + 1 = 0$

(c) $yy'' + (y')^2 - 1 = 0$

(d) $yy'' + 2(y')^2 + 1 = 0$

5. Let $f(x)$ be a quadratic expression which is positive for all the real values of x . If $g(x) = f(x) + f'(x) + f''(x)$, then for any real x , [1990 - 2 Marks]

(a) $g(x) < 0$

(b) $g(x) > 0$

(c) $g(x) = 0$

(d) $g(x) \geq 0$

6. If $y^2 = P(x)$, a polynomial of degree 3, then

$2 \frac{d}{dx} \left(y^3 \frac{d^2y}{dx^2} \right)$ equals [1988 - 2 Marks]

(a) $P'''(x) + P'(x)$

(b) $P'(x)P'''(x)$

(c) $P(x)P'''(x)$

(d) a constant



2 Integer Value Answer/ Non-Negative Integer

7. For a polynomial $g(x)$ with real coefficients, let m_g denote the number of distinct real roots of $g(x)$. Suppose S is the set of polynomials with real coefficients defined by

$$S = \{(x^2 - 1)^2(a_0 + a_1x + a_2x^2 + a_3x^3) : a_0, a_1, a_2, a_3 \in \mathbb{R}\}.$$

For a polynomial f , let f' and f'' denote its first and second order derivatives, respectively. Then the minimum possible value of $(m_{f'} + m_{f''})$, where $f \in S$, is ____ [Adv 2019]



6 MCQs with One or More Than One Correct

8. Let S be the set of all twice differentiable functions f

from \mathbb{R} to \mathbb{R} such that $\frac{d^2f}{dx^2}(x) > 0$ for all $x \in (-1, 1)$.

For $f \in S$, let X_f be the number of points $x \in (-1, 1)$ for which $f(x) = x$. Then which of the following statements is(are) true? [Adv. 2023]

(a) There exists a function $f \in S$ such that $X_f = 0$

(b) For every function $f \in S$, we have $X_f \leq 2$

(c) There exists a function $f \in S$ such that $X_f = 2$

(d) There does NOT exist any function f in S such that $X_f = 1$

9. Let $f: (0, \pi) \rightarrow \mathbb{R}$ be a twice differentiable function such that

$$\lim_{t \rightarrow x} \frac{f(x) \sin t - f(t) \sin x}{t - x} = \sin^2 x \text{ for all } x \in (0, \pi).$$

If $f\left(\frac{\pi}{6}\right) = -\frac{\pi}{12}$, then which of the following statement(s) is(are) TRUE? [Adv. 2018]

(a) $f\left(\frac{\pi}{4}\right) = \frac{\pi}{4\sqrt{2}}$

(b) $f(x) < \frac{x^4}{6} - x^2$ for all $x \in (0, \pi)$

(c) There exists $\alpha \in (0, \pi)$ such that $f'(\alpha) = 0$

(d) $f''\left(\frac{\pi}{2}\right) + f\left(\frac{\pi}{2}\right) = 0$



9 Assertion and Reason / Statement Type Questions

10. Let f and g be real valued functions defined on interval $(-1, 1)$ such that $g''(x)$ is continuous, $g(0) \neq 0$, $g'(0) = 0$, $g''(0) \neq 0$, and $f(x) = g(x) \sin x$

STATEMENT - 1 : $\lim_{x \rightarrow 0} [g(x) \cot x - g(0) \operatorname{cosec} x] = f''(0)$ and

STATEMENT - 2 : $f'(0) = g(0)$ [2008]

- (a) Statement - 1 is True, Statement - 2 is True; Statement - 2 is a correct explanation for Statement - 1

(b) Statement - 1 is True, Statement - 2 is True; Statement - 2 is NOT a correct explanation for Statement - 1

(c) Statement - 1 is True, Statement - 2 is False

(d) Statement - 1 is False, Statement - 2 is True



10 Subjective Problems

11. Let f be a twice differentiable function such that

$$f''(x) = -f(x), \text{ and } f'(x) = g(x),$$

$$h(x) = [f(x)]^2 + [g(x)]^2$$

Find $h(10)$ if $h(5) = 11$

[1982 - 3 Marks]



Answer Key

Topic-1 : Continuity

1. (d) 2. (c) 3. $1 - \{-1, 0\}$ 4. (7) 5. (a, c, d) 6. (a, d) 7. (b, d) 8. (b, c) 9. (b)

Topic-2 : Differentiability

1. (b) 2. (b) 3. (c) 4. (a) 5. (b) 6. (a) 7. (d) 8. (d) 9. (d) 10. (a)
 11. (d) 12. (b) 13. (a) 14. (c) 15. (d) 16. (4) 17. (3) 18. (6) 19. $\mathbb{R} - \{0\}$
 20. (0) 21. (a, c, d) 22. (a, b) 23. (a, c)
 24. (a, c, d) 25. (b, c) 26. (b, c) 27. (a, b) 28. (a, d) 29. (a, b, c, d) 30. (b, c) 31. (a, d) 32. (a, c, d)
 33. (a, c) 34. (a, b) 35. (b, c, d) 36. (a, b, c) 37. (a) 38. (a, b, d) 39. (b, d, e) 40. (b) 41. (a, b, d)
 42. (c) 43. (d) 44. (d) 45. (A)-p, q, r; (B)-p, s; (C)-s, r; (D)-p, q 46. (A)-p; (B)-r

Topic-3 : Chain Rule of Differentiation, Differentiation of Explicit & Implicit Functions, Parametric & Composite Functions, Logarithmic & Exponential Functions, Inverse Functions, Differentiation by Trigonometric Substitution

1. (a) 2. (a) 3. (a) 4. (1) 5. (2) 6. $\frac{dy}{dx} = 1$ 7. $g'(x) = -4$
 8. $\left. \frac{du}{dv} \right|_{x=\frac{1}{2}} = 4$ 9. $\frac{1}{e}$ 10. (0) 11. $\frac{2+2x-2x^2}{(x^2+1)^2} \sin\left(\frac{2x-1}{x^2+1}\right)^2$ 12. True 13. (d) 14. (a, b, d)
 15. (b, c) 16. (b)

Topic-4 : Differentiation of Infinite Series, Successive Differentiation, nth Derivative of Some Standard Functions, Leibnitz's Theorem, Rolle's Theorem, Lagrange's Mean Value Theorem

1. (a) 2. (d) 3. (d) 4. (b) 5. (b) 6. (c) 7. (5.00) 8. (a, b, c) 9. (b, c, d) 10. (a)

Hints & Solutions



Topic-1: Continuity

1. (d) We have $f(x) = [x]^2 - [x^2]$
At $x = 0$,
L.H.L. = $\lim_{h \rightarrow 0} f(-h) = \lim_{h \rightarrow 0} ([-h]^2 - [(-h)^2])$
= $\lim_{h \rightarrow 0} ((-1)^2 - [h^2]) = \lim_{h \rightarrow 0} (1 - 0) = 1$
R.H.L. = $\lim_{h \rightarrow 0} f(h) = \lim_{h \rightarrow 0} ([h]^2 - [h^2]) = \lim_{h \rightarrow 0} (0 - 0) = 0$
 \therefore L.H.L. \neq R.H.L.
 $\therefore f(x)$ is not continuous at $x = 0$.
At $x = 1$
L.H.L. = $\lim_{h \rightarrow 0} f(1-h) = \lim_{h \rightarrow 0} ([1-h]^2 - [(1-h)^2])$
= $\lim_{h \rightarrow 0} (0 - 0) = 0$
R.H.L. = $\lim_{h \rightarrow 0} f(1+h) = \lim_{h \rightarrow 0} ([1+h]^2 - [(1+h)^2])$
= $\lim_{h \rightarrow 0} (1 - 1) = 0$
 $f(1) = [1]^2 - [1^2] = 1 - 1 = 0$
 \therefore L.H.L. = R.H.L. = $f(1)$
 $\therefore f(x)$ is continuous at $x = 1$.
Clearly $f(x)$ is not continuous at other integral points.
2. (c) $f(x) = [x] \cos\left(\frac{2x-1}{2}\right)\pi$
When x is not an integer, both the functions $[x]$ and $\cos\left(\frac{2x-1}{2}\right)\pi$ are continuous.
 $\therefore f(x)$ is continuous on all non integral points.
For $x = n \in \mathbb{I}$
LHL = $\lim_{x \rightarrow n^-} f(x) = \lim_{x \rightarrow n^-} [x] \cos\left(\frac{2x-1}{2}\right)\pi$
= $(n-1) \cos\left(\frac{2n-1}{2}\right)\pi = 0$
RHL = $\lim_{x \rightarrow n^+} f(x) = \lim_{x \rightarrow n^+} [x] \cos\left(\frac{2x-1}{2}\right)\pi$
= $n \cos\left(\frac{2n-1}{2}\right)\pi = 0$
Also $f(n) = n \cos\left(\frac{2n-1}{2}\right)\pi = 0$
Thus LHL = RHL = $f(x)$
 $\therefore f$ is continuous at all integral point.
Hence, f is continuous everywhere.
3. Clearly the given function is not defined for those values of x for which $[x+1] = 0$.
i.e., $0 \leq x+1 < 1 \Rightarrow -1 \leq x < 0$
 \therefore Required domain is $\mathbb{R} - [-1, 0)$

We know that $[x]$ is discontinuous at all integral value of x and

$\sin\left(\frac{\pi}{[x+1]}\right)$ is discontinuous for $[x+1] = 0$

$$\Rightarrow 0 \leq x+1 < 1 \Rightarrow -1 \leq x < 0$$

i.e., $[-1, 0)$

Also domain of $f = \mathbb{R} - [-1, 0)$

Hence points of discontinuity of f in their domain
= $\mathbb{I} - \{-1, 0\}$

4. $f(x)$ will be continuous at $x = 2$, if

$$\lim_{x \rightarrow 2} f(x) = f(2) \Rightarrow \lim_{x \rightarrow 2} \frac{x^3 + x^2 - 16x + 20}{(x-2)^2} = k$$

$$\Rightarrow k = \lim_{x \rightarrow 2} \frac{(x-2)^2(x+5)}{(x-2)^2} = \lim_{x \rightarrow 2} (x+5) = 7$$

5. (a, c, d) Given : $f(x) = x \cos(\pi(x + [x]))$

Let $x = n$ be any integer not equal to zero.

$$\text{Then } \lim_{x \rightarrow n^-} x \cos(\pi(x + [x])) = n \cos(\pi(n + n - 1))$$

$$= n \cos(2n - 1)\pi = -n$$

$$\text{and } \lim_{x \rightarrow n^+} x \cos(\pi(x + [x])) = n \cos(\pi(n + [n])) = n \cos 2n\pi = n$$

$$\therefore \text{LHL} \neq \text{RHL}$$

$$\Rightarrow f \text{ is discontinuous at } x = -1, 1, 2$$

$$\text{At } x = 0, \text{LHL} = \text{RHL} = 0 = f(0)$$

$$\therefore f \text{ is continuous at } x = 0.$$

6. (a, d) Let f and g be maximum at c_1 and c_2 respectively,

$$c_1, c_2 \in [0, 1]$$

$$\text{Then, } f(c_1) = g(c_2)$$

$$\text{Let } h(x) = f(x) - g(x)$$

$$\text{Then, } h(c_1) = f(c_1) - g(c_1) > 0$$

$$\text{and } h(c_2) = f(c_2) - g(c_2) < 0$$

$$\therefore h(x) = 0 \text{ has atleast one root in } [c_1, c_2]$$

$$\text{i.e. } f(c) = g(c) \text{ for } c \in [c_1, c_2],$$

which shows that options (a) and (d) are correct.

7. (b, d) Given : $f(x) = \begin{cases} a_n + \sin \pi x, & x \in [2n, 2n+1] \\ b_n + \cos \pi x, & x \in (2n-1, 2n) \end{cases}$

$\therefore f$ is continuous for all n

$$\therefore \text{At } x = 2n, \text{LHL} = \text{RHL} = f(2n)$$

$$\Rightarrow b_n + \cos 2\pi n = a_n + \sin 2\pi n = a_n + \sin 2\pi n$$

$$\Rightarrow b_n + 1 = a_n \Rightarrow a_n - b_n = 1, \therefore \text{option (b) is correct.}$$

$$\text{Also at } x = 2n+1, \text{LHL} = \text{RHL} = f(2n+1)$$

$$\Rightarrow \lim_{h \rightarrow 0} a_n + \sin \pi(2n+1-h)$$

$$= \lim_{h \rightarrow 0} b_{n+1} + \cos \pi(2n+1-h) = a_n + \sin(2n+1)\pi$$

$$\Rightarrow a_n = b_{n+1} - 1 = a_n \Rightarrow a_n - b_{n+1} = -1$$

$$\therefore \text{option (c) is incorrect.}$$

$$\Rightarrow a_{n-1} - b_n = -1, \therefore \text{option (d) is correct.}$$

8. (b, c) On $(0, \pi)$

$$(a) f(x) = \tan x$$

We know that $\tan x$ is discontinuous at $x = \pi/2$

$$(b) f(x) = \int_0^x t \sin\left(\frac{1}{t}\right) dt$$

$\Rightarrow f'(x) = x \sin \left(\frac{1}{x} \right)$, which exists on $(0, \pi)$

$\therefore f(x)$ is differentiable, on $(0, \pi)$, therefore it is continuous on $(0, \pi)$.

$$(c) f(x) = \begin{cases} 1, & 0 < x \leq 3\pi/4 \\ 2\sin \frac{2x}{9}, & 3\pi/4 < x < \pi \end{cases}$$

Clearly $f(x)$ may or may not be continuous at $x = \frac{3\pi}{4}$ but it is

continuous on $(0, \pi)$ except at $x = \frac{3\pi}{4}$.

$$\text{L.H.L.} = \lim_{h \rightarrow 0} f\left(\frac{3\pi}{4} - h\right) = \lim_{x \rightarrow 0} 1 = 1$$

$$\text{R.H.L.} = \lim_{h \rightarrow 0} f\left(\frac{3\pi}{4} + h\right) = \lim_{x \rightarrow 0} 2\sin \frac{2}{9}\left(\frac{3\pi}{4} + h\right)$$

$$= \lim_{h \rightarrow 0} 2\sin \left(\frac{\pi}{6} + \frac{2h}{9} \right) = 2\sin \frac{\pi}{6} = 1$$

$$\text{Also } f\left(\frac{3\pi}{4}\right) = 1$$

$$\therefore \text{L.H.L.} = \text{R.H.L.} = f\left(\frac{3\pi}{4}\right)$$

$\therefore f(x)$ is continuous at $x = \frac{3\pi}{4}$ and hence it is continuous on $(0, \pi)$

$$(d) f(x) = \begin{cases} x \sin x, & 0 < x \leq \pi/2 \\ \frac{\pi}{2} \sin(\pi + x), & \frac{\pi}{2} < x < \pi \end{cases}$$

Clearly $f(x)$ may or may not be continuous at

$x = \frac{\pi}{2}$ but it is continuous on $(0, \pi)$ except at $x = \pi/2$.

$$\text{At } x = \pi/2, \text{ L.H.L.} = \lim_{h \rightarrow 0} f\left(\frac{\pi}{2} - h\right)$$

$$= \lim_{h \rightarrow 0} \left(\frac{\pi}{2} - h \right) \sin \left(\frac{\pi}{2} - h \right) = \frac{\pi}{2} \sin \frac{\pi}{2} = \frac{\pi}{2}$$

$$\text{R.H.L.} = \lim_{h \rightarrow 0} f\left(\frac{\pi}{2} + h\right) = \lim_{h \rightarrow 0} \frac{\pi}{2} \sin \left(\pi + \frac{\pi}{2} + h \right)$$

$$= \frac{\pi}{2} \sin \left(\pi + \frac{\pi}{2} \right) = \frac{-\pi}{2} \sin \frac{\pi}{2} = -\frac{\pi}{2}$$

Thus, L.H.L. \neq R.H.L., $\therefore f(x)$ is not continuous on $(0, \pi)$.

$$9. (b) f(x) = \frac{x}{2} - 1$$

$$\therefore [f(x)] = \left[\frac{x}{2} - 1 \right] = -1, \text{ if } 0 \leq x < 2$$

$$\tan[f(x)] = \begin{cases} \tan(-1), & 0 \leq x < 2 \\ 0, & 2 \leq x \leq \pi \end{cases}$$

\therefore The function $\tan[f(x)]$ is discontinuous at $x = 2$.

$$\text{Also the function } \frac{1}{f(x)} = \frac{1}{\frac{x}{2} - 1} = \frac{2}{x - 2} \text{ is}$$

discontinuous at $x = 2$.

Thus both the given functions $\tan[f(x)]$ as well as $\frac{1}{f(x)}$ are discontinuous on the interval $[0, \pi]$.

$$\text{Now, } f^{-1}(x) = y \Rightarrow x = f(y) = \frac{y}{2} - 1 \Rightarrow y = 2(x + 1)$$

$\therefore f^{-1}(x) = 2(x + 1)$ is continuous on $[0, \pi]$

$$10. \text{ Given : } f(x) = \begin{cases} (1 + |\sin x|)^{\frac{a}{|\sin x|}}, & -\frac{\pi}{6} < x < 0 \\ b, & x = 0 \\ \frac{\tan 2x}{e^{\tan 3x}}, & 0 < x < \frac{\pi}{6} \end{cases}$$

is continuous at $x = 0$

$$\therefore \lim_{h \rightarrow 0} f(0 - h) = f(0) = \lim_{h \rightarrow 0} f(0 + h)$$

$$\text{Now, } \lim_{h \rightarrow 0} f(0 - h) = \lim_{h \rightarrow 0} [1 + |\sin(-h)|]^{\frac{a}{|\sin(-h)|}}$$

$$= \lim_{h \rightarrow 0} [1 + \sin h]^{\frac{a}{\sin h}} \Rightarrow \lim_{h \rightarrow 0} \frac{a}{\sin h} \log(1 + \sin h) = e^a$$

$$\text{and } \lim_{h \rightarrow 0} f(0 + h) = \lim_{h \rightarrow 0} \frac{\tan 2h}{e^{\tan 3h}}$$

$$= \lim_{h \rightarrow 0} \frac{\tan 2h}{2h} \times \frac{3h}{\tan 3h} \times \frac{2}{3} = e^{\frac{2}{3}}$$

Also $f(0) = b$

$$\therefore e^a = b = e^{\frac{2}{3}} \Rightarrow a = \frac{2}{3} \text{ and } b = e^{\frac{2}{3}}$$

$$11. \text{ Given : } f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2}, & x < 0 \\ a, & x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x} - 4}}, & x > 0 \end{cases}$$

Since $f(x)$ is continuous at $x = 0$,

\therefore L.H.L. at $(x = 0) = f(0)$

$$= \lim_{h \rightarrow 0} \frac{1 - \cos 4(0 - h)}{(0 - h)^2} = a \Rightarrow \lim_{h \rightarrow 0} \frac{1 - \cos 4h}{h^2} = a$$

$$= \lim_{h \rightarrow 0} \frac{2 \sin^2 2h}{4h^2} \cdot 4 = a \Rightarrow 8 = a$$

12. Given :

$$f(x) = \begin{cases} x + a\sqrt{2} \sin x, & 0 \leq x < \frac{\pi}{4} \\ 2x \cot x + b, & \frac{\pi}{4} \leq x \leq \frac{\pi}{2} \\ a \cos 2x - b \sin x, & \frac{\pi}{2} < x \leq \pi \end{cases}$$

is continuous for $0 \leq x \leq \pi$.

$$\therefore f(x) \text{ must be continuous at } x = \frac{\pi}{4} \text{ and } x = \frac{\pi}{2}$$

$$\Rightarrow \lim_{x \rightarrow \left(\frac{\pi}{4}\right)^-} f(x) = f\left(\frac{\pi}{4}\right)$$

$$\Rightarrow \lim_{h \rightarrow 0} f\left(\frac{\pi}{4} - h\right) = \frac{2\pi}{4} \cot \frac{\pi}{4} + b$$

$$\Rightarrow \lim_{h \rightarrow 0} \left(\frac{\pi}{4} - h\right) + a\sqrt{2} \lim_{h \rightarrow 0} \sin\left(\frac{\pi}{4} - h\right) = \frac{\pi}{2} + b$$

$$\Rightarrow \frac{\pi}{4} + a = \frac{\pi}{2} + b \Rightarrow a - b = \frac{\pi}{4} \quad \dots(i)$$

Also, $\lim_{x \rightarrow \left(\frac{\pi}{2}\right)^+} f(x) = f\left(\frac{\pi}{2}\right)$

$$\Rightarrow \lim_{h \rightarrow 0} f\left(\frac{\pi}{2} + h\right) = 2 \cdot \frac{\pi}{2} \cot \frac{\pi}{2} + b$$

$$\Rightarrow \lim_{h \rightarrow 0} a \cos 2\left(\frac{\pi}{2} + h\right) - b \sin\left(\frac{\pi}{2} + h\right) = b$$

$$\Rightarrow a \cos \pi - b \sin \frac{\pi}{2} = b \Rightarrow -a - b = b$$

$$\Rightarrow a + 2b = 0 \quad \dots(ii)$$

On solving (i) and (ii), we get $a = \frac{\pi}{6}$ and $b = \frac{-\pi}{12}$.

13. Let $h(x) = f(x) + g(x)$ be continuous.
 $\Rightarrow g(x) = h(x) - f(x)$
 Now, $h(x)$ and $f(x)$ both are continuous functions.
 $\therefore h(x) - f(x)$ must also be continuous. But it contradicts the given statement that $g(x)$ is discontinuous. Therefore our assumption that $f(x) + g(x)$ a continuous function is wrong and hence $f(x) + g(x)$ is discontinuous.

14. $f(x) = \begin{cases} 1+x, & 0 \leq x \leq 2 \\ 3-x, & 2 < x \leq 3 \end{cases}$

$$f(f(x)) = \begin{cases} 1+f(x) & 0 \leq f(x) \leq 2 \\ 3-f(x) & 2 < f(x) \leq 3 \end{cases}$$

$$\text{Now } 0 \leq x < 2 \Rightarrow 1 \leq x+1 \leq 3$$

$$\Rightarrow 1 \leq f(x) \leq 3$$

$$2 < x \leq 3 \Rightarrow -3 \leq -x < -2$$

$$\Rightarrow 0 \leq 3-x < 1 \Rightarrow 0 \leq f(x) < 1$$

$$\Rightarrow 0 \leq x \leq 1 \Rightarrow 1 \leq f(x) \leq 2$$

$$1 < x \leq 2 \Rightarrow 2 < f(x) \leq 3$$

$$2 < x \leq 3 \Rightarrow 0 \leq f(x) < 1$$

$$\Rightarrow f(f(x)) = \begin{cases} 2+x & 0 \leq x \leq 1 \\ 2-x & 1 < x \leq 2 \\ 4-x & 2 < x \leq 3 \end{cases}$$

$$\text{At } x = 1, \text{ R.H.L.} = \lim_{h \rightarrow 0} g(1+h) = \lim_{h \rightarrow 0} 2 - (1+h) = 1$$

$g(1) = 3$, \therefore discontinuous at $x = 1$

$$\text{At } x = 2, \text{ R.H.L.} = \lim_{h \rightarrow 0} g(2+h) = \lim_{h \rightarrow 0} 4 - (2+h) = 2$$

$$g(2) = 0, \therefore \text{discontinuous at } x = 2$$

15. Given: $f(x+y) = f(x) + f(y) + x, y$
 As $f(x)$ is continuous at $x = 0$, we have
 LHL = RHL = $f(0)$

$$\Rightarrow \lim_{h \rightarrow 0} f(0-h) = \lim_{h \rightarrow 0} f(0+h) = f(0)$$

$$\Rightarrow f(0) + \lim_{h \rightarrow 0} f(-h) = f(0) + \lim_{h \rightarrow 0} f(h) = f(0)$$

$$\Rightarrow \lim_{h \rightarrow 0} f(-h) = \lim_{h \rightarrow 0} f(h) = 0 \quad \dots(i)$$

Now let $x = a$ be any arbitrary point then at $x = a$,

$$\text{LHL} = \lim_{h \rightarrow 0} f(a-h) = \lim_{h \rightarrow 0} [f(a) + f(-h)]$$

$$= f(a) + \lim_{h \rightarrow 0} f(-h) = f(a) \quad [\text{using (i)}]$$

$$\text{Similarly, R.H.L.} = \lim_{h \rightarrow 0} f(a+h) = f(a)$$

$$\therefore \lim_{h \rightarrow 0} f(a-h) = \lim_{h \rightarrow 0} f(a+h) = f(a)$$

Hence, f is continuous at $x = a$. Since a is any arbitrary point,

$\therefore f$ is continuous $\forall x \in R$.



Topic-2: Differentiability

1. (b) Given that, $\lim_{t \rightarrow x} \frac{t^{10} f(x) - x^{10} f(t)}{t^9 - x^9} = 1$

By L-Hospital Rule

$$\lim_{t \rightarrow x} \frac{10t^9 f(x) - f'(t)x^{10}}{9t^8} = 1$$

$$\Rightarrow 10x^9 f(x) - f'(x)x^{10} = 9x^8$$

$$\Rightarrow f'(x) - \frac{10}{x} f(x) = -\frac{9}{x^2}$$

$$\text{IF} = e^{-\int \frac{10}{x} dx} = \frac{1}{x^{10}}$$

\therefore Solution is

$$\frac{y}{x^{10}} = \int -\frac{9}{x^{10}} \times \frac{1}{x^2} dx$$

$$= -9 \int x^{-12} dx$$

$$\frac{y}{x^{10}} = \frac{9}{11} x^{-11} + C$$

$$\text{put } x = 1 \text{ and } y = 2, \text{ we get } C = \frac{13}{11}$$

$$\Rightarrow y = \frac{9}{11x} + \frac{13}{11} x^{10}$$

2. (b) $f'(0^+) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$

$$= \lim_{h \rightarrow 0} \frac{h^2 \left| \cos \frac{\pi}{h} \right|}{h} = \lim_{h \rightarrow 0} h \left| \cos \frac{\pi}{h} \right|$$

$$= 0 \times \text{some finite value} = 0$$

$$\text{and } f'(0^-) = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{h^2 \left| \cos \frac{\pi}{-h} \right|}{-h}$$

$$= \lim_{h \rightarrow 0} -h \left| \cos \frac{\pi}{h} \right| = 0 \times \text{some finite value} = 0$$

$\therefore f'(0^+) = f'(0^-) \therefore f$ is differentiable at $x = 0$

$$\text{Now } f'(2^+) = \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2 \left| \cos \frac{\pi}{2+h} \right| - 4 \left| \cos \frac{\pi}{2} \right|}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2 \left(\cos \frac{\pi}{2+h} \right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2}{h} \sin \left(\frac{\pi}{2} - \frac{\pi}{2+h} \right)$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2}{h} \sin \left(\frac{\pi h}{2(2+h)} \right)$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2}{h} \times \frac{\sin \left(\frac{\pi h}{2(2+h)} \right)}{\left(\frac{\pi h}{2(2+h)} \right)} \times \frac{\pi h}{2(2+h)} = \pi$$

$$\text{and } f'(2^-) = \lim_{h \rightarrow 0} \frac{f(2-h) - f(2)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{(2-h)^2 \left| \cos \left(\frac{\pi}{2-h} \right) \right| - 4}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{-(2-h)^2 \cos \left(\frac{\pi}{2-h} \right)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{(2-h)^2 \sin \left(\frac{\pi}{2} - \frac{\pi}{2-h} \right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2-h)^2}{h} \times \frac{\sin \left(\frac{-\pi h}{2(2-h)} \right)}{\left(\frac{-\pi h}{2(2-h)} \right)} \times \left(\frac{-\pi h}{2(2-h)} \right) = -\pi$$

$\therefore f'(2^+) \neq f'(2^-), \therefore f$ is not differentiable at $x = 2$.

3. (c) $\therefore p$ = left hand derivative of $|x-1|$ at $x = 1 \Rightarrow p = -1$

Now $\lim_{x \rightarrow 1^+} g(x) = p$, where

$$g(x) = \frac{(x-1)^n}{\log \cos^m(x-1)}, \quad 0 < x < 2,$$

m, n are integers, $m \neq 0, n > 0$

$$\therefore \lim_{x \rightarrow 1^+} \frac{(x-1)^n}{\log \cos^m(x-1)} = -1$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^n}{\log \cos^m h} = -1$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^n}{m(\log \cosh h)} = -1 \quad (\text{using LH rule})$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{n h^{n-1} \cosh h}{m(-\sin h)} = -1 \quad (\text{using LH rule})$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{n h^{n-2} \cosh h}{m \left(\frac{\sin h}{h} \right)} = 1 \Rightarrow n = 2 \text{ and } m = 2$$

4. (a) Given : $f(x)$ is differentiable on $(0, \infty)$ such that

$$f(1) = 1 \text{ and } \lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1 \text{ for each } x > 0$$

$$\Rightarrow \lim_{t \rightarrow x} \frac{2t f(x) - x^2 f'(t)}{1} = 1 \quad (\text{using L'H rule})$$

$$\Rightarrow 2xf(x) - x^2 f'(x) = 1 \Rightarrow f'(x) - \frac{2}{x} f(x) = -\frac{1}{x^2}$$

(Linear differential equation) Integrating factor,

$$e^{\int -\frac{2}{x} dx} = e^{-2 \log x} = e^{\log 1/x^2} = \frac{1}{x^2}$$

$$\therefore \text{Solution : } f(x) \times \frac{1}{x^2} = \int \left(-\frac{1}{x^2} \right) \times \frac{1}{x^2} dx$$

$$\Rightarrow \frac{f(x)}{x^2} = \frac{1}{3x^3} + c \Rightarrow f(x) = cx^2 + \frac{1}{3x}$$

$$\therefore f(1) = 1,$$

$$\therefore 1 = c + \frac{1}{3} \Rightarrow c = 2/3$$

$$\therefore f(x) = \frac{2}{3}x^2 + \frac{1}{3x}$$

5. (b) Given : $f(x)$ is a continuous and differentiable function and

$$f\left(\frac{1}{n}\right) = 0, \forall n \geq 1 \text{ and } n \in \mathbb{I}$$

$$\therefore f(0^+) = f\left(\frac{1}{\infty}\right) = 0$$

$$\therefore \text{R.H.L.} = 0,$$


$$\therefore f(0) = 0 \text{ for } f(x) \text{ to be continuous.}$$

$$\text{Also } f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h - 0} = \lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$$

= 0 (using $f(0) = 0$ and $f(0^+) = 0$)

$$\therefore f(0) = 0, f'(0) = 0$$

6. (a) Graph of $y = ||x| - 1|$ is as follows :



The graph has sharp turnings at $x = -1, 0$. Therefore given function is not differentiable at $x = -1, 0, 1$.

7. (d) $f(x) = \begin{cases} \tan^{-1} x, & \text{if } |x| \leq 1 \\ \frac{1}{2}(|x|-1), & \text{if } |x| > 1 \end{cases}$

$$\Rightarrow f(x) = \begin{cases} \frac{1}{2}(-x-1), & \text{if } x < -1 \\ \tan^{-1} x, & \text{if } -1 \leq x \leq 1 \\ \frac{1}{2}(x-1), & \text{if } x > 1 \end{cases}$$

Clearly L.H.L. at $(x = -1) = \lim_{h \rightarrow 0} f(-1-h) = 0$

R.H.L. at $(x = -1) = \lim_{h \rightarrow 0} f(-1+h)$
 $= \lim_{h \rightarrow 0} \tan^{-1}(-1+h) = 3\pi/4$

\therefore At $x = -1$, L.H.L. \neq R.H.L.

$\therefore f(x)$ is discontinuous at $x = -1$

Also we can prove in the same way, that $f(x)$ is discontinuous at $x = 1$

$\therefore f'(x)$ can not be found for $x = \pm 1$

Hence, domain of $f'(x) = R - \{-1, 1\}$

8. (d) Let us test each of four options :

(a) $f(x) = \cos |x| + |x| = \begin{cases} \cos x - x, & x < 0 \\ \cos x + x, & x \geq 0 \end{cases}$

$$f'(x) = \begin{cases} -\sin x - 1, & x < 0 \\ -\sin x + 1, & x \geq 0 \end{cases}$$

At $x = 0$, LHD = -1, RHD = 1

$\therefore f(x)$ is not differentiable.

(b) $f(x) = \cos |x| - |x| = \begin{cases} \cos x + x, & x < 0 \\ \cos x - x, & x \geq 0 \end{cases}$

$\therefore f(x)$ is not differentiable at $x = 0$

(c) $f(x) = \sin |x| + |x| = \begin{cases} -\sin x - x, & x < 0 \\ \sin x + x, & x \geq 0 \end{cases}$

$\therefore f(x)$ is not differentiable at $x = 0$

(d) $f(x) = \sin |x| - |x| = \begin{cases} -\sin x + x, & x < 0 \\ \sin x - x, & x \geq 0 \end{cases}$

$$f'(x) = \begin{cases} -\cos x + 1, & x < 0 \\ \cos x - 1, & x \geq 0 \end{cases}$$


At $x = 0$, LHD = 0, RHD = 0

$\therefore f(x)$ is differentiable at $x = 0$.

9. (d) $f(x) = \max. \{x, x^3\}$

$$= \begin{cases} x; & x < -1 \\ x^3; & -1 \leq x \leq 1 \\ x; & 0 \leq x \leq 1 \\ x^3; & x \geq 1 \end{cases}$$

Graph of $f(x) = \max. \{x, x^3\}$ is as shown with solid lines.



We know that a continuous function $f(x)$ is not differentiable at $x = a$ if graphically it takes a sharp turn at $x = a$. Since, in the graph there are sharp turns at $x = -1, 0, 1$; $\therefore f(x)$ is not differentiable at $x = -1, 0, 1$.

10. (a) LHD = $\lim_{h \rightarrow 0} \frac{f(k) - f(k-h)}{h}$ ($k = \text{integer}$)

$$= \lim_{h \rightarrow 0} \frac{[k] \sin k\pi - [k-h] \sin(k-h)\pi}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-(k-1) \sin(k-h)\pi}{h} \quad [\because \sin k\pi = 0]$$

$$= \lim_{h \rightarrow 0} \frac{-(k-1) \sin(k\pi - h\pi)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-(k-1)(-1)^{k-1} \sin h\pi}{h\pi} \times \pi$$

$$= \pi(k-1)(-1)^k \quad [\because \sin(k\pi - \theta) = (-1)^{k-1} \sin \theta]$$

11. (d) Since $|x| = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$

$$|x^2 - 3x + 2| = |(x-1)(x-2)| = \begin{cases} (x-1)(x-2), & \text{if } x < 1 \\ (x-1)(2-x), & \text{if } 1 \leq x < 2 \\ (x-1)(x-2), & \text{if } x \geq 2 \end{cases}$$

and $\cos(-\theta) = \cos \theta \Rightarrow \cos |x| = \cos x$

$$\therefore f(x) = \begin{cases} (x^2-1)(x-1)(x-2) + \cos x, & \text{if } x \leq 1 \\ -(x^2-1)(x-1)(x-2) + \cos x, & \text{if } 1 \leq x < 2 \\ (x^2-1)(x-1)(x-2) + \cos x, & \text{if } x \geq 2 \end{cases}$$

This function may or may not be differentiable at $x = 1$ and $x = 2$ but is differentiable at all points except at $x = 1$ and $x = 2$. Let us check the differentiability at $x = 1$ and $x = 2$.

$$\text{Lf}'(1) = \left\{ \frac{d}{dx} [(x^2-1)(x-1)(x-2) + \cos x] \right\}_{x=1} = -\sin 1$$

$$\text{Rf}'(1) = \left\{ \frac{d}{dx} [-(x^2-1)(x-1)(x-2) + \cos x] \right\}_{x=1} = -\sin 1$$

$\therefore \text{Lf}'(1) = \text{Rf}'(1)$
 $\therefore f$ is differentiable at $x = 1$.

$$\text{Lf}'(2) = \left\{ \frac{d}{dx} [-(x^2-1)(x-1)(x-2) + \cos x] \right\}_{x=2} = -3 - \sin 2$$

$$Rf'(2) = \left\{ \frac{d}{dx} ((x^2 - 1)(x - 1)(x - 2) + \cos x) \right\}_{x=2}$$

$$= 3 - \sin 2$$

$$\therefore Lf'(2) \neq Rf'(2)$$

$\therefore f$ is not differentiable at $x = 2$.

12. (b) $f(x) = [\tan^2 x]$

$\tan x$ is an increasing function for $-\frac{\pi}{4} < x < \frac{\pi}{4}$

$$\therefore \tan\left(-\frac{\pi}{4}\right) < \tan x < \tan\left(\frac{\pi}{4}\right)$$

$$\Rightarrow -1 < \tan x < 1 \Rightarrow 0 \leq \tan^2 x < 1$$

$$\Rightarrow [\tan^2 x] = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} [\tan^2 x] = 0, f(0) = 0$$

$\therefore f(x)$ is continuous at $x = 0$

13. (a) $f: R \rightarrow R$ is a differentiable function and $f(1) = 4$

$$\lim_{x \rightarrow 1} \int_4^{f(x)} \frac{2t}{x-1} dt = \lim_{x \rightarrow 1} \left[\frac{t^2}{x-1} \right]_4^{f(x)}$$

$$= \lim_{x \rightarrow 1} \frac{(f(x))^2 - 16}{x-1} = \lim_{x \rightarrow 1} \frac{f(x)-4}{x-1} \cdot \lim_{x \rightarrow 1} (f(x)+4)$$

$$= \lim_{x \rightarrow 1} f'(x) \cdot \lim_{x \rightarrow 1} (f(x)+4)$$

$$= f'(1) \cdot (f(1)+4) = 8f'(1) \quad [\because f(1)=4]$$

14. (c) $\lim_{x \rightarrow a} \frac{g(x)f(a) - g(a)f(x)}{x-a}$

$$= \lim_{h \rightarrow 0} \frac{g(a+h)f(a) - g(a)f(a+h)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{g(a+h)f(a) - g(a)f(a) + g(a)f(a) - g(a)f(a+h)}{h}$$

$$= \lim_{h \rightarrow 0} f(a) \left[\frac{g(a+h) - g(a)}{h} \right] - \lim_{h \rightarrow 0} g(a) \left[\frac{f(a+h) - f(a)}{h} \right]$$

$$= f(a)g'(a) - g(a)f'(a) = 2 \times 2 - (-1) \times 1 = 5$$

15. (d) Given: $f(x) = \frac{\tan(\pi[x - \pi])}{1 + [x]^2}$

Clearly $[x - \pi]$ is an integer whatever be the value of x .

$\therefore \pi[x - \pi]$ is an integral multiple of π .

Consequently $\tan(\pi[x - \pi]) = 0, \forall x$.

Also $1 + [x]^2 \neq 0$ for any x .

$$\therefore f(x) = 0.$$

Hence, $f(x)$ is constant function and therefore, it is continuous and differentiable any number of times, that is $f'(x), f''(x), f'''(x), \dots$ all exist for every x , their value being 0 at every point x . Hence, out of all the alternatives only (d) is correct.


16. (4) Given that $f(x) = |2x - 1| + |2x + 1|$

$$\text{and } g(x) = x - [x] = \{x\}$$

$$\therefore (\text{fog})(x) = |2\{x\} - 1| + |2\{x\} + 1|$$

$$\Rightarrow (\text{fog})(x) = \begin{cases} |2x+1| + |2x+3|, & x \in \left(-1, -\frac{1}{2}\right] \\ |2x+1| + |2x+3|, & x \in \left(-\frac{1}{2}, 0\right] \\ |2x-1| + |2x+1|, & x \in \left(0, \frac{1}{2}\right] \\ |2x-1| + |2x+1|, & x \in \left(\frac{1}{2}, 1\right] \end{cases}$$

$$\Rightarrow (\text{fog})(x) = \begin{cases} 2, & x \in \left[-1, -\frac{1}{2}\right] \\ 4x+4, & x \in \left(-\frac{1}{2}, 0\right] \\ 2, & x \in \left(0, \frac{1}{2}\right] \\ 4x, & x \in \left(\frac{1}{2}, 1\right] \end{cases}$$



$\therefore f(g(x))$ is discontinuous at $x = 0$.

$$\therefore c = 1$$

$$\text{Now, } (\text{fog})(x) = \begin{cases} 0, & x \in \left(-1, \frac{1}{2}\right) \\ 4, & x \in \left(-\frac{1}{2}, 0\right) \\ 0, & x \in \left(0, \frac{1}{2}\right) \\ 4, & x \in \left(\frac{1}{2}, 1\right) \end{cases}$$


$\therefore f(g(x))$ is non-differentiable at $x = -\frac{1}{2}, 0, \frac{1}{2}$

$$\therefore d = 3$$

$$\text{Hence, } c + d = 4$$

17. (3) $f(x) = |x| + 1 = \begin{cases} x+1, & x \geq 0 \\ -x+1, & x < 0 \end{cases}$

$$g(x) = x^2 + 1$$



From graph, it is clear that there are 3 points at which $h(x)$ is not differentiable.

18. (6) $6 \int_1^x f(t) dt = 3xf(x) - x^3$
On differentiating, we get $6f(x) = 3f(x) + 3xf'(x) - 3x^2$
 $\Rightarrow f'(x) - \frac{1}{x}f(x) = x$, I.F. = $\frac{1}{x}$

$$\therefore \text{Solution is } f(x) \cdot \frac{1}{x} = \int 1 \cdot dx = x + c$$

$$\therefore f(x) = x^2 + cx$$

But $f(1) = 2 \Rightarrow c = 1$, $\therefore f(x) = x^2 + x$
 $\Rightarrow f(2) = 4 + 2 = 6$

Note: Putting $x = 1$ in given integral equation, we get

$$f(1) = \frac{1}{3} \text{ while given } f(1) = 2.$$

\therefore Data given in the question is inconsistent.

19. We have,

$$f(x) = x|x| = \begin{cases} -x^2, & x < 0 \\ x^2, & x \geq 0 \end{cases}$$

$$f'(x) = \begin{cases} -2x, & x < 0 \\ 2x, & x \geq 0 \end{cases} \Rightarrow f''(x) = \begin{cases} -2, & x < 0 \\ 2, & x \geq 0 \end{cases}$$

Thus $f''(x)$ exists at each point except at $x = 0$
 $\therefore f(x)$ is twice differentiable on $\mathbb{R} - \{0\}$.

20. Given: $f(x) = \begin{cases} (x-1)^2 \sin \frac{1}{x-1} - |x|, & x \neq 1 \\ -1, & x = 1 \end{cases}$

Since $|x|$ is not differentiable at $x = 0$

$$\therefore (x-1)^2 \sin \frac{1}{x-1} - |x| \text{ is not differentiable at } x = 0.$$

At all other values of x , $f(x)$ is differentiable.

\therefore Required set of points is $\{0\}$.

21. (a, c, d) Let domain and codomain of function $y = f(x)$ are S and T respectively.

(a) There are infinitely many elements in domain sets S and four elements in codomain set T .

So, there are infinitely many function from S to T .

Hence, option (a) is correct

(b) If number of elements in domain is greater than number of elements in co-domain, then number of strictly increasing function is zero. Hence, option (b) is incorrect.

(c) Since, every subset $(0, 1)$, $(1, 2)$, $(3, 4)$ has four choices.

\therefore Maximum number of continuous functions

$$= 4 \times 4 \times 4 = 64 [\because 64 < 120]$$

Hence, option (c) is correct.

(d) Since, every continuous function is piece wise constant functions. $f'(x) = 0$, so, $f(x)$ is differentiable.

Hence option (d) is correct

$$22. \text{ (a, b) } f(x) = \begin{cases} 0 & ; 0 < x < \frac{1}{4} \\ \left(x - \frac{1}{4}\right)^2 \left(x - \frac{1}{2}\right) & ; \frac{1}{4} \leq x < \frac{1}{2} \\ 2\left(x - \frac{1}{4}\right)^2 \left(x - \frac{1}{2}\right) & ; \frac{1}{2} \leq x < \frac{3}{4} \\ 3\left(x - \frac{1}{4}\right)^2 \left(x - \frac{1}{2}\right) & ; \frac{3}{4} \leq x < 1 \end{cases}$$

$$f\left(\frac{3}{4}^-\right) = \frac{1}{8} \text{ and } f\left(\frac{3}{4}^+\right) = \frac{3}{16}$$

So, $f(x)$ is discontinuous at $x = \frac{3}{4}$ only and

$$f'(x) = \begin{cases} 0 & ; 0 < x < \frac{1}{4} \\ 2\left(x - \frac{1}{4}\right)\left(x - \frac{1}{2}\right) + \left(x - \frac{1}{4}\right)^2 & ; \frac{1}{4} < x < \frac{1}{2} \\ 4\left(x - \frac{1}{4}\right)\left(x - \frac{1}{2}\right) + 2\left(x - \frac{1}{4}\right)^2 & ; \frac{1}{2} < x < \frac{3}{4} \\ 6\left(x - \frac{1}{4}\right)\left(x - \frac{1}{2}\right) + 3\left(x - \frac{1}{4}\right)^2 & ; \frac{3}{4} < x < 1 \end{cases}$$

$$\therefore f'\left(\frac{1}{2}^-\right) \neq f'\left(\frac{1}{2}^+\right) \text{ and } f'\left(\frac{3}{4}^-\right) \neq f'\left(\frac{3}{4}^+\right)$$

$f(x)$ is non-differentiable at $x = \frac{1}{2}$ and $\frac{3}{4}$ and minimum

values of $f(x)$ occur at $x = \frac{5}{12}$ whose values is $-\frac{1}{432}$

23. (a, c) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = (x^2 + \sin x)(x - 1)$

$$\text{Then, } f(1^+) = f(1^-) = f(1) = 0$$

$$\text{Let } (fg): \mathbb{R} \rightarrow \mathbb{R} \text{ is defined by } (fg)(x) = f(x) \cdot g(x)$$

$$\text{Let } fg(x) = h(x) = f(x) \cdot g(x) \text{ then } h: \mathbb{R} \rightarrow \mathbb{R}$$

$$h'(x) = f'(x)g(x) + f(x)g'(x)$$

If g is differentiable at $x = 1$

$$h'(1) = f'(1)g(1) + 0, \quad [\because f(1) = 0]$$

\Rightarrow if $g(x)$ is differentiable then $h(x)$ is also differentiable (true)

\Rightarrow if $g(x)$ is differentiable at $x = 1$, then fg is also differentiable at $x = 1$

If $g(x)$ is continuous at $x = 1$, then $g(1^+) = g(1^-) = g(1)$

$$h'(1^+) = \lim_{h \rightarrow 0^+} \frac{h(1+h) - h(1)}{h}$$

$$h'(1^+) = \lim_{h \rightarrow 0^+} \frac{f(1+h)g(1+h) - 0}{h} = f'(1)g(1)$$

$$h'(1^-) = \lim_{h \rightarrow 0^+} \frac{f(1-h)g(1-h) - 0}{-h} = f'(1)g(1)$$

$\Rightarrow h(x) = f(x)g(x)$ is differentiable at $x = 1$ (True)

So, if g is continuous at $x = 1$, then fg is differentiable at $x = 1$.

option (b) (d) $h'(1^+) = \lim_{h \rightarrow 0^+} \frac{h(1+h) - h(1)}{-h}$

$$h'(1^+) = \lim_{h \rightarrow 0^+} \frac{f(1+h)g(1+h)}{h} = f'(1)g(1^+)$$

$$h'(1^-) = \lim_{h \rightarrow 0^+} \frac{f(1-h)g(1-h)}{-h} = f'(1)g(1^-)$$

$$\Rightarrow g(1^+) = g(1^-)$$

So, it does not mean that if fg is differentiable at $x = 1$, then g is continuous or differentiable at $x = 1$

24. (a, c, d)

$$f(x) = \begin{cases} (x^5 + 5x^4 + 10x^3 + 10x^2 + 5x + 1) - 2x, & x < 0 \\ x^2 - 2 \times \frac{1}{2} \times x + \frac{1}{4} + \frac{3}{4}, & 0 \leq x < 1 \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3 \\ (x-2)\log_e(x-2) - x + \frac{10}{3}, & x \geq 3 \end{cases}$$

$$= \begin{cases} (x+1)^5 - 2x, & x < 0 \\ \left(x - \frac{1}{2}\right)^2 + \frac{3}{4}, & 0 \leq x < 1 \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3 \\ (x-2)\log_e(x-2) - x + \frac{10}{3}, & x \geq 3 \end{cases}$$

For $x = 0$, $f(x) = 1$

For $x < 0$, $f(x) = (x+1)^5 - 2x$

It decreases to $-\infty$.

$\therefore f(x) \in (-\infty, 1]$ for $x \leq 0$

For $x = 3$, $f(x) = \frac{1}{3}$

For $x \geq 3$, $f(x)$ increases to ∞

$\therefore f(x) \in \left[\frac{1}{3}, \infty\right)$ for $x \geq 3$

On combining the two $f(x) \in R \Rightarrow f$ is onto.

\therefore option (d) is correct.

$$f'(x) = \begin{cases} 5(x+1)^4 - 2, & x < 0 \\ 2x - 1, & 0 \leq x < 1 \\ 2(x-2)^2 - 1, & 1 \leq x < 3 \\ \log_e(x-2), & x \geq 3 \end{cases}$$

$Lf''(1) = 2$, $Rf''(1) = -4 \Rightarrow f'$ is not differentiable at $x = 1$

\therefore option (c) is correct.

For $x < 0$, $f(x) = 5(x+1)^4 - 1$


$$\text{Now, } f'(x) = 0 \Rightarrow (x+1)^4 = \frac{1}{5} \Rightarrow x = -1 \pm \left(\frac{1}{5}\right)^{1/4}$$

$\Rightarrow f$ is changing its nature at $x = -1 - \left(\frac{1}{5}\right)^{1/4}$

$\therefore f$ is not increasing on $(-\infty, 0) \therefore$ option (b) is incorrect.

$$f'(x) = \begin{cases} 2x - 1, & 0 \leq x < 1 \\ 2(x-2)^2 - 1, & 1 \leq x < 3 \end{cases}$$

From its graph $f'(x)$ has local maxima at $x = 1$.



\therefore option (a) is correct.

25. (b, c) Given: $f'(x) = e^{(f(x)-g(x))} \cdot g'(x) \quad \forall x \in R$

$$\Rightarrow e^{-f(x)} f'(x) = e^{-g(x)} g'(x)$$

Integrating both sides, we get

$$-e^{-f(x)} = -e^{-g(x)} + c \Rightarrow -e^{-f(x)} + e^{-g(x)} = c$$

$$\Rightarrow -e^{-f(1)} + e^{-g(1)} = -e^{-f(2)} + e^{-g(2)}$$


$$\therefore -e^{-1} + e^{-g(1)} = -e^{-f(2)} + e^{-1} \quad [\because f(1) = g(2) = 1]$$

$$\Rightarrow e^{-f(2)} + e^{-g(1)} = \frac{2}{e} \Rightarrow e^{-f(2)} < \frac{2}{e} \text{ and } e^{-g(1)} < \frac{2}{e}$$

$$\Rightarrow -f(2) < \ln 2 - 1 \text{ and } -g(1) < \ln 2 - 1$$

$$\Rightarrow f(2) > 1 - \ln 2 \text{ and } g(1) > 1 - \ln 2$$

$$26. \text{ (b, c) } f(x) = [x^2 - 3] = [x^2] - 3 = \begin{cases} -3, & -1/2 \leq x < 1 \\ -2, & 1 \leq x < \sqrt{2} \\ -1, & \sqrt{2} \leq x < \sqrt{3} \\ 0, & \sqrt{3} \leq x < 2 \\ 1, & x = 2 \end{cases}$$



Clearly, $f(x)$ is discontinuous at 4 points. Option (b) is correct.


$$\text{and } g(x) = |x|f(x) + |4x-7|f(x)$$

$$= (|x| + |4x-7|)f(x)$$

$$= (|x| + |4x-7|)[x^2 - 3]$$

$$= \begin{cases} (-x-4x-7)(-3), & -1/2 \leq x < 0 \\ (x-4x+7)(-3), & 0 \leq x < 1 \\ (x-4x+7)(-2), & 1 \leq x < \sqrt{2} \\ (x-4x+7)(-1), & \sqrt{2} \leq x < \sqrt{3} \\ (x-4x+7)(0), & \sqrt{3} \leq x < 7/4 \\ (x+4x-7)(0), & 7/4 \leq x < 2 \\ (x+4x-7)(1), & x = 2 \end{cases}$$

$$\therefore g(x) = \begin{cases} 15x+21, & -1/2 \leq x < 0 \\ 9x-21, & 0 \leq x < 1 \\ 6x-14, & 1 \leq x < \sqrt{2} \\ 3x-7, & \sqrt{2} \leq x < \sqrt{3} \\ 0, & \sqrt{3} \leq x < 2 \\ 5x-7, & x = 2 \end{cases}$$



Clearly, $g(x)$ is not differentiable at 4 points, when $x \in (-1/2, 2)$.

\therefore Option (c) is correct.

27. (a, b) $f(x) = a \cos(|x^3 - x|) + b |x| \sin(|x^3 + x|)$

(a) If $a = 0, b = 1$

$$\Rightarrow f(x) = |x| \sin |x^3 + x| \\ = x \sin(x^3 + x)$$

Which is differentiable every where.

- (b), (c) If $a = 1, b = 0 \Rightarrow f(x) = \cos(|x^3 - x|) = \cos(x^3 - x)$

Which is differentiable every where.

- (d) When $a = 1, b = 1, f(x) = \cos(x^3 - x) + x \sin(x^3 + x)$

Which is differentiable at $x = 1$

Hence only options (a) and (b) are the correct options.

28. (a, d) $f(x) = \begin{cases} \frac{x}{|x|} g(x), & x \neq 0 \\ 0, & x = 0 \end{cases} = \begin{cases} -g(x), & x < 0 \\ 0, & x = 0 \\ g(x), & x > 0 \end{cases}$

$$f'(x) = \begin{cases} -g'(x), & x < 0 \\ 0, & x = 0 \\ g'(x), & x > 0 \end{cases}$$

$$\therefore Lf'(0) = -g'(0) = 0 \text{ and } Rf'(0) = g'(0) = 0$$

$\therefore f$ is differentiable at $x = 0$

$$h(x) = e^{|x|} = \begin{cases} e^{-x}, & x < 0 \\ e^x, & x \geq 0 \end{cases}$$

$$h'(x) = \begin{cases} -e^{-x}, & x < 0 \\ e^x, & x \geq 0 \end{cases}$$

$$\therefore Lh'(0) = -1, Rh'(0) = 1$$

$\therefore h$ is not differentiable at $x = 0$

$$f \circ h(x) = f(h(x)) = f(e^{|x|})$$

$$= \begin{cases} g(e^{-x}) & \text{if } x < 0 \\ g(1) & \text{if } x = 0 \\ g(e^x) & \text{if } x > 0 \end{cases}$$

$$f'[h(x)] = \begin{cases} -g'(e^{-x}) \cdot e^{-x}, & x < 0 \\ 0, & x = 0 \\ g'(e^x) \cdot e^x, & x > 0 \end{cases}$$

$$Lf'(h(0)) = -g'(1) \cdot 1, Rf'(h(0)) = g'(1)$$

$$\therefore g'(1) \neq 0, \therefore Lf'(h(0)) \neq Rf'(h(0))$$

$\therefore f \circ h$ is not differentiable at $x = 0$.

$$hof(x) = \begin{cases} e^{|f(x)|}, & x \neq 0 \\ 1, & x = 0 \end{cases}$$

$$Lh'(f(0)) = \lim_{k \rightarrow 0} \frac{h(f(0)) - h(f(0-k))}{k}$$

$$= \lim_{k \rightarrow 0} \frac{1 - e^{|g(-k)|}}{k} = \lim_{k \rightarrow 0} \frac{1 - e^{|g(-k)|}}{|g(-k)|} \times \frac{|g(-k)|}{k}$$

$$= 1 \times 0 = 0 \quad \left(\because g'(0) = 0 \Rightarrow \lim_{k \rightarrow 0} \frac{g(-k)}{k} = \lim_{k \rightarrow 0} \frac{g(k)}{k} = 0 \right)$$

$$Rh'(f(0)) = \lim_{k \rightarrow 0} \frac{h(f(0+k)) - h(f(0))}{k}$$

$$= \lim_{k \rightarrow 0} \frac{e^{|g(k)|} - 1}{k} = \lim_{k \rightarrow 0} \frac{e^{|g(k)|} - 1}{|g(k)|} \times \frac{|g(k)|}{k} = 0$$

$$\therefore Lh'(f(0)) = Rh'(f(0)) = 0$$

$\therefore hof$ is differentiable at $x = 0$.

29. (a, b, c, d) At $x = -\frac{\pi}{2}$, $LHL = \lim_{x \rightarrow \frac{\pi}{2}^-} -x - \frac{\pi}{2} = 0$

and $RHL = \lim_{x \rightarrow \frac{\pi}{2}^+} -\cos x = 0$ and $f\left(-\frac{\pi}{2}\right) = 0$

$$\therefore LHL = RHL = f\left(-\frac{\pi}{2}\right)$$

$$\therefore f(x) \text{ is continuous at } x = -\frac{\pi}{2}$$

$$\text{At } x = 0, Lf'(0) = \sin 0 = 0 \text{ and } Rf'(0) = 1 - 0 = 1$$

$$\therefore Lf'(0) \neq Rf'(0)$$

$\therefore f$ is not differentiable at $x = 0$

$$\text{At } x = 1, Lf'(1) = Rf'(1)$$

$\therefore f$ is differentiable at $x = 1$.

$$\text{At } x = \frac{-3}{2}, f(x) = -\cos x, \text{ which is differentiable.}$$

Hence, all four options are correct.

30. (b, c) Given: $f(x+y) = f(x) + f(y) \quad \forall x, y \in \mathbb{R}$

\therefore On putting $x = y = 0$, we get $f(0) = 0$

$$\text{Also } f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$


$$= \lim_{h \rightarrow 0} \frac{f(h)}{h} = f'(0) = k \text{ (say)} \Rightarrow f(x) = kx + c$$

$$\text{But } f(0) = 0 \Rightarrow c = 0, \therefore f(x) = kx$$

Which is continuous $\forall x \in \mathbb{R}$.


Also $f'(x) = k$, a constant.

31. (a, d) From graph, $f(x)$ is continuous everywhere but not differentiable at $x = 1$ as there is sharp turns in the graph at $x = 1$.




32. (a, c, d) From the figure it is clear that

$$h(x) = \begin{cases} x, & \text{if } x \leq 0 \\ x^2, & \text{if } 0 < x < 1 \\ x, & \text{if } x \geq 1 \end{cases}$$



From the graph it is clear that h is continuous for all $x \in \mathbb{R}$, $h'(x) = 1$ for all $x > 1$ and h is not differentiable at $x = 0$ and 1 as there are sharp turns at $x = 0$ and 1 .

33. (a, c)



From graph it is clear that $f(x)$ is continuous everywhere and also differentiable everywhere except at $x = 1$ and -1 .

34. (a, b) $g(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$

$$\text{If } x \neq 0, g'(x) = x^2 \cos\left(\frac{1}{x}\right) \left(-\frac{1}{x^2}\right) + 2x \sin\left(\frac{1}{x}\right)$$

$$= -\cos\left(\frac{1}{x}\right) + 2x \sin\left(\frac{1}{x}\right), \text{ which exists for } \forall x \neq 0.$$

$$\text{If } x = 0, g'(0) = \lim_{x \rightarrow 0} \frac{g(x) - g(0)}{x - 0}$$

$$= \lim_{x \rightarrow 0} \frac{x^2 \sin(1/x) - 0}{x - 0} = \lim_{x \rightarrow 0} x \sin\left(\frac{1}{x}\right) = 0$$

$$\therefore g'(x) = \begin{cases} -\cos\left(\frac{1}{x}\right) + 2x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

At $x = 0$, $\cos\left(\frac{1}{x}\right)$ is not continuous, therefore $g'(x)$ is not continuous at $x = 0$.

$$\text{At } x = 0, Lf' = \lim_{x \rightarrow 0} \frac{0 - (-x) \sin\left(-\frac{1}{x}\right)}{x} = -\sin\left(\frac{1}{x}\right),$$


which does not exist.

$\therefore f$ is not differentiable at $x = 0$.

35. (b, c, d) $f(x) = \begin{cases} 0, & x < 0 \\ x^2, & x \geq 0 \end{cases} \therefore f'(x) = \begin{cases} 0, & x < 0 \\ 2x, & x \geq 0 \end{cases}$

Clearly $f'(x)$ may or may not exist at $x = 0$ but it exists $\forall x$ except at $x = 0$.

Now $Lf'(0) = 0 = Rf'(0) \Rightarrow f$ is differentiable at $x = 0$



Thus, $f(x)$ is differentiable for all values of x and hence it is continuous also for all values of x .

From graph of $f'(x)$, it is clear that $f'(x)$ is continuous but not differentiable at $x = 0$ as there is sharp turns at $x = 0$ in the graph.

36. (a, b, c) $f(x) = \begin{cases} |x-3|, & x \geq 1 \\ \frac{x^2}{4} - \frac{3x}{2} + \frac{13}{4}, & x < 1 \end{cases}$

$$= \begin{cases} \frac{x^2}{4} - \frac{3x}{2} + \frac{13}{4}, & x < 1 \\ 3 - x, & 1 \leq x < 3 \\ x - 3, & x \geq 3 \end{cases}$$

$$Lf'(1) = 1 \text{ and } Rf'(1) = -1.$$

$$\therefore Lf'(1) \neq Rf'(1)$$

Hence, f is differentiable at $x = 1$ and therefore continuous at $x = 1$.

$$\text{Now, } Lf'(3) = -1 \text{ and } Rf'(3) = 1$$

$$\therefore Lf'(3) \neq Rf'(3)$$

Hence, f is not differentiable at $x = 3$

$$\text{Now, L.H.L.} = \lim_{h \rightarrow 0} f(3-h) = \lim_{h \rightarrow 0} [3-(3-h)] = 0$$

$$\text{R.H.L.} = \lim_{h \rightarrow 0} f(3+h) = \lim_{h \rightarrow 0} [3+h-3] = 0$$

$$\text{and } f(3) = 0, \therefore \text{LHL} = \text{RHL} = f(3)$$

Hence, f is continuous at $x = 3$

$$37. \quad (a) \quad f(x) = \frac{x}{1+|x|} = \begin{cases} \frac{x}{1-x}, & x < 0 \\ \frac{x}{1+x}, & x \geq 0 \end{cases}$$

Clearly $f(x)$ may or may not be differentiable at $x = 0$ but $f(x)$ differentiable at each pair in $(-\infty, \infty)$ except at $x = 0$

$$Lf'(0) = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{\frac{-h}{1+h} - 0}{-h} = 1$$

$$Rf'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{\frac{h}{1+h} - 0}{h} = 1$$

$$\therefore Lf'(0) = Rf'(0) \Rightarrow f \text{ is differentiable at } x = 0$$

Thus, f is differentiable in $(-\infty, \infty)$.

$$38. \quad (a, b, d) \text{ If } -1 \leq x \leq 1, \text{ then } 0 \leq x \sin \pi x \leq 1/2$$


$$\therefore f(x) = [x \sin \pi x] = 0$$

$$\text{Also } f(x) = [x \sin \pi x] = -1, \text{ when } 1 < x < 1+h$$

Thus $f(x)$ is constant and equal to 0 in the closed interval $[-1, 1]$ and so $f(x)$ is continuous and differentiable in the open interval $(-1, 1)$.

At $x = 1$, $f(x)$ is clearly discontinuous, since $f(1-0) = 0$, $f(1+0) = -1$ and $f(x)$ is non-differentiable at $x = 1$.

$$39. \quad (b, d, e) \text{ Graph of } f(x) = 1 + |\sin x| \text{ is as follows:}$$



From graph it is clear that function is continuous every where but not differentiable at integral multiples of π because at these points curve has sharp turnings.

$$40. \quad (b) \quad f(x) = x(\sqrt{x} - \sqrt{x+1})$$

$$Rf'(0) = \lim_{h \rightarrow 0} \frac{(0-h)[\sqrt{0-h} - \sqrt{0-h+1}] - 0}{-h}$$

$$= \lim_{h \rightarrow 0} [\sqrt{-h} - \sqrt{-h+1}] = 0 - \sqrt{1} = -1$$

$$Lf'(0) = \lim_{h \rightarrow 0} \frac{(0+h)[\sqrt{0+h} - \sqrt{0+h+1}] - 0}{h}$$

$$= \lim_{h \rightarrow 0} [\sqrt{h} - \sqrt{h+1}] = -1$$

$$\text{Since } Lf'(0) = Rf'(0)$$

$\therefore f$ is differentiable at $x = 0$.

$$41. \quad (a, b, d) \text{ Given : } x + |y| = 2y$$

$$\text{If } y < 0 \text{ then } x - y = 2y$$

$$\Rightarrow y = x/3 \Rightarrow x < 0$$

$$\text{If } y = 0 \text{ then } x = 0. \text{ If } y > 0 \text{ then } x + y = 2y$$

$$\Rightarrow y = x \Rightarrow x > 0$$

$$\therefore f(x) = y = \begin{cases} x/3, & x < 0 \\ x, & x \geq 0 \end{cases}$$

Continuity at $x = 0$

$$\text{LHL} = \lim_{h \rightarrow 0} f(0-h) = \lim_{h \rightarrow 0} (-h/3) = 0$$

$$\text{RHL} = \lim_{h \rightarrow 0} f(0+h) = \lim_{h \rightarrow 0} h = 0$$

$$f(0) = 0$$

$$\therefore \text{LHL} = \text{RHL} = f(0)$$

$\therefore f(x)$ is continuous at $x = 0$

Differentiability at $x = 0$

$$Lf' = 1/3; Rf' = 1$$

As $Lf' \neq Rf' \Rightarrow f(x)$ is not differentiable at $x = 0$

$$\text{But for } x < 0, \frac{dy}{dx} = \frac{1}{3}$$

$$42. \quad (c) \text{ Given that } f(x) = \begin{cases} x|x| \sin \frac{1}{x} & ; x \neq 0 \\ 0 & ; x = 0 \end{cases}$$

$$\text{and } g(x) = \begin{cases} 1-2x & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$

$$\therefore g\left(\frac{1}{2}-x\right) = \begin{cases} 2x & ; 0 \leq \frac{1}{2}-x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$

$$= \begin{cases} 2x & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$

$$\text{Now, } g(x) + g\left(\frac{1}{2}-x\right) = \begin{cases} 1 & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$

$$(P) \text{ Let } a = 0, b = 1, c = 0, d = 0$$

$$\therefore h(x) = g(x) + g\left(\frac{1}{2}-x\right) = \begin{cases} 1 & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$

Hence range of $h(x)$ is $\{0, 1\}$

$$(Q) \text{ Let } a = 1, b = 0, c = 0, d = 0$$

$$h(x) = f(x) = \begin{cases} x|x| \sin \frac{1}{x} & ; x \neq 0 \\ 0 & ; x = 0 \end{cases}$$


$$\text{RHD} = \lim_{x \rightarrow 0} \frac{x^2 \sin \frac{1}{x} - 0}{x} = 0$$

$$\text{LHD} = \lim_{x \rightarrow 0} \frac{-x^2 \sin \frac{1}{x} - 0}{x} = 0$$

Hence $h(x)$ is differentiable on \mathbb{R}

$$(R) \text{ Let } a = 0, b = 0, c = 1, d = 0$$


$$h(x) = x - g(x) = \begin{cases} 3x-1 & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$



$\therefore h(x)$ is ONTO

(S) Let $a=0, b=0, c=0, d=1$

$$h(x) = g(x) = \begin{cases} 1-2x & ; 0 \leq x \leq \frac{1}{2} \\ 0 & ; \text{otherwise} \end{cases}$$



Range of $h(x)$ is $[0, 1]$

$$\begin{aligned} 43. \text{ (d) (i) } f'_1(0) &= \lim_{h \rightarrow 0} \left[\frac{\sin \sqrt{1-e^{-h^2}} - 0}{h} \right] \\ &= \lim_{h \rightarrow 0} \left[\frac{\sin \sqrt{1-e^{-h^2}}}{\sqrt{1-e^{-h^2}}} \times \frac{\sin \sqrt{1-e^{-h^2}}}{h^2} \times \frac{|h|}{h} \right] \\ &= \lim_{h \rightarrow 0} \left[1 \times 1 \times \frac{|h|}{h} \right] = \lim_{h \rightarrow 0} \frac{|h|}{h} \quad \left[\because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \end{aligned}$$

which does not exist.

\therefore for (P), (2) is correct.

$$\begin{aligned} \text{(ii) } \lim_{x \rightarrow 0} f_2(x) &= \lim_{x \rightarrow 0} \left[\frac{|\sin x|}{\tan^{-1} x} \right] \\ &= \lim_{x \rightarrow 0} \left[\frac{|\sin x|}{|x|} \times \frac{x}{\tan^{-1} x} \times \frac{|x|}{x} \right] \\ &= \lim_{x \rightarrow 0} \left[1 \times 1 \times \frac{|x|}{x} \right] = \lim_{x \rightarrow 0} \frac{|x|}{x} \quad \left[\because \lim_{x \rightarrow \infty} \frac{x}{\tan^{-1} x} = 1 \right] \end{aligned}$$

which does not exist, so for Q, (1) is correct.

$$\text{(iii) } \lim_{x \rightarrow 0} f_3(x) = \lim_{x \rightarrow 0} [\sin(\log_e(x+2))]$$

$$\text{if } x \rightarrow 0 \Rightarrow (x+2) \rightarrow 2 \Rightarrow \log_e(x+2) \rightarrow \log_e 2 < 1$$

$$\Rightarrow 0 < \lim_{x \rightarrow 0} \sin(\log_e(x+2)) < \sin 1$$

$$\Rightarrow \lim_{x \rightarrow 0} [\sin(\log_e(x+2))] = 0$$

$$f_3(x) = 0 \quad \forall x \in [-1, e^{\pi/2} - 2]$$

$$\Rightarrow f'_3(x) = 0 \quad \forall x \in (-1, e^{\pi/2} - 2)$$

$$\Rightarrow f''_3(x) = 0 \quad \forall x \in (-1, e^{\pi/2} - 2)$$

\therefore for (R), (4) is correct.

$$\text{(iv) } \lim_{x \rightarrow 0} f_4(x) = \lim_{x \rightarrow 0} \left(x^2 \sin \frac{1}{x} \right) = \lim_{x \rightarrow 0} x^2 \left(\sin \frac{1}{x} \right) = 0$$

$$f'_4(0) = \lim_{x \rightarrow 0} \frac{h^2 \sin\left(\frac{1}{h}\right) - 0}{h} = \lim_{x \rightarrow 0} h \sin\left(\frac{1}{h}\right) = 0$$


$$f'_4(x) = -\cos \frac{1}{x} + 2x \sin \frac{1}{x}, \quad x \neq 0$$

$$\lim_{x \rightarrow 0} f'_4(x) = \lim_{x \rightarrow 0} \left[-\cos \frac{1}{x} + 2x \sin \frac{1}{x} \right] = -\lim_{x \rightarrow 0} \cos \frac{1}{x}$$

which does not exist

So for (S), (3) is correct.

$$44. \text{ (d) } P(1): f_4(x) = \begin{cases} x^2, & x < 0 \\ e^{2x} - 1, & x \geq 0 \end{cases}$$




Range of $f_4 = [0, \infty)$

$\therefore f_4$ is onto.


From graph f_4 is not one one.

$$Q(3): f_3(x) = \begin{cases} \sin x, & x < 0 \\ x, & x \geq 0 \end{cases}$$




From graph f is differentiable but not one one.

$$R(2): f_2 \circ f_1(x) = \begin{cases} x^2, & x < 0 \\ e^{2x}, & x \geq 0 \end{cases}$$



From graph $f_2 \circ f_1$ is neither continuous nor one one.

$$S(4): f_2(x) = x^2, x \in [0, \infty)$$




It is continuous and one one.

45. $A \rightarrow (p, q, r); B \rightarrow (p, s); C \rightarrow (s, r); D \rightarrow (p, q)$

$$(A) y = x|x| = \begin{cases} -x^2, & \text{if } x < 0 \\ x^2, & \text{if } x \geq 0 \end{cases}$$

Graph is as follows :




From graph, $y = x|x|$ is continuous in $(-1, 1)$ (p)
differentiable in $(-1, 1)$ (q)
and strictly increasing in $(-1, 1)$. (r)

$$(B) y = \sqrt{|x|} = \begin{cases} \sqrt{-x}, & \text{if } x < 0 \\ \sqrt{x}, & \text{if } x \geq 0 \end{cases}$$

$$\Rightarrow y^2 = -x, x < 0 \text{ [where } y \text{ can take only +ve values]}$$

$$\text{and } y^2 = x, x \geq 0$$

\therefore Graph is as follows :




From graph $y = \sqrt{|x|}$ is continuous in $(-1, 1)$ (p)

and not differentiable at $x = 0$ (s)

$$(C) y = x + [x] = \begin{cases} x-1, & -1 \leq x < 0 \\ x, & 0 \leq x < 1 \\ x+1, & 1 \leq x < 2 \end{cases}$$

Graph of $y = x + [x]$ is as follows :




From graph, $y = x + [x]$ is neither continuous, nor differentiable at $x = 0$ and hence in $(-1, 1)$. (s)

Also it is strictly increasing in $(-1, 1)$ (r)

$$(D) y = |x-1| + |x+1| = \begin{cases} -2x, & x < -1 \\ 2, & -1 \leq x < 1 \\ 2x, & x \geq 1 \end{cases}$$

Graph of function is as follows :



From graph, $y = f(x)$ is continuous (p) and differentiable (q) in $(-1, 1)$ but not strictly increasing in $(-1, 1)$.

46. $A \rightarrow (p); B \rightarrow (r)$

$$(A) \sin(\pi[x]) = 0, \forall x \in \mathbb{R}$$


\therefore Differentiable everywhere.

$$\therefore (A) \rightarrow (p)$$

$$(B) \sin(\pi(x - [x])) = f(x)$$

$$\text{We know that } x - [x] = \begin{cases} x, & \text{if } 0 \leq x < 1 \\ x-1, & \text{if } 1 \leq x < 2 \\ x-2, & \text{if } 2 \leq x < 3 \end{cases}$$

It's graph is, as shown in figure, which is discontinuous at $\forall x \in \mathbb{Z}$.



Clearly $x - [x]$ and hence $\sin(\pi(x - [x]))$ is not differentiable $\forall x \in \mathbb{Z}$

(B) $\rightarrow r$

47. Given : $f(x-y) = f(x)g(y) - f(y)g(x)$
Put $y = x$ and we get $f(0) = 0$
put $y = 0$ and we get $g(0) = 1$

$$\text{R.H.D. of } f(x) : f'(0^+) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

($h \in \mathbb{R}^+$ and tending to zero)

$$= \lim_{h \rightarrow 0} \frac{f(0)g(-h) - g(0)f(-h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-f(-h)}{h} = f(0^-) = \text{L.H.D.} \quad \dots(i)$$

and L.H.D. of $f(x) : f'(0^-)$

$$= \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = f(0^+) = \text{R.H.D.} \quad \dots(ii)$$

Hence from (i) and (ii) we get $f'(0)$ is exist. So it is finite.

Put $y = x$ in given condition

$$g(x-y) = g(x)g(y) + f(x)f(y)$$

$$\Rightarrow g(0) = g^2(x) + f^2(x)$$

$$\Rightarrow g^2(x) + f^2(x) = 1 \Rightarrow g^2(x) = 1 - f^2(x)$$

On diff. w.r.t.x, we get

$$2g(x)g'(x) + 2f(x)f'(x) = 0 \Rightarrow g'(0) = 0$$

[Note : g is differentiable at zero because f is diff. at 0 and $g^2(x) = 1 - f^2(x)$]

48. Given : $f(x)$ is differentiable at $x = 0$.
 $\therefore f(x)$ will also be continuous at $x = 0$

$$\Rightarrow \lim_{h \rightarrow 0} f(0+h) = f(0) \Rightarrow \lim_{h \rightarrow 0} \frac{e^{\frac{ah}{2}} - 1}{h} = \frac{1}{2}$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{e^{\frac{ah}{2}} - 1}{\frac{ah}{2}} \times \frac{a}{2} = \frac{1}{2} \Rightarrow a = 1$$

Since $f(x)$ is differentiable at $x = 0$,

$$Lf'(0) = Rf'(0)$$

$$\therefore \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{b \sin^{-1}\left(\frac{c-h}{2}\right) - \frac{1}{2}}{-h} = \lim_{h \rightarrow 0} \frac{e^{\frac{h}{2}} - 1 - \frac{1}{2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2e^{\frac{h}{2}} - 2 - h}{2h^2} \quad \left[\frac{0}{0} \text{ form}\right]$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{\frac{b}{\sqrt{1 - \left(\frac{c-h}{2}\right)^2}} \cdot \left(-\frac{1}{2}\right)}{-1} \quad [\text{using LH rule}]$$

$$= \lim_{h \rightarrow 0} \frac{2e^{\frac{h}{2}} \cdot \frac{a}{2} - 1}{4h} = \lim_{h \rightarrow 0} \frac{e^{\frac{h}{2}} - 1}{8\left(\frac{h}{2}\right)} \quad [\because a=1]$$

$$\Rightarrow \frac{b}{\sqrt{1 - \frac{c^2}{4}}} = \frac{1}{8} \Rightarrow 64b^2 = 4 - c^2$$

49. Given that $f : [-2a, 2a] \rightarrow \mathbb{R}$ is an odd function.

$$\therefore f(-x) = -f(x)$$

Lf' at $x = a$ is 0.

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h} = 0$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{h} = 0 \quad \dots(i)$$

$Lf'(-a)$ is given by

$$\lim_{h \rightarrow 0} \frac{f(-a-h) - f(-a)}{-h} = \lim_{h \rightarrow 0} \frac{-f(a+h) + f(a)}{-h} \quad [\because f(-x) = -f(x)]$$

$$= \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \quad \dots(ii)$$

Now, for $x \in [a, 2a]$, $f(x) = f(2a-x)$

$$\therefore f(a+h) = f(2a-a-h) = f(a-h)$$

Substituting this values in equation (ii), we get

$$Lf'(-a) = \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{h} = 0 \quad [\text{using equation (i)}]$$

$$50. \quad g \circ f(x) = \begin{cases} f(x) + 1, & \text{if } f(x) < 0 \\ \{f(x) - 1\}^2 + b, & \text{if } f(x) \geq 0 \end{cases}$$

$$= \begin{cases} x + a + 1, & \text{if } x < -a \\ (x + a - 1)^2 + b, & \text{if } -a \leq x < 0 \\ (|x - 1| - 1)^2 + b, & \text{if } x \geq 0 \end{cases}$$

As $g \circ f(x)$ is continuous at $x = -a$

$$g \circ f(-a) = g \circ f(-a^+) = g \circ f(-a^-)$$

$$\Rightarrow 1 + b = 1 + b = 1 \Rightarrow b = 0$$

Also, $g \circ f(x)$ is continuous at $x = 0$

$$\Rightarrow g \circ f(0) = g \circ f(0^+) = g \circ f(0^-)$$

$$\Rightarrow b = b = (a-1)^2 + b \Rightarrow a = 1$$

$$\text{Hence, } g \circ f(x) = \begin{cases} x + 2 & \text{if } x < -1 \\ x^2, & \text{if } -1 \leq x < 0 \\ (|x - 1| - 1)^2, & \text{if } x \geq 0 \end{cases}$$

In the neighbourhood of $x = 0$, $g \circ f(x) = x^2$, which is differentiable at $x = 0$.

51. (I) Given : g is continuous at α and
 $f(x) - f(\alpha) = g(x)(x - \alpha), \forall x \in \mathbb{R}$
 \Rightarrow Since g is continuous at $x = \alpha$
and $g(x) = \frac{f(x) - f(\alpha)}{x - \alpha}$
 $\therefore \lim_{x \rightarrow \alpha} g(x) = g(\alpha)$

$$\Rightarrow \lim_{x \rightarrow \alpha} \frac{f(x) - f(\alpha)}{x - \alpha} = g(\alpha) \Rightarrow f'(\alpha) = g(\alpha)$$

(II) $f'(x)$ is differentiable at $x = \alpha$ (Given)

$$\therefore \lim_{x \rightarrow \alpha} \frac{f(x) - f(\alpha)}{x - \alpha} = f'(\alpha) \text{ exists and is finite.}$$

$$\text{Let us define, } g(x) = \begin{cases} \frac{f(x) - f(\alpha)}{x - \alpha}, & x \neq \alpha \\ f'(\alpha), & x = \alpha \end{cases}$$

Then, $f(x) - f(\alpha) = (x - \alpha)g(x)$, $\forall x \neq \alpha$.

Now for continuity of $g(x)$ at $x = \alpha$

$$\lim_{x \rightarrow \alpha} g(x) = \lim_{x \rightarrow \alpha} \frac{f(x) - f(\alpha)}{x - \alpha} = f'(\alpha) = g(\alpha)$$

$\therefore g$ is continuous at $x = \alpha$.

$$52. \text{ Given : } f(x) = \begin{cases} 1 - x, & x < 1 \\ (1 - x)(2 - x), & 1 \leq x \leq 2 \\ 3 - x, & x > 2 \end{cases}$$

It is clear that the function f is continuous and differentiable at all points except possibility at $x = 1$ and $x = 2$.

Continuity at $x = 1$:

$$\text{L.H.L.} = \lim_{h \rightarrow 0} [1 - (1 - h)] = \lim_{h \rightarrow 0} h = 0$$

$$\text{R.H.L.} = \lim_{h \rightarrow 0} [1 - (1 + h)][2 - (1 + h)] = 0$$

and $f(1) = 0$, \therefore L.H.L. = R.H.L. = $f(1) = 0$

Hence, f is continuous at $x = 1$

Differentiability at $x = 1$.

$$Lf'(1) = \lim_{h \rightarrow 0} \frac{f(1 - h) - f(1)}{-h} = \lim_{h \rightarrow 0} \frac{(1 - (1 - h)) - 0}{-h} = -1$$

$$\text{and } Rf'(1) = \lim_{h \rightarrow 0} \frac{f((1 + h) - f(1))}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\{1 - (1 - h)\} \{2 - (1 - h)\} - 0}{h} = \lim_{h \rightarrow 0} \frac{-h(1 - h)}{h} = -1$$

$$\therefore Lf'(1) = Rf'(1)$$

$\therefore f$ is differentiable at $x = 1$

Continuity at $x = 2$:

$$\text{L.H.L.} = \lim_{h \rightarrow 0} [1 - (2 - h)][2 - (2 - h)] = 0$$

$$\text{and R.H.L.} = \lim_{h \rightarrow 0} [3 - (2 + h)] = 1$$

\therefore L.H.L. \neq R.H.L., $\therefore f$ is not continuous at $x = 2$ and hence f cannot be differentiable at $x = 2$.

$\therefore f$ is continuous and differentiable at all points except at $x = 2$.

$$53. \text{ Given : } f\left(\frac{x+y}{2}\right) = \frac{f(x) + f(y)}{2} \quad \dots(i)$$

On putting $y = 0$ in (i), we get

$$f\left(\frac{x}{2}\right) = \frac{1}{2}[f(x) + 1] \quad [\because f(0) = 1]$$

$$\therefore f(x) = 2f\left(\frac{x}{2}\right) - 1 \quad \dots(ii)$$

$$\text{Now, } f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[\frac{f(2x) + f(2h)}{2} - f(x) \right], \quad [\text{using (i)}]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[\frac{(2f(x) - 1) + (2f(h) - 1)}{2} - f(x) \right], \quad [\text{using (ii)}]$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} [f(h) - 1] = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} \quad [\because f(0) = 1]$$

$$= f'(0) = -1 \quad [\because f'(0) = -1]$$

On integrating both sides w.r.t. x , we get

$f(x) = -x + c$. On putting $x = 0$, we get

$$f(0) = c = 1 \quad [\because f(0) = 0] \quad \therefore f(x) = 1 - x$$

$$\Rightarrow f(2) = 1 - 2 = -1$$

$$54. f(x+y) = f(x)f(y), \quad \forall x, y \in R$$

Hence, for $x = y = 0$, $f(0+0) = f(0)f(0)$

$$\Rightarrow f(0) = [f(0)]^2 \Rightarrow f(0) = 1 \quad [\because f(x) \neq 0, \text{ for any } x]$$

$$\text{Again } f'(0) = 2$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = 2 \Rightarrow \lim_{h \rightarrow 0} \frac{f(0)f(h) - f(0)}{h} = 2$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(0)[f(h) - 1]}{h} = 2$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(h) - 1}{h} = 2 \quad \dots(i) \quad [\because f(0) = 1]$$

$$\text{Now, } f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(x)f(h) - f(x)}{h} = \lim_{h \rightarrow 0} f(x) \left(\frac{f(h) - 1}{h} \right)$$

$$= f(x) \lim_{h \rightarrow 0} \left[\frac{f(h) - 1}{h} \right] \Rightarrow f'(x) = f(x) \cdot 2 \quad [\text{using eq. (i)}]$$

$$\text{Also, } \frac{f'(x)}{f(x)} = 2$$

On integrating both sides with respect to x , we get

$$\log |f(x)| = 2x + c$$


$$\text{At } x = 0, \log f(0) = c \Rightarrow c = \log 1 = 0$$

$$\Rightarrow \log |f(x)| = 2x \Rightarrow f(x) = e^{2x}$$

$$55. \text{ Given : } y = [x] + |1 - x|, \quad -1 \leq x \leq 3$$

$$\Rightarrow y = \begin{cases} -1 + 1 - x, & -1 \leq x < 0 \\ 0 + 1 - x, & 0 \leq x < 1 \\ 1 - 1 + x, & 1 \leq x < 2 \\ 2 - 1 + x, & 2 \leq x < 3 \\ 3 - 1 + x, & x = 3 \end{cases}$$

$$\Rightarrow y = \begin{cases} -x, & -1 \leq x < 0 \\ 1 - x, & 0 \leq x < 1 \\ x, & 1 \leq x < 2 \\ 1 + x, & 2 \leq x < 3 \\ 2 + x, & x = 3 \end{cases}$$



From graph we can say that given function is not differentiable at $x = 0, 1, 2, 3$.

56. Given : $f(x)$ is a function satisfying

$$f(-x) = f(x), \forall x \in R$$

Also $f'(0)$ exists

$$\Rightarrow f'(0) = Rf'(0) = Lf'(0)$$

$$\text{Now, } Rf'(0) = Lf'(0)$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h}$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{f(-h) - f(0)}{-h}$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{-h}$$

$$\Rightarrow 2 \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = 0$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = 0 \Rightarrow f'(0) = 0$$

57. Given $f(x) = \begin{cases} -1, & -2 \leq x \leq 0 \\ x-1, & 0 < x \leq 2 \end{cases}$

and $g(x) = f(|x| + |f(x)|)$

Here $g(x)$ involves $|x|$ and $|x-1|$ and $|-1| = 1$

Therefore, we should divide the given interval $[-2, 2]$ into the following intervals.

I_1	I_2	I_3
$[-2, 0)$	$[0, 1)$	$[1, 2]$
$x = -ve$	$+ve$	$+ve$
$ x = -x$	x	x
$f(x) = -1$	$x-1$	$x-1$
$f(x) = -1$	$= x-1$	$= x-1$
$ f(x) = -1 $	$ x-1 $	$ x-1 $
$= 1$	$= -(x-1)$	$= x-1$

\therefore Using above, we get

$$g(x) = f(|x| + |f(x)|)$$

$$\Rightarrow g(x) = \begin{cases} -1+1=0 & \text{in } I_1 \\ x-1-(x-1)=0 & \text{in } I_2 \\ x-1+x-1=2(x-1) & \text{in } I_3 \end{cases}$$

Hence, $g(x)$ is defined as follows :

$$g(x) = \begin{cases} 0, & -2 \leq x < 1 \\ 2(x-1), & 1 \leq x \leq 2 \end{cases}$$

$$Lg'(1) = 0; Rg'(1) = 2$$

$\therefore g(x)$ is not differentiable at $x = 1$.

58. Here, $f(x) = x^3 - x^2 + x + 1$

$$\Rightarrow f'(x) = 3x^2 - 2x + 1 \text{ which is positive } \forall x \in R$$

Hence, $f(x)$ is strictly increasing in $(0, 2)$

$$g(x) = \begin{cases} \max \{f(t)\}, & 0 \leq t \leq x, 0 \leq x \leq 1 \\ 3-x, & 1 < x \leq 2 \end{cases}$$

As $f(x)$ is increasing function

$$\text{So, } \max \{f(t)\}, 0 \leq t \leq x, 0 \leq x \leq 1 = f(x)$$

$$\therefore g(x) = \begin{cases} x^3 - x^2 + x + 1 & 0 \leq x \leq 1 \\ 3-x, & 1 < x \leq 2 \end{cases}$$

$$\text{Clearly } g(1) = g(1^-) = g(1^+) = 2$$

Hence, $g(x)$ is continuous for all $x \in [0, 2]$

$$\text{Also, } g'(x) = \begin{cases} 3x^2 - 2x + 1 & 0 < x < 1 \\ -1 & 1 < x < 2 \end{cases}$$

$$\text{At } x = 1, \text{ L.H.D.} = 2, \text{ but R.H.D.} = -1$$

Thus $g(x)$ is not differentiable of $x = 1$

59. Given : $f(x) = \begin{cases} \frac{x^2}{2}, & 0 \leq x < 1 \\ 2x^2 - 3x + \frac{3}{2}, & 1 \leq x \leq 2 \end{cases}$

Clearly $f(x)$ may or may not be continuous at $x = 1$ but it is continuous everywhere on $[0, 2]$ except at $x = 1$

$$\Rightarrow \text{At } x = 1, Lf' = \frac{2}{2} \times 1 = 1; Rf' = 4 \times 1 - 3 = 1$$

$\Rightarrow f$ is differentiable and hence continuous at $x = 1$

$\therefore f(x)$ is continuous on $[0, 2]$

$$f'(x) = \begin{cases} x, & 0 \leq x < 1 \\ 4x-3, & 1 \leq x \leq 2 \end{cases}$$

At $x = 1$,

$$\lim_{x \rightarrow 1^-} f'(x) = \lim_{h \rightarrow 0} f'(1-h) = \lim_{h \rightarrow 0} (1-h) = 1$$

$$\lim_{x \rightarrow 1^+} f'(x) = \lim_{h \rightarrow 0} f'(1+h) = \lim_{h \rightarrow 0} 4(1+h) - 3 = 1$$

$$f'(1) = 4 - 3 = 1$$

$$\text{Thus } \lim_{x \rightarrow 1^-} f'(x) = \lim_{x \rightarrow 1^+} f'(x) = f'(1)$$

$\therefore f'$ is continuous at $x = 1$

Hence, f' is continuous on $[0, 2]$

$$f''(x) = \begin{cases} 1, & 0 \leq x < 1 \\ 4, & 1 \leq x \leq 2 \end{cases}$$

Clearly $f''(x)$ is discontinuous at $x = 1$,

$\therefore f''(x)$ is discontinuous on $[0, 2]$.

60. Given : $f(x) = \begin{cases} \frac{x-1}{2x^2-7x+5}, & x \neq 1 \\ -\frac{1}{3}, & x = 1 \end{cases}$

$$\therefore f'(x)|_{x=1} = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$

$$= \lim_{h \rightarrow 0} \left[\frac{\frac{1+h-1}{2(1+h)^2-7(1+h)+5} + \frac{1}{3}}{h} \right] = \lim_{h \rightarrow 0} \frac{h}{2h^2-3h+\frac{1}{3}}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{2h-3} + \frac{1}{3}}{h} = \lim_{h \rightarrow 0} \frac{2}{3(2h-3)} = -2/9$$

Topic-3: Chain Rule of Differentiation, Differentiation of Explicit & Implicit Functions, Parametric & Composite Functions, Logarithmic & Exponential Functions, Inverse Functions, Differentiation by Trigonometric Substitution

1. (a) Given : $\log(x+y) = 2xy$
Clearly, when $x = 0$ then $y = 1$
On differentiating w.r.t. x , we get
- $$\frac{1}{x+y} \left[1 + \frac{dy}{dx} \right] = 2y + \frac{2xdy}{dx}$$
- $$\Rightarrow \frac{dy}{dx} = \frac{\frac{1}{x+y} - 2y}{2x - \frac{1}{x+y}} \Rightarrow y'(0) = \frac{1-2}{0-1} = 1$$
- [\because when $x = 0$, then $y = 1$]

2. (a) $y = (\sin x)^{\tan x} \Rightarrow \log y = \tan x \cdot \log \sin x$
On differentiating w.r.t. x , we get

$$\frac{1}{y} \frac{dy}{dx} = \sec^2 x \log \sin x + \tan x \cdot \frac{1}{\sin x} \cdot \cos x$$

$$\Rightarrow \frac{dy}{dx} = (\sin x)^{\tan x} [1 + \sec^2 x \log \sin x]$$

3. (a) $f(x) = e^{-x}$ is one such function.
Here $f(0) = 1, f'(0) = -1, f(x) > 0, \forall x$.
 $\therefore f''(x) > 0 \forall x$

4. (1) $f(\theta) = \sin \left(\tan^{-1} \left(\frac{\sin \theta}{\sqrt{\cos 2\theta}} \right) \right)$
- $$= \sin \left[\sin^{-1} \left(\frac{\sin \theta}{\sqrt{\sin^2 \theta + \cos 2\theta}} \right) \right] \left[\because \tan^{-1} \frac{x}{y} = \sin^{-1} \frac{x}{\sqrt{x^2 + y^2}} \right]$$
- $$= \sin \left[\sin^{-1} \left(\frac{\sin \theta}{\cos \theta} \right) \right] = \tan \theta$$
- $$\Rightarrow \frac{df(\theta)}{d \tan \theta} = 1.$$

5. (2) Given : $f(x) = x^3 + e^{x/2}$ and $g(x) = f^{-1}(x)$
therefore we should have $gof(x) = x$

$$\therefore g(f(x)) = x \Rightarrow g(x^3 + e^{x/2}) = x$$

On differentiating both sides w.r.t. x , we get

$$g'(x^3 + e^{x/2}) \cdot \left(3x^2 + e^{x/2} \cdot \frac{1}{2} \right) = 1$$

$$\Rightarrow g'(x^3 + e^{x/2}) = \frac{1}{3x^2 + e^{x/2} \cdot \frac{1}{2}}$$

For $x = 0$, we get $g'(1) = 2$

6. Given : $xe^{xy} = y + \sin^2 x$
Differentiating both sides w. r.to x , we get

$$e^{xy} \cdot 1 + xe^{xy} \left(y + x \frac{dy}{dx} \right) = \frac{dy}{dx} + 2 \sin x \cos x$$

$$\text{On putting } x = 0, \text{ we get } 1 + 0 = \frac{dy}{dx} + 0 \Rightarrow \frac{dy}{dx} = 1$$

7. $f(x) = |x - 2|$
 $\Rightarrow g(x) = f(f(x)) = |f(x) - 2|$ for $x > 20$
 $= ||x - 2| - 2| = |x - 2 - 2|$ for $x > 20$
 $= |x - 4| = x - 4$ for $x > 20$
 $\therefore g'(x) = 1 \Rightarrow g'(x) = -4$

8. Let $u = \sec^{-1} \left(\frac{1}{2x^2 - 1} \right) \Rightarrow u = \cos^{-1}(2x^2 - 1) = 2 \cos^{-1} x$

$$\text{and } v = \sqrt{1 - x^2}$$

$$\therefore \frac{du}{dx} = \frac{-2}{\sqrt{1 - x^2}} \text{ and } \frac{dv}{dx} = \frac{-x}{\sqrt{1 - x^2}}$$

$$\therefore \frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} = \frac{\frac{-2}{\sqrt{1 - x^2}}}{\frac{-x}{\sqrt{1 - x^2}}} = \frac{2}{x} \Rightarrow \frac{du}{dv} \Big|_{x=\frac{1}{2}} = 4$$

9. Given that, $f(x) = \log_x(\ln x) = \frac{\log_e(\log_e x)}{(\log_e x)}$

$$f'(x) = \frac{\frac{1}{\log_e x} \times \frac{1}{x} \times \log_e x - \frac{1}{x} \log_e(\log_e x)}{(\log_e x)^2}$$

$$= \frac{\frac{1}{x} [1 - \log_e(\log_e x)]}{(\log_e x)^2}$$

$$f'(e) = \frac{\frac{1}{e} [1 - \log_e(\log_e e)]}{(\log_e e)^2} = \frac{\frac{1}{e} [1 - \log_e 1]}{(1)^2} = \frac{1}{e} (1 - 0) = \frac{1}{e}$$

10. Given that $F(x) = \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix} \dots (i)$

where $f_r(x), g_r(x), h_r(x), r = 1, 2, 3$, are polynomials in x and hence differentiable and

$$f_r(a) = g_r(a) = h_r(a), r = 1, 2, 3 \dots (ii)$$

On differentiating equation (i) with respect to x , we get

$$F'(x) = \begin{vmatrix} f_1'(x) & f_2'(x) & f_3'(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix}$$

$$+ \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ g_1'(x) & g_2'(x) & g_3'(x) \\ h_1(x) & h_2(x) & h_3(x) \end{vmatrix} + \begin{vmatrix} f_1(x) & f_2(x) & f_3(x) \\ g_1(x) & g_2(x) & g_3(x) \\ h_1'(x) & h_2'(x) & h_3'(x) \end{vmatrix}$$

$$\Rightarrow F'(a) = \begin{vmatrix} f_1'(a) & f_2'(a) & f_3'(a) \\ g_1(a) & g_2(a) & g_3(a) \\ h_1(a) & h_2(a) & h_3(a) \end{vmatrix}$$

$$+ \begin{vmatrix} f_1(a) & f_2(a) & f_3(a) \\ g_1'(a) & g_2'(a) & g_3'(a) \\ h_1(a) & h_2(a) & h_3(a) \end{vmatrix} + \begin{vmatrix} f_1(a) & f_2(a) & f_3(a) \\ g_1(a) & g_2(a) & g_3(a) \\ h_1'(a) & h_2'(a) & h_3'(a) \end{vmatrix}$$

$$F'(a) = D_1 + D_2 + D_3$$

Using equation (ii) and the property of determinants that $D = 0$, if two rows in D are identical, we get $D_1 = D_2 = D_3 = 0$

$$\therefore F'(a) = 0.$$

11. Given: $y = f\left(\frac{2x-1}{x^2+1}\right)$; $f'(x) = \sin x^2$

$$\begin{aligned}\therefore \frac{dy}{dx} &= f'\left(\frac{2x-1}{x^2+1}\right) \cdot \frac{d}{dx}\left(\frac{2x-1}{x^2+1}\right) \\ &= \left[\sin\left(\frac{2x-1}{x^2+1}\right)\right] \cdot \left[\frac{2(x^2+1) - 2x(2x-1)}{(x^2+1)^2}\right] \\ &= \frac{2+2x-2x^2}{(x^2+1)^2} \sin\left(\frac{2x-1}{x^2+1}\right)\end{aligned}$$

12. (True) Consider $g(x) = \frac{f(x) + f(-x)}{2}$, which is an even function

$$\therefore g'(x) = \frac{f'(x) - f'(-x)}{2} = h(x), \text{ let}$$

$$\text{Now } h(-x) = \frac{f'(-x) - f'(x)}{2} = -h(x),$$

$\therefore h$ is an odd function.

Hence, derivative of an even function is an odd function.

13. (d) $f_n(x) = \sum_{j=1}^n \tan^{-1}\left(\frac{1}{1+(x+j)(x+j-1)}\right)$

$$\begin{aligned}&= \sum_{j=1}^n \tan^{-1}\left[\frac{(x+j) - (x+j-1)}{1+(x+j)(x+j-1)}\right] \\ &= \sum_{j=1}^n [\tan^{-1}(x+j) - \tan^{-1}(x+j-1)] \\ &\Rightarrow f_n(x) = \tan^{-1}(x+n) - \tan^{-1}(x) \\ &= \tan^{-1}\left(\frac{n}{1+x(n+x)}\right) \Rightarrow f'_n(x) = \frac{1}{1+(x+n)^2} - \frac{1}{1+x^2}\end{aligned}$$

$$\text{and } f'_n(0) = \tan^{-1}(n), \therefore \tan^2(\tan^{-1}n) = n^2$$

Here $x=0$ is not in the given domain, i.e., $x \in (0, \infty)$.

\therefore Options (a) & (b) are not correct options.

(c) $\lim_{x \rightarrow \infty} \tan(f'_n(x)) = \lim_{x \rightarrow \infty} \left(\frac{n}{1+x(n+x)}\right) = 0$

(d) $\lim_{x \rightarrow \infty} \sec^2(f'_n(x)) = \lim_{n \rightarrow \infty} 1 + \tan^2(f'_n(x))$
 $= 1 + \lim_{x \rightarrow \infty} \tan^2(f'_n(x)) = 1$

14. (a, b, d)

(a) $f(x)$ being twice differentiable, it is continuous but can't be constant throughout the domain.

Hence we can find $x \in (r, s)$ such that $f(x)$ is one one.

\therefore (a) is true.

(b) By Lagrange's Mean Value theorem for $f(x)$ in $[-4, 0]$, there exists

$$x_0 \in (-4, 0) \text{ such that } f'(x_0) = \frac{f(0) - f(-4)}{0 - (-4)}$$

$$\Rightarrow |f'(x_0)| = \left| \frac{f(0) - f(-4)}{4} \right|$$

$$\therefore -2 \leq f(x) \leq 2, \therefore -4 \leq f(0) - f(-4) \leq 4$$

$$\Rightarrow |f'(x_0)| \leq 1, \therefore \text{(b) is true.}$$

(c) If we consider $f(x) = \sin(\sqrt{85}x)$ then $f(x)$ satisfies the given condition $[f(0)]^2 + [f'(0)]^2 = 1$

But $\lim_{x \rightarrow \infty} (\sin \sqrt{85}x)$ does not exist

\therefore (c) is false.

(d) Let us consider $g(x) = [f(x)]^2 + [f'(x)]^2$

By Lagrange's Mean Value theorem $|f'(x)| \leq 1$

Also $|f(x_1)| \leq 2$ as $f(x) \in [-2, 2]$

$\therefore g(x_1) \leq 5$, for $x_1 \in (-4, 0)$

Similarly $g(x_2) \leq 5$, for $x_2 \in (0, 4)$

Also $g(0) = 85$

Hence $g(x)$ has maxima in (x_1, x_2) say at α such that

$$g'(\alpha) = 0 \text{ and } g(\alpha) \geq 85$$

$$g'(\alpha) = 0 \Rightarrow 2f(\alpha)f'(\alpha) + 2f'(\alpha)f''(\alpha) = 0$$

$$\Rightarrow 2f'(\alpha)[f(\alpha) + f''(\alpha)] = 0$$

$$\text{If } f'(\alpha) = 0 \Rightarrow g(\alpha) = [f(\alpha)]^2 \text{ and } [f(\alpha)]^2 \leq 4$$

$$\therefore g(\alpha) \geq 85 \text{ (is not possible.)}$$

$$\Rightarrow f(\alpha) + f''(\alpha) = 0 \text{ for } \alpha \in (x_1, x_2) \in (-4, 4)$$

Hence, (d) is true.

15. (b, c) Given: $f(x) = x^3 + 3x + 2 \Rightarrow f'(x) = 3x^2 + 3$

$$\therefore f(0) = 2, f(1) = 6, f(2) = 16, f(3) = 38, f(6) = 236$$

Also given $g(f(x)) = x \Rightarrow g(2) = 0, g(6) = 1, g(16) = 2,$

$$g(3, 8) = 3, g(236) = 6$$

(a) $g(f(x)) = x \Rightarrow g'(f(x)) \cdot f'(x) = 1$

$$\text{For } g'(2), f(x) = 2 \Rightarrow x = 0$$

On putting $x = 0$, we get $g'(f(0)) \cdot f'(0) = 1$

$$\Rightarrow g'(2) = \frac{1}{3}$$

(b) $h(g(g(x))) = x \Rightarrow h'(g(g(x))) \cdot g'(g(x)) \cdot g'(x) = 1$

For $h'(1)$, we need $g(g(x)) = 1$

$$\Rightarrow g(x) = 6 \Rightarrow x = 236$$

On putting $x = 236$, we get

$$h'[g(g(236))] = \frac{1}{g'(g(236)) \cdot g'(236)}$$

$$\Rightarrow h'(g(6)) = \frac{1}{g'(6) \cdot g'(236)}$$

$$\Rightarrow h'(1) = \frac{1}{g'(f(1)) \cdot g'(f(6))}$$

$$= f'(1) \cdot f'(6) = 6 \times 111 = 666$$

(c) $h[g(g(x))] = x$

For $h(0)$, $g(g(x)) = 0 \Rightarrow g(x) = 2 \Rightarrow x = 16$

On putting $x = 16$, we get

$$h(g(g(16))) = 16 \Rightarrow h(0) = 16$$

(d) $h[g(g(x))] = x$

For $h(g(3))$, we need $g(x) = 3 \Rightarrow x = 38$

On putting $x = 38$, we get

$$h[g(g(38))] = 38 \Rightarrow h(g(3)) = 38$$

16. (b) Given that $f(x) = 2 + \cos x$ which is continuous and differentiable every where.

$$\text{Also } f'(x) = -\sin x, \therefore f'(x) = 0 \Rightarrow x = n\pi$$

\Rightarrow There exists $c \in [t, t + \pi]$ for $t \in \mathbb{R}$ such that

$$f'(c) = 0$$

\therefore Statement-1 is true.

Also $f(x)$ being periodic of period 2π , statement-2 is true, but statement-2 is not a correct explanation of statement-1.

$$\begin{aligned}
 17. \quad y &= \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1 \\
 &= \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{x}{x-c} \\
 &= \frac{ax^2}{(x-a)(x-b)(x-c)} + \left(\frac{b}{x-b} + 1\right) \frac{x}{x-c} \\
 &= \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{x^2}{(x-b)(x-c)} \\
 &= \left(\frac{a}{x-a} + 1\right) \frac{x^2}{(x-b)(x-c)} = \frac{x^3}{(x-a)(x-b)(x-c)}
 \end{aligned}$$

Taking log on both sides, we get

$$\log y = 3 \log x - \log(x-a) - \log(x-b) - \log(x-c)$$

$$\begin{aligned}
 \Rightarrow \frac{y'}{y} &= \frac{3}{x} - \frac{1}{x-a} - \frac{1}{x-b} - \frac{1}{x-c} \\
 &= \left(\frac{1}{x} - \frac{1}{x-a}\right) + \left(\frac{1}{x} - \frac{1}{x-b}\right) + \left(\frac{1}{x} - \frac{1}{x-c}\right) \\
 &= \frac{a}{x(a-x)} + \frac{b}{x(b-x)} + \frac{c}{x(c-x)} \\
 &= \frac{1}{x} \left[\frac{a}{a-x} + \frac{b}{b-x} + \frac{c}{c-x} \right]
 \end{aligned}$$

$$18. \quad (\sin y)^{\sin\left(\frac{\pi x}{2}\right)} + \frac{\sqrt{3}}{2} \sec^{-1}(2x) + 2^x \tan[\ln(x+2)] = 0 \quad \dots (i)$$

Put $x = -1$, in equation (i), we get

$$(\sin y)^{\sin\left(-\frac{\pi}{2}\right)} + \frac{\sqrt{3}}{2} \sec^{-1}(-2) + 2^{-1} \tan[\ln(-1+2)] = 0$$

$$\Rightarrow (\sin y)^{-1} + \frac{\sqrt{3}}{2} \left(\frac{2\pi}{3}\right) + \frac{1}{2} \tan 0 = 0$$

$$\Rightarrow \sin y = -\frac{\sqrt{3}}{\pi}, \text{ when } x = -1 \quad \dots (ii)$$

$$\text{Let } u = (\sin y)^{\sin\left(\frac{\pi x}{2}\right)}$$

Taking ln on both sides, we get

$$\ln u = \sin\left(\frac{\pi x}{2}\right) \ln \sin y$$

Differentiating both sides with respect to x , we get

$$\frac{1}{u} \frac{du}{dx} = \frac{\pi}{2} \cos\left(\frac{\pi x}{2}\right) \ln \sin y + \cot y \sin\left(\frac{\pi x}{2}\right) \frac{dy}{dx}$$

$$\begin{aligned}
 \Rightarrow \frac{du}{dx} &= (\sin y)^{\sin\left(\frac{\pi x}{2}\right)} \\
 &\times \left[\frac{\pi}{2} \cos\left(\frac{\pi x}{2}\right) \ln \sin y + \sin\left(\frac{\pi x}{2}\right) \cot y \frac{dy}{dx} \right] \quad \dots (iii)
 \end{aligned}$$

Now differentiating equation (i), we get

$$\frac{d}{dx} \left[(\sin y)^{\sin\left(\frac{\pi x}{2}\right)} \right] + \frac{\sqrt{3}}{2} \frac{1}{2x\sqrt{4x^2-1}} \cdot 2$$

$$\begin{aligned}
 &+ 2^x (\ln 2) \tan[\ln(x+2)] \\
 &+ 2^x \sec^2[\ln(x+2)] \frac{1}{x+2} = 0
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow (\sin y)^{\sin\left(\frac{\pi x}{2}\right)} &\left[\frac{\pi}{2} \cos\left(\frac{\pi x}{2}\right) \ln \sin y \right. \\
 &\left. + \sin\left(\frac{\pi x}{2}\right) \cot y \frac{dy}{dx} \right]
 \end{aligned}$$

$$\begin{aligned}
 &+ \frac{\sqrt{3}}{2x\sqrt{4x^2-1}} + 2^x \ln 2 \tan(\ln(x+2)) \\
 &+ \frac{2^x \sec^2[\ln(x+2)]}{x+2} = 0 \quad \text{[using (iii)]}
 \end{aligned}$$

$$\text{At } x = -1 \text{ and } \sin y = -\frac{\sqrt{3}}{\pi}, \text{ we get}$$

$$\left(-\frac{\sqrt{3}}{\pi}\right)^{-1} \left[0 - (-1) \sqrt{\frac{\pi^2}{3}} - 1 \left(\frac{dy}{dx}\right)_{x=-1} \right]$$

$$+ \frac{\sqrt{3}}{-2\sqrt{3}} + 0 + 2^{-1} = 0$$

$$\Rightarrow -\frac{\pi}{\sqrt{3}\sqrt{3}} \sqrt{\pi^2-3} \left(\frac{dy}{dx}\right)_{x=-1} - \frac{1}{2} + \frac{1}{2} = 0$$

$$\Rightarrow \frac{dy}{dx} \Big|_{x=-1} = 0$$

$$19. \quad \text{Given : } x = \sec \theta - \cos \theta, y = \sec^n \theta - \cos^n \theta$$

$$\Rightarrow \frac{dx}{d\theta} = \sec \theta \tan \theta + \sin \theta$$

$$= \sec \theta \tan \theta + \tan \theta \cos \theta = \tan \theta (\sec \theta + \cos \theta)$$

$$\text{and } \frac{dy}{d\theta} = n \sec^{n-1} \theta \sec \theta \tan \theta - n \cos^{n-1} \theta (-\sin \theta)$$

$$= n \tan \theta (\sec^n \theta + \cos^n \theta)$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{n \tan \theta (\sec^n \theta + \cos^n \theta)}{\tan \theta (\sec \theta + \cos \theta)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{n(\sec^n \theta + \cos^n \theta)}{(\sec \theta + \cos \theta)} \quad \dots (i)$$

$$\begin{aligned}
 \text{Now } x^2 + 4 &= (\sec \theta - \cos \theta)^2 + 4 \\
 &= \sec^2 \theta + \cos^2 \theta - 2 \sec \theta \cos \theta + 4 \\
 &= \sec^2 \theta + \cos^2 \theta + 2 = (\sec \theta + \cos \theta)^2 \quad \dots (ii)
 \end{aligned}$$

$$\begin{aligned}
 \text{and } y^2 + 4 &= (\sec^n \theta - \cos^n \theta)^2 + 4 \\
 &= \sec^{2n} \theta + \cos^{2n} \theta - 2 \sec^n \theta \cos^n \theta + 4 \\
 &= \sec^{2n} \theta + \cos^{2n} \theta + 2 = (\sec^n \theta + \cos^n \theta)^2 \quad \dots (iii) \\
 &= n^2 (y^2 + 4)
 \end{aligned}$$

$$\begin{aligned}
 \text{Now } (x^2 + 4) \left(\frac{dy}{dx}\right)^2 &= (\sec \theta + \cos \theta)^2 \cdot \frac{n^2 (\sec^n \theta + \cos^n \theta)^2}{(\sec \theta + \cos \theta)^2} \\
 &= n^2 (\sec^n \theta + \cos^n \theta)^2 \quad \text{[using (i) and (ii)]}
 \end{aligned}$$

$$\Rightarrow (x^2 + 4) \left(\frac{dy}{dx}\right)^2 = n^2 (y^2 + 4) \quad \text{[From (iii)]}$$

20. Let $F(x) = \begin{vmatrix} A(x) & B(x) & C(x) \\ A(\alpha) & B(\alpha) & C(\alpha) \\ A'(x) & B'(x) & C'(x) \end{vmatrix}$ (i)

Since α is a repeated root of quadratic equation $f(x) = 0$

\therefore We must have $f(x) = k(x - \alpha)^2$; where k is a non-zero real number.

Put $x = \alpha$ on both sides of equation (i); we get

$$F(\alpha) = \begin{vmatrix} A(\alpha) & B(\alpha) & C(\alpha) \\ A(\alpha) & B(\alpha) & C(\alpha) \\ A'(\alpha) & B'(\alpha) & C'(\alpha) \end{vmatrix} = 0$$

[$\because R_1$ and R_2 are identical]

$\therefore (x - \alpha)$ is a factor of $F(x)$

Differentiating equation (i) w.r. to x , we get

$$F'(x) = \begin{vmatrix} A'(x) & B'(x) & C'(x) \\ A(\alpha) & B(\alpha) & C(\alpha) \\ A'(x) & B'(x) & C'(x) \end{vmatrix}$$

Putting $x = \alpha$, we get

$$F'(\alpha) = \begin{vmatrix} A'(\alpha) & B'(\alpha) & C'(\alpha) \\ A(\alpha) & B(\alpha) & C(\alpha) \\ A'(\alpha) & B'(\alpha) & C'(\alpha) \end{vmatrix} = 0$$

[$\because R_1$ and R_3 are identical]

$\Rightarrow (x - \alpha)$ is a factor of $F'(x)$.

$\Rightarrow (x - \alpha)^2$ is a factor of $F(x)$.

$\therefore F(x)$ is divisible by $f(x)$.

21. Given: $y = e^{x \sin x^3} + (\tan x)^x$

Here y is the sum of two functions and in the second function base as well as power are functions of x . Therefore, here we will use logarithmic differentiation.

Let $y = u + v$, where $u = e^{x \sin x^3}$ and $v = (\tan x)^x$

$$\therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots (i)$$

$$\text{Now, } \frac{du}{dx} = e^{x \sin x^3} \cdot \frac{d}{dx}(x \sin x^3)$$

$$= e^{x \sin x^3} \cdot [3x^3 \cdot \cos x^3 + \sin x^3]$$

$$\text{Now } v = (\tan x)^x \Rightarrow \log v = x \log \tan x$$

Now, differentiating the both sides with respect to x , then

$$\frac{1}{v} \frac{dv}{dx} = x \cdot \frac{1}{\tan x} \cdot \sec^2 x + 1 \cdot \log \tan x$$

$$\therefore \frac{dv}{dx} = (\tan x)^x \left[\frac{2x}{\sin 2x} + \log \tan x \right]$$

Now substituting the value of $\frac{du}{dx}$ and $\frac{dv}{dx}$ in (i), we get

$$\frac{dy}{dx} = e^{x \sin x^3} [\sin x^3 + 3x^3 \cos x^3]$$

$$+ (\tan x)^x \left[\frac{2x}{\sin 2x} + \log \tan x \right]$$

22. Given: $y = \frac{5x}{3|1-x|} + \cos^2(2x+1)$

[Clearly y is not defined at $x = 1$]

$$\Rightarrow y = \begin{cases} \frac{5x}{3(1-x)} + \cos^2(2x+1), & x < 1 \\ \frac{5x}{3(x-1)} + \cos^2(2x+1), & x > 1 \end{cases}$$

$$\Rightarrow \frac{dy}{dx} = \begin{cases} \frac{5}{3} \left(\frac{(1-x) - x(-1)}{(1-x)^2} \right) - 2 \sin(4x+2), & x < 1 \\ \frac{5}{3} \left(\frac{(x-1) - x}{(x-1)^2} \right) - 2 \sin(4x+2), & x > 1 \end{cases}$$

$$\text{or } \frac{dy}{dx} = \begin{cases} \frac{5}{3} \frac{1}{(1-x)^2} - 2 \sin(4x+2), & x < 1 \\ -\frac{5}{3} \frac{1}{(x-1)^2} - 2 \sin(4x+2), & x > 1 \end{cases}$$

Topic-4: Differentiation of Infinite Series, Successive Differentiation, n th Derivative of Some Standard Functions, Leibnitz's Theorem, Rolle's Theorem, Lagrange's Mean Value Theorem

1. (a) Given: $g(x) = \log f(x) \Rightarrow g(x+1) = \log f(x+1)$

$$\Rightarrow g(x+1) = \log x f(x) \quad [\because f(x+1) = x f(x)]$$

$$\Rightarrow g(x+1) = \log x + \log f(x) \Rightarrow g(x+1) - g(x) = \log(x)$$

$$\Rightarrow g'(x+1) - g'(x) = \frac{1}{x} \Rightarrow g''(x+1) - g''(x) = -\frac{1}{x^2}$$

On putting, $x = x - \frac{1}{2}$, we get

$$\Rightarrow g''\left(x + \frac{1}{2}\right) - g''\left(x - \frac{1}{2}\right) = -\frac{1}{\left(x - \frac{1}{2}\right)^2} = \frac{-2^2}{(2x-1)^2}$$

On putting $x = 1, 2, 3, \dots, N$; we get

$$g''\left(\frac{3}{2}\right) - g''\left(\frac{1}{2}\right) = -\frac{2^2}{1^2} \quad \dots (i)$$

$$g''\left(\frac{5}{2}\right) - g''\left(\frac{3}{2}\right) = \frac{-2^2}{3^2} \quad \dots (ii)$$

$$g''\left(\frac{7}{2}\right) - g''\left(\frac{5}{2}\right) = \frac{-2^2}{5^2} \quad \dots (iii)$$

$$g''\left(N + \frac{1}{2}\right) - g''\left(N - \frac{1}{2}\right) = -\frac{2^2}{(2N-1)^2} \quad \dots (N)$$

On adding all the above equations, we get

$$g''\left(N + \frac{1}{2}\right) - g''\left(\frac{1}{2}\right) = -4 \left[1 + \frac{1}{3^2} + \frac{1}{5^2} + \dots + \frac{1}{(2N-1)^2} \right]$$

$$2. \quad (d) \quad \frac{d^2x}{d^2y} = \frac{d}{dy} \left(\frac{dx}{dy} \right) = \frac{d}{dx} \left(\frac{dx}{dy} \right) \times \frac{dx}{dy}$$

$$= \left\{ \frac{d}{dx} \left[\frac{1}{\left(\frac{dy}{dx} \right)} \right] \right\} \times \frac{1}{\frac{dy}{dx}} = -\frac{1}{\left(\frac{dy}{dx} \right)^2} \times \frac{d^2y}{dx^2} \times \frac{1}{\left(\frac{dy}{dx} \right)}$$

$$= -\left(\frac{dy}{dx} \right)^{-3} \frac{d^2y}{dx^2}$$

$$3. \quad (d) \quad \text{Let us consider the function } g(x) = f(x) - x^2$$

$$\text{such that } g(1) = f(1) - 1^2 = 1 - 1 = 0$$

$$g(2) = f(2) - 2^2 = 4 - 4 = 0$$

$$g(3) = f(3) - 3^2 = 9 - 9 = 0$$

Since $f(x)$ is twice differentiable, therefore we can say $g(x)$ is continuous and differentiable everywhere and

$$g(1) = g(2) = g(3) = 0$$

\therefore By Rolle's theorem, $g'(c) = 0$ for some $c \in (1, 2)$ and

$g'(d) = 0$ for some $d \in (2, 3)$

Again by Rolle's theorem,

$$g''(e) = 0 \text{ for some } e \in (c, d) \Rightarrow e \in (1, 3)$$

$$\Rightarrow f''(e) - 2 = 0 \text{ or } f''(e) = 2 \text{ for some } x \in (1, 3)$$

$$\therefore f''(x) = 2 \text{ for some } x \in (1, 3)$$

$$4. \quad (b) \quad x^2 + y^2 = 1 \Rightarrow 2x + 2yy' = 0 \Rightarrow x + yy' = 0$$

$$\Rightarrow 1 + yy'' + (y')^2 = 0 \Rightarrow yy'' + (y')^2 + 1 = 0$$

$$5. \quad (b) \quad \text{Let } f(x) = ax^2 + bx + c \text{ and } f(x) > 0, \forall x \in \mathbb{R}$$

$$\therefore a > 0 \text{ and } D < 0$$

$$\Rightarrow a > 0 \text{ and } b^2 - 4ac < 0 \quad \dots (i)$$

$$\begin{aligned} \text{Now, } g(x) &= f(x) + f'(x) + f''(x) \\ &= ax^2 + bx + c + 2ax + b + 2a \\ &= ax^2 + (2a + b)x + (2a + b + c) \end{aligned}$$

$$\begin{aligned} \text{Here, } D &= (2a + b)^2 - 4a(2a + b + c) \\ &= 4a^2 + b^2 + 4ab - 8a^2 - 4ab - 4ac \\ &= b^2 - 4a^2 - 4ac = -4a^2 + b^2 - 4ac \\ &= (-ve) + (-ve) = -ve \quad [\text{using (i)}] \end{aligned}$$

Also from (i), $a > 0$

$$\therefore g(x) > 0, \forall x \in \mathbb{R}$$

$$6. \quad (c) \quad \text{Given : } y^2 = P(x), \text{ where } P(x) \text{ is a polynomial of degree 3 and hence thrice differentiable.}$$

$$\therefore 2y \frac{dy}{dx} = P'(x) \quad \dots (i)$$

Again differentiating with respect to x , we get

$$2 \left(\frac{dy}{dx} \right)^2 + 2y \frac{d^2y}{dx^2} = P''(x)$$

$$\Rightarrow \frac{[P'(x)]^2}{2y^2} + 2y \frac{d^2y}{dx^2} = P''(x) \quad [\text{using (i)}]$$

$$\Rightarrow 4y^3 \frac{d^2y}{dx^2} = 2y^2 P''(x) - [P'(x)]^2$$

$$\Rightarrow 4y^3 \frac{d^2y}{dx^2} = 2P(x)P''(x) - [P'(x)]^2 \quad [\because y^2 = P(x)]$$

$$\Rightarrow 2y^3 \frac{d^2y}{dx^2} = P(x)P''(x) - \frac{1}{2}[P'(x)]^2$$

$$\text{Again on differentiating w.r. to } x, \text{ we get } 2 \frac{d}{dx} \left(y^3 \frac{d^2y}{dx^2} \right)$$

$$= P'''(x)P(x) + P''(x)P'(x) - P'(x)P''(x) = P'''(x)P(x)$$

$$7. \quad (5.00) \quad g(x) = (x^2 - 1)^2 h(x); h(x) = a_0 + a_1x + a_2x^2 + a_3x^3$$

$$\therefore f(1) = f(-1) = 0$$

$$\Rightarrow f(x) \text{ has two roots } x = 1 \text{ and } x = -1$$

$$\Rightarrow g(x) \text{ has atleast 3 roots } x = 1, x = -1 \text{ and } x = \alpha$$

Then by Rolle's theorem

$$\Rightarrow g'(\alpha) = 0, \alpha \in (-1, 1)$$

$$g'(1) = g'(-1) = 0 \Rightarrow g'(x) = 0 \text{ has atleast 3 root,}$$

$$\Rightarrow g''(x) = 0 \text{ will have at least 2 root, say } \beta, \gamma \text{ such that}$$

Then by Rolle's theorem

$$-1 < \beta < \alpha < \gamma < 1$$


$$\text{So, } \min(m_{f''}) = 2 \text{ and we find } (m_f + m_{f''}) = 5$$

$$8. \quad (a, b, c) \quad \text{Given, } S = \text{Set of twice differentiable functions } f: \mathbb{R} \rightarrow \mathbb{R}$$

$$\frac{d^2f}{dx^2} > 0 \text{ in } (-1, 1)$$

Graph 'f' is Concave upward.

Number of solutions of $f(x) = x \rightarrow x_f$



\Rightarrow Graph of $y = f(x)$ can intersect graph of $y = x$ at atmost two points $\Rightarrow 0 \leq x_f \leq 2$

$$9. \quad (b, c, d) \quad \lim_{t \rightarrow x} \frac{f(x) \sin t - f(t) \sin x}{t - x} = \sin^2 x$$

$$\Rightarrow \lim_{t \rightarrow x} \frac{f(x) \cos t - f'(t) \sin x}{1} = \sin^2 x \quad (\text{using LH Rule})$$

$$\Rightarrow f(x) \cos x - f'(x) \sin x = \sin^2 x$$

$$\Rightarrow -\left(\frac{f'(x) \sin x - f(x) \cos x}{\sin^2 x}\right) = 1$$

$$\Rightarrow -d\left(\frac{f(x)}{\sin x}\right) = 1 \Rightarrow \frac{f(x)}{\sin x} = -x + c$$

$$\text{Put } x = \frac{\pi}{6}$$

$$\therefore \frac{\frac{12}{1}}{\frac{1}{2}} = -\frac{\pi}{6} + c \quad \left[\because f\left(\frac{\pi}{6}\right) = -\frac{\pi}{12} \right]$$

$$\Rightarrow \frac{-\pi}{12} = -\frac{\pi}{12} + c \Rightarrow c = 0 \Rightarrow f(x) = -x \sin x$$

$$(a) \quad f\left(\frac{\pi}{4}\right) = -\frac{\pi}{4} \cdot \frac{1}{\sqrt{2}}$$

$$(b) \quad f(x) = -x \sin x$$


$$\therefore \sin x > x - \frac{x^3}{6} \quad \forall x \in (0, \pi)$$

$$\therefore -x \sin x < -x^2 + \frac{x^4}{6} \Rightarrow f(x) < -x^2 + \frac{x^4}{6} \quad \forall x \in (0, \pi)$$

$$(c) \quad f'(x) = -\sin x - x \cos x$$

$$\text{Now } f'(x) = 0 \Rightarrow \tan x = -x$$

$$\therefore \text{There exist } \alpha \in (0, \pi) \text{ for which } f'(\alpha) = 0$$



$$(d) \quad \text{Here, } f''(x) = -2 \cos x + x \sin x$$

$$\therefore f''\left(\frac{\pi}{2}\right) = \frac{\pi}{2} \text{ and } f\left(\frac{\pi}{2}\right) = -\frac{\pi}{2} \Rightarrow f''\left(\frac{\pi}{2}\right) + f\left(\frac{\pi}{2}\right) = 0$$

$$10. \quad (a) \quad \text{We have } f(x) = g(x) \sin x$$

$$\Rightarrow f'(x) = g'(x) \sin x + g(x) \cos x$$

$$\Rightarrow f'(0) = g'(0) \times 0 + g(0) = g(0) \quad [\because g'(0) = 0]$$

\therefore Statement 2 is correct.

$$\text{Also } f''(0) = \lim_{x \rightarrow 0} \frac{f'(x) - f'(0)}{x}$$

$$= \lim_{x \rightarrow 0} \frac{g(x) \cos x + g'(x) \sin x - g(0)}{x}$$

$$= \lim_{x \rightarrow 0} \frac{g(x) \cos x - g(0)}{x} + \lim_{x \rightarrow 0} \frac{g'(x) \sin x}{x}$$

$$= \lim_{x \rightarrow 0} \frac{g(x) \cos x - g(0)}{x \times \frac{\sin x}{x}} + \lim_{x \rightarrow 0} g'(x)$$

$$= \lim_{x \rightarrow 0} \frac{g(x) \cos x - g(0)}{\sin x} + g'(0)$$

$$= \lim_{x \rightarrow 0} [g(x) \cot x - g(0) \operatorname{cosec} x] + 0$$

$$= \lim_{x \rightarrow 0} [g(x) \cot x - g(0) \operatorname{cosec} x]$$

\therefore Statement 1 is also true and statement 2 is a correct explanation for statement 1.

$$11. \quad \text{Given : } f \text{ is twice differentiable such that}$$

$$f'''(x) = -f(x) \text{ and } f'(x) = g(x)$$

$$h(x) = [f'(x)]^2 + [g(x)]^2$$

$$\Rightarrow h'(x) = 2f'f'' + 2gg' = 2f(x)g(x) + 2g(x)f''(x)$$

$$[\because g(x) = f'(x) \Rightarrow g'(x) = f''(x)]$$

$$= 2f(x)g(x) + 2g(x)(-f(x)) \quad [\because f''(x) = -f(x)]$$

$$= 2f(x)g(x) - 2f(x)g(x) = 0$$

$$\therefore h'(x) = 0, \text{ for all } x \Rightarrow h \text{ is a constant function}$$

$$\therefore h(5) = 11 \Rightarrow h(10) = 11.$$