

Chapter 16

Probability

Miscellaneous Exercise

Q. 1 A box contains 10 red marbles, 20 blue marbles and 30 green marbles. 5 marbles are drawn from the box, what is the probability that

(i) all will be blue? (ii) at least one will be green?

Answer:

No. of red marbles = 10

No. of blue marbles = 20

No. of green marbles = 30

Total number of marbles = $10 + 20 + 30 = 60$

Number of ways of drawing 5 marbles from 60 marbles = $60C_5$

(i) All the drawn marbles will be blue if we draw 5 marbles out of 20 blue marbles.

Number of ways of drawing 5 blue marbles from 20 blue marbles = $20C_5$

Probability that all marbles will be blue = $20C_5 / 60C_5$

(ii) Number of ways in which the drawn marble is not green = $(20+10)C_5$

Probability that no marble is green = $30C_5 / 60C_5$

Probability that at least one marble is green = $1 - 30C_5 / 60C_5$

Q. 2 4 cards are drawn from a well – shuffled deck of 52 cards. What is the probability of obtaining 3 diamonds and one spade?

Answer:

Number of ways of drawing 4 cards from 52 cards = $52C_4$

In a deck of 52 cards, there are 13 diamonds and 13 spades.

Number of ways of drawing 3 diamonds and one spade = $13C_3 \times 13C_1$

Thus, the probability of obtaining 3 diamonds and one spade

$$= (13C_3 \times 13C_1) / 52C_4$$

Q. 3 A die has two faces each with number '1', three faces each with number '2' and one face with number '3'. If die is rolled once, determine

(i) P (2) (ii) P (1 or 3) (iii) P (not 3)

Answer:

Total number of faces = 6

(i) Number faces with number '2' = 3

$$P(2) = \frac{3}{6} = \frac{1}{2}$$

(ii) $P(1 \text{ or } 3) = P(\text{not } 2) = 1 - P(2)$

$$P(1 \text{ or } 3) = 1 - \frac{1}{2} = \frac{1}{2}$$

(iii) Number of faces with number '3' = 1

$$P(3) = \frac{1}{6}$$

$$P(\text{not } 3) = 1 - P(3) = 1 - \frac{1}{6} = \frac{5}{6}$$

Q. 4 In a certain lottery 10,000 tickets are sold and ten equal prizes are awarded. What is the probability of not getting a prize if you buy?

(a) one ticket

(b) two tickets

(c) 10 tickets.

Answer:

Total number of tickets sold = 10,000

Number prizes awarded = 10

(i) If we buy one ticket, then

$$P(\text{getting a prize}) = \frac{10}{10000} = \frac{1}{1000}$$

$$P(\text{not getting a prize}) = 1 - \frac{1}{1000} = \frac{999}{1000}$$

(ii) If we buy two tickets, then

Number of tickets not awarded = $10,000 - 10 = 9990$

$$P(\text{not getting a prize}) = \frac{9990C_2}{10000C_2}$$

(iii) If we buy 10 tickets, then

Number of tickets not awarded = $10,000 - 10 = 9990$

$$P(\text{not getting a prize}) = \frac{9990C_{10}}{10000C_{10}}$$

Q. 5 Out of 100 students, two sections of 40 and 60 are formed. If you and your friend are among the 100 students, what is the probability that

(a) you both enter the same section?

(b) you both enter the different sections?

Answer:

My friend and I are among the 100 students.

Total number of ways of selecting 2 students out of 100 students =

(a) Let S = The two of us will enter the same section if both of us are among 40 students or among 60 students.

Number of ways in which both of us enter the same section = $P(S)$

$$P(S) = (40C_2 + 60C_2) / 100C_2$$

$$P(S) = \frac{\frac{40!}{2! \times 38!} + \frac{60!}{2! \times 58!}}{\frac{100!}{2! \times 98!}} = \frac{39 \times 40 + 59 \times 60}{99 \times 100} = \frac{5100}{9900} = \frac{17}{33}$$

(b) $P(\text{we enter different sections}) = 1 - P(\text{we enter the same section})$

$$P(\text{we enter different sections}) = 1 - \frac{17}{33} = \frac{16}{33}$$

Q. 6 Three letters are dictated to three persons and an envelope is addressed to each of them, the letters are inserted into the envelopes at random so that each envelope contains exactly one letter. Find the probability that at least one letter is in its proper envelope.

Answer:

Let L_1, L_2, L_3 be three letters and $E_1, E_2,$ and E_3 be their corresponding envelopes respectively.

Let L_1E_1 denote with letter is inserted in with envelope

\therefore Sample space is

$L_1E_1, L_2E_3, L_3E_2,$

$L_2E_2, L_1E_3, L_3E_1,$

$L_3E_3, L_1E_2, L_2E_1,$

$L_1E_1, L_2E_2, L_3E_3,$

$L_1E_2, L_2E_3, L_3E_1,$

$L_1E_3, L_2E_1, L_3E_2,$

\therefore there are 6 ways of inserting 3 letters in 3 envelopes.

And there are 4 ways in which at least one letter is inserted in proper envelope. (first 4 rows of sample space)

$$\text{Probability that at least one letter is inserted in proper envelope} = \frac{4}{6} = \frac{2}{3}$$

Q. 7 An A and B are two events such that $P(A) = 0.54$, $P(B) = 0.69$ and $P(A \cap B) = 0.35$.

Find (i) $P(A \cup B)$ (ii) $P(A \cap B)$ (iii) $P(A \cap B)$ (iv) $P(B \cap A)$

Answer:

It is given that $P(A) = 0.54$, $P(B) = 0.69$, $P(A \cap B) = 0.35$

(i) We know that $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$$\Rightarrow P(A \cup B) = 0.54 + 0.69 - 0.35 = 0.88$$

$$\therefore P(A \cup B) = 0.88$$

(ii) $A' \cap B' = (A \cup B)'$ [by De Morgan's law]

$$\text{So, } P(A' \cap B') = P(A \cup B)' = 1 - P(A \cup B) = 1 - 0.88 = 0.12$$

$$\therefore P(A' \cap B') = 0.12$$

(iii) $P(A \cap B') = P(A) - P(A \cap B) = 0.54 - 0.35 = 0.19$

$$\therefore P(A \cap B') = 0.19$$

(iv) We know that: $P(B \cap A') = P(B) - P(A \cap B)$

$$\Rightarrow P(B \cap A') = 0.69 - 0.35$$

$$\therefore P(B \cap A') = 0.34$$

Q. 8 From the employees of a company, 5 persons are selected to represent them in the managing committee of the company. Particulars of five persons are as follows:

S. No.	Name	Sex	Age in years
1	Harish	M	30
2	Rohan	M	33
3	Sheetal	F	46
4	Alis	F	28
5	Salim	M	41

A person is selected at random from this group to act as a spokesperson. What is the probability that the spokesperson will be either male or over 35 years?

Answer:

Given Data

S. No.	Name	Sex	Age in years
1	Harish	M	30
2	Rohan	M	33
3	Sheetal	F	46
4	Alis	F	28
5	Salim	M	41

Total no. of person = 5

No. of spokesperson who are male = 3

No. of spokesperson who are 35 years of age = 2

Let, E be the event in which the spokesperson will be a male and F be the event in which the spokesperson will be over 35 years of age.

Accordingly, $P(E) = 3/5$ and $P(F) = 2/5$

Since there is only one male who is over 35 years of age,

$$\therefore P(E \cap F) = \frac{1}{5}$$

We know that: $P(E \cup F) = P(E) + P(F) - P(E \cap F)$

$$\Rightarrow P(E \cap F) = \frac{3}{5} + \frac{2}{5} - \frac{1}{5}$$

$$\therefore P(E \cup F) = 4/5$$

Thus, the probability that the spokesperson will either be a male or over 35 years of age is $4/5$.

Q. 9 If 4-digit numbers greater than 5,000 are randomly formed from the digits 0, 1, 3, 5, and 7, what is the probability of forming a number divisible by 5 when,

(i) the digits are repeated? (ii) the repetition of digits is not allowed?

Answer:

(i) When the digits are repeated

Since four-digit numbers greater than 5000 are formed,

The thousand's place digit is either 7 or 5. The remaining 3 places can be filled by any of the digits 0, 1, 3, 5, or 7 as repetition of digits is allowed.

Total number of 4-digit numbers greater than 5000 $= 2 \times 5 \times 5 \times 5 - 1$
 $= 250 - 1 = 249$

[In this case, 5000 cannot be counted; so, 1 is subtracted]

A number is divisible by 5 if the digit at its unit's place is either 0 or 5.

\therefore Total number of 4-digit numbers greater than 5000 that are divisible by 5 $= 2 \times 5 \times 5 \times 2 - 1 = 100 - 1 = 99$ Thus, the probability of forming a number divisible by 5 when the digits are repeated is $= P$ (number divisible by 5 when digits repeated)

$$\Rightarrow P \text{ (number divisible by 5 when digits repeated)} = \frac{99}{249} = \frac{33}{83}$$

(ii) When repetition of digits is not allowed

The thousands place can be filled with either of the two digits 5 or 7 i.e. by 2 ways.

The remaining 3 places can be filled with any of the remaining 4 digits.

Total number of 4-digit numbers greater than 5000 $= 2 \times 4 \times 3 \times 2 = 48$

When the digit at the thousands place is 5, the unit place can be filled only with 0 and the tens and hundreds places can be filled with any two of the remaining 3 digits. Here, number of 4-digit numbers starting with 5 and divisible by 5 = $1 \times 3 \times 2 \times 1 = 6$

When the digit at the thousands place is 7, the unit place can be filled in two ways (0 or 5) and the tens and hundreds of places can be filled with any two of the remaining 3 digits. Here, number of 4-digit numbers starting with 7 and divisible by 5 = $1 \times 2 \times 3 \times 2 = 12$

Total number of 4-digit numbers greater than 5000 that are divisible by 5

$$= 6 + 12 = 18$$

Thus, the probability of forming a number divisible by 5 when the repetition of digits is not allowed

= P (number divisible by 5 when digits are not repeated)

$$\Rightarrow P (\text{number divisible by 5 when digits repeated}) = \frac{18}{48} = \frac{3}{8}$$

Q. 10 The number lock of a suitcase has 4 wheels, each labelled with ten digits i.e., from 0 to 9. The lock opens with a sequence of four digits with no repeats. What is the probability of a person getting the right sequence to open the suitcase?

Answer:

The number lock has 4 wheels, each labelled with ten digits i.e., from 0 to 9.

Number of ways of selecting 4 different digits out of the 10 digits = ${}^{10}C_4$

Now, each combination of 4 different digits can be arranged in 4! Ways.

Number of four digits with no repetitions = $4! \times {}^{10}C_4 = 5040$

There is only one number that can open the suitcase.

Thus, the required probability is $\frac{1}{5040}$