


NCERT Intext Questions

Ans. Glucose contains five —OH groups and sucrose (molecular mass = 342) contains eight —OH groups. These —OH groups form hydrogen bonds with water. Because of this extensive intermolecular hydrogen-bonding, glucose and sucrose are soluble in water.

[CBSE 2019 (56/2/3)]

$$\text{C}_{12}\text{H}_{22}\text{O}_{11} + \text{H}_2\text{O} \xrightarrow[\text{or Lactase}]{\text{H}_3\text{O}^+} \underset{\text{D-(+) - Glucose}}{\text{C}_6\text{H}_{12}\text{O}_6} + \underset{\text{D-(+) - Galactose}}{\text{C}_6\text{H}_{12}\text{O}_6}$$

Ans. The cyclic hemiacetal form of glucose contains an —OH group at C-1 which gets hydrolysed in the aqueous solution to produce the open chain aldehydic form which then reacts with NH_4OH to form the corresponding oxime. Therefore, glucose contains an aldehydic group. On the other hand, when glucose is reacted with acetic anhydride, the —OH group at C-1, along with the four other —OH groups at C-2, C-3, C-4 and C-6 form a pentaacetate. As the pentaacetate of glucose does not contain a free —OH group at C-1, it cannot get hydrolysed in aqueous solution to produce the open chain aldehydic form and thus glucose pentaacetate does not react with NH_4OH to form glucose oxime.



Q. 4. The melting points and solubility in water of amino acids are generally higher than that of the corresponding halo acids. Explain.

Ans. The amino acids exist as zwitter ions, $\text{H}_3\text{N}^+-\text{CHR}-\text{COO}^-$. Because of this dipolar salt-like character, they have strong dipole-dipole attractions. So, their melting points are higher than halo acids which do not have salt-like character. Moreover, due to this salt-like character, they interact strongly with H_2O . Thus, solubility in water of amino acids is higher than that of the corresponding halo acids which do not have salt-like character.

Q. 5. Where does the water present in the egg go after boiling the egg?

Ans. On boiling the egg, the proteins undergo denaturation and the water present in the egg gets absorbed or adsorbed in the denaturated proteins probably through H-bonding.

Q. 6. Why cannot vitamin C be stored in our body?

Ans. Vitamin C is soluble in water, hence, it is readily excreted in urine and thus cannot be stored in the body.

Q. 7. What products would be formed when a nucleotide from DNA containing thymine is hydrolysed?

Ans. Besides thymine, the two products are 2-deoxy-D-ribose and phosphoric acid.

Q. 8. When RNA is hydrolysed, there is no relationship among the quantities of different bases obtained. What does this fact suggest about the structure of RNA? [HOTS]

Ans. A DNA molecule has two strands in which the four complementary bases pair each other, viz., cytosine (C) always pairs with guanine (G) while thymine (T) always pairs with adenine (A). Therefore, when a DNA molecule is hydrolysed, the molar amounts of cytosine is always equal to that of guanine and that of adenine is always equal to that of thymine. RNA also contains four bases, the first three are same as in DNA but the fourth one is uracil (U).

As in RNA, there is no relationship between the quantities of four bases (C, G, A and U) obtained, therefore, the base-pairing principle, viz., A pairs with U and C pairs with G is not followed. So, unlike DNA, RNA has a single strand.

NCERT Exercises

Q. 1. What are monosaccharides?

Ans. Monosaccharides are carbohydrates which cannot be further hydrolysed to simpler molecules. The general formula is $(\text{CH}_2\text{O})_n$ where $n = 3 - 7$.

Q. 2. What are reducing sugars?

Ans. Carbohydrates which reduce Fehling's solution to red precipitate of Cu_2O or Tollens' reagent to metallic Ag are called reducing sugars. All monosaccharides (both aldoses and ketoses) and disaccharides except sucrose are reducing sugars.

Q. 3. Write two main functions of carbohydrates in plants.

Ans. (i) **Structural material for cell walls:** The polysaccharide, cellulose acts as the chief structural material of the cell walls.

(ii) **Reserve food material:** The polysaccharide, starch is the major reserve food material in the plants.

Q. 4. Classify the following into monosaccharides and disaccharides: Ribose, 2-deoxyribose, maltose, galactose, fructose and lactose.

Ans. Monosaccharides: Ribose, 2-deoxyribose, galactose and fructose.

Disaccharides: Maltose and lactose.

Q. 5. What do you understand by the term glycosidic linkage?

Ans. The linkage between two monosaccharide units in an oligosaccharide or a polysaccharide through oxygen atom is called glycosidic linkage.


Q. 6. What is glycogen? How is it different from starch?

[CBSE (F) 2012]

Ans. Glycogen is a polymer of α -D-glucose. The carbohydrates are stored in animal body as glycogen. Starch is also a polymer of α -D-glucose and consist of two components amylose and amylopectin. Amylose is linear chain polymer of α -D-glucose. Both glycogen and amylopectin are branched chain polymers of α -D-glucose but glycogen is more highly branched than amylopectin. Starch is the main storage polysaccharide of plants.

Q. 7. What are the hydrolysis products of (i) sucrose (ii) lactose? [CBSE (AI) 2010, 2013, 2014]

Ans. Both sucrose and lactose are disaccharides. Sucrose on hydrolysis gives one molecule each of glucose and fructose. Lactose on hydrolysis gives one molecule each of glucose and galactose.



Q. 8. What is the basic structural difference between starch and cellulose? [CBSE 2019 (56/2/2)]
OR

Starch and cellulose both contain glucose units as monomer, yet they are structurally different.
Explain. [CBSE 2020 (56/1/2)]

Ans. Starch consists of amylose and amylopectin.

Amylose is a long unbranched chain with α -D-(+)-glucose units that are held together with C1–C4 glycosidic linkage. Amylopectin is a highly branched chain polymer of α -D-glucose units in which glycosidic linkage is present between C1–C4 atoms and branching occurs through C1–C6 glycosidic linkage. For figures refer to Points to remember 10(a).

Cellulose is a straight chain polysaccharide composed only of β -D-glucose units which are joined by glycosidic linkage between C1 of one glucose units and C4 of next glucose units. For figure refer to Points to remember 10(b).

Q. 9. What happens when D-glucose is treated with the following reagents? [CBSE 2019 (56/5/2)]
(i) HI (ii) Bromine water (iii) HNO_3

Ans. Refer to Points to remember 7.

Q. 10. Enumerate the reactions of D-glucose which cannot be explained by its open chain structure.

Ans. The following reactions cannot be explained by its open chain structure proposed by Baeyer:

- Despite having an aldehydic group, glucose does not give Schiff's test and it does not react with sodium bisulphite and ammonia.
- The pentaacetate of glucose does not react with hydroxylamine indicating absence of —CHO group.
- When D-glucose is treated with methyl alcohol in presence of dry hydrogen chloride gas, it gives two isomeric mono methyl derivatives known as α -D-glucoside and methyl β -D-glucoside. These glucosides does not reduce Fehling's solution and also do not react with hydrogen cyanide or hydroxylamine indicating the absence of free —CHO group.

Q. 11. What are essential and non-essential amino acids? Give two examples of each type.

[CBSE Delhi 2010]

Ans. α -Amino acids which are required for health and growth of human beings but are not synthesised by the human body are known as essential amino acids. Examples: valine, leucine, phenylalanine, etc. On the other hand, α -amino acids which are needed for health and growth of human beings and are synthesised by the human body are called non-essential amino acids. Examples: glycine, alanine, aspartic acid, etc.

Q. 12. Define the following terms as related to proteins:

- (i) Peptide linkage (ii) Primary structure (iii) Denaturation

[CBSE (AI) 2014; 2019 (56/5/2)]

Ans. (i) A **peptide linkage** is an amide ($\text{—}\overset{\text{O}}{\parallel}\text{C—NH—}$) linkage formed between —COOH group of one α -amino acid and —NH_2 group of other α -amino acid by loss of a water molecule.

(ii) The specific sequence in which various α -amino acids present in a protein are linked to one another is called its **primary structure**. Any change in its primary structure creates a new protein.

(iii) When a protein in its native form is subjected to a change, such as change in temperature or change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called **denaturation of protein**. During denaturation, 2° and 3° structures are destroyed but 1° structure remains intact, e.g., coagulation of egg white on boiling, curdling of milk, etc.

Q. 13. What are common types of secondary structure of proteins?

Ans. The conformation which the polypeptide chains assume as a result of hydrogen bonding is known as the secondary structure of the proteins. The two types of secondary structures are α -helix and β -pleated sheet structure.

Q. 14. What type of bonding helps in stabilising the α -helix structure of proteins?

Ans. The α -helix structure of proteins is stabilised by intramolecular H-bonding between C=O of one amino acid residue and the N—H of the fourth amino acid residue in the chain.

Q. 15. Differentiate between globular and fibrous proteins.


[CBSE 2019 (56/5/2), 2020 (56/5/1)]

Ans. Refer to Points to remember 12(a).

Q. 16. How do you explain the amphoteric behaviour of amino acids?

Ans. Amino acids contain both acidic (carboxyl group) and basic (amino group) groups in the same molecule. In aqueous solution, the carboxyl group can lose a proton and amino group can accept a proton, giving rise to a dipolar ion known as zwitter ion. This is neutral but contains both positive and negative charges.

In zwitter ionic form, amino acids show amphoteric behaviour as they react with both acids and bases.



Q. 17. What are enzymes?

Ans. Enzymes are biological catalysts. Each biological system requires a different enzyme. So, as compared to conventional catalysts, enzymes are specific and efficient in their action. They are required in only small quantity and work at optimum temperature (310 K) and pH (7.4) under 1 atmospheric pressure. Chemically, they are globular proteins. However, some enzymes are also associated with some non-protein components called the co-factors for their activity. Cofactors are of two types:

(i) **Inorganic ions** such as Zn^{2+} , Mg^{2+} , Mn^{2+} , Fe^{2+} , Cu^{2+} , Co^{2+} , etc.

(ii) **Organic molecules:** These are of two types:

(a) **Coenzymes:** These are usually derived from vitamins such as thymine, riboflavin, niacin, etc. They are loosely held to the protein and can be easily separated by dialysis.

(b) **Prosthetic group:** They are also derived from vitamins such as biotin but are tightly held to the protein molecule by covalent bonds. They can be separated only by careful hydrolysis.

Q. 18. What is the effect of denaturation on the structure of proteins?

Ans. During denaturation, 2° and 3° structures of proteins are destroyed but 1° structure remains intact. Due to denaturation, the globular proteins (soluble in H_2O) are converted into fibrous proteins (insoluble in H_2O) and their biological activity is lost. For example, boiled egg which contains coagulated proteins cannot be hatched.

Q. 19. How are vitamins classified? Name the vitamin responsible for the coagulation of blood.

Ans. Vitamins are classified into two groups depending upon their solubility in water or fat.

(i) **Water-soluble vitamins:** These include vitamin B-complex (B_1 , B_2 , B_3 , i.e., nicotinic acid, B_6 , B_{12} , pantothenic acid, biotin, i.e., vitamin H and folic acid) and vitamin C.

(ii) **Fat-soluble vitamins:** These include vitamin A, D, E and K. These are stored in liver and adipose tissues (fat storing tissues).

Vitamin K is responsible for coagulation of blood.

Q. 20. Why are vitamin A and vitamin C essential to us? Give their important sources.

Ans. Vitamin A is essential for us because its deficiency can cause xerophthalmia (hardening of cornea of eye) and night blindness.

Sources: Carrots, fish liver oil, butter and milk.

Vitamin C: Vitamin C is essential for us because its deficiency causes scurvy (bleeding gums) and pyorrhea (loosening and bleeding of teeth).

Sources: Amla, citrus fruits and green leafy vegetables.

Q. 21. What are nucleic acids? Mention their two important functions.

Ans. Nucleic acids are biomolecules which are found in the nuclei of all living cells in the form of nucleoproteins or chromosomes (proteins containing nucleic acids as the prosthetic group).

These are of two types: deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). The two main functions of nucleic acids are:


- DNA is responsible for transmission of hereditary effects from one generation to another. This is because of the unique property of replication during cell division and the transfer of two identical DNA strands to the daughter cells.
- DNA and RNA are responsible for synthesis of all proteins essential for the growth and maintenance of our body. Actually, the proteins are synthesised by various RNA molecules (*rRNA*, *mRNA* and *tRNA*) in the cell but the message for the synthesis of a particular protein is present in DNA.

Q. 22. What is the difference between a nucleoside and a nucleotide?

[CBSE Delhi 2010; (F) 2013]

Ans. A nucleoside is formed when 1-position of pyrimidine (cytosine, thymine or uracil) or 9-position of purine (guanine or adenine) base is connected to C-1 of sugar (ribose or deoxyribose) by a β -linkage. Hence, in general, nucleosides may be represented as: Sugar–Base.



A nucleotide contains all the three basic compounds of nucleic acids, *i.e.*, a phosphoric acid group, a pentose sugar and a nitrogenous base. These are obtained by esterification of $C_5 - OH$ group of the pentose sugar by phosphoric acid. Thus, in general, a nucleotide is represented as:



For figures refer to Points to remember 14(a).

Q. 23. The two strands in DNA are not identical but are complementary. Explain. [CBSE 2020 (56/1/2)]

Ans. The two strands in DNA molecule are held together through hydrogen bonds between purine base of one strand and pyrimidine base of the other and vice versa. Because of different sizes and geometries of the base, the only possible pairing in DNA are G (guanine) and C (cytosine) through three H-bonds (*i.e.*, $C \equiv G$) and between A (adenine) and T (thymine) through two H-bonds (*i.e.*, $A = T$). Due to this base-pairing principle, the sequence of bases in one strand automatically fixes the sequence of bases in the other strand. Thus, the two strands are not identical but are complementary.



Q. 24. Write the important structural and functional differences between DNA and RNA.

Ans.

Structural differences		
	DNA	RNA
(i)	The sugar present in DNA is 2-deoxy-D-(–)-ribose.	The sugar present in RNA is D-(–)-ribose.
(ii)	DNA contains cytosine and thymine as pyrimidine bases.	RNA contains cytosine and uracil as pyrimidine bases.
(iii)	DNA has a double stranded α -helix structure.	RNA has a single stranded α -helix structure.
(iv)	DNA molecules are very large; their molecular mass may vary from $6 \times 10^6 - 16 \times 10^6$ u.	RNA molecules are much smaller with molecular mass ranging from 20,000 to 40,000 u.
Functional differences		
(i)	DNA has unique property of replication.	RNA usually does not replicate.
(ii)	DNA controls the transmission of hereditary effects.	RNA controls the synthesis of proteins.

Q. 25. What are the different types of RNA found in the cell?

[CBSE Delhi 2013]

Ans. There are three types of RNAs:




- (i) Ribosomal RNA (rRNA)
- (ii) Messenger RNA (mRNA)
- (iii) Transfer RNA (tRNA)

Multiple Choice Questions

Choose and write the correct option(s) in the following questions.




1. Three cyclic structures of monosaccharides are given below which of these are anomers.

[NCERT Exemplar]



- (a) I and II (b) II and III (c) I and III (d) III is anomer of I and II
2. Optical rotations of some compounds along with their structures are given below which of them have D configuration.

[NCERT Exemplar]



- (a) I, II, III (b) II, III (c) I, II

(d) III

3. Which of the following statements is not true about glucose? [NCERT Exemplar]
 (a) It is an aldohexose. (b) On heating with HI it forms n-hexane.
 (c) It is present in furanose form. (d) It does not give 2,4-DNP test.


4. Which of the following reactions of glucose can be explained only by its cyclic structure? [NCERT Exemplar]

- (a) Glucose forms pentaacetate.
 (b) Glucose reacts with hydroxylamine to form an oxime.
 (c) Pentaacetate of glucose does not react with hydroxylamine.
 (d) Glucose is oxidised by nitric acid to gluconic acid.

5. The glycosidic linkage involved in linking the glucose units in amylose part of starch is: [CBSE 2023 (56/1/1)]

- (a) $C_1 - C_6$ α linkage (b) $C_1 - C_6$ β linkage
 (c) $C_1 - C_4$ α linkage (d) $C_1 - C_4$ β linkage

6. Which of the following pairs represents anomers? [NCERT Exemplar]



7. Nucleosides are composed of [CBSE 2022 (56/3/4)]
 (a) a pentose sugar and phosphoric acid.
 (b) a nitrogenous base and phosphoric acid.
 (c) a nitrogenous base and a pentose sugar.
 (d) a nitrogenous base, a pentose sugar and phosphoric acid.

8. Carbohydrates are classified on the basis of their behaviour on hydrolysis and also as reducing or non-reducing sugar. Sucrose is a _____. [NCERT Exemplar]
 (a) monosaccharide (b) polysaccharide
 (c) reducing sugar (d) non-reducing sugar

9. Disaccharides that are reducing in nature are: [CBSE Sample Paper 2021]
 (a) sucrose and lactose (b) sucrose and maltose
 (c) lactose and maltose (d) sucrose, lactose and maltose

10. An α -helix is a structural feature of [CBSE 2023(56/2/1)]
 (a) Sucrose (b) Polypeptides
 (c) Nucleotides (d) Starch

11. Cellulose is not digestible by human beings due to absence of cellulose hydrolysing enzyme called
 (a) cellulase (b) invertase
 (c) zymase (d) urease

12. Glycogen is a branched chain polymer of α -D-glucose units in which chain is formed by C1—C4 glycosidic linkage whereas branching occurs by the formation of C1—C6 glycosidic linkage. Structure of glycogen is similar to _____. [NCERT Exemplar]
 (a) Amylose (b) Amylopectin
 (c) Cellulose (d) Glucose
13. Which of the following polymer is stored in the liver of animals? [NCERT Exemplar]
 (a) Amylose (b) Cellulose
 (c) Amylopectin (d) Glycogen
14. Which of the following naturally occurring α -amino acids is optically inactive?
 (a) Glycine (b) Alanine (c) Leucine (d) Valine
15. Curdling of milk is an example of [CBSE Sample Paper 2021]
 (a) breaking of peptide linkage. (b) hydrolysis of lactose.
 (c) breaking of protein into amino acids. (d) denaturation of protein.
16. Each polypeptide in a protein has amino acids linked with each other in a specific sequence. This sequence of amino acids is said to be _____. [NCERT Exemplar]
 (a) primary structure of proteins. (b) secondary structure of proteins.
 (c) tertiary structure of proteins. (d) quaternary structure of proteins.
17. Which parts of amino acids molecules are linked through hydrogen bonds in the secondary structure of proteins? [CBSE 2020 (56/3/2)]
 (a) NH_2 group (c) COOH group
 (b) $-\text{C}-$ and $-\text{NH}-$ groups (d) None of the above

$$\begin{array}{c} \text{O} \\ || \\ -\text{C}- \end{array}$$
18. Proteins can be classified into two types on the basis of their molecular shape i.e., fibrous proteins and globular proteins. Examples of globular proteins are : [NCERT Exemplar]
 (a) Insulin (b) Fibrinogen
 (c) Albumin (d) All of these
19. Proteins are found to have two different types of secondary structures viz. α -helix and β -pleated sheet structure. α -helix structure of protein is stabilised by : [NCERT Exemplar]
 (a) Peptide bonds (b) van der Waals forces
 (c) Hydrogen bonds (d) Dipole-dipole interactions
20. Which of the following B group vitamins can be stored in our body? [NCERT Exemplar]
 (a) Vitamin B₁ (b) Vitamin B₂
 (c) Vitamin B₆ (d) Vitamin B₁₂
21. Which of the following acids is a vitamin? [NCERT Exemplar]
 (a) Aspartic acid (b) Ascorbic acid
 (c) Adipic acid (d) Saccharic acid
22. Dinucleotide is obtained by joining two nucleotides together by phosphodiester linkage. Between which carbon atoms of pentose sugars of nucleotides are these linkages present? [NCERT Exemplar]
 (a) 5' and 3' (b) 1' and 5'
 (c) 5' and 5' (d) 3' and 3'
23. Which of the following statements is not true about glucose? [CBSE 2023 (56/5/2)]
 (a) It is an aldohexose. (b) On heating with HI it forms *n*-hexane.
 (c) It is present in pyranose form. (d) It gives 2, 4 DNP test.
24. Which one is the complementary base of cytosine in one strand to that in other strand of DNA? [CBSE 2020 (56/4/3)]
 (a) Adenine (b) Guanine (c) Thymine (d) Uracil
25. On hydrolysis, which of the following carbohydrates gives glucose and fructose? [CBSE 2023 (56/4/2)]
 (a) Sucrose (b) Starch (c) Lactose (d) Maltose

26. Match the following:

Column-I	Column-II
(i) Globular protein	A. Protein with the tertiary structure in which it normally occurs in living systems.
(ii) Zwitter ion	B. Protein that is usually water soluble having a hydrophilic exterior and hydrophobic interior and an overall rounded shape.
(iii) Fibrous protein	C. A protein that is water insoluble, is very tough and has a long shape.
(iv) Native protein	D. Overall a neutral molecule but migrates to cathode or anode depending on pH.
(v) Simple protein	E. A protein that produces only amino acids on hydrolysis.

(a) (i) — B, (ii) — D, (iii) — C, (iv) — A, (v) — E

(b) (i) — D, (ii) — B, (iii) — A, (iv) — C, (v) — E

(c) (i) — C, (ii) — B, (iii) — A, (iv) — D, (v) — E

(d) (i) — E, (ii) — A, (iii) — C, (iv) — D, (v) — B

Answers

1. (a) 2. (a) 3. (c) 4. (c) 5. (c) 6. (c) 7. (c) 8. (d) 9. (c) 10. (b)
 11. (a) 12. (b) 13. (d) 14. (a) 15. (d) 16. (a) 17. (c) 18. (d) 19. (c) 20. (d)
 21. (b) 22. (a) 23. (d) 24. (b) 25. (a) 26. (a)

Assertion-Reason Questions

In the following questions, two statements are given—one labeled Assertion (A) and the other labeled Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below:

- (a) Both Assertion (A) and Reason (R) are correct statements, and Reason (R) is the correct explanation of the Assertion (A).
 (b) Both Assertion (A) and Reason (R) are correct statements, but Reason (R) is not the correct explanation of the Assertion (A).
 (c) Assertion (A) is correct, but Reason (R) is incorrect statement.
 (d) Assertion (A) is incorrect, but Reason (R) is correct statement.
- Assertion (A) : Deoxyribose, $C_5H_{10}O_4$ is a carbohydrate.
Reason (R) : Carbohydrates are hydrates of carbon so compounds which follow $C_x(H_2O)_y$ formula are carbohydrates.
 - Assertion (A) : D (+) – Glucose is dextrorotatory in nature.
Reason (R) : 'D' represents its dextrorotatory nature.
 - Assertion (A) : Glucose reacts with hydroxylamine to form an oxime and also adds a molecule of hydrogen cyanide to give cyanohydrin. [CBSE Sample Paper 2020]
Reason (R) : The carbonyl group is present in the open chain structure of glucose.
 - Assertion (A) : Reducing sugars undergo mutarotation.
Reason (R) : During mutarotation, one pure anomer is converted into an equilibrium mixture of two anomers.
 - Assertion (A) : Sucrose is a non-reducing sugar.
Reason (R) : Sucrose has glycosidic linkage. [CBSE 2020 (56/1/1)]
 - Assertion (A) : A solution of sucrose in water is dextrorotatory but on hydrolysis in presence of little hydrochloric acid, it becomes laevorotatory.
Reason (R) : Sucrose on hydrolysis gives unequal amounts of glucose and fructose as a result of which change in sign of rotation is observed.

OR

Define the following as related to carbohydrates :

(a) Anomers

(b) Glycosidic linkage

Answers

1. It indicates the absence of free $\text{—}\overset{\text{O}}{\parallel}\text{C—H}$ group.
2. Vitamin C is soluble in water therefore it is readily excreted in urine and hence cannot be stored in body.
3. (a) A **peptide linkage** is an amide ($\text{—}\overset{\text{O}}{\parallel}\text{C—NH—}$) linkage formed between —COOH group of one α -amino acid and —NH_2 group of other α -amino acid by loss of a water molecule.
- (b) When a protein in its native form is subjected to a change, such as change in temperature or change in pH, the hydrogen bonds are disturbed. Due to this, globules unfold and helix get uncoiled and protein loses its biological activity. This is called **denaturation of protein**. During denaturation, 2° and 3° structures are destroyed but 1° structure remains intact, e.g., coagulation of egg while on boiling, curdling of milk, etc.

OR

- (a) Carbohydrates which differ in their configuration at the glycosidic carbon (*i.e.*, C_1 in aldoses and C_2 in ketoses) are called anomers *e.g.*, α -D glucose and β -D glucose.
- (b) The linkage between two monosaccharides through oxygen atom in an oligosaccharide or a polysaccharide is known as glycosidic linkage.

PASSAGE-2

Living systems are made up of various complex biomolecule, like carbohydrates, proteins, nucleic acids, lipids, etc. Carbohydrates are optically active polyhydroxy aldehydes or ketones or molecules which provide such units on hydrolysis. They are broadly classified into three groups - monosaccharides, oligosaccharides and polysaccharides. Monosaccharides are held together by glycosidic linkages to form disaccharides like sucrose, maltose or polysaccharides like starch and cellulose.

Another biomolecule: proteins are polymers of α -amino acids which are linked by peptide bonds. Ten amino acids are called essential amino acids. Structure and shape of proteins can be studied at four different levels *i.e.*, primary, secondary, tertiary and quaternary, each level being more complex than the previous one.

Answer the following questions:

[CBSE 2023(56/4/2)]

- What is the difference between a glycosidic linkage and peptide linkage?
- Which amino acids are called essential amino acids?
- What are the common types of secondary structures of proteins? Write any two forces which stabilise the secondary and tertiary structures of protein.

OR

Define denaturation of protein with an example. During denaturation which structures of protein lose their biological activity?

Answers

1.

Glycosidic Linkage	Peptide Linkage
Joins two monosaccharides	Joins two amino acids
The monomer units are joined by an oxide linkage called glycosidic linkage	Peptide linkage is an amide formed between —COOH and —NH_2 group of amino acids.

- The amino acid that cannot be synthesised in the body and must be obtained through food are known as essential amino acid. Example– Valine, Leucine, Arginine.
- Secondary structure of the protein refers to the shape in which the polypeptide chain can exist. The shapes are due to the interaction between the atoms in the polypeptide chain.

The common types of 2° proteins are:

- α -helix:** twisted like right handed screw because of Hydrogen bonding.
- β -pleated sheet structure:** In this all the peptide chains are stretched to maximum extent and laid side by side, which are held together by hydrogen bonds.

The two forces that stabilise the 2° and 3° structure of proteins are:

- Hydrogen bonds
- Disulphide linkages.

OR

The hydrogen bonds in the native protein are disturbed or broken when the protein molecules are subjected to physical stress (like temperature change) or chemical changes like change in pH. Owing to this, the proteins lose their biological activity, which is known as denaturation of protein.

Example – The egg protein undergoes coagulation when subjected to high temperature (boiling point). During denaturation only the secondary and tertiary structures are destroyed, the primary structure remains intact.

CONCEPTUAL QUESTIONS


Q. 1. Why are carbohydrates generally optically active?

Ans. Carbohydrates have chiral or asymmetric carbon atom.

Q. 2. What happens when glucose is treated with bromine water?

[CBSE (F) 2010]


Ans. When glucose is treated with bromine water it forms gluconic acid.



Q. 3. What happens when glucose reacts with nitric acid?

[CBSE (F) 2010]


Ans. Glucose gets oxidised to succinic acid



Q. 4. Write a reaction which shows that all the carbon atoms in glucose are linked in a straight chain.

[CBSE (AI) 2012]

Ans. On prolonged heating with HI, glucose gives *n*-hexane.



Q. 5. Which component of starch is a branched polymer of α -glucose and insoluble in water?

[CBSE Delhi 2014]

Ans. Amylopectin

Q. 6. Which of the two components of starch is water soluble?

[CBSE Delhi 2014]

Ans. Amylose is water soluble whereas amylopectin is water insoluble component.

Q. 7. Name two α -amino acids which form a dipeptide which is 100 times more sweet than cane sugar.

[NCERT Exemplar]

Ans. Aspartic acid and phenylalanine.


Q. 8. What type of bonding helps in stabilising the α -helix structure of proteins?

[CBSE Delhi 2013]

Ans. The α -helix structure of proteins is stabilised by intramolecular H-bonding between C=O of one amino acid residue and the N—H of the fourth amino acid residue in the chain.

Q. 9. How do you explain the presence of five —OH groups in glucose molecule? [NCERT Exemplar] [HOTS]

Ans. Glucose gives pentaacetate derivative on acetylation with acetic anhydride. This confirms the presence of five —OH groups.



Q. 10. Name the species formed when an aqueous solution of amino acid is dissolved in water?

[CBSE Sample Paper 2020]

Ans. Zwitter ion

Q. 11. What are biocatalysts? Give an example.

[CBSE (F) 2014]

Ans. Enzymes are termed as biocatalysts as they catalyse numerous reactions that occur in the bodies of animals and plants to maintain life process e.g., invertase, pepsin, urease.

Q. 12. What are three types of RNA molecules which perform different functions?

[CBSE Delhi 2013]

Ans. There are three types of RNAs:

(i) Ribosomal RNA (rRNA)

(ii) Messenger RNA (mRNA)

(iii) Transfer RNA (tRNA)

Short Answer Questions–I

Each of the following questions are of 2 marks.

Q. 1. Write the reactions involved when D-glucose is treated with the following reagents:



(i) HCN

(ii) Br₂ water

[CBSE (F) 2013]

Ans. (i) HCN


(ii) Br₂ water




Q. 2. How do you explain the presence of an aldehydic group in a glucose molecule? [NCERT Exemplar]

Ans. Glucose reacts with hydroxylamine to form a monoxime and adds one molecule of hydrogen cyanide to give cyanohydrin so it contains a carbonyl group which can be an aldehyde or a ketone. On mild oxidation with bromine water, glucose gives gluconic acid which is a six carbon carboxylic acid. This indicates that carbonyl group present in glucose is an aldehydic group.

Q. 3. Label the glucose and fructose units in the following disaccharide and identify anomeric carbon atoms in these units. Is the sugar reducing in nature? Explain. [NCERT Exemplar] [HOTS]




Ans. C-1 of glucose unit and C-2 of fructose unit are anomeric carbon atoms in the given disaccharide. The disaccharide is non-reducing sugar because —OH groups attached to anomeric carbon atoms are involved in the formation of glycosidic bond.



Q. 4. Amino acids behave like salts rather than simple amines or carboxylic acids. Explain.

[NCERT Exemplar]

Ans. In aqueous solution, the carboxyl group loses a proton and amino group accepts a proton to form a zwitter ion.




Q. 5. Explain what is meant by the following:

- peptide linkage
- pyranose structure of glucose.

[CBSE (AI) 2011; (F) 2011]

Ans. (i) Refer to NCERT Exercises, Q. 12(i).

(ii) The six-membered cyclic structure of glucose is called pyranose structure (α - or β -), in analogy with heterocyclic compound pyran.



Q. 6. Describe what you understand by primary structure and secondary structure of proteins.

[CBSE Delhi 2011; (F) 2011]

Ans. Primary structure: The specific sequence in which the various α -amino acids present in a protein are linked to one another is called its primary structure. Any change in the primary structure creates a different protein.

Secondary structure: The conformation which the polypeptide chain assumes as a result of hydrogen bonding is known as secondary structure. The two types of secondary structures are α -helix and β -pleated sheet structures. In α -helix structure, the polypeptide chain forms all the possible hydrogen bonds by twisting into a right-handed screw (helix) with the —NH group of each amino acid residue hydrogen bonded to the $>\text{C}=\text{O}$ groups of an adjacent turn of the helix. In β -pleated structure, all peptide chains are stretched out to nearly maximum extension and then laid side by side and are held together by hydrogen bonds.

Q. 7. (i) What type of linkage is present in nucleic acids?

(ii) Give one example each for fibrous protein and globular protein.

[CBSE Central 2016]

- Ans.** (i) Phosphodiester linkage.
 (ii) **Fibrous protein:** Myosin, keratin, collagen, etc.
Globular protein: Insulin, haemoglobin, etc.

1

1

[CBSE Marking Scheme 2016]

Q. 8. Name the bases present in RNA. Which one of these is not present in DNA?

[CBSE Delhi 2011, 2020 (56/3/3)]

Ans. The bases present in RNA are adenine (A), guanine (G), cytosine (C) and uracil (U). Uracil is not present in DNA.

Q. 9. If one strand of a DNA has the sequence —ATGCTTCA—, what is the sequence of the bases in the complementary strand?

Ans. As we know that in DNA molecule, adenine (A) always pairs with thymine (T) and cytosine (C) always pairs with guanine (G). Thus,

Sequence of bases in one strand:

A T G C T T C A

Sequence of bases in the complementary strand:


T A C G A A G T

Q. 10. (i) Write chemical reaction to show that open structure of D-glucose contains the straight chain.

(ii) What type of linkage is responsible for the formation of protein?

[CBSE 2023 (56/1/1)]

Ans. (i) On prolonged heating with HI, glucose gives *n*-hexane.



(ii) Peptide linkage $\left(\begin{array}{c} \text{O} \\ || \\ -\text{C}-\text{NH}- \end{array} \right)$

Q. 11. Write two differences between DNA and RNA.

[CBSE 2023 (56/2/1)]

Ans.

Structural differences		
	DNA	RNA
(i)	The sugar present in DNA is 2-deoxy-D-(–)-ribose.	The sugar present in RNA is D-(–)-ribose.
(ii)	DNA contains cytosine and thymine as pyrimidine bases.	RNA contains cytosine and uracil as pyrimidine bases.
(iii)	DNA has a double stranded α -helix structure.	RNA has a single stranded α -helix structure.
(iv)	DNA molecules are very large; their molecular mass may vary from $6 \times 10^6 - 16 \times 10^6$ u.	RNA molecules are much smaller with molecular mass ranging from 20,000 to 40,000 u.
Functional differences		
(i)	DNA has unique property of replication.	RNA usually does not replicate.
(ii)	DNA controls the transmission of hereditary effects.	RNA controls the synthesis of proteins.

(Any two)

Short Answer Questions–II

Each of the following questions are of 3 marks.

Q. 1. Define the following terms:

(i) Glycosidic linkage

(ii) Invert sugar

[CBSE (AI) 2014]

[CBSE 2020 (56/3/2)]

(iii) Oligosaccharides

- Ans.** (i) The linkage between two monosaccharides through oxygen atom in an oligosaccharide or a polysaccharide is known as glycosidic linkage.
- (ii) Sucrose is dextrorotatory (+66.5°) but after hydrolysis it gives an equimolar mixture of D-(+)-glucose and D-(-)-fructose, which is laevorotatory. This change of specific rotation from dextrorotation to laevorotation is called inversion of sugar and the mixture obtained is called invert sugar.
- (iii) Carbohydrates which on hydrolysis give two to ten molecules of monosaccharides are called oligosaccharides e.g., sucrose.

Q. 2. (i) Which of the following biomolecules is insoluble in water? Justify.


Insulin, Haemoglobin, Keratin.

(ii) Draw the Haworth structure for α -D-Glucopyranose.


(iii) Write chemical reaction to show that glucose contains aldehyde as carbonyl group.

[CBSE Sample Paper 2015]

Ans. (i) Keratin being a fibrous protein insoluble in water.



(iii) Glucose reduces Tollens' reagent to metallic silver.



Q. 3. (i) Give one structural difference between amylose and amylopectin

(ii) Name the protein and its shape present in oxygen carrier in human body.

(iii) Name two fat storing tissues in human body.

[CBSE Sample Paper 2014]


Ans. (i) Amylose is a long unbranched chain polymer of α -D(+) glucose.

Amylopectin is a branched chain polymer of α -D glucose.

(ii) Globular protein and its shape is spherical.

(iii) Liver and adipose tissue.

Q. 4. (i) A non-reducing disaccharide 'A' on hydrolysis with dilute acid gives an equimolar mixture of D-(+)-glucose and D-(-)-fructose.



Identify A. What is the mixture of D-(+)-glucose and D-(-)-fructose known as? Name the linkage that holds the two units in the disaccharide.

(ii) α -amino acids have relatively higher melting points than the corresponding halo acids. Explain.

[CBSE Sample Paper 2016] [HOTS]

Ans. (i) ● A = $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ (sucrose).

● Invert sugar.

● Glycosidic linkage.

(ii) α -Amino acids act as zwitter ions, ($\text{H}_3\text{N}^+-\text{CHR}-\text{COO}^-$) or dipolar ions. Due to this dipolar salt-like structure, they have strong dipole-dipole interactions. Therefore, their melting points are higher than the corresponding halo acids which do not exist as zwitter ions.


Q. 8. Give reasons for any 3 of the following observations:

- (i) Penta-acetate of glucose does not react with hydroxylamine.
- (ii) Amino acids behave like salts.
- (iii) Water soluble vitamins must be taken regularly in diet.
- (iv) The two strands in DNA are complimentary to each other.

[CBSE 2023 (56/5/2)]

Ans.

- (i) When glucose is reacted with acetic anhydride the —OH groups at C-1, C-2, C-3, C-4 and C-6 form a pentaacetate. As the pentaacetate of glucose does not contain a free —OH group at C-1, it cannot get hydrolyzed in aqueous solution to produce the open chain aldehydic form and thus glucose pentaacetate does not react with $\text{NH}_2\text{—OH}$ to form oxime.
- (ii) In aqueous solution, the carboxyl group loses a proton and amino group accepts a proton to form a zwitter ion.



- (iii) Water soluble vitamins must be supplied regularly in diet because they are readily excreted in urine and cannot be stored (except vitamin B12) in our body.
- (iv) The two strands in DNA molecule are held together through hydrogen bonds between purine base of one strand and pyrimidine base of the other and vice versa. Because of different sizes and geometries of the base, the only possible pairing in DNA are G (guanine) and C (cytosine) through three H-bonds (i.e., $\text{C}=\text{G}$) and between A (adenine) and T (thymine) through two H-bonds (i.e., $\text{A}=\text{T}$). Due to this base-pairing principle, the sequence of bases in one strand automatically fixes the sequence of bases in the other strand. Thus, the two strands are not identical but are complementary. (Any three)

Long Answer Questions

Each of the following questions are of 5 marks.

Q. 1. (i) (a) What is the difference between native protein and denatured protein?

(b) Which one of the following is a disaccharide?

Glucose, Lactose, Amylose, Fructose

(c) Write the name of the vitamin responsible for the coagulation of blood. [CBSE 2019 (56/4/3)]

(ii) Define the following terms:

(a) Native protein

[CBSE 2020 (56/3/2)]

(b) Nucleotide

[CBSE 2019 (56/2/2)]

Ans.

(i) (a) Protein found in a biological system with unique three-dimensional structure and biological activity is called native protein.

When a protein in its native form is subjected to change such as change in temperature, change in pH, its 2° and 3° structures are destroyed and it loses its biological activity. The protein thus formed is called denatured protein.

(b) Lactose

(c) Vitamin K.

(ii) (a) Protein found in biological system with unique three dimensional structure and biological activity is called native protein.

(b) A unit formed by the combination of nitrogenous base, pentose sugar and phosphate.

Questions for Practice

Choose and write the correct answer for each of the following.

- Which of the following reaction confirms the presence of carbonyl group ($>C=O$) in glucose?
(a) Reaction with HI
(b) Reaction with hydroxylamine
(c) Reaction with HCN
(d) Both (b) and (c)
- Amino acids are best represented as:
(a) dipolar ions
(b) isoelectric ions
(c) amphoteric ions
(d) zwitter ions
- Fructose reduces Tollens' reagent due to
(a) Primary alcoholic group
(b) Asymmetric carbons
(c) Secondary alcoholic groups
(d) Enolisation of fructose followed by conversion to aldehyde by base
- Which of the following reaction confirms the presence of carbonyl group ($>C=O$) in glucose?
(a) Reaction with HI
(b) Reaction with hydroxylamine
(c) Reaction with HCN
(d) Both (b) and (c)

In the following questions, two statements are given—one labeled Assertion (A) and the other labeled Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below:


- Both Assertion (A) and Reason (R) are correct statements, and Reason (R) is the correct explanation of the Assertion (A).
 - Both Assertion (A) and Reason (R) are correct statements, but Reason (R) is not the correct explanation of the Assertion (A).
 - Assertion (A) is correct, but Reason (R) is incorrect statement.
 - Assertion (A) is incorrect, but Reason (R) is correct statement.
- Assertion (A) : In a course of curdling of milk protein present in milk is denatured.
Reason (R) : Denaturation is caused by lactic acid formed from lactose by the bacteria present in milk.
 - Assertion (A) : Vitamin D can be stored in our body.
Reason (R) : Vitamin D is fat soluble vitamin.
 - Assertion (A) : Fructose is a ketohexose
Reason (R) : Fructose reduces Tollen's reagent.
 - Assertion (A) : Vitamin C is called ascorbic acid.
Reason (R) : It contains a $-COOH$ group.
 - Assertion (A) : Maltose is a reducing sugar, one molecule of which gives two molecules of D-glucose on hydrolysis.
Reason (R) : Maltose has a 1, 4 β -glycosidic linkage.

Answer the following questions:

- Write the reactions showing the presence of following in the open structure of glucose:
(i) five $-OH$ groups
(ii) a carbonyl group

[CBSE 2020 (56/4/3)]

11. (i) Give the reaction of glucose with acetic anhydride. Presence of which group is confirmed by this reaction? [CBSE 2020 (56/5/2)]
 (ii) What is essentially the difference between α -form of glucose and β -form of glucose? [CBSE Delhi 2011]
12. The Fischer projection of D-Glucose is given alongside.
 (i) Give the Fischer projection of L-Glucose.
 (ii) What happens when L-Glucose is treated with Tollens' reagent?



13. Define the following terms:
 (i) Polysaccharides
 (ii) Amino acids
 (iii) Enzymes [CBSE (F) 2014]
14. (i) Name the three major classes of carbohydrates and give an example of each of these classes.
 (ii) What type of linkage is responsible for the primary structure of proteins?
 (iii) Name the location where protein synthesis occurs in our body.
15. What is meant by the following:
 (i) Anomers
 (ii) Polypeptide
 (iii) Nucleotide
16. Define the following:
 (i) Denaturation of protein
 (ii) Inversion of sugar
 (iii) Essential amino acid
17. (i) Write about the following on protein synthesis:
 (a) Name the location where protein synthesis occurs.
 (b) How do 64 codons code for only 20 amino acids?
 (c) Which of the two bases of the codon are more important for coding?
 (ii) What deficiency diseases are caused due to lack of vitamins A, B₁, B₆ and K in human diet?
18. (i) Comment on the specificity of enzyme action. What is the most important reason for their specificity?
 (ii) (a) What are the products of hydrolysis of sucrose and lactose?
 (b) Name the vitamin whose deficiency causes pernicious anaemia.

[CBSE 2019 (56/2/2)]

Answers

1. (d) 2. (d) 3. (d) 4. (c) 5. (a) 6. (a) 7. (b) 8. (c) 9. (c)

