

Topic : Binomial Theorem

Type of Questions

M.M., Min.

Single choice Objective (no negative marking) Q.1 to 12

(3 marks, 3 min.)

[36, 36]

Multiple choice objective (no negative marking) Q.13, 14, 15

(5 marks, 4 min.)

[15, 12]

- In the expansion of $\left(x^3 - \frac{1}{x^2}\right)^n$, $n \in \mathbb{N}$, if the sum of the coefficients of x^5 and x^{10} is 0, then n is :
(A) 25 (B) 20 (C) 15 (D) None of these
- The sum of the coefficients of all the integral powers of x in the expansion of $(1 + 2\sqrt{x})^{40}$ is :
(A) $3^{40} + 1$ (B) $3^{40} - 1$ (C) $\frac{1}{2}(3^{40} - 1)$ (D) $\frac{1}{2}(3^{40} + 1)$
- The coefficient of the term independent of x in the expansion of $\left(\frac{x+1}{x^{\frac{2}{3}} - x^{\frac{1}{3}} + 1} - \frac{x-1}{x - x^{\frac{1}{2}}}\right)^{10}$ is :
(A) 70 (B) 112 (C) 105 (D) 210
- Coefficient of x^{n-1} in the expansion of, $(x+3)^n + (x+3)^{n-1}(x+2) + (x+3)^{n-2}(x+2)^2 + \dots + (x+2)^n$ is :
(A) ${}^{n+1}C_2(3)$ (B) ${}^{n-1}C_2(5)$ (C) ${}^{n+1}C_2(5)$ (D) ${}^nC_2(5)$
- Let $f(n) = 10^n + 3 \cdot 4^{n+2} + 5$, $n \in \mathbb{N}$. The greatest value of the integer which divides $f(n)$ for all n is :
(A) 27 (B) 9 (C) 3 (D) None of these
- If $\{x\}$ denotes the fractional part of ' x ', then $\left\{\frac{3^{1001}}{82}\right\} =$
(A) $9/82$ (B) $81/82$ (C) $3/82$ (D) $1/82$
- The sum $\sum_{r=0}^n (r+1) C_r^2$ is equal to :
(A) $\frac{(n+2)(2n-1)!}{n!(n-1)!}$ (B) $\frac{(n+2)(2n+1)!}{n!(n-1)!}$ (C) $\frac{(n+2)(2n+1)!}{n!(n+1)!}$ (D) $\frac{(n+2)(2n-1)!}{n!(n+1)!}$
- If $a_n = \sum_{r=0}^n \frac{1}{{}^nC_r}$, the value of $\sum_{r=0}^n \frac{n-2r}{{}^nC_r}$ is :
(A) $\frac{n}{2} a_n$ (B) $\frac{1}{4} a_n$ (C) na_n (D) 0
- The sum of the series $\sum_{r=1}^n (-1)^{r-1} \cdot {}^nC_r(a-r)$ is equal to :
(A) $n \cdot 2^{n-1} + a$ (B) 0 (C) a (D) None of these

10. The sum of: $3 \cdot {}^nC_0 - 8 \cdot {}^nC_1 + 13 \cdot {}^nC_2 - 18 \cdot {}^nC_3 + \dots$ upto $(n+1)$ terms is :
 (A) zero (B) 1 (C) 2 (D) none of these
11. The number of terms in the expansion of $\left(x^2 + 1 + \frac{1}{x^2}\right)^n$, $n \in \mathbb{N}$, is :
 (A) $2n$ (B) $3n$ (C) $2n + 1$ (D) $3n + 1$
12. If $(1 + x + 2x^2)^{20} = a_0 + a_1x + a_2x^2 + \dots + a_{40}x^{40}$, then $a_0 + a_2 + a_4 + \dots + a_{38}$ is equal to :
 (A) $2^{19}(2^{30} + 1)$ (B) $2^{19}(2^{20} - 1)$
 (C) $2^{20}(2^{19} - 1)$ (D) none of these
13. In the expansion of $\left(\sqrt[3]{4} + \frac{1}{\sqrt[4]{6}}\right)^{20}$
 (A) the number of irrational terms is 19 (B) middle term is irrational
 (C) the number of rational terms is 2 (D) 9th term is rational
14. If $(9 + \sqrt{80})^n = I + f$, where I, n are integers and $0 < f < 1$, then :
 (A) I is an odd integer (B) I is an even integer
 (C) $(I + f)(1 - f) = 1$ (D) $1 - f = (9 - \sqrt{80})^n$
15. $7^9 + 9^7$ is divisible by :
 (A) 16 (B) 24 (C) 64 (D) 72

Answers Key

1. (C) 2. (D) 3. (D) 4. (C)
 5. (B) 6. (C) 7. (A) 8. (D)
 9. (C) 10. (A) 11. (C) 12. (B)
 13. (A, B, C, D) 14. (A, C, D) 15. (A, C)