

Ordinary Thinking

Objective Questions

Atomic Structure

1. If in nature there may not be an element for which the principal quantum number $n > 4$, then the total possible number of elements will be

[IIT 1983; MP PET 1999; RPMT 1999; RPET 2001]

- (a) 60 (b) 32
(c) 4 (d) 64

2. In the Bohr's hydrogen atom model, the radius of the stationary orbit is directly proportional to ($n =$ principle quantum number) [MNR 1988; SCRA 1994;

CBSE PMT 1996; AIIMS 1999; DCE 2002]

- (a) n^{-1} (b) n
(c) n^{-2} (d) n^2

3. In the n orbit, the energy of an electron $E_n = -\frac{13.6}{n^2} eV$ for hydrogen atom. The energy required to take the electron from first orbit to second orbit will be

[MP PMT 1987; CPMT 1991, 97; RPMT 1999; DCE 2001; Kerala PMT 2004]

- (a) 10.2 eV (b) 12.1 eV
(c) 13.6 eV (d) 3.4 eV

4. In the following atoms and molecules for the transition from $n=2$ to $n=1$, the spectral line of minimum wavelength will be produced by [IIT 1983]

- (a) Hydrogen atom (b) Deuterium atom
(c) Uni-ionized helium (d) di-ionized lithium

5. The Lyman series of hydrogen spectrum lies in the region

[MNR 1993; MP PMT 1995; UPSEAT 2002]

- (a) Infrared (b) Visible
(c) Ultraviolet (d) Of X - rays

6. The size of an atom is of the order of

[CPMT 1990; MP PMT 1984; KCET 1994]

- (a) $10^{-8} m$ (b) $10^{-10} m$
(c) $10^{-12} m$ (d) $10^{-14} m$


7. Which one of the series of hydrogen spectrum is in the visible region [RPMT 1999; MP PET 1990; MP PMT 1994;

AFMC 1998; CBSE PMT 1990; MH CET 2004]

- (a) Lyman series (b) Balmer series
(c) Paschen series (d) Bracket series

8. The energy levels of the hydrogen spectrum is shown in figure. There are some transitions A, B, C, D and E. Transition A, B and C respectively represent

[CPMT 1986, 88]



- (a) First member of Lyman series, third spectral line of Balmer series and the second spectral line of Paschen series
(b) Ionization potential of hydrogen, second spectral line of Balmer series and third spectral line of Paschen series
(c) Series limit of Lyman series, third spectral line of Balmer series and second spectral line of Paschen series
(d) Series limit of Lyman series, second spectral line of Balmer series and third spectral line of Paschen series

9. In the above figure D and E respectively represent

[CPMT 1986, 88]

- (a) Absorption line of Balmer series and the ionization potential of hydrogen
(b) Absorption line of Balmer series and the wavelength lesser than lowest of the Lyman series
(c) Spectral line of Balmer series and the maximum wavelength of Lyman series
(d) Spectral line of Lyman series and the absorption of greater wavelength of limiting value of Paschen series

10. The Rutherford α -particle experiment shows that most of the α -particles pass through almost unscattered while some are scattered through large angles. What information does it give about the structure of the atom [AFMC 1997]

- (a) Atom is hollow
(b) The whole mass of the atom is concentrated in a small centre called nucleus
(c) Nucleus is positively charged
(d) All the above

11. Which of the following is true [MP PET 1993]

- (a) Lyman series is a continuous spectrum
(b) Paschen series is a line spectrum in the infrared
(c) Balmer series is a line spectrum in the ultraviolet
(d) The spectral series formula can be derived from the Rutherford model of the hydrogen atom

12. The energy required to knock out the electron in the third orbit of a hydrogen atom is equal to [DPMT 1987]

- (a) 13.6 eV (b) $+\frac{13.6}{9} eV$
(c) $-\frac{13.6}{3} eV$ (d) $-\frac{3}{13.6} eV$

13. An electron has a mass of $9.1 \times 10^{-31} kg$. It revolves round the nucleus in a circular orbit of radius $0.529 \times 10^{-10} metre$ at a speed of $2.2 \times 10^6 m/s$. The magnitude of its linear momentum in this motion is

[AFMC 1988]

[Roorkee 1993]

- (a) $1.1 \times 10^{-34} \text{ kg} - \text{m} / \text{s}$ (b) $2.0 \times 10^{-24} \text{ kg} - \text{m} / \text{s}$
 (c) $4.0 \times 10^{-24} \text{ kg} - \text{m} / \text{s}$ (d) $4.0 \times 10^{-31} \text{ kg} - \text{m} / \text{s}$

14. In a beryllium atom, if a_0 be the radius of the first orbit, then the radius of the second orbit will be in general

[CBSE PMT 1992; Roorkee 1993; BHU 1998]

- (a) na_0 (b) a_0
 (c) $n^2 a_0$ (d) $\frac{a_0}{n^2}$

15. The ionization potential for second He electron is

- (a) 13.6 eV (b) 27.2 eV
 (c) 54.4 eV (d) 100 eV

16. The energy required to remove an electron in a hydrogen atom from $n = 10$ state is [MP PMT 1993]

- (a) 13.6 eV (b) 1.36 eV
 (c) 0.136 eV (d) 0.0136 eV

17. Every series of hydrogen spectrum has an upper and lower limit in wavelength. The spectral series which has an upper limit of wavelength equal to 18752 \AA is [MP PMT 1993]

- (a) Balmer series (b) Lyman series
 (c) Paschen series (d) Pfund series

(Rydberg constant $R = 1.097 \times 10^7 \text{ per metre}$)

18. The kinetic energy of the electron in an orbit of radius r in hydrogen atom is ($e =$ electronic charge) [MP PMT 1987]

- (a) $\frac{e^2}{r^2}$ (b) $\frac{e^2}{2r}$
 (c) $\frac{e^2}{r}$ (d) $\frac{e^2}{2r^2}$

19. Ionization potential of hydrogen atom is 13.6 V.


Hydrogen atoms in the ground state are excited by monochromatic radiation of photon energy 12.1 eV. The spectral lines emitted by hydrogen atoms according to Bohr's theory will be

[CBSE PMT 1996; MP PMT 1999; AMU (Med.) 2002]

- (a) One (b) Two
 (c) Three (d) Four

20. Energy levels A, B, C of a certain atom corresponding to increasing values of energy i.e. $E_A < E_B < E_C$. If $\lambda_1, \lambda_2, \lambda_3$ are the wavelengths of radiations corresponding to the transitions C to B, B to A and C to A respectively, which of the following statements is correct

[AIIMS 1995; CBSE PMT 1990, 2005]



- (a) $\lambda_3 = \lambda_1 + \lambda_2$ (b) $\lambda_3 = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$
 (c) $\lambda_1 + \lambda_2 + \lambda_3 = 0$ (d) $\lambda_3^2 = \lambda_1^2 + \lambda_2^2$

21. The angular momentum of electron in n orbit is given by

- (a) nh (b) $\frac{h}{2\pi m}$
 (c) $n \frac{h}{2\pi}$ (d) $n^2 \frac{h}{2\pi}$

22. The ratio of the energies of the hydrogen atom in its first to second excited state is [CPMT 1978]

- (a) 1/4 (b) 4/9
 (c) 9/4 (d) 4

23. An electron jumps from the 4th orbit to the 2nd orbit of hydrogen atom. Given the Rydberg's constant $R = 10^5 \text{ cm}^{-1}$. The frequency in Hz of the emitted radiation will be

- (a) $\frac{3}{16} \times 10^5$ (b) $\frac{3}{16} \times 10^{15}$
 (c) $\frac{9}{16} \times 10^{15}$ (d) $\frac{3}{4} \times 10^{15}$

24. The ionisation potential of hydrogen atom is 13.6 volt. The energy required to remove an electron in the $n = 2$ state of the hydrogen atom is [NCERT 1983; MP PET 2005]

- (a) 27.2 eV (b) 13.6 eV
 (c) 6.8 eV (d) 3.4 eV

25. The ionisation energy of 10 times ionised sodium atom is


[DPMT 1991]

- (a) 13.6 eV (b) $13.6 \times 11 \text{ eV}$
 (c) $\frac{13.6}{11} \text{ eV}$ (d) $13.6 \times (11)^2 \text{ eV}$

26. If the wavelength of the first line of the Balmer series of hydrogen is 6561 \AA , the wavelength of the second line of the series should be [CPMT 1984; DPMT 2004]

- (a) 13122 \AA (b) 3280 \AA
 (c) 4860 \AA (d) 2187 \AA

27. The following diagram indicates the energy levels of a certain atom when the system moves from $2E$ level to E , a photon of wavelength λ is emitted. The wavelength of photon produced during its transition from $\frac{4E}{3}$ level to E is




[CPMT 1989]


- (a) $\lambda/3$ (b) $3\lambda/4$
 (c) $4\lambda/3$ (d) 3λ

28. A beam of fast moving alpha particles were directed towards a thin film of gold. The parts A', B' and C' of the transmitted and reflected beams corresponding to the incident parts A, B and C of the beam, are shown in the adjoining diagram. The number of alpha particles in

[CPMT 1986, 88; RPET 2000]



- (a) B' will be minimum and in C' maximum
 (b) A' will be maximum and in B' minimum
 (c) A' will be minimum and in B' maximum
 (d) C' will be minimum and in B' maximum
29. According to Bohr's theory the radius of electron in an orbit described by principal quantum number n and atomic number Z is proportional to [CPMT 1988]
- (a) $Z^2 n^2$ (b) $\frac{Z^2}{n^2}$
 (c) $\frac{Z^2}{n}$ (d) $\frac{n^2}{Z}$
30. The radius of electron's second stationary orbit in Bohr's atom is R . The radius of the third orbit will be [EAMCET 1992; DPMT 1999]
- (a) $3R$ (b) $2.25R$
 (c) $9R$ (d) $\frac{R}{3}$
31. If m is mass of electron, v its velocity, r the radius of stationary circular orbit around a nucleus with charge Ze , then from Bohr's first postulate, the kinetic energy $K = \frac{1}{2}mv^2$ of the electron in C.G.S. system is equal to [NCERT 1977]
- (a) $\frac{1}{2} \frac{Ze^2}{r}$ (b) $\frac{1}{2} \frac{Ze^2}{r^2}$
 (c) $\frac{Ze^2}{r}$ (d) $\frac{Ze}{r^2}$
32. Consider an electron in the n orbit of a hydrogen atom in the Bohr model. The circumference of the orbit can be expressed in terms of the de Broglie wavelength λ of that electron as
- (a) $(0.259)n\lambda$ (b) $\sqrt{n}\lambda$
 (c) $(13.6)\lambda$ (d) $n\lambda$
33. In any Bohr orbit of the hydrogen atom, the ratio of kinetic energy to potential energy of the electron is [MP PET 1994]
- (a) $1/2$ (b) 2
 (c) $-1/2$ (d) -2
34. The spectral series of the hydrogen spectrum that lies in the ultraviolet region is the [CPMT 1990; MP PET 1994; MP PMT 2000]
- (a) Balmer series (b) Pfund series
 (c) Paschen series (d) Lyman series
35. Figure shows the energy levels P, Q, R, S and G of an atom where G is the ground state. A red line in the emission spectrum of the atom can be obtained by an energy level change from Q to S . A blue line can be obtained by following energy level change




- (a) P to Q (b) Q to R
 (c) R to S (d) R to G

36. A hydrogen atom (ionisation potential 13.6 eV) makes a transition from third excited state to first excited state. The energy of the photon emitted in the process is [MNR 1995]

- (a) 1.89 eV (b) 2.55 eV
 (c) 12.09 eV (d) 12.75 eV

37. The figure indicates the energy level diagram of an atom and the origin of six spectral lines in emission (e.g. line no. 5 arises from the transition from level B to A). The following spectral lines will also occur in the absorption spectrum [CBSE PMT 1995]



- (a) 1, 4, 6 (b) 4, 5, 6
 (c) 1, 2, 3 (d) 1, 2, 3, 4, 5, 6

38. When a hydrogen atom is raised from the ground state to an excited state [CBSE PMT 1995; AMU (Med.) 1999]

- (a) P.E. increases and K.E. decreases
 (b) P.E. decreases and K.E. increases
 (c) Both kinetic energy and potential energy increase
 (d) Both K.E. and P.E. decrease

39. An electron makes a transition from orbit $n = 4$ to the orbit $n = 2$ of a hydrogen atom. The wave number of the emitted radiations ($R = \text{Rydberg's constant}$) will be [CBSE PMT 1990]

- (a) $\frac{16}{3R}$ (b) $\frac{2R}{16}$
 (c) $\frac{3R}{16}$ (d) $\frac{4R}{16}$

40. In Bohr model of the hydrogen atom, the lowest orbit corresponds to [Manipal MEE 1995]

- (a) Infinite energy (b) The maximum energy
 (c) The minimum energy (d) Zero energy

41. The ratio of the kinetic energy to the total energy of an electron in a Bohr orbit is [Roorkee 1995; BHU 2002]

- (a) -1 (b) 2
 (c) $1:2$ (d) None of these

42. An electron in the $n = 1$ orbit of hydrogen atom is bound by 13.6 eV . If a hydrogen atom is in the $n = 3$ state, how much energy is required to ionize it [MP PMT 1994]

- (a) 13.6 eV (b) 4.53 eV
 (c) 3.4 eV (d) 1.51 eV

43. Which of the following statements about the Bohr model of the hydrogen atom is false [MP PMT 1995]
- (a) Acceleration of electron in $n = 2$ orbit is less than that in $n = 1$ orbit
 (b) Angular momentum of electron in $n = 2$ orbit is more than that in $n = 1$ orbit
 (c) Kinetic energy of electron in $n = 2$ orbit is less than that in $n = 1$ orbit
 (d) Potential energy of electron in $n = 2$ orbit is less than that in $n = 1$ orbit
44. If an electron jumps from 1st orbital to 3rd orbital, then it will
- (a) Absorb energy (b) Release energy
 (c) No gain of energy (d) None of these
45. The ratio of the frequencies of the long wavelength limits of Lyman and Balmer series of hydrogen spectrum is [KCEE 1996]
- (a) 27 : 5 (b) 5 : 27
 (c) 4 : 1 (d) 1 : 4
46. Which of the following transitions in a hydrogen atom emits photon of the highest frequency [MP PET 1996; DPMT 2001]
- (a) $n = 1$ to $n = 2$ (b) $n = 2$ to $n = 1$
 (c) $n = 2$ to $n = 6$ (d) $n = 6$ to $n = 2$
47. In terms of Rydberg's constant R , the wave number of the first Balmer line is [MP PMT 1996]
- (a) R (b) $3R$
 (c) $\frac{5R}{36}$ (d) $\frac{8R}{9}$
48. If the ionisation potential of helium atom is 24.6 volt, the energy required to ionise it will be [MP PMT 1996]
- (a) 24.6 eV (b) 24.6 V
 (c) 13.6 V (d) 13.6 eV
49. Which of the transitions in hydrogen atom emits a photon of lowest frequency ($n =$ quantum number) [BHU 1999]
- (a) $n = 2$ to $n = 1$ (b) $n = 4$ to $n = 3$
 (c) $n = 3$ to $n = 1$ (d) $n = 4$ to $n = 2$
50. According to Bohr's theory, the expressions for the kinetic and potential energy of an electron revolving in an orbit is given respectively by
- (a) $+\frac{e^2}{8\pi\epsilon_0 r}$ and $-\frac{e^2}{4\pi\epsilon_0 r}$ (b) $+\frac{8\pi\epsilon_0 e^2}{r}$ and $-\frac{4\pi\epsilon_0 e^2}{r}$
 (c) $-\frac{e^2}{8\pi\epsilon_0 r}$ and $-\frac{e^2}{4\pi\epsilon_0 r}$ (d) $+\frac{e^2}{8\pi\epsilon_0 r}$ and $+\frac{e^2}{4\pi\epsilon_0 r}$
51. In a hydrogen atom, which of the following electronic transitions would involve the maximum energy change [MP PET 1997]
- (a) From $n = 2$ to $n = 1$ (b) From $n = 3$ to $n = 1$
 (c) From $n = 4$ to $n = 2$ (d) From $n = 3$ to $n = 2$
52. In the lowest energy level of hydrogen atom, the electron has the angular momentum [MP PET 1997; BCECE 2003]
- (a) π/h (b) h/π
 (c) $h/2\pi$ (d) $2\pi/h$
53. The minimum energy required to excite a hydrogen atom from its ground state is [EAMCET (Engg.) 1995; MP PMT 1997; CPMT 1999; DCE 1999]
- (a) 13.6 eV (b) -13.6 eV
 (c) 3.4 eV (d) 10.2 eV
54. Ratio of the wavelengths of first line of Lyman series and first line of Balmer series is [AFMC 1996]
- (a) 1 : 3 (b) 27 : 5
 (c) 5 : 27 (d) 4 : 9
55. The Rydberg constant R for hydrogen is [MP PMT/PET 1998]
- (a) $R = -\left(\frac{1}{4\pi\epsilon_0}\right) \cdot \frac{2\pi^2 m e^2}{ch^2}$ (b) $R = \left(\frac{1}{4\pi\epsilon_0}\right) \cdot \frac{2\pi^2 m e^4}{ch^2}$
 (c) $R = \left(\frac{1}{4\pi\epsilon_0}\right)^2 \cdot \frac{2\pi^2 m e^4}{c^2 h^2}$ (d) $R = \left(\frac{1}{4\pi\epsilon_0}\right)^2 \cdot \frac{2\pi^2 m e^4}{ch^3}$
56. The wavelength of the first line of Balmer series is 6563 Å. The Rydberg constant for hydrogen is about [MP PMT/PET 1998]
- (a) 1.09×10^7 per m (b) 1.09×10^8 per m
 (c) 1.09×10^9 per m (d) 1.09×10^5 per m
57. According to Bohr's theory the moment of momentum of an electron revolving in second orbit of hydrogen atom will be [MP PET 1999; KCET 2003]
- (a) $2\pi h$ (b) πh
 (c) $\frac{h}{\pi}$ (d) $\frac{2h}{\pi}$
58. The velocity of an electron in the second orbit of sodium atom (atomic number = 11) is v . The velocity of an electron in its fifth orbit will be [MP PET 1999]
- (a) v (b) $\frac{22}{5}v$
 (c) $\frac{5}{2}v$ (d) $\frac{2}{5}v$
59. The absorption transitions between the first and the fourth energy states of hydrogen atom are 3. The emission transitions between these states will be [MP PET 1999]
- (a) 3 (b) 4
 (c) 5 (d) 6
60. The ratio of longest wavelength and the shortest wavelength observed in the five spectral series of emission spectrum of hydrogen is [MP PET 1999]
- (a) $\frac{4}{3}$ (b) $\frac{525}{376}$
 (c) 25 (d) $\frac{900}{11}$

1466 Atomic and Nuclear Physics



61. In the Bohr model of a hydrogen atom, the centripetal force is furnished by the coulomb attraction between the proton and the electron. If a_0 is the radius of the ground state orbit, m is the mass, e is the charge on the electron and ϵ_0 is the vacuum permittivity, the speed of the electron is [RPET 1997]
- [CBSE PMT 1998]
- (a) 0 (b) $\frac{e}{\sqrt{\epsilon_0 a_0 m}}$
- (c) $\frac{e}{\sqrt{4\pi\epsilon_0 a_0 m}}$ (d) $\frac{\sqrt{4\pi\epsilon_0 a_0 m}}{e}$
62. The electron in a hydrogen atom makes a transition $n_1 \rightarrow n_2$, where n_1 and n_2 are the principal quantum numbers of the two states. Assume the Bohr model to be valid. The time period of the electron in the initial state is eight times that in the final state. The possible values of n and n are [IIT 1998; KCET 2005]
- (a) $n_1 = 4, n_2 = 2$ (b) $n_1 = 8, n_2 = 2$
- (c) $n_1 = 8, n_2 = 1$ (d) $n_1 = 6, n_2 = 3$
63. As per Bohr model, the minimum energy (in eV) required to remove an electron from the ground state of doubly ionized Li atom ($Z = 3$) is [IIT 1997 Re-Exam; MH CET 2000]
- (a) 1.51 (b) 13.6
- (c) 40.8 (d) 122.4
64. Which one of these is non-divisible [KCET 1994]
- (a) Nucleus (b) Photon
- (c) Proton (d) Atom
65. In Bohr's model of hydrogen atom, let PE represents potential energy and TE the total energy. In going to a higher level
- (a) PE decreases, TE increases
- (b) PE increases, TE increases
- (c) PE decreases, TE decreases
- (d) PE increases, TE decreases
66. According to Bohr's model, the radius of the second orbit of helium atom is [Bihar MEE 1995]
- (a) 0.53 Å (b) 1.06 Å
- (c) 2.12 Å (d) 0.265 Å
67. The fact that photons carry energy was established by [ISM Dhanbad 1994]
- (a) Doppler's effect (b) Compton's effect
- (c) Bohr's theory (d) Diffraction of light
68. An ionic atom equivalent to hydrogen atom has wavelength equal to 1/4 of the wavelengths of hydrogen lines. The ion will be
- (a) He^+ (b) Li^{++}
- (c) Ne^{9+} (d) Na^{10+}
69. The extreme wavelengths of Paschen series are
- (a) 0.365 μm and 0.565 μm (b) 0.818 μm and 1.89 μm
- (c) 1.45 μm and 4.04 μm (d) 2.27 μm and 7.43 μm
70. The third line of Balmer series of an ion equivalent to hydrogen atom has wavelength of 108.5 nm. The ground state energy of an electron of this ion will be [RPET 1997]
- (a) 3.4 eV (b) 13.6 eV
- (c) 54.4 eV (d) 122.4 eV
71. An electron in the $n = 1$ orbit of hydrogen atom is bound by 13.6 eV energy is required to ionize it is [MP PMT 2003]
- (a) 13.6 eV (b) 6.53 eV
- (c) 5.4 eV (d) 1.51 eV
72. Ionization energy of hydrogen is 13.6 eV. If $h = 6.6 \times 10^{-34} \text{ J-sec}$, the value of R will be of the order of [RPMT 1997]
- (a) 10^{10} m^{-1} (b) 10^7 m^{-1}
- (c) 10^4 m^{-1} (d) 10^{-7} m^{-1}
73. To explain his theory, Bohr used [CBSE PMT 1993; MP PET 2002]
- (a) Conservation of linear momentum
- (b) Conservation of angular momentum
- (c) Conservation of quantum frequency
- (d) Conservation of energy
74. The ionisation energy of hydrogen atom is 13.6 eV. Following Bohr's theory, the energy corresponding to a transition between the 3rd and the 4th orbit is [CBSE PMT 1992; DPMT 2000; RPMT 1999; AMU (Med.) 2001]
- [KCET 1994]
- (a) 3.40 eV (b) 1.51 eV
- (c) 0.85 eV (d) 0.66 eV
75. Hydrogen atoms are excited from ground state of the principal quantum number 4. Then the number of spectral lines observed will be [CBSE PMT 1993]
- (a) 3 (b) 6
- (c) 5 (d) 2
76. Hydrogen atom emits blue light when it changes from $n = 4$ energy level to the $n = 2$ level. Which colour of light would the atom emit when it changes from the $n = 5$ level to the $n = 2$ level
- (a) Red (b) Yellow
- (c) Green (d) Violet
77. In Rutherford scattering experiment, what will be the correct angle for α scattering for an impact parameter $b = 0$ [CBSE PMT 1994; JIPMER 2000]
- [RPET 1997]
- (a) 90° (b) 270°
- (c) 0° (d) 180°
78. The radius of hydrogen atom in its ground state is $5.3 \times 10^{-11} \text{ m}$. After collision with an electron it is found to have a radius of

- $21.2 \times 10^{-11} m$. What is the principal quantum number n of the final state of the atom
[CBSE PMT 1994; CPMT 2001; MH CET 2000]
- (a) $n = 4$ (b) $n = 2$
(c) $n = 16$ (d) $n = 3$
79. The splitting of line into groups under the effect of electric or magnetic field is called [AFMC 1995]
- (a) Zeeman's effect (b) Bohr's effect
(c) Heisenberg's effect (d) Magnetic effect
80. The energy of a hydrogen atom in its ground state is $-13.6 eV$. The energy of the level corresponding to the quantum number $n = 2$ (first excited state) in the hydrogen atom is [CBSE PMT 1996; CBSE PMT 1997, 2001; MP PET 2000; AFMC 2000, 01, 02; BCECE 2003]
- (a) $-2.72 eV$ (b) $-0.85 eV$
(c) $-0.54 eV$ (d) $-3.4 eV$
81. The first line of Balmer series has wavelength 6563 \AA . What will be the wavelength of the first member of Lyman series [RPMT 1996]
- (a) 1215.4 \AA (b) 2500 \AA
(c) 7500 \AA (d) 600 \AA
82. The wavelength of Lyman series is [BHU 1997]
- (a) $\frac{4}{3 \times 10967} cm$ (b) $\frac{3}{4 \times 10967} cm$
(c) $\frac{4 \times 10967}{3} cm$ (d) $\frac{3}{4} \times 10967 cm$
83. When hydrogen atom is in its first excited level, its radius is its ground state radius [CBSE PMT 1997]
- (a) Half (b) Same
(c) Twice (d) Four times
84. Hydrogen atom excites energy level from fundamental state to $n = 3$. Number of spectrum lines according to Bohr, is [CPMT 1997]
- (a) 4 (b) 3
(c) 1 (d) 2
85. Number of spectral lines in hydrogen atom is [CPMT 1997]
- (a) 3 (b) 6
(c) 15 (d) Infinite
86. In Bohr's model, the atomic radius of the first orbit is r_0 , then the radius of the third orbit is [AIIMS 1997; CPMT 2001; KCET (Engg./Med.) 1999; Pb. PMT 2004]
- (a) $\frac{r_0}{9}$ (b) r_0
(c) $9r_0$ (d) $3r_0$
87. The wavelength of the energy emitted when electron come from fourth orbit to second orbit in hydrogen is $20.397 cm$. The wavelength of energy for the same transition in He^+ is [AIIMS 1997; JIPMER 2000]
- (a) $5.099 cm^{-1}$ (b) $20.497 cm^{-1}$
(c) $40.994 cm^{-1}$ (d) $81.988 cm^{-1}$
88. Minimum excitation potential of Bohr's first orbit in hydrogen atom is [BHU 1998; JIPMER 2001, 02; Pb. PMT 2004]
- (a) $13.6 V$ (b) $3.4 V$
(c) $10.2 V$ (d) $3.6 V$
89. Which of the following statements are true regarding Bohr's model of hydrogen atom
- (I) Orbiting speed of electron decreases as it shifts to discrete orbits away from the nucleus
(II) Radii of allowed orbits of electron are proportional to the principal quantum number
(III) Frequency with which electrons orbits around the nucleus in discrete orbits is inversely proportional to the principal quantum number
(IV) Binding force with which the electron is bound to the nucleus increases as it shifts to outer orbits
- Select correct answer using the codes given below
- Codes : [SCRA 1998]
- (a) I and III (b) II and IV
(c) I, II and III (d) II, III and IV
90. The wavelength of radiation emitted is λ_0 when an electron jumps from the third to the second orbit of hydrogen atom. For the electron jump from the fourth to the second orbit of the hydrogen atom, the wavelength of radiation emitted will be
- (a) $\frac{16}{25} \lambda_0$ (b) $\frac{20}{27} \lambda_0$
(c) $\frac{27}{20} \lambda_0$ (d) $\frac{25}{16} \lambda_0$
91. For electron moving in n orbit of H -atom the angular velocity is proportional to [RPET 1999]
- (a) n (b) $1/n$
(c) n (d) $1/n$
92. The energy of electron in first excited state of H -atom is $-3.4 eV$ its kinetic energy is [RPET 1999; CBSE PMT 2005]
- (a) $-3.4 eV$ (b) $+3.4 eV$
(c) $-6.8 eV$ (d) $6.8 eV$
93. The energy required to excite an electron from the ground state of hydrogen atom to the first excited state, is [Pb. PMT 1999]
- (a) $1.602 \times 10^{-14} J$ (b) $1.619 \times 10^{-16} J$
(c) $1.632 \times 10^{-18} J$ (d) $1.656 \times 10^{-20} J$
94. Which of the following phenomena suggests the presence of electron energy levels in atoms [JIPMER 1999]
- (a) Radio active decay
(b) Isotopes
(c) Spectral lines
(d) α -particles scattering
95. Which of the following spectral series in hydrogen atom give spectral line of 4860 \AA [Roorkee 1999]

1468 Atomic and Nuclear Physics

- (a) Lyman (b) Balmer
(c) Paschen (d) Brackett
96. If scattering particles are 56° for 90° angle then this will be at 60° angle [RPMT 2000]
(a) 224 (b) 256
(c) 98 (d) 108
97. When an electron in hydrogen atom is excited, from its 4th to 5th stationary orbit, the change in angular momentum of electron is (Planck's constant: $h = 6.6 \times 10^{-34} \text{ J-s}$) [AFMC 2000; Pb. PET 2001]
(a) $4.16 \times 10^{-34} \text{ J-s}$ (b) $3.32 \times 10^{-34} \text{ J-s}$
(c) $1.05 \times 10^{-34} \text{ J-s}$ (d) $2.08 \times 10^{-34} \text{ J-s}$
98. Energy of electron in a orbit of H-atom is [RPET 2000]
(a) Positive (b) Negative
(c) Zero (d) Nothing can be said
99. The concept of stationary orbits was proposed by [Pb. PMT 2000]
(a) Neil Bohr (b) J.J. Thomson
(c) Rutherford (d) I. Newton
100. In a hydrogen atom, the distance between the electron and proton is $2.5 \times 10^{-11} \text{ m}$. The electrical force of attraction between them will be [Pb. PMT 2000]
(a) $2.8 \times 10^{-7} \text{ N}$ (b) $3.7 \times 10^{-7} \text{ N}$
(c) $6.2 \times 10^{-7} \text{ N}$ (d) $9.1 \times 10^{-7} \text{ N}$
101. If λ_{max} is 6563 Å, then wave length of second line for Balmer series will be [RPMT 2000]
(a) $\lambda = \frac{16}{3R}$ (b) $\lambda = \frac{36}{5R}$
(c) $\lambda = \frac{4}{3R}$ (d) None of the above
102. What will be the angular momentum of a electron, if energy of this electron in H-atom is 1.5 eV (in J-sec) [RPMT 2000]
(a) 1.05×10^{-34} (b) 2.1×10^{-34}
(c) 3.15×10^{-34} (d) -2.1×10^{-34}
103. Who discovered spin quantum number [RPMT 2000]
(a) Unlenbeck and Goudsmit
(b) Nell's Bohr
(c) Zeeman
(d) Sommerfield
104. The time of revolution of an electron around a nucleus of charge Ze in n Bohr orbit is directly proportional to [MP PET 2003]
(a) n (b) $\frac{n^3}{Z^2}$
- (c) $\frac{n^2}{Z}$ (d) $\frac{Z}{n}$
105. In Bohr's model, if the atomic radius of the first orbit is r_0 , then the radius of the fourth orbit is [CBSE PMT 2000]
(a) r_0 (b) $4r_0$
(c) $r_0/16$ (d) $16r_0$
106. If R is the Rydberg's constant for hydrogen the wave number of the first line in the Lyman series will be [KCET 2000]
(a) $\frac{R}{4}$ (b) $\frac{3R}{4}$
(c) $\frac{R}{2}$ (d) 2R
107. In hydrogen atom, if the difference in the energy of the electron in $n = 2$ and $n = 3$ orbits is E, the ionization energy of hydrogen atom is [EAMCET (Med.) 2000]
(a) 13.2 E (b) 7.2 E
(c) 5.6 E (d) 3.2 E
108. The first member of the Paschen series in hydrogen spectrum is of wavelength 18,800 Å. The short wavelengths limit of Paschen series is [EAMCET (Med.) 2000]
(a) 1215 Å (b) 6560 Å
(c) 8225 Å (d) 12850 Å
109. The ratio of the largest to shortest wavelengths in Lyman series of hydrogen spectra is [EAMCET (Med.) 2000]
(a) $\frac{25}{9}$ (b) $\frac{17}{6}$
(c) $\frac{9}{5}$ (d) $\frac{4}{3}$
110. In Bohr model of hydrogen atom, the ratio of periods of revolution of an electron in $n = 2$ and $n = 1$ orbits is [EAMCET (Engg.) 2000]
(a) 2 : 1 (b) 4 : 1
(c) 8 : 1 (d) 16 : 1
111. The ratio of the longest to shortest wavelengths in Brackett series of hydrogen spectra is [EAMCET (Engg.) 2000]
(a) $\frac{25}{9}$ (b) $\frac{17}{6}$
(c) $\frac{9}{5}$ (d) $\frac{4}{3}$
112. The electron in a hydrogen atom makes a transition from an excited state to the ground state. Which of the following statements is true
(a) Its kinetic energy increases and its potential and total energies decrease
(b) Its kinetic energy decreases, potential energy increases and its total energy remains the same

- (c) Its kinetic and total energies decrease and its potential energy increases
 (d) Its kinetic, potential and total energies decreases
113. The ratio of minimum to maximum wavelength in Balmer series is
 (a) 5 : 9 (b) 5 : 36
 (c) 1 : 4 (d) 3 : 4
114. The radius of the Bohr orbit in the ground state of hydrogen atom is 0.5 \AA . The radius of the orbit of the electron in the third excited state of He^+ will be [MP PMT 2000]
 (a) 8 \AA (b) 4 \AA
 (c) 0.5 \AA (d) 0.25 \AA
115. The ratio of the speed of the electron in the first Bohr orbit of hydrogen and the speed of light is equal to (where e , h and c have their usual meanings) [MP PMT 2000]
 (a) $2\pi hc/e^2$ (b) $e^2 h/2\pi c$
 (c) $e^2 c/2\pi h$ (d) $2\pi e^2/hc$
116. According to the Rutherford's atomic model, the electrons inside the atom are [KCET (Med.) 2000]
 (a) Stationary (b) Not stationary
 (c) Centralized (d) None of these
117. The energy of hydrogen atom in its ground state is -13.6 eV . The energy of the level corresponding to the quantum number n is equal to 5 is [KCET (Engg./Med.) 2001]
 (a) -5.40 eV (b) -2.72 eV
 (c) -0.85 eV (d) -0.54 eV
118. According to classical theory, the circular path of an electron in Rutherford atom is [BHU 2001]
 (a) Spiral (b) Circular
 (c) Parabolic (d) Straight line
119. Rutherford's α -particle experiment showed that the atoms have
 (a) Proton (b) Nucleus
 (c) Neutron (d) Electrons
120. Orbital acceleration of electron is [RPET 2001]
 (a) $\frac{n^2 h^2}{4\pi^2 m^2 r^3}$ (b) $\frac{n^2 h^2}{2\pi^2 r^3}$
 (c) $\frac{4n^2 h^2}{\pi^2 m^2 r^3}$ (d) $\frac{4n^2 h^2}{4\pi^2 m^2 r^3}$
121. Which of the following is true for number of spectral lines in going from Lyman series to Pfund series [RPET 2001]
 (a) Increases
 (b) Decreases
 (c) Unchanged
 (d) May decrease or increase
122. The wavelength of yellow line of sodium is 5896 \AA . Its wave number will be [MP PET 2001]
 (a) $50883 \times 10^3 \text{ per second}$
 (b) 16961 per cm
 (c) 17581 per cm

- (d) 50883 per cm
123. Radius of the first orbit of the electron in a hydrogen atom is 0.53 \AA . So, the radius of the third orbit will be [Kerala (Engg.) 2001]
 [MP PET 2000]
 (a) 2.12 \AA (b) 4.77 \AA
 (c) 1.06 \AA (d) 1.59 \AA
124. The first line in the Lyman series has wavelength λ . The wavelength of the first line in Balmer series is [MH CET (Med.) 2001]
 (a) $\frac{2}{9} \lambda$ (b) $\frac{9}{2} \lambda$
 (c) $\frac{5}{27} \lambda$ (d) $\frac{27}{5} \lambda$
125. In hydrogen atom which quantity is integral multiple of $\frac{h}{2\pi}$ [DCE 2001]
 (a) Angular momentum (b) Angular velocity
 (c) Angular acceleration (d) Momentum
126. In the following transitions, which one has higher frequency [UPSEAT 2001]
 (a) $3 - 2$ (b) $4 - 3$
 (c) $4 - 2$ (d) $3 - 1$
127. The diagram shows the path of four α -particles of the same energy being scattered by the nucleus of an atom simultaneously. Which of these are/is not physically possible [AMU (Med.) 2001]
- 
- (a) 3 and 4 (b) 2 and 3
 (c) 1 and 4 (d) 4 only
128. An electron jumps from 5th orbit to 4th orbit of hydrogen atom. Taking the Rydberg constant as 10^7 per metre . What will be the frequency of radiation emitted [AFMC 2001] [Pb. PMT 2001]
 (a) $6.75 \times 10^{12} \text{ Hz}$ (b) $6.75 \times 10^{14} \text{ Hz}$
 (c) $6.75 \times 10^{13} \text{ Hz}$ (d) None of these
129. For principal quantum number $n = 3$, the possible values of orbital quantum number ' l ' are [MP PET 2001; MP PMT 2001]
 (a) 1, 2, 3 (b) 0, 1, 2, 3
 (c) 0, 1, 2 (d) -1, 0, +1
130. Four lowest energy levels of H-atom are shown in the figure. The number of possible emission lines would be [MP PMT 2001]
- 

1470 Atomic and Nuclear Physics

- (a) 3 (b) 4
(c) 5 (d) 6
- 131.** The order of the size of nucleus and Bohr radius of an atom respectively are [MP PET 2001; MP PMT 2001]
(a) $10^{-14} m, 10^{-10} m$ (b) $10^{-10} m, 10^{-8} m$
(c) $10^{-20} m, 10^{-16} m$ (d) $10^{-8} m, 10^{-6} m$
- 132.** Energy of an electron in an excited hydrogen atom is $-3.4 eV$. Its angular momentum will be: $h = 6.626 \times 10^{-34} J - s$ [UPSEAT 1999; Kerala PET 2002]
(a) $1.11 \times 10^{34} J \text{ sec}$ (b) $1.51 \times 10^{-31} J \text{ sec}$
(c) $2.11 \times 10^{-34} J \text{ sec}$ (d) $3.72 \times 10^{-34} J \text{ sec}$
- 133.** The ratio of the wavelengths for $2 \rightarrow 1$ transition in *Li*, *He* and *H* is
(a) 1 : 2 : 3 (b) 1 : 4 : 9
(c) 4 : 9 : 36 (d) 3 : 2 : 1
- 134.** The wavelength of light emitted from second orbit to first orbits in a hydrogen atom is [Pb. PMT 2002]
(a) $1.215 \times 10^{-7} m$ (b) $1.215 \times 10^{-5} m$
(c) $1.215 \times 10^{-4} m$ (d) $1.215 \times 10^{-3} m$
- 135.** Energy of the electron in *n* orbit of hydrogen atom is given by $E_n = -\frac{13.6}{n^2} eV$. The amount of energy needed to transfer electron from first orbit to third orbit is [MH CET 2002; Kerala PMT 2002]
(a) 13.6 eV (b) 3.4 eV
(c) 12.09 eV (d) 1.51 eV
- 136.** The ratio of speed of an electron in ground state in Bohrs first orbit of hydrogen atom to velocity of light in air is [MH CET 2002]
(a) $\frac{e^2}{2\epsilon_0 hc}$ (b) $\frac{2e^2 \epsilon_0}{hc}$
(c) $\frac{e^3}{2\epsilon_0 hc}$ (d) $\frac{2\epsilon_0 hc}{e^2}$
- 137.** Whenever a hydrogen atom emits a photon in the Balmer series [KCET 2002]
(a) It need not emit any more photon
(b) It may emit another photon in the Paschen series
(c) It must emit another photon in the Lyman series
(d) It may emit another photon in the Balmer series
- 138.** The de-Broglie wavelength of an electron in the first Bohr orbit is
(a) Equal to one fourth the circumference of the first orbit
(b) Equal to half the circumference of the first orbit
(c) Equal to twice the circumference of the first orbit
(d) Equal to the circumference of the first orbit
- 139.** In hydrogen atom, when electron jumps from second to first orbit, then energy emitted is [AIEEE 2002]
(a) $-13.6 eV$ (b) $-27.2 eV$
(c) $-6.8 eV$ (d) None of these
- 140.** Minimum energy required to takeout the only one electron from ground state of He^+ is [CPMT 2002]
(a) 13.6 eV (b) 54.4 eV
(c) 27.2 eV (d) 6.8 eV
- 141.** The frequency of $H\gamma$ line of Balmer series in H_2 atom is ν_0 . The frequency of line emitted by singly ionised He atom is [CPMT 2002]
(a) $2\nu_0$ (b) $4\nu_0$
(c) $\nu_0/2$ (d) $\nu_0/4$
- 142.** When the electron in the hydrogen atom jumps from 2^{nd} orbit to 1^{st} orbit, the wavelength of radiation emitted is λ . When the electrons jump from 3^{rd} orbit to 1^{st} orbit, the wavelength of emitted radiation would be [MP PMT 2002]
(a) $\frac{27}{32} \lambda$ (b) $\frac{32}{27} \lambda$
(c) $\frac{2}{3} \lambda$ (d) $\frac{3}{2} \lambda$
- 143.** The possible quantum number for $3d$ electron are [MP PMT 2002]
(a) $n = 3, l = 1, m_l = +1, m_s = -\frac{1}{2}$
(b) $n = 3, l = 2, m_l = +2, m_s = -\frac{1}{2}$
(c) $n = 3, l = 1, m_l = -1, m_s = +\frac{1}{2}$
(d) $n = 3, l = 0, m_l = +1, m_s = -\frac{1}{2}$
- 144.** The radius of the first (lowest) orbit of the hydrogen atom is a_0 . The radius of the second (next higher) orbit will be [MP PET 2002; MP PMT 2004]
(a) $4a_0$ (b) $6a_0$
(c) $8a_0$ (d) $10a_0$
- 145.** Which of the following transition will have highest emission wavelength [BHU 2003]
(a) $n = 2$ to $n = 1$ (b) $n = 1$ to $n = 2$
(c) $n = 2$ to $n = 5$ (d) $n = 5$ to $n = 2$
- 146.** When the wave of hydrogen atom comes from infinity into the first orbit then the value of wave number is [RPET 2003]
(a) 109700 cm^{-1} (b) 1097 cm^{-1}
(c) 109 cm^{-1} (d) None of these
- 147.** [KCET 2002] With the increase in principle quantum number, the energy difference between the two successive energy levels [RPET 2003]
(a) Increases
(b) Decreases
(c) Remains constant
(d) Sometimes increases and sometimes decreases
- 148.** In which of the following systems will the radius of the first orbit ($n = 1$) be minimum

[Kerala PET 2002; CBSE PMT 2003]

[Pb. PMT 2004]

- (a) Single ionized helium
(b) Deuterium atom
(c) Hydrogen atom
(d) Doubly ionized lithium
149. If the binding energy of the electron in a hydrogen atom is 13.6 eV , the energy required to remove the electron from the first excited state of Li^{++} is [AIEEE 2003]
(a) 122.4 eV (b) 30.6 eV
(c) 13.6 eV (d) 3.4 eV
150. Which of the following is quantised according to Bohr's theory of hydrogen atom [MP PMT 2004]
(a) Linear momentum of electron
(b) Angular momentum of electron
(c) Linear velocity of electron
(d) Angular velocity of electron
151. The shortest wavelength in the Lyman series of hydrogen spectrum is 912 \AA corresponding to a photon energy of 13.6 eV . The shortest wavelength in the Balmer series is about [MP PMT 2004]
(a) 3648 \AA (b) 8208 \AA
(c) 1228 \AA (d) 6566 \AA
152. Energy E of a hydrogen atom with principal quantum number n is given by $E = \frac{-13.6}{n^2} \text{ eV}$. The energy of a photon ejected when the electron jumps from $n = 3$ state to $n = 2$ state of hydrogen is approximately [CBSE PMT 2004]
(a) 1.5 eV (b) 0.85 eV
(c) 3.4 eV (d) 1.9 eV
153. The Bohr model of atoms [CBSE PMT 2004]
(a) Assumes that the angular momentum of electrons is quantized
(b) Uses Einstein's photo-electric equation
(c) Predicts continuous emission spectra for atoms
(d) Predicts the same emission spectra for all types of atoms
154. The colour of the second line of Balmer series is [J & K CET 2004]
(a) Blue (b) Yellow
(c) Red (d) Violet
155. Which state of triply ionised Beryllium (Be^{+++}) has the same orbital radius as that of the ground state of hydrogen [KCET 2004]
(a) $n = 4$ (b) $n = 3$
(c) $n = 2$ (d) $n = 1$
156. The ratio of areas within the electron orbits for the first excited state to the ground state for hydrogen atom is [BCECE 2004]
(a) $16 : 1$ (b) $18 : 1$
(c) $4 : 1$ (d) $2 : 1$
157. The kinetic energy of an electron revolving around a nucleus will be
(a) Four times of P.E. (b) Double of P.E.
(c) Equal to P.E. (d) Half of its P.E.
158. Taking Rydberg's constant $R_H = 1.097 \times 10^7 \text{ m}^{-1}$ first and second wavelength of Balmer series in hydrogen spectrum is
(a) 2000 \AA , 3000 \AA (b) 1575 \AA , 2960 \AA
(c) 6529 \AA , 4280 \AA (d) 6552 \AA , 4863 \AA
159. The kinetic energy of electron in the first Bohr orbit of the hydrogen atom is [Pb. PET 2000]
(a) -6.5 eV (b) -27.2 eV
(c) 13.6 eV (d) -13.6 eV
160. In the spectrum of hydrogen atom, the ratio of the longest wavelength in Lyman series to the longest wavelength in the Balmer series is [UPSEAT 2004]
(a) $5/27$ (b) $1/93$
(c) $4/9$ (d) $3/2$
161. In Bohr's model of hydrogen atom, which of the following pairs of quantities are quantized [UPSEAT 2004]
(a) Energy and linear momentum
(b) Linear and angular momentum
(c) Energy and angular momentum
(d) None of the above
162. The energy of the highest energy photon of Balmer series of hydrogen spectrum is close to [UPSEAT 2004]
(a) 13.6 eV (b) 3.4 eV
(c) 1.5 eV (d) 0.85 eV
163. Energy of an electron in n orbit of hydrogen atom is $\left(k = \frac{1}{4\pi\epsilon_0} \right)$
(a) $-\frac{2\pi^2 k^2 m e^4}{n^2 h^2}$ (b) $-\frac{4\pi^2 m k e^2}{n^2 h^2}$
(c) $-\frac{n^2 h^2}{2\pi k m e^4}$ (d) $-\frac{n^2 h^2}{4\pi^2 k m e^2}$
164. Which one of the relation is correct between time period and number of orbits while an electron is revolving in a orbit [DPMT 2003]
(a) n^2 (b) $\frac{1}{n^2}$
(c) n^3 (d) $\frac{1}{n}$
165. An electron changes its position from orbit $n = 4$ to the orbit $n = 2$ of an atom. The wavelength of the emitted radiation's is ($R =$ Rydberg's constant) [BHU 2004]
(a) $\frac{16}{R}$ (b) $\frac{16}{3R}$
(c) $\frac{16}{5R}$ (d) $\frac{16}{7R}$
166. If the energy of a hydrogen atom in n th orbit is E_n , then energy in the n th orbit of a singly ionized helium atom will be [BCECE 2004]
(a) $4E_n$ (b) $E_n/4$
(c) $2E_n$ (d) $E_n/2$

1472 Atomic and Nuclear Physics

167. What is the ratio of wavelength of radiations emitted when an electron in hydrogen atom jump from fourth orbit to second orbit and from third orbit to second orbit

[MH CET 2004]

- (a) 27 : 25 (b) 20 : 27
(c) 20 : 25 (d) 25 : 27

168. The energy of electron in the n th orbit of hydrogen atom is expressed as $E_n = \frac{-13.6}{n^2} eV$. The shortest and longest wavelength of Lyman series will be [Pb. PET 2003]


- (a) 910 Å, 1213 Å (b) 5463 Å, 7858 Å
(c) 1315 Å, 1530 Å (d) None of these

169. The ground state energy of hydrogen atom is $-13.6 eV$. What is the potential energy of the electron in this state

[AIIMS 2005]

- (a) 0 eV (b) $-27.2 eV$
(c) 1 eV (d) 2 eV

170. The diagram shows the energy levels for an electron in a certain atom. Which transition shown represents the emission of a photon with the most energy [AIEEE 2005]



- (a) I (b) II
(c) III (d) IV

171. As the electron in Bohr orbit of Hydrogen atom passes from state $n = 2$ to $n = 1$, the kinetic energy K and potential energy U change as [MP PET 2005]

- (a) K two-fold, U four-fold
(b) K four-fold, U two-fold
(c) K four-fold, U also four-fold
(d) K two-fold, U also two-fold

172. The magnetic moment (μ) of a revolving electron around the nucleus varies with principal quantum number n as

[AIIMS 2005]

- (a) $\mu \propto n$ (b) $\mu \propto 1/n$
(c) $\mu \propto n^2$ (d) $\mu \propto 1/n^2$

173. Bohr's atom model assumes [KCET 2005]

- (a) The nucleus is of infinite mass and is at rest
(b) Electrons in a quantized orbit will not radiate energy
(c) Mass of electron remains constant
(d) All the above conditions

174. Radius of first Bohr orbit is r . What is the radius of 2nd Bohr orbit?

- (a) $8r$ (b) $2r$
(c) $4r$ (d) $2\sqrt{2}r$

1. Which of the following particles are constituents of the nucleus [CBSE PMT 1999]

- (a) Protons and electrons (b) Protons and neutrons
(c) Neutrons and electrons (d) Neutrons and positrons

2. The particles which can be added to the nucleus of an atom without changing its chemical properties are called

[NCERT 1979]

- (a) Electrons (b) Protons
(c) Neutrons (d) None of the above

3. The neutron was discovered by

[MP PMT 1992; RPMT 1996]

- (a) Marie Curie (b) Pierre Curie
(c) James Chadwick (d) Rutherford

4. The mass number of a nucleus is

[IIT 1986; ISM Dhanbad 1994;

MP PMT 1997; CBSE PMT 2003; MH CET (Med.) 2001]

- (a) Always less than its atomic number
(b) Always more than its atomic number
(c) Always equal to its atomic number
(d) Sometimes more than and sometimes equal to its atomic number

5. The energy equivalent of 1 kilogram of matter is about

[MP PET/PMT 1988; MNR 1987]

- (a) $10^{-15} J$ (b) 1 J
(c) $10^{-12} J$ (d) $10^{17} J$

6. Nuclear binding energy is equivalent to [MP PET/PMT 1988]

- (a) Mass of proton (b) Mass of neutron
(c) Mass of nucleus (d) Mass defect of nucleus

7. If the binding energy of the deuterium is 2.23 MeV.

The mass defect given in *a.m.u* is [MP PET 1993]

- (a) -0.0024 (b) -0.0012
(c) 0.0012 (d) 0.0024

8. Which of the following has the mass closest in value to that of the positron [AFMC 1993]

- (a) Proton (b) Electron
(c) Photon (d) Neutrino

(1 *a.m.u* = 931 MeV)

9. Size of nucleus is of the order of

[CPMT 1983; MP PET 2002, 03]


- (a) $10^{-10} m$ (b) $10^{-15} m$
(c) $10^{-12} m$ (d) $10^{-19} m$

[BHU 2005]


1500 Atomic and Nuclear Physics

4. A radioactive sample consists of two distinct species having equal number of atoms initially. The mean life time of one species is τ and that of the other is 5τ . The decay products in both cases are stable. A plot is made of the total number of radioactive nuclei as a function of time. Which of the following figures best represents the form of this plot


[IIT-JEE (Screening) 2001]




5. Radioactive element decays to form a stable nuclide, then the rate of decay of reactant $\left(\frac{dN}{dt}\right)$ will vary with time (t) as shown in figure



6. A radioactive sample has N_0 active atoms at $t = 0$. If the rate of disintegration at any time is R and the number of atoms is N , then the ratio R/N varies with time as




7. The count rate of 10g of radioactive material was measured at different times and this has been shown in the figure. The half life of material and the total counts (approximately) in the first half life period, respectively are [CPMT 1986]



- (a) 4h, 9000 (b) 3h, 14000
(c) 3h, 235 (d) 3h, 50


8. The fraction f of radioactive material that has decayed in time t , varies with time t . The correct variation is given by the curve

- (a) A
(b) B
(c) C
(d) D




9. Binding energy per nucleon versus mass number curve for nuclei is shown in the figure. W, X, Y and Z are four nuclei indicated on the curve. The process that would release energy is

[DCE 2000]




- (a) $Y \rightarrow 2Z$
(c) $W \rightarrow 2Y$


10. The plot of the number (N) of decayed atoms versus activity (A) of a radioactive substance is




11. If in hydrogen atom, radius of n^{th} Bohr orbit is r_n^D , frequency of revolution of electron in n^{th} orbit is f_n , choose the correct option




12. The graph between the instantaneous concentration (N) of a radioactive element and time (t) is




13. In Fig. X represents time and Y represent activity of a radioactive sample. Then the activity of sample, varies with time according to the curve




14. The graph which represents the correct variation of logarithm of activity ($\log A$) versus time, in figure is



15. The charge density in a nucleus varies with distance from the centre of the nucleus according to the curve in Fig.




16. The graph between $\log R$ and $\log A$ where R is the nuclear radius and A is the mass number is




(c)

(d)


17. The curve between the activity A of a radioactive sample and the number of active atoms N is



18. The graph between number of decayed atoms N' of a radioactive element and time t is



19. The figure shows a graph between $\ln \frac{A_n}{A_1}$ and $\ln |n|$, where A_n is the area enclosed by the n th orbit in a hydrogen like atom. The correct curve is



Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below:

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- (c) If assertion is true but reason is false.
- (d) If the assertion and reason both are false.
- (e) If assertion is false but reason is true.

1. Assertion : It is not possible to use ^{35}Cl as the fuel for fusion energy.

Reason : The binding energy of ^{35}Cl is too small.

[AIIMS 2005]

1502 Atomic and Nuclear Physics

2. Assertion : ${}^{90}\text{Sr}$ from the radioactive fall out from a nuclear bomb ends up in the bones of human beings through the milk consumed by them. It causes impairment of the production of red blood cells.
Reason : The energetics β -particles emitted in the decay of ${}^{90}\text{Sr}$ damage the bone marrow. [AIIMS 2004]
3. Assertion : Neutrons penetrate matter more readily as compared to protons.
Reason : Neutrons are slightly more massive than protons. [AIIMS 2003]
4. Assertion : Bohr had to postulate that the electrons in stationary orbits around the nucleus do not radiate.
Reason : According to classical physics all moving electrons radiate. [AIIMS 2003]
5. Assertion : Radioactive nuclei emit β^{-1} particles.
Reason : Electrons exist inside the nucleus. [AIIMS 2003]
6. Assertion : ${}_Z X^A$ undergoes 2α -decays, 2β -decays and 2γ -decays and the daughter product is ${}_{Z-2} Y^{A-8}$.
Reason : In α -decay the mass number decreases by 4 and atomic number decreases by 2. In β -decay the mass number remains unchanged, but atomic number increases by 1 only. [AIIMS 2001]
7. Assertion : Density of all the nuclei is same.
Reason : Radius of nucleus is directly proportional to the cube root of mass number. [AIIMS 2000]
8. Assertion : Isobars are the element having same mass number but different atomic number.
Reason : Neutrons and protons are present inside nucleus. [AIIMS 1997]
9. Assertion : The force of repulsion between atomic nucleus and α -particle varies with distance according to inverse square law.
Reason : Rutherford did α -particle scattering experiment.
10. Assertion : The positively charged nucleus of an atom has a radius of almost $10^{-15} m$.
Reason : In α -particle scattering experiment, the distance of closest approach for α -particles is $\approx 10^{-15} m$.
11. Assertion : According to classical theory, the proposed path of an electron in Rutherford atom model will be parabolic.
Reason : According to electromagnetic theory an accelerated particle continuously emits radiation.
12. Assertion : Electrons in the atom are held due to coulomb forces.
Reason : The atom is stable only because the centripetal force due to Coulomb's law is balanced by the centrifugal force.
13. Assertion : The electron in the hydrogen atom passes from energy level $n=4$ to the $n=1$ level. The maximum and minimum number of photon that can be emitted are six and one respectively.
Reason : The photons are emitted when electron make a transition from the higher energy state to the lower energy state.
14. Assertion : Hydrogen atom consists of only one electron but its emission spectrum has many lines.
Reason : Only Lyman series is found in the absorption spectrum of hydrogen atom whereas in the emission spectrum, all the series are found.
15. Assertion : It is essential that all the lines available in the emission spectrum will also be available in the absorption spectrum.
Reason : The spectrum of hydrogen atom is only absorption spectrum.
16. Assertion : For the scattering of α -particles at a large angles, only the nucleus of the atom is responsible.
Reason : Nucleus is very heavy in comparison to electrons.
17. Assertion : All the radioactive elements are ultimately converted in lead.
Reason : All the elements above lead are unstable.
18. Assertion : Amongst alpha, beta and gamma rays, α -particle has maximum penetrating power.
Reason : The alpha particle is heavier than beta and gamma rays.
19. Assertion : The ionising power of β -particle is less compared to α -particles but their penetrating power is more.
Reason : The mass of β -particle is less than the mass of α -particle.
20. Assertion : The mass of β -particles when they are emitted is higher than the mass of electrons obtained by other means.
Reason : β -particle and electron, both are similar particles.
21. Assertion : Radioactivity of 10^8 undecayed radioactive nuclei of half life of 50 days is equal to that of 1.2×10^8 number of undecayed nuclei of some other material with half life of 60 days
Reason : Radioactivity is proportional to half-life.
22. Assertion : Fragments produced in the fission of U^{235} are radioactive.
Reason : The fragments have abnormally high proton to neutron ratio.
23. Assertion : Electron capture occurs more often than positron emission in heavy elements.
Reason : Heavy elements exhibit radioactivity.
24. Assertion : The mass of a nucleus can be either less than or more than the sum of the masses of nucleons present in it.
Reason : The whole mass of the atom is considered in the nucleus.

Answers

Atomic Structure

1	a	2	d	3	a	4	d	5	c
6	b	7	b	8	c	9	a	10	d
11	b	12	b	13	b	14	c	15	c
16	c	17	c	18	b	19	c	20	b
21	c	22	c	23	c	24	d	25	d
26	c	27	d	28	b	29	d	30	b
31	a	32	d	33	c	34	d	35	d
36	b	37	c	38	a	39	c	40	c
41	a	42	d	43	d	44	a	45	a
46	a	47	c	48	a	49	b	50	a
51	b	52	c	53	d	54	c	55	d
56	a	57	c	58	d	59	d	60	d
61	c	62	ad	63	d	64	b	65	b
66	b	67	c	68	a	69	b	70	c
71	a	72	b	73	b	74	d	75	b
76	d	77	d	78	b	79	a	80	d
81	a	82	a	83	b	84	b	85	d
86	c	87	a	88	c	89	a	90	b
91	d	92	b	93	c	94	c	95	b
96	a	97	c	98	b	99	a	100	b
101	a	102	c	103	a	104	b	105	d
106	b	107	b	108	c	109	d	110	c
111	a	112	a	113	a	114	b	115	d
116	b	117	d	118	a	119	b	120	a
121	b	122	b	123	b	124	d	125	a
126	d	127	d	128	c	129	c	130	d
131	a	132	c	133	c	134	a	135	c
136	a	137	c	138	d	139	d	140	b
141	b	142	a	143	b	144	a	145	d
146	a	147	b	148	d	149	b	150	b
151	a	152	d	153	a	154	a	155	c
156	d	157	d	158	d	159	c	160	a
161	c	162	b	163	a	164	c	165	b
166	a	167	b	168	a	169	b	170	c
171	c	172	a	173	d	174	c		

46	c	47	d	48	b	49	a	50	b
51	d	52	b	53	d	54	c	55	a
56	a	57	d	58	ad	59	c	60	a
61	c	62	b	63	a	64	a	65	d
66	c	67	a	68	b	69	b	70	d
71	b	72	d	73	bc	74	d	75	c
76	d	77	d	78	c	79	b	80	b
81	c	82	d	83	c	84	b	85	d
86	b	87	a	88	d	89	d	90	a
91	c	92	b	93	a	94	c	95	a
96	b	97	c	98	d	99	d	100	b
101	c	102	b	103	a	104	d	105	c
106	b	107	a	108	b	109	d	110	c
111	c	112	c	113	b	114	b	115	d
116	a	117	b	118	a	119	c	120	b
121	d	122	d	123	a	124	a	125	c
126	c	127	b	128	d	129	c	130	a
131	c	132	a	133	a	134	b	135	b
136	c	137	a	138	c	139	b	140	a
141	b	142	b	143	b	144	d	145	d
146	a	147	b	148	b	149	d	150	a
151	c	152	b	153	d	154	c	155	c
156	a	157	b	158	a	159	c	160	c
161	a	162	a	163	b	164	b	165	c
166	b	167	d	168	d	169	a	170	b
171	b	172	a	173	c	174	b	175	a
176	b	177	a	178	c	179	b		

Radioactivity

1	a	2	a	3	d	4	c	5	a
6	c	7	c	8	d	9	c	10	c
11	b	12	c	13	c	14	c	15	a
16	c	17	a	18	c	19	b	20	a
21	a	22	c	23	a	24	d	25	d
26	d	27	c	28	b	29	a	30	c
31	c	32	c	33	d	34	c	35	c
36	b	37	b	38	d	39	d	40	d
41	a	42	b	43	c	44	d	45	b
46	b	47	d	48	d	49	b	50	a
51	b	52	c	53	a	54	d	55	c
56	d	57	b	58	d	59	d	60	b
61	a	62	d	63	a	64	d	65	b
66	a	67	b	68	c	69	d	70	c
71	d	72	a	73	a	74	d	75	c
76	d	77	d	78	c	79	a	80	d
81	d	82	b	83	a	84	a	85	b

Nucleus, Nuclear Reaction

1	b	2	c	3	c	4	d	5	d
6	d	7	d	8	b	9	b	10	c
11	a	12	b	13	d	14	c	15	c
16	c	17	c	18	c	19	d	20	c
21	b	22	d	23	a	24	b	25	c
26	b	27	c	28	c	29	a	30	a
31	b	32	a	33	d	34	c	35	c
36	b	37	a	38	a	39	d	40	c
41	b	42	a	43	c	44	a	45	c

1504 Atomic and Nuclear Physics

86	c	87	d	88	d	89	b	90	a
91	b	92	d	93	c	94	c	95	a
96	d	97	d	98	a	99	b	100	c
101	a	102	d	103	b	104	b	105	b
106	d	107	a	108	d	109	c	110	b
111	c	112	c	113	d	114	d	115	c
116	b	117	a	118	a	119	d	120	a
121	c	122	d	123	a	124	d	125	d
126	d	127	c	128	d	129	c	130	b
131	d	132	b	133	c	134	a	135	a
136	b	137	ac	138	b	139	c	140	c
141	d	142	c	143	a	144	d	145	c
146	b	147	d	148	b	149	b	150	c
151	c	152	a	153	b	154	b	155	d
156	b	157	c	158	c	159	d	160	c
161	a	162	d	163	c	164	c	165	d
166	d	167	c	168	c	169	b	170	d
171	b	172	c	173	b	174	a	175	c
176	d								

16	a	17	c	18	d	19	b	20	b
21	c	22	c	23	b	24	e		

Critical Thinking Questions

1	c	2	c	3	b	4	a	5	a
6	a	7	a	8	d	9	b	10	d
11	a	12	a	13	a	14	d	15	c
16	d	17	c	18	d	19	c	20	d
21	b	22	c	23	d	24	a	25	d
26	d	27	c	28	c	29	cd	30	a
31	a	32	c	33	a	34	a	35	b
36	b	37	b	38	c	39	b	40	a
41	a	42	b	43	c	44	c	45	d
46	a	47	b	48	d	49	a	50	b
51	d	52	a	53	b	54	b	55	a
56	a	57	b	58	b	59	b	60	a
61	a	62	a	63	b	64	c	65	a

Graphical Questions

1	a	2	a	3	c	4	d	5	c
6	d	7	b	8	b	9	c	10	d
11	d	12	d	13	b	14	d	15	c
16	a	17	b	18	c	19	a		


Assertion and Reason

1	c	2	a	3	b	4	b	5	c
6	a	7	a	8	b	9	b	10	a
11	e	12	c	13	b	14	b	15	d

AS Answers and Solutions

Atomic Structure

1. (a) For $n=1$, maximum number of states $= 2n^2 = 2$ and for $n = 2, 3, 4$, maximum number of states would be 8, 18, 32 respectively. Hence number of possible elements $= 2 + 8 + 18 + 32 = 60$.
2. (d) Bohr radius $r = \frac{\epsilon_0 n^2 h^2}{\pi Z m e^2}$; $\therefore r \propto n^2$
3. (a)
- $$n=2 \quad \text{-----} \quad E_2 = -\frac{13.6}{(2)^2} = -3.4 \text{ eV}$$
- $$\uparrow$$
- $$n=1 \quad \text{-----} \quad E_1 = -13.6 \text{ eV}$$
- $$E_{1 \rightarrow 2} = -3.4 - (-13.6) = +10.2 \text{ eV}$$
4. (d) $\frac{1}{\lambda} = RZ^2 \left(\frac{1}{1^2} - \frac{1}{2^2} \right)$
- For di-ionised lithium the value of Z is maximum.
5. (c) Lyman series lies in the UV region.
6. (b) The size of the atom is of the order of $1\text{\AA} = 10^{-10} \text{ m}$.
7. (b) Balmer series lies in the visible region.
8. (c) Transition A ($n = \infty$ to 1): Series limit of Lyman series
 Transition B ($n = 5$ to $n = 2$): Third spectral line of Balmer series
 Transition C ($n = 5$ to $n = 3$): Second spectral line of Paschen series
9. (a) D is excitation of electron from 2^{nd} orbit corresponding to absorption line in Balmer series and E is the energy released to bring the electron from ∞ to ground state *i.e.* ionisation potential.
10. (d)
11. (b) Paschen series lies in the infrared region.
12. (b) Energy required to knock out the electron in the n orbit $= +\frac{13.6}{n^2} \text{ eV} \Rightarrow E_3 = +\frac{13.6}{9} \text{ eV}$.
13. (b) Linear momentum $= mv = 9.1 \times 10^{-31} \times 2.2 \times 10^6$
 $= 2.0 \times 10^{-24} \text{ kg-m/s}$
14. (c) $r \propto n^2 \Rightarrow r_n = n^2 a_0$ ($\because r_1 = a_0$)
15. (c) For the ionization of second He electron. He^+ will act as hydrogen like atom.
 Hence ionization potential
 $= Z^2 \times 13.6 \text{ volt} = (2)^2 \times 13.6 = 54.4 \text{ V}$
16. (c) Energy required $= \frac{13.6}{n^2} = \frac{13.6}{10^2} = 0.136 \text{ eV}$

17. (c) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \Rightarrow \frac{1}{n_1^2} - \frac{1}{n_2^2} = \frac{1}{R\lambda}$
 $= \frac{1}{1.097 \times 10^7 \times 18752 \times 10^{-10}} = 0.0486 = \frac{7}{144}$. But
 $\frac{1}{3^2} - \frac{1}{4^2} = \frac{7}{144} \Rightarrow n_1 = 3$ and $n_2 = 4$ (Paschen series)
18. (b) Potential energy of electron in n orbit of radius r in H -atom
 $U = -\frac{e^2}{r}$ (in CGS)
 $\therefore \text{K.E.} = \frac{1}{2} |P.E.| \Rightarrow K = \frac{e^2}{2r}$
19. (c) Final energy of electron $= -13.6 + 12.1 = -1.51 \text{ eV}$. which is corresponds to third level *i.e.* $n = 3$. Hence number of spectral lines emitted $= \frac{n(n-1)}{2} = \frac{3(3-1)}{2} = 3$
20. (b) Let the energy in A, B and C state be E_1, E_2 and E_3 , then from the figure
- 
- $$(E_C - E_B) + (E_B - E_A) = (E_C - E_A) \text{ or } \frac{hc}{\lambda_1} + \frac{hc}{\lambda_2} = \frac{hc}{\lambda_3}$$
- $$\Rightarrow \lambda_3 = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$$
21. (c) According to Bohr's second postulate.
22. (c) First excited state *i.e.* second orbit ($n = 2$)
 Second excited state *i.e.* third orbit ($n = 3$)
 $\therefore E = -\frac{13.6}{n^2} \Rightarrow \frac{E_2}{E_3} = \left(\frac{3}{2} \right)^2 = \frac{9}{4}$
23. (c) $\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{4^2} \right) = \frac{3R}{16} \Rightarrow \lambda = \frac{16}{3R} = \frac{16}{3} \times 10^{-5} \text{ cm}$
 Frequency $n = \frac{c}{\lambda} = \frac{3 \times 10^{10}}{\frac{16}{3} \times 10^{-5}} = \frac{9}{16} \times 10^{15} \text{ Hz}$
24. (d) Energy required to remove electron in the $n = 2$ state $= +\frac{13.6}{(2)^2} = +3.4 \text{ eV}$
25. (d) $(E)_\infty = Z^2 (E_{ion})_H = (11)^2 13.6 \text{ eV}$
26. (c) The wavelength of spectral line in Balmer series is given by
 $\frac{1}{\lambda} = R \left[\frac{1}{2^2} - \frac{1}{n^2} \right]$
 For first line of Balmer series, $n = 3$

$$\Rightarrow \frac{1}{\lambda_1} = R \left[\frac{1}{2^2} - \frac{1}{3^2} \right] = \frac{5R}{36}; \text{ For second line } n = 4.$$

$$\Rightarrow \frac{1}{\lambda_2} = R \left[\frac{1}{2^2} - \frac{1}{4^2} \right] = \frac{3R}{16}$$

$$\therefore \frac{\lambda_2}{\lambda_1} = \frac{20}{27} \Rightarrow \lambda_1 = \frac{20}{27} \times 6561 = 4860 \text{ \AA}$$

27. (d) $2E - E = \frac{hc}{\lambda} \Rightarrow E = \frac{hc}{\lambda}$

$$\frac{4E}{3} - E = \frac{hc}{\lambda'} \Rightarrow \frac{E}{3} = \frac{hc}{\lambda'} \therefore \frac{\lambda'}{\lambda} = 3 \Rightarrow \lambda' = 3\lambda$$

28. (b) Because atom is hollow and whole mass of atom is concentrated in a small centre called nucleus.


29. (d) $r = \frac{\epsilon_0 n^2 h^2}{\pi Z m e^2}; \therefore r \propto \frac{n^2}{Z}$

30. (b) $r \propto n^2 \Rightarrow \frac{r_{(n=2)}}{r_{(n=3)}} = \frac{4}{9} \Rightarrow r_{(n=3)} = \frac{9}{4} R = 2.25 R$

31. (a) In the revolution of electron, coulomb force provides the necessary centripetal force

$$\Rightarrow \frac{z e^2}{r^2} = \frac{m v^2}{r} \Rightarrow m v^2 = \frac{z e^2}{r}$$

$$\therefore \text{K.E.} = \frac{1}{2} m v^2 = \frac{z e^2}{2r}$$



32. (d) According to Bohr's theory $mvr = n \frac{h}{2\pi}$

$$\Rightarrow \text{Circumference } 2\pi r = n \left(\frac{h}{mv} \right) = n\lambda$$

33. (c) $K.E. = \frac{kZe^2}{2r}$ and $P.E. = -\frac{kZe^2}{r}; \therefore \frac{K.E.}{P.E.} = -\frac{1}{2}$.

34. (d) Lyman series lies in the UV region.

35. (d) If E is the energy radiated in transition

$$\text{then } E_{R \rightarrow G} > E_{Q \rightarrow S} > E_{R \rightarrow S} > E_{Q \rightarrow R} > E_{P \rightarrow Q}$$

For getting blue line energy radiated should be maximum

$$\left(E \propto \frac{1}{\lambda} \right). \text{ Hence (d) is the correct option.}$$

36. (b) Energy released = $13.6 \left[\frac{1}{(2)^2} - \frac{1}{(4)^2} \right] = 2.55 \text{ eV}$

37. (c) The absorption lines are obtained when the electron jumps from ground state ($n = 1$) to the higher energy states. Thus only 1, 2 and 3 lines will be obtained.

38. (a) $P.E. \propto -\frac{1}{r}$ and $K.E. \propto \frac{1}{r}$

As r increases so K.E. decreases but P.E. increases.

39. (c) Wave number $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] = R \left[\frac{1}{4} - \frac{1}{16} \right] = \frac{3R}{16}$

40. (c) In hydrogen atom, the lowest orbit ($n = 1$) corresponds to minimum energy (-13.6 eV).

41. (a) $K.E. = -(\text{T.E.})$

42. (d) Required energy $E_3 = \frac{+13.6}{3^2} = 1.51 \text{ eV}$

43. (d) As n increases P.E. also increases.

44. (a) When an electron jumps from the orbit of lower energy ($n=1$) to the orbit of higher energy ($n=3$), energy is absorbed.

45. (a) For Lyman series

$$V_{\text{Lyman}} = \frac{c}{\lambda_{\text{max}}} = Rc \left[\frac{1}{(1)^2} - \frac{1}{(2)^2} \right] = \frac{3RC}{4}$$

For Balmer series


$$V_{\text{Balmer}} = \frac{c}{\lambda_{\text{max}}} = Rc \left[\frac{1}{(2)^2} - \frac{1}{(3)^2} \right] = \frac{5RC}{36}$$

$$\therefore \frac{V_{\text{Lyman}}}{V_{\text{Balmer}}} = \frac{27}{5}$$

46. (a) $\therefore E_1 > E_2$

$$\therefore v_1 > v_2$$

i.e. photons of higher frequency will be emitted if transition takes place from $n = 2$ to 1.



47. (c) Wave number


$$= \frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

For first Balmer line $n_1 = 2, n_2 = 3$

$$\therefore \text{Wave number} = R \left(\frac{1}{2^2} - \frac{1}{3^2} \right) = R \left(\frac{9-4}{9 \times 4} \right) = \frac{5R}{36}$$

48. (a) Energy required to ionise helium atom = 24.6 eV

49. (b) From diagram



$$E_1 = -13.6 - (-3.4) = -10.2 \text{ eV}$$

$$E_2 = -13.6 - (-1.51) = -12.09 \text{ eV}$$

$$E_3 = -1.51 - (-0.85) = -0.66 \text{ eV}$$

$$E_4 = -3.4 - (-1.51) = -1.89 \text{ eV}$$

E_3 is least *i.e.* frequency is lowest.

50. (a) $P.E. = -\frac{ke^2}{r} = -\frac{e^2}{4\pi\epsilon_0 r}; \text{ K.E.} = -\frac{1}{2}(\text{P.E.}) = \frac{e^2}{8\pi\epsilon_0 r}$

51. (b) Similar to Q. 49

52. (c) $mvr = \frac{nh}{2\pi}$, for $n=1$ it is $\frac{h}{2\pi}$

53. (d) Minimum energy required to excite from ground state

$$= 13.6 \left[\frac{1}{1^2} - \frac{1}{2^2} \right] = 10.2 \text{ eV}$$

54. (c) $\frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$

For first line of Lyman series $n_1 = 1$ and $n_2 = 2$
 For first line of Balmer series $n_1 = 2$ and $n_2 = 3$

So, $\frac{\lambda_{\text{Lyman}}}{\lambda_{\text{Balmer}}} = \frac{5}{27}$

55. (d) $R = \frac{2\pi^2 k^2 e^4 m}{ch^3} = \left(\frac{1}{4\pi\epsilon_0}\right)^2 \frac{2\pi^2 me^4}{ch^3}$
56. (a) $\frac{1}{\lambda} = R\left[\frac{1}{4} - \frac{1}{9}\right] = \frac{5R}{36}$
 $\therefore R = \frac{36}{5\lambda} = \frac{36}{5 \times 6563 \times 10^{-10}} = 1.09 \times 10^7 \text{ m}^{-1}$
57. (c) Angular momentum $L = n\left(\frac{h}{2\pi}\right)$
 For this case $n=2$, hence $L = 2 \times \frac{h}{2\pi} = \frac{h}{\pi}$
58. (d) $v \propto \frac{1}{n} \Rightarrow \frac{v_5}{v_2} = \frac{2}{5} \Rightarrow v_5 = \frac{2}{5}v_2 = \frac{2}{5}v$
59. (d) By using $N_E = \frac{n(n-1)}{2} \Rightarrow N_E = \frac{4(4-1)}{2} = 6$
60. (d) Shortest wavelength comes from $n_1 = \infty$ to $n_2 = 1$ and longest wavelength comes from $n_1 = 6$ to $n_2 = 5$ in the given case. Hence $\frac{1}{\lambda_{\text{min}}} = R\left(\frac{1}{1^2} - \frac{1}{\infty^2}\right) = R$
 $\frac{1}{\lambda_{\text{max}}} = R\left(\frac{1}{5^2} - \frac{1}{6^2}\right) = R\left(\frac{36-25}{25 \times 36}\right) = \frac{11}{900}R$
 $\therefore \frac{\lambda_{\text{max}}}{\lambda_{\text{min}}} = \frac{900}{11}$
61. (c) $\frac{mv^2}{a_0} = \frac{1}{4\pi\epsilon_0} \frac{e^2}{a_0^2} \Rightarrow v = \frac{e}{\sqrt{4\pi\epsilon_0 a_0 m}}$
62. (a,d) $T \propto n^3$. Given $T_{n_1} = 8 T_{n_2}$, hence $n_1 = 2n_2$
 Therefore, option (a) and (d) both are correct.
63. (d) $E = -Z^2 \times 13.6 \text{ eV} = -9 \times 13.6 \text{ eV} = -122.4 \text{ eV}$
 So ionisation energy = + 122.4 eV.
64. (b)
65. (b) As n increases P.E. increases and K.E. decreases.
66. (b) $r = \frac{n^2}{Z}(r_0)$; $\Rightarrow r_{(n=2)} = \frac{(2)^2}{2} \times 0.53 = 1.06 \text{ \AA}$
67. (c)
68. (a) $\bar{v} \propto \frac{1}{\lambda} \propto Z^2 \Rightarrow \lambda Z^2 = \text{constant} \Rightarrow \lambda = \frac{\lambda}{4} Z^2 \Rightarrow Z = 2$
69. (b) In Paschen series $\frac{1}{\lambda_{\text{max}}} = R\left[\frac{1}{(3)^2} - \frac{1}{(4)^2}\right]$
 $\Rightarrow \lambda_{\text{max}} = \frac{144}{7R} = \frac{144}{7 \times 1.1 \times 10^7} = 1.89 \times 10^{-6} \text{ m} = 1.89 \text{ \mu m}$
 Similarly $\lambda_{\text{min}} = \frac{9}{R} = \frac{9}{1.1 \times 10^7} = 0.818 \text{ \mu m}$

70. (c) For third line of Balmer series $n_1 = 2, n_2 = 5$
 $\therefore \frac{1}{\lambda} = RZ^2\left[\frac{1}{n_1^2} - \frac{1}{n_2^2}\right]$ gives $Z^2 = \frac{n_1^2 n_2^2}{(n_2^2 - n_1^2)\lambda R}$
 On putting values $Z = 2$
 From $E = -\frac{13.6Z^2}{n^2} = \frac{-13.6(2)^2}{(1)^2} = -54.4 \text{ eV}$
71. (a) Ionization energy = Binding energy.
72. (b) $E = -Rch \Rightarrow R = -\frac{E}{ch} = -\frac{13.6 \times 1.6 \times 10^{-19}}{3 \times 10^8 \times 6.6 \times 10^{-34}}$
 $= 1.098 \times 10^7 \text{ per m}$
73. (b) Bohr postulated that the angular momentum of the electron is conserved.
74. (d) $E_3 = -\frac{13.6}{9} = -1.51 \text{ eV}; E_4 = -\frac{13.6}{16} = -0.85 \text{ eV}$
 $\therefore E_4 - E_3 = 0.66 \text{ eV}$
75. (b) Number of spectral lines $N_E = \frac{n(n-1)}{2} = \frac{4(4-1)}{2} = 6$
76. (d) In the transition from orbit $5 \rightarrow 2$, more energy is liberated as compared to transition from $4 \rightarrow 2$.
77. (d) Impact parameter $b \propto \cot \frac{\theta}{2}$
 Here $b = 0$, hence $\theta = 180^\circ$
78. (b) $r \propto n^2$ i.e. $\frac{r_f}{r_i} = \left(\frac{n_f}{n_i}\right)^2$
 $\Rightarrow \frac{21.2 \times 10^{-11}}{5.3 \times 10^{-11}} = \left(\frac{n}{1}\right)^2 \Rightarrow n^2 = 4 \Rightarrow n = 2$
79. (a)
80. (d) $E_n = \frac{-13.6}{n^2} = \frac{-13.6}{4} = -3.4 \text{ eV}$
81. (a) $\frac{1}{\lambda_{\text{Balmer}}} = R\left[\frac{1}{2^2} - \frac{1}{3^2}\right] = \frac{5R}{36}$, $\frac{1}{\lambda_{\text{Lyman}}} = R\left[\frac{1}{1^2} - \frac{1}{2^2}\right] = \frac{3R}{4}$
 $\therefore \lambda_{\text{Lyman}} = \lambda_{\text{Balmer}} \times \frac{5}{27} = 1215.4 \text{ \AA}$
82. (a) $\frac{1}{\lambda} = R_H\left[\frac{1}{n_1^2} - \frac{1}{n_2^2}\right]$. For Lyman series $n=1$ and $n=2, 3, 4$,
 When $n=2$, we get $\lambda = \frac{4}{3R_H} = \frac{4}{3 \times 10967} \text{ cm}$
83. (b) $r \propto n^2$. For ground state $n=1$ and for first excited state $n=2$.
84. (b) No. of lines $N_E = \frac{n(n-1)}{2} = \frac{3(3-1)}{2} = 3$
85. (d) Infinitely large transitions are possible (in principle) for the hydrogen atom.
86. (c) $r_n \propto n^2$
87. (a) $E\left(\frac{hc}{\lambda}\right) \propto \frac{Z^2}{n^2} \Rightarrow \lambda \propto \frac{1}{Z^2}$

1508 Atomic and Nuclear Physics

Hence $\lambda_{He^+} = \frac{20.397}{4} = 5.099 \text{ cm}$

88. (c) Excitation potential = $\frac{\text{Excitation energy}}{e}$

Minimum excitation energy corresponds to excitation from $n = 1$ to $n = 2$

\therefore Minimum excitation energy in hydrogen atom = $-3.4 - (-13.6) = +10.2 \text{ eV}$

so minimum excitation potential = 10.2 eV .

89. (a) Orbital speed varies inversely as the radius of the orbit. Energy increases with the increase in quantum number.

90. (b) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \Rightarrow \frac{1}{\lambda_{3 \rightarrow 2}} = R \left[\frac{1}{(2)^2} - \frac{1}{(3)^2} \right] = \frac{5R}{36}$

and $\frac{1}{\lambda_{4 \rightarrow 2}} = R \left[\frac{1}{(2)^2} - \frac{1}{(4)^2} \right] = \frac{3R}{16}$

$\therefore \frac{\lambda_{4 \rightarrow 2}}{\lambda_{3 \rightarrow 2}} = \frac{20}{27} \Rightarrow \lambda_{4 \rightarrow 2} = \frac{20}{27} \lambda_0$

91. (d)

92. (b) Kinetic energy = |Total energy|

93. (c) Energy to excite the e^- from $n = 1$ to $n = 2$

$\xrightarrow{\hspace{10em}}$ First excited state $n = 2$ (-3.4 eV)

\uparrow
 Ground state $n = 1$ (-13.6 eV)

(For H_2 - atom)
 $E = -3.4 - (-13.6) = 10.2 \text{ eV} = 10.2 \times 1.6 \times 10^{-19}$

$= 1.632 \times 10^{-18} \text{ J}$

94. (c)

95. (b)

96. (a) According to scattering formula

$$N \propto \frac{1}{\sin^4(\theta/2)} \Rightarrow \frac{N_2}{N_1} = \left[\frac{\sin(\theta_1/2)}{\sin(\theta_2/2)} \right]^4$$

$$\Rightarrow \frac{N_2}{N_1} = \left[\frac{\sin \frac{90^\circ}{2}}{\sin \frac{60^\circ}{2}} \right]^4 = \left[\frac{\sin 45^\circ}{\sin 30^\circ} \right]^4$$

$\Rightarrow N_2 = (\sqrt{2})^4 \times N_1 = 4 \times 56 = 224$

97. (c) Change in the angular momentum

$$\Delta L = L_2 - L_1 = \frac{n_2 h}{2\pi} - \frac{n_1 h}{2\pi} \Rightarrow \Delta L = \frac{h}{2\pi} (n_2 - n_1)$$

$$= \frac{6.6 \times 10^{-34}}{2 \times 3.14} (5 - 4) = 1.05 \times 10^{-34} \text{ J-S}$$

98. (b) $E_n = -\frac{13.6}{n^2} \text{ eV}$

99. (a)

100. (b) $F = \frac{9 \times 10^9 \times 1.6 \times 10^{-19} \times 1.6 \times 10^{-19}}{(2.5 \times 10^{-11})^2} = 3.7 \times 10^{-7} \text{ N}$

101. (a) For Balmer series $\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{n^2} \right)$ where $n = 3, 4, 5$

For second line $n = 4$

So $\frac{1}{\lambda} = R \left(\frac{1}{2^2} - \frac{1}{4^2} \right) = \frac{3}{16} R \Rightarrow \lambda = \frac{16}{3R}$

102. (c) Energy of electron in H atom $E_n = \frac{-13.6}{n^2} \text{ eV}$

$\Rightarrow -1.5 = \frac{-13.6}{n^2} \Rightarrow n^2 = \frac{13.6}{1.5} = 3$

Now angular momentum

$$p = n \frac{h}{2\pi} = \frac{3 \times 6.6 \times 10^{-34}}{2 \times 3.14} = 3.15 \times 10^{-34} \text{ J} \times \text{sec}$$

103. (a)

104. (b) $T = \frac{2\pi r}{v}$; $r =$ radius of n orbit = $\frac{n^2 h^2}{\pi m Z e^2}$

$v =$ speed of e^- in n orbit = $\frac{ze^2}{2\epsilon_0 n h}$

$\therefore T = \frac{4\epsilon_0^2 n^3 h^3}{m Z^2 e^4} \Rightarrow T \propto \frac{n^3}{Z^2}$

105. (d) $r_n \propto n^2 \Rightarrow \frac{r_4}{r_1} = \left(\frac{4}{1} \right)^2 = \frac{16}{1} \Rightarrow r_4 = 16 r_1 \Rightarrow r_4 = 16 r_0$

106. (b) For Lyman series

$\bar{\nu} = \frac{1}{\lambda} = R \left(\frac{1}{1^2} - \frac{1}{n^2} \right)$ here $n = 2, 3, 4, 5, \dots$

For first line

$\bar{\nu} = R \left(\frac{1}{1^2} - \frac{1}{2^2} \right) \Rightarrow \bar{\nu} = R \left(1 - \frac{1}{4} \right) = \frac{3R}{4}$

107. (b) Energy $E = K \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$ ($K =$ constant)

$n = 2$ and $n = 3$, so $E = K \left[\frac{1}{2^2} - \frac{1}{3^2} \right] = K \left[\frac{5}{36} \right]$

For removing an electron $n = 1$ to $n_2 = \infty$

Energy $E_1 = K[1] = \frac{36}{5} E = 7.2 E$

\therefore Ionization energy = $7.2 E$

108. (c) For Paschen series $\bar{\nu} = \frac{1}{\lambda} = R \left[\frac{1}{3^2} - \frac{1}{n^2} \right]$; $n = 4, 5, 6, \dots$

For first member of Paschen series $n = 4$

$\frac{1}{\lambda_1} = R \left[\frac{1}{3^2} - \frac{1}{4^2} \right] \Rightarrow \frac{1}{\lambda_1} = \frac{7R}{144}$

$\Rightarrow R = \frac{144}{7\lambda_1} = \frac{144}{7 \times 18800 \times 10^{-10}} = 1.1 \times 10^{-7}$

For shortest wave length $n = \infty$

So $\frac{1}{\lambda} = R \left[\frac{1}{3^2} - \frac{1}{\infty^2} \right] = \frac{R}{9}$

$$\Rightarrow \lambda = \frac{9}{R} = \frac{9}{1.1 \times 10^{-7}} = 8.225 \times 10^{-7} m = 8225 \text{ \AA}$$

109. (d) For Lyman series $\frac{1}{\lambda_{\max}} = R \left[\frac{1}{1^2} - \frac{1}{2^2} \right] = \frac{3}{4} R$ and

$$\frac{1}{\lambda_{\min}} = R \left[\frac{1}{1^2} - \frac{1}{\infty^2} \right] = \frac{R}{1} \Rightarrow \frac{\lambda_{\max}}{\lambda_{\min}} = \frac{4}{3}$$

110. (c) $T \propto n^3 \Rightarrow \frac{T_2}{T_1} = \frac{2^3}{1^3} = \frac{8}{1}$

111. (a) For Bracket series $\frac{1}{\lambda_{\max}} = R \left[\frac{1}{4^2} - \frac{1}{5^2} \right] = \frac{9}{25 \times 16} R$

and $\frac{1}{\lambda_{\min}} = R \left[\frac{1}{4^2} - \frac{1}{\infty^2} \right] = \frac{R}{16} \Rightarrow \frac{\lambda_{\max}}{\lambda_{\min}} = \frac{25}{9}$

112. (a) For hydrogen and hydrogen like atoms $E_n = -13.6 \frac{z^2}{n^2} eV$

$$U_n = 2E_n = -27.2 \frac{z^2}{n^2} eV \text{ and } K_n = |E_n| = 13.6 \frac{z^2}{n^2} eV$$

From these three relations we can see that as n decreases, K will increase but E and U will decrease.

113. (a) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \Rightarrow \frac{\lambda_{\min}}{\lambda_{\max}} = \frac{\left[\frac{1}{2^2} - \frac{1}{3^2} \right]}{\left[\frac{1}{2^2} - \frac{1}{\infty} \right]} = \frac{5}{9}$

114. (b) By using $r_n = r_0 \frac{n^2}{Z}$; Where r_n = Radius of the Bohr orbit in the ground state atom. So for He^+ third excited state $n = 4, Z = 2, r_0 = 0.5 \text{ \AA} \Rightarrow r_4 = 0.5 \times \frac{4^2}{2} = 4 \text{ \AA}$

115. (d) Speed of electron in n orbit (in CGS) $v_n = \frac{2\pi Ze^2}{nh} (k=1)$

For first orbit $H_2; n=1$ and $Z=1$

$$\text{So } v = \frac{2\pi e^2}{h} \Rightarrow \frac{v}{c} = \frac{2\pi e^2}{hc}$$

116. (b)

117. (d) $E_n = -\frac{13.6}{n^2} eV \Rightarrow E_5 = \frac{-13.6}{5^2} = \frac{-13.6}{25} = -0.54 eV$

118. (a)

119. (b)

120. (a) $mvr = \frac{nh}{2\pi} \Rightarrow v = \frac{nh}{2\pi mr} \Rightarrow \frac{v^2}{r} = \frac{n^2 h^2}{4\pi^2 m^2 r^3}$

121. (b) Maximum number of spectral lines are observed in Lyman series.

122. (b) Wave number $\bar{\nu} = \frac{1}{\lambda} = \frac{1}{5896 \times 10^{-8}} = 16961 \text{ per cm}$

123. (b) $r_n \propto n^2 \Rightarrow \frac{r_3}{r_1} = \frac{3^2}{1} \Rightarrow r_3 = 9r_1 = 9 \times 0.53 = 4.77 \text{ \AA}$

124. (d) For first line in Lyman series $\lambda_{L_1} = \frac{4}{3R}$ (i)

For first line in Balmer series $\lambda_{B_1} = \frac{36}{5R}$ (ii)

From equation (i) and (ii)

$$\frac{\lambda_{B_1}}{\lambda_{L_1}} = \frac{27}{5} \Rightarrow \lambda_{B_1} = \frac{27}{5} \lambda_{L_1} \Rightarrow \lambda_{B_1} = \frac{27}{5} \lambda$$

125. (a)

126. (d) $3-1$ transition has higher energy so it has higher frequency $\left(\nu = \frac{E}{h} \right)$

127. (d) α -particles cannot be attracted by the nucleus.

128. (c) By using $\nu = RC \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$

$$\Rightarrow \nu = 10^7 \times (3 \times 10^8) \left[\frac{1}{4^2} - \frac{1}{5^2} \right] = 6.75 \times 10^8 \text{ Hz}$$

129. (c) For M shell ($n=3$), orbital quantum number $l=0, 1, 2$.

130. (d) Number of possible emission lines = $\frac{n(n-1)}{2}$

Where $n=4$; Number = $\frac{4(4-1)}{2} = 6$.

131. (a) Diameter of nucleus is of the order of $10^{-14} m$ and radius of first Bohr orbit of hydrogen atom $r = 0.53 \times 10^{-10} m$.

132. (c) The electron is in the second orbit ($n=2$)

Hence $L = \frac{nh}{2\pi} = \frac{2h}{2\pi} = \frac{6.6 \times 10^{-34}}{\pi} = 2.11 \times 10^{-34} \text{ J-sec}$

133. (c) $\frac{1}{\lambda} = RZ^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \Rightarrow \lambda \propto \frac{1}{Z^2}$

$$\lambda_{Li^{++}} : \lambda_{He^+} : \lambda_H = 4 : 9 : 36$$

134. (a) Energy radiated $E = 10.2 eV = 10.2 \times 1.6 \times 10^{-19} J$

$$\Rightarrow E = \frac{hc}{\lambda} \Rightarrow \lambda = 1.215 \times 10^{-7} m$$

135. (c) For $n=1, E_1 = -\frac{13.6}{(1)^2} = -13.6 eV$

and for $n=3, E_3 = -\frac{13.6}{(3)^2} = -1.51 eV$

So required energy

$$= E_3 - E_1 = -1.51 - (-13.6) = 12.09 eV$$

136. (a) Similar to Q. 115

137. (c) Since in spectral series of hydrogen atom, Lyman series lies lower Balmer series.

138. (d) $mvr_n = \frac{nh}{2\pi} \Rightarrow pr_n = \frac{nh}{2\pi} \Rightarrow \frac{h}{\lambda} \times r_n = \frac{nh}{2\pi}$

$$\Rightarrow \lambda = \frac{2\pi r_n}{n}, \text{ for first orbit } n=1 \text{ so } \lambda = 2\pi r_1$$

= circumference of first orbit

139. (d) $E_{n_1 \rightarrow n_2} = -13.6 \left[\frac{1}{n_2^2} - \frac{1}{n_1^2} \right]; n_1 = 2 \text{ \& } n_2 = 1$

$$\Rightarrow E_{II} \rightarrow E_I = -13.6 \times \frac{3}{4} = -10.2 eV$$

140. (b) $E_n = -\frac{13.6z^2}{n^2} eV \Rightarrow E_1 = -\frac{13.6 \times (2)^2}{(1)^2} = -54.4 eV$

141. (b) $v \propto Z^2 \Rightarrow \frac{v_{H_2}}{v_{He}} = \left(\frac{1}{2}\right)^2 = \frac{1}{4} \Rightarrow v_{He} = 4v_{H_2} = 4V$

142. (a) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$


First condition $\frac{1}{\lambda} = R \left[\frac{1}{1^2} - \frac{1}{2^2} \right] \Rightarrow R = \frac{4}{3\lambda}$

Second condition $\frac{1}{\lambda'} = R \left[\frac{1}{1^2} - \frac{1}{3^2} \right]$
 $\Rightarrow \lambda' = \frac{9}{8R} \Rightarrow \lambda' = \frac{9}{8 \times \frac{4}{3\lambda}} = \frac{27\lambda}{32}$

143. (b)

144. (a) $r_n \propto n^2$


145. (d) $\because E_2 < E_1 \Rightarrow \lambda_2 > \lambda_1$



146. (a) Wave number $\bar{\nu} = \frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]; n_2 = \infty$ and $n_1 = 1$

$\Rightarrow \bar{\nu} = R = 1.097 \times 10^7 m^{-1} = 109700 cm^{-1}$

147. (b) $E_1 > E_2 > E_3$



148. (d) $r \propto \frac{1}{Z}$, for double ionized lithium $Z(=3)$ will be maximum.

So r will be minimum

149. (b) $E_n = \frac{13.6}{n^2} \times Z^2$. For first excited state $n = 2$ and for

$Li^{++}, z = 3 \Rightarrow E = \frac{13.6}{4} \times 9 = 30.6 eV$

150. (b)

151. (a) In Lyman series $(\lambda_{min})_L = \frac{1}{R}$ and $(\lambda_{min})_B = \frac{4}{R}$

$\Rightarrow (\lambda_{min})_B = 4 \times (\lambda_{min})_L = 4 \times 912 = 3648 \text{ \AA}$

152. (d)



$E_{3 \rightarrow 2} = -3.4 - (-1.51) = -1.89 eV \Rightarrow |E_{3 \rightarrow 2}| \approx 1.9 eV$

153. (a)

154. (a)

155. (c) Radius of n orbit for any hydrogen like atom

$r_n = r_0 \left(\frac{n^2}{Z} \right)$ ($r_0 =$ radius of first orbit of H_2 -atom)

If $r_n = r_0 \Rightarrow n = \sqrt{Z}$. For $Be, Z = 4 \Rightarrow n = 2$.

156. (d) $r_n \propto n^4 \Rightarrow A_n \propto n^4 \Rightarrow \frac{A_1}{A_0} = \left(\frac{2}{1}\right)^4 = \frac{16}{1}$

157. (d)

158. (d) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$. For first wavelength, $n_1 = 2, n_2 = 3$

$\Rightarrow \lambda_1 = 6563 \text{ \AA}$. For second wavelength, $n_1 = 2, n_2 = 4$
 $\Rightarrow \lambda_2 = 4861 \text{ \AA}$

159. (c) K.E. = - (Total energy) = - (-13.6 eV) = + 13.6 eV

160. (a) In Lyman series $\lambda_{max} = \frac{4}{3R}$

In Balmer series $\lambda_{max} = \frac{36}{5R}$. So required ratio = $\frac{5}{27}$

161. (c)

162. (b) $E = 13.6 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$. For highest energy in Balmer series

$n_1 = 2$ and $n_2 = \infty \Rightarrow E = 13.6 \left[\frac{1}{(2)^2} - \frac{1}{(\infty)^2} \right] = 3.4 eV$

163. (a)

164. (c) $T \propto n^3$

165. (b) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] = R \left[\frac{1}{(2)^2} - \frac{1}{(4)^2} \right] \Rightarrow \lambda = \frac{16}{3R}$

166. (a) $E_n \propto Z^2 \Rightarrow \frac{(E_n)_{He}}{(E_n)_H} = \frac{Z_{He}^2}{Z_H^2} = 4 \Rightarrow (E_n)_{He} = 4 \times (E_n)_H$

167. (b) By using $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$

168. (a) $\frac{1}{\lambda_{max}} = R \left[\frac{1}{(1)^2} - \frac{1}{(2)^2} \right] \Rightarrow \lambda_{max} = \frac{4}{3R} \approx 1213 \text{ \AA}$

and $\frac{1}{\lambda_{min}} = R \left[\frac{1}{(1)^2} - \frac{1}{\infty} \right] \Rightarrow \lambda_{min} = \frac{1}{R} \approx 910 \text{ \AA}$

169. (b) P.E. = $2 \times$ Total energy = $2 \times (-13.6) = -27.2 eV$

170. (c) Emitted energy $\Delta E = \frac{hc}{\lambda} \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$.

171. (c) $U = 2E, K = -E$ and $E = -\frac{13.6}{n^2} = eV$.

172. (a)

173. (d)

174. (c) $r \propto n^2$

Nucleus, Nuclear Reaction

1. (b)

2. (c) Neutrons are neutral particles.

3. (c) James Chadwick discovered the neutron.

4. (d) In hydrogen, atomic number and mass number are equal.
5. (d) $E = mc^2 = 1 \times (3 \times 10^8)^2 = 9 \times 10^{16} \approx 10^{17} J$
6. (d) $B.E. = \Delta m \text{ amu} = \Delta m \times 931 \text{ MeV}$.
7. (d) Mass defect $\Delta m = \frac{2.23}{931} = 0.0024$.
8. (b) Positron is the antiparticle of electron.
9. (b)
10. (c)
11. (a) $B.E. = \Delta mc^2 = [2(1.0087 + 1.0073) - 4.0015] = 28.4 \text{ MeV}$
12. (b) $\frac{\text{Binding energy}}{\text{Nucleon}} = \frac{0.0303 \times 931}{4} \approx 7$
13. (d) Energy / day = $200 \times 10^6 \times 24 \times 3600$
 $= 2 \times 2.4 \times 3.6 \times 10^{12} = 1728 \times 10^{10} J$
14. (c) $E = \Delta mc^2 = 10^{-6} \times (3 \times 10^8)^2 = 9 \times 10^{10} J$
15. (c)
16. (c) Mass of ${}_1H^2 = 2.01478 \text{ a.m.u.}$
 Mass of ${}_2He^4 = 4.00388 \text{ a.m.u.}$
 Mass of two deuterium = $2 \times 2.01478 = 4.02956$
 Energy equivalent to $2{}_1H^2$
 $= 4.02956 \times 1.112 \text{ MeV} = 4.48 \text{ MeV}$
 Energy equivalent to ${}_2He^4$
 $= 4.00388 \times 7.047 \text{ MeV} = 28.21 \text{ MeV}$
 Energy released = $28.21 - 4.48 = 23.73 \text{ MeV} = 24 \text{ MeV}$
17. (c) Energy released while forming a nucleus is known as binding energy (by definition).
18. (c) Nuclear force is stronger than coulomb force.
19. (d)
20. (c)
21. (b) $Q = 4(x - x)$
22. (d)
23. (a) Rest energy of an electron = $m_e c^2$
 Here $m_e = 9.1 \times 10^{-31} \text{ kg}$ and $c = \text{velocity of light}$
 $\therefore \text{Rest energy} = 9.1 \times 10^{-31} \times (3 \times 10^8)^2 \text{ joule}$
 $= \frac{9.1 \times 10^{-31} \times (3 \times 10^8)^2}{1.6 \times 10^{-19}} \text{ eV} = 510 \text{ keV}$
24. (b) ${}_z X^A = {}_{88} Ra^{226}$
 Number of protons = $Z = 88$
 Number of neutrons = $A - Z = 226 - 88 = 138$.
25. (c) Out side the Nucleus, neutron is unstable (life $\approx 932 \text{ sec}$).
26. (b) The order of magnitude of mass and volume of uranium nucleus will be
 $m \approx A(1.67 \times 10^{-27} \text{ kg})$ (A is atomic number)
 $V = \frac{4}{3} \pi r^3 \approx \frac{4}{3} \pi [(1.25 \times 10^{-15} \text{ m})A^{1/3}]^3$
 $\approx (8.2 \times 10^{-45} \text{ m}^3)A$

Hence, $\rho = \frac{m}{V} = \frac{A(1.67 \times 10^{-27} \text{ kg})}{(8.2 \times 10^{-45} \text{ m}^3)A}$
 $\approx 2.0 \times 10^{17} \text{ kg/m}^3$.

27. (c) We have $r \propto A^{1/3} \Rightarrow \frac{r_2}{r_1} = \left(\frac{A_2}{A_1}\right)^{1/3} = \left(\frac{206}{4}\right)^{1/3}$
 $\therefore r_2 = 3\left(\frac{206}{4}\right)^{1/3} = 11.6 \text{ Fermi}$.
28. (c) Nucleus does not contains electron.
29. (a) Let the percentage of B^{10} atoms be x , then Average atomic weight
 $= \frac{10x + 11(100 - x)}{100} = 10.81 \Rightarrow x = 19 \therefore \frac{N_{B^{10}}}{N_{B^{11}}} = \frac{19}{81}$
30. (a)
31. (b)
32. (a) Nuclear force is charge independent, it also acts between two neutrons.
33. (d) $p \rightarrow \pi^+ + n$, $n \rightarrow p + \pi^-$ and $n \rightarrow n' + \pi^0$
34. (c) Helium nucleus $\rightarrow {}_2He^4$.
 Number of protons = $Z = 2$
 Number of Neutrons = $A - Z = 2$.
35. (c)
36. (b) Binding energy per nucleon increases with atomic number and is maximum for iron. After that it decrease.
-
37. (a) For isotopes Z is same and A is different. Therefore the number of neutrons $A - Z$ will also be different.
38. (a) This is due to mass defect because a part of mass is used in keeping the neutrons and protons bound as α -particle.
39. (d) $B.E. \text{ of } Li^7 = 39.20 \text{ MeV}$ and $He^4 = 28.24 \text{ MeV}$
 Hence binding energy of $2He^4 = 56.48 \text{ MeV}$
 Energy of reaction = $56.48 - 39.20 = 17.28 \text{ MeV}$.
40. (c) $r \propto (A)^{1/3}$
41. (b) $r \propto A^{1/3}$
42. (a) $E = mc^2 = (1 \times 10^{-3})(3 \times 10^8)^2 = 9 \times 10^{13} J$.
43. (c) $\Delta E = 8.5 \times 234 - 7.6 \times 236 = 195.4 \text{ MeV} = 200 \text{ MeV}$.
44. (a) $N = M - Z = \text{Total no. of nucleons} - \text{no. of protons}$.
45. (c)
46. (c) Both coulomb and nuclear force act inside the nucleus.
47. (d) For stability in case of lighter nuclei $\frac{N}{Z} = 1$ and for heavier nuclei $\frac{N}{Z} > 1$.
48. (b) Nuclear forces are charge independent.

1512 Atomic and Nuclear Physics

49. (a) Actual mass of the nucleus is always less than total mass of nucleons so

$$M < (NM_n + Zm_p).$$
50. (b) Mass of H_1 nucleus = mass of proton = 1 amu energy equivalent to 1 amu is 931 MeV so correct option is (b).
51. (d) $R = R_1 A \Rightarrow R \propto A.$
52. (b)
53. (d) Number of neutrons = $A - Z = 23 - 11 = 12.$
54. (c) For ${}_6C^{12}$, $p = 6, e = 6, n = 6$
 For ${}_6C^{14}$, $p = 6, e = 6, n = 8$
55. (a)
56. (a) The mass of nucleus formed is always less than the sum of the masses of the constituent protons and neutrons i.e.

$$m < (A - Z)m_n + Zm_p.$$
57. (d) $E = \Delta m.c^2 \Rightarrow E = \frac{0.3}{1000} \times (3 \times 10^8)^2 = 2.7 \times 10^{13} J$

$$= \frac{2.7 \times 10^{13}}{3.6 \times 10^6} = 7.5 \times 10^6 kWh.$$
58. (a, d)
59. (c) ${}_5B^{10} + {}_0n^1 \rightarrow {}_3Li^7 + {}_2He^4.$
60. (a)
61. (c) ${}_{92}U^{235}$ is normally fissionable.
62. (b)
63. (a)
64. (a) In atom bomb nuclear fission takes place with huge temperature.
65. (d) The given equation is ${}_2He^4 + {}_zX^A \rightarrow {}_{z+2}Y^{A+3} + A$
 Applying charge and mass conservation

$$4 + A = A + 3 + x \Rightarrow x = 1 \Rightarrow 2 + z = z + 2 + n \Rightarrow n = 0$$

 Hence A is a neutron.
66. (c) Energy of stars is due to the fusion of light hydrogen nuclei into He. In this process much energy is released.
67. (a) ${}_1H^2 + {}_1H^2 \rightarrow {}_2He^4 + 24 MeV.$
68. (b) Energy $\propto c^2$; \therefore Decrease in energy $\propto \frac{4}{9}.$
69. (b) Fusion reaction requires a very high temperature
 = (10 K).
70. (d) ${}_4Be^9 + {}_2He^4 \rightarrow {}_6C^{12} + {}_0n^1.$
71. (b)
72. (d)
73. (b, c)
74. (d)
75. (c) Cadmium rods absorb the neutrons so they are used to control the chain reaction process.
76. (d)
77. (d) No energy and mass enters or goes out of the system of the reaction and no external force is assumed to act.
78. (c)
79. (b)
80. (b) Energy of γ -ray photon = $0.5 + 0.5 + 0.78 = 1.78 MeV.$
81. (c)
82. (d)
83. (c) When fast moving neutrons pass through a moderator, they collide with the molecules of the moderator. As a result of this the neutrons are in thermal equilibrium with the surrounding molecules of moderator. These neutrons are called thermal neutrons.
84. (b) $E_b + E_c > E_a$
85. (d) Because sound waves require medium to travel through and there is no medium (air) on moon's surface.
86. (b) Heavy water is used as moderators in nuclear reactions to slow down the neutrons.
87. (a) $m = \frac{E}{c^2} = \frac{931 \times 1.6 \times 10^{-13}}{(3 \times 10^8)^2} = 1.66 \times 10^{-27} kg.$
88. (d) $E = \Delta mc^2, \Delta m = \frac{0.1}{100} = 10^{-3} kg$

$$\therefore E = 10^{-3} \times (3 \times 10^8)^2 = 10^{-3} \times 9 \times 10^{16} = 9 \times 10^{13} J.$$
89. (d) Energy released by γ -rays for pair production must be greater than 1.02 MeV.
90. (a) ${}_8O^{18} + {}_1H^1 \rightarrow {}_9F^{18} + {}_0n^1$
91. (c) Power = 1000 kW = $10^6 J/s$

$$\text{Rate of nuclear fission} = \frac{10^6}{200 \times 1.6 \times 10^{-13}} = 3.125 \times 10^9.$$
92. (b) $A = 238 - 4 = 234$ and $Z = 92 - 2 = 90.$
93. (a) $P = n \left(\frac{E}{t} \right) \Rightarrow 1000 = \frac{n \times 200 \times 10^6 \times 1.6 \times 10^{-19}}{t}$

$$\Rightarrow \frac{n}{t} = 3.125 \times 10^{13}.$$
94. (c) Due to the production of neutrons, a chain of nuclear fission is established which continues until the whole of the source substance is consumed.
95. (a) ${}_{92}U^{235} + {}_0n^1 \rightarrow {}_{38}Sr^{90} + {}_{54}Xe^{143} + 3{}_0n^1$
96. (b) ${}_1H^2 + {}_1H^2 \rightarrow {}_2He^4 + Q.$
97. (c) Fast neutrons can escape from the reaction. So as to proceed the chain reaction. Slow neutrons are best.
98. (d) ${}_1H^2 + {}_1H^2 \rightarrow {}_1H^3 + {}_1H^1$
99. (d)
100. (b) $\Delta m = 1 - 0.993 = 0.007 gm$

$$\therefore E = (\Delta m)c^2 = (0.007 \times 10^{-3})(3 \times 10^8)^2 = 63 \times 10^{10} J.$$
101. (c) ${}_{85}X^{297} \rightarrow {}_{77}Y^{281} + 4({}_2He^4)$
102. (b) $x+1 = 24 + 4 \Rightarrow x = 27.$
103. (a)
104. (d) ${}_6C^{11} \rightarrow {}_5B^{11} + \beta^+ + \gamma$ because $\beta^+ = {}_1e^0$
105. (c)
106. (b) $\frac{\text{Energy}}{\text{Fission}} = 200 MeV = 200 \times 10^6 \times 1.6 \times 10^{-19} J$

$$\text{Fission rate} = \frac{5}{200 \text{ MeV}} = 1.56 \times 10^{11} \text{ fission/sec.}$$

107. (a)
108. (b) Energy is released in the sun due to fusion.
109. (d)
110. (c) In nuclear fission, neutrons are released.
111. (c) ${}_1H^1 + {}_1H^1 + {}_1H^2 \rightarrow {}_2He^4 + {}_{+1}e^0 + \text{energy.}$
112. (c)
113. (b) ${}_0n^1 = {}_1p^1 + {}_{-1}e^0 + \bar{\nu}$
Antineutrino is required for conservation of spin.
114. (b)
115. (d) Fusion is the main process of energy production in the sun.
116. (a)
117. (b) Mass of proton = mass of antiproton
 $= 1.67 \times 10^{-27} \text{ kg} = 1 \text{ amu}$
Energy equivalent to $1 \text{ amu} = 931 \text{ MeV}$
So energy equivalent to $2 \text{ amu} = 2 \times 931 \text{ MeV}$
 $= 1862 \times 10^6 \times 1.6 \times 10^{-19} = 2.97 \times 10^{-10} \text{ J} = 3 \times 10^{-10} \text{ J.}$
118. (a) In fusion reaction, two lighter nuclei combines.
119. (c)
120. (b) Hydrogen bomb is based on nuclear fusion.
121. (d) ${}_{92}U^{235} + {}_0n^1 \rightarrow {}_{92}U^{236}$ and
 ${}_{92}U^{236} \rightarrow {}_{56}Ba^{144} + {}_{36}Kr^{89} + 3{}_0n^1 + Q.$
122. (d) Fusion reaction of deuterium is
 ${}_1H^2 + {}_1H^2 \rightarrow {}_2He^3 + {}_0n^1 + 3.27 \text{ MeV}$
So $E = \frac{6.02 \times 10^{23} \times 10^3 \times 3.27 \times 1.6 \times 10^{-13}}{2 \times 2} = 7.8 \times 10^{13} \text{ J}$
 $= 8 \times 10^{13} \text{ J.}$
123. (a)
124. (a)
125. (c)
126. (c)
127. (b)
128. (d) Energy released in the fission of one nucleus = 200 MeV
 $= 200 \times 10^6 \times 1.6 \times 10^{-19} \text{ J} = 3.2 \times 10^{-11} \text{ J}$
 $P = 16 \text{ KW} = 16 \times 10^3 \text{ Watt}$
Now, number of nuclei required per second
 $n = \frac{P}{E} = \frac{16 \times 10^3}{3.2 \times 10^{-11}} = 5 \times 10^{14}.$
129. (c)
130. (a) Number of fissions per second
 $= \frac{\text{Power output}}{\text{Energy released per fission}}$
 $= \frac{3.2 \times 10^6}{200 \times 10^6 \times 1.6 \times 10^{-19}} = 1 \times 10^{17}$

$$\Rightarrow \text{Number of fission per minute} = 60 \times 10^{17} = 6 \times 10^{18}$$


131. (c)
132. (a) $X(n, \alpha) {}_3Li^7 \Rightarrow {}_Z X^A + {}_0n^1 \rightarrow 3Li^7 + {}_2He^4$
 $Z = 3 + 2 = 5$ and $A = 7 + 4 - 1 = 10$
 $\therefore {}_5X^{10} = {}_5B^{10}$
133. (a)
134. (b) Mass of electron = mass of positron = $9.1 \times 10^{-31} \text{ kg}$
Energy released $E = (2m).c^2$
 $= 2 \times 9.1 \times 10^{-31} \times (3 \times 10^8)^2 = 1.6 \times 10^{-13} \text{ J.}$
135. (b) ${}_1H^2 + {}_1H^2 \rightarrow {}_2He^4 + \text{energy}$
Binding energy of a $({}_1H^2)$ deuterium nuclei
 $= 2 \times 1.1 = 2.2 \text{ MeV}$
Total binding energy of two deuterium nuclei
 $= 2.2 \times 2 = 4.4 \text{ MeV}$
Binding energy of a $({}_2He^4)$ nuclei = $4 \times 7 = 28 \text{ MeV}$
So, energy released in fusion = $28 - 4.4 = 23.6 \text{ MeV}$
136. (c) Mass of a uranium nucleus
 $= 92 \times 1.6725 \times 10^{-27} + 143 \times 1.6747 \times 10^{-27}$
 $= 393.35 \times 10^{-27} \text{ kg}$
Number of nuclei in the given mass
 $= \frac{1}{393.35 \times 10^{-27}} = 2.542 \times 10^{24}$
Energy released = $200 \times 2.542 \times 10^{24} \text{ MeV}$
 $= 5.08 \times 10^{26} \text{ MeV} = 8.135 \times 10^{13} \text{ J} = 8.2 \times 10^{13} \text{ J}$
137. (a)
138. (c)
139. (b) In a material medium, when a positron meets an electron both the particles annihilate leading to the emission of two γ ray photons. This process forms the basis of an important diagnostic procedure called PET.
140. (a) Total mass of reactants
 $= (2.0141) \times 2 = 4.0282 \text{ amu}$
Total mass of products = 4.0024 amu
Mass defect = $4.0282 \text{ amu} - 4.0024 \text{ amu}$
 $= 0.0258 \text{ amu}$
 \therefore Energy released $E = 931 \times 0.0258 = 24 \text{ MeV}$
141. (b) ${}_7N^{14} + {}_2He^4 \rightarrow {}_8O^{17} + {}_1H^1$
142. (b)
143. (b)
144. (d) ${}_8O^{16} + {}_1H^2 \rightarrow {}_7N^{14} + {}_2He^4$

1514 Atomic and Nuclear Physics

145. (d)
 146. (a)
 147. (b) Nuclear fusion takes place in stars which results in joining of nuclei accompanied by release of tremendous amount of energy.
 148. (b)
 149. (d) $B.E.$ per nucleon \propto stability.
 150. (a) Nuclei of different elements having the same mass number are called isotones e.g., ${}_4Be^9$ and ${}_5B^{10}$
 151. (c)
 152. (b)
 153. (d) Packing fraction = $\frac{M - A}{A}$
 154. (c) $B = [ZM_p + NM_n - M(N, Z)]c^2$
 $\Rightarrow M(M, Z) = ZM_p + NM_n - B/c^2$
 155. (c)
 156. (a) In nuclear reactor, nuclear fission can be carried out through a sustained and a controlled chain reaction.
 157. (b) ${}_6C^{12} + {}_0n^1 \rightarrow {}_7N^{13} + {}_{-1}\beta^0$
 158. (a) ${}_0n^1 + {}_{92}U^{235} \rightarrow {}_{56}Ba^{144} + {}_{36}Kr^{89} + 3{}_0n^1$
 159. (c) The energy released in sun and hydrogen bomb are due to nuclear fusion.
 160. (c)
 161. (a)
 162. (a) ${}_{84}Po^{210} \rightarrow {}_{82}X^{206} + {}_2He^4$
 Using conservation of linear moments
 $206v' + 4v = 0 \Rightarrow v' = -\frac{4v}{206} \Rightarrow |v'| = \frac{4v}{206}$
 163. (b) Power $P = \frac{\text{Energy}}{\text{time}} = \frac{mc^2}{t}$, $= 1 \times 10 \times (3 \times 10)^4$
 $= 9 \times 10^4 W = 9 \times 10^4 kW$
 164. (b)
 165. (c)
 166. (b) The elements high on the $B.E.$ versus mass number plot are very tightly bound and hence, are stable. And the elements those are lower on this plot, are less tightly bound and hence, are unstable.
 Since helium nucleus shows a peak on this plot so, it is very stable.
 167. (d)
 168. (d) $E = \Delta mc^2 = 1 \times (3 \times 10^8)^2 = 9 \times 10^{10} J$
 $\Rightarrow E = \frac{9 \times 10^{16}}{1.6 \times 10^{-19}} = 5.625 \times 10^{35} eV = 5.625 \times 10^{29} MeV$
 169. (a) $E = \Delta mc^2 = 0.5 \times 10^{-3} \times (3 \times 10^8)^2 = 4.5 \times 10^{13} j$

$$\Rightarrow E = \frac{4.5 \times 10^{13}}{3.6 \times 10^6} = 1.25 \times 10^7 kWh.$$

170. (b) ${}_0n^1 + {}_{92}U^{235} \rightarrow {}_{36}Kr^{94} + {}_{56}Ba^{139} + 3{}_0n^1$
 171. (b)
 172. (a) Number of protons = $2 + 2 + 6 + 2 + 6 = 18$
 Number of neutrons = $40 - 18 = 22$.
 173. (c) Neutrons are unstable and having mean life time of 32 sec, decaying by emitting an electron and antineutrino to become proton.
 174. (b) During fusion binding energy of daughter nucleus is always greater than the total energy of the parent nuclei so energy released = $c - (a + b) = c - a - b$
 175. (a) These nuclei having different Z and A but equal $(A - Z)$ are called isotones.



177. (a)
 178. (c) $r \propto A^{1/3} \Rightarrow \frac{r_1}{r_2} = \left(\frac{A_1}{A_2}\right)^{1/3}$
 $\Rightarrow \frac{3.6}{r_2} = \left(\frac{27}{125}\right)^{1/3} = \frac{3}{5} \Rightarrow r_2 = 6 \text{ Fermi}$
 179. (b)

Radioactivity

1. (a)
 2. (a) By formula $N = N_0 \left(\frac{1}{2}\right)^{t/T}$ or $10^4 = 8 \times 10^4 \left(\frac{1}{2}\right)^{t/3}$
 or $\left(\frac{1}{8}\right) = \left(\frac{1}{2}\right)^{t/3}$ or $\left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{t/3} \Rightarrow 3 = \frac{t}{3}$
 Hence $t = 9 \text{ years}$.
 3. (d) Fraction = $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{\frac{6400}{1600}} = \left(\frac{1}{2}\right)^4 = \frac{1}{16}$
 4. (c) Negative β -decay is expressed by the equation
 $n = p^+ + e^- + \nu^-$
 5. (a) No radioactive substance emit both α and β particles simultaneously. Some substances emit α -particles and some other emits β -particles. γ -rays are emitted along with both α and β -particles.
 6. (c) γ -rays are highly penetrating.

7. (c) Average life $\frac{1}{\lambda} = \frac{1600}{0.693} = 2308 \approx 2319$ years.
8. (d) Fraction of atoms remains after five half lives

$$\frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T} = \left(\frac{1}{2}\right)^{5T/T} = \frac{1}{32}$$

$$\Rightarrow \text{Percentage atom remains} = \frac{1}{32} \times 100 = 3.125\%$$
9. (c) β -rays emitted from nucleus and they carry negative charge.
10. (c)
11. (b) ${}_Z X^A \xrightarrow{-1\beta^0} {}_{Z+1} Y^A \xrightarrow{2He^4(\alpha)} {}_{Z-1} K^{A-4} \xrightarrow{-0e^0} {}_{Z-1} K^{A-4}$
12. (c) $N_t = N_0 \left(\frac{1}{2}\right)^{t/T} = 50000 \left(\frac{1}{2}\right)^{10/5} = 12500$
13. (c) ${}_Z X^A \xrightarrow{-1\beta^0} {}_{Z+1} X^A$
14. (c) $N = N_0 \left(\frac{1}{2}\right)^{t/T} \Rightarrow \frac{N_0}{64} = N_0 \left(\frac{1}{2}\right)^{30/T} \Rightarrow T = \frac{30}{6} = 5$ sec
15. (a)
16. (c)
17. (a) Average life $T = \frac{\text{Sum of all lives of all the atom}}{\text{Total number of atoms}} = \frac{1}{\lambda}$

$$\Rightarrow T\lambda = 1$$
18. (c) Fraction remains after n half lives $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n = \left(\frac{1}{2}\right)^{t/T}$

$$\therefore \frac{N}{N_0} = \left(\frac{1}{2}\right)^{T/2} = \left(\frac{1}{2}\right)^{1/2} = \frac{1}{\sqrt{2}}$$
19. (b)
20. (a) Penetration power of γ is 100 times of β , while that of β is 100 times of α .
21. (a) $\frac{N_0}{32} = N_0 \left(\frac{1}{2}\right)^{60/T} \Rightarrow 5 = \frac{60}{T} \Rightarrow T = 12$ days
22. (c) By using $N = N_0 \left(\frac{1}{2}\right)^{t/T}$; where $N = \left(1 - \frac{7}{8}\right) N_0 = \frac{1}{8} N_0$
 So $\frac{1}{8} N_0 = N_0 \left(\frac{1}{2}\right)^{t/T} \Rightarrow \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{t/5} \Rightarrow t = 15$ days.
23. (a) ${}_{72}A^{180} \xrightarrow{\alpha} {}_{70}A_1^{176} \xrightarrow{\beta} {}_{71}A_2^{176} \xrightarrow{\alpha} {}_{69}A_3^{172} \xrightarrow{\gamma} {}_{69}A_4^{172}$
24. (d)
25. (d) Half life of a substance doesn't depend upon Amount, temperature and pressure. It depends upon the nature of the substance.
26. (d) $T = \frac{0.6931 \times 1}{\lambda} = \frac{0.6931 \times 1}{4.28 \times 10^{-4}} \text{ year} = 1620$ years
27. (c) In fusion two lighter nuclei combine, it is not the radioactive decay.
28. (b) $n_\alpha = \frac{A - A'}{4} = \frac{232 - 208}{4} = 6$
 and $n_\beta = (2n_\alpha - Z + Z') = (2 \times 6 - 90 + 82) = 4$
29. (a) Remaining amount

$$= 16 \times \left(\frac{1}{2}\right)^{32/2} = 16 \times \left(\frac{1}{2}\right)^{16} = \left(\frac{1}{2}\right)^{12} < 1 \text{ mg}$$
30. (c) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{15/5} = \frac{1}{8} \Rightarrow \text{Decayed fraction} = 1 - \frac{1}{8} = \frac{7}{8}$
31. (c)
32. (c) By using $n_\alpha = \frac{A - A'}{4}$ and $n_\beta = 2n_\alpha - Z + Z'$

$$\Rightarrow A' = A - 4n_\alpha = 236 - 4 \times 3 = 224$$

 and $Z' = (n_\beta - 2n_\alpha + Z) = (1 - 2 \times 3 + 88) = 83$
33. (d) Uncertain, because it is infinite. No radioactive element can be disintegrated fully.
34. (c) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/140} \Rightarrow \frac{1}{16} = \left(\frac{1}{2}\right)^{t/140}$

$$\Rightarrow \frac{t}{140} = 4 \Rightarrow t = 560 \text{ days}$$
35. (c) $\frac{C_{14}}{C_{12}} = \frac{1}{4} = \left(\frac{1}{2}\right)^{t/5700} \Rightarrow \frac{t}{5700} = 2 \Rightarrow t = 11400$ years
36. (b) Ionising property depends upon the charge and mass.
37. (b) $R = \frac{dN}{dt} \propto N \Rightarrow \frac{R_2}{R_1} = \frac{N_2}{N_1}$
 But $\frac{N_2}{N_1} = \left(\frac{1}{2}\right)^{t/2} \Rightarrow \frac{25}{200} = \frac{1}{8} = \left(\frac{1}{2}\right)^3 \Rightarrow \frac{t}{t_{1/2}} = 3$

$$\therefore t_{1/2} = \frac{t}{3} = \frac{3}{3} = 1 \text{ hour} = 60 \text{ minutes}$$
38. (d) $t_{1/2} = \frac{0.6931}{0.01} = 69.31$ seconds.
39. (d) Because radioactivity is a spontaneous phenomenon.
40. (d) Undecayed isotope $= 1 - \frac{7}{8} = \frac{1}{8}$

$$\therefore \frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T} \Rightarrow \left(\frac{1}{8}\right) = \left(\frac{1}{2}\right)^{t/15} \Rightarrow \frac{t}{15} = 3$$

 or $t = 45$ hours
41. (a) Mean life $= \frac{\text{Half life}}{0.6931} = \frac{10}{0.6931} = 14.4$ hours

1516 Atomic and Nuclear Physics

42. (b) 20 gm substance reduces to 10 gm (i.e. becomes half in 4 min.
So $T_{1/2} = 4 \text{ min}$. Again $M = M_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$
 $\Rightarrow 10 = 20 \left(\frac{1}{2}\right)^{t/4} \Rightarrow \frac{1}{2} = \left(\frac{1}{2}\right)^{t/4} \Rightarrow t = 4 \text{ min}$
43. (c) $N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow 1 = 16 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow T_{1/2} = \frac{1}{2} \text{ hour}$
44. (d)
45. (b) β -decay from nuclei based on this process only.
46. (b) $A = A_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow 5 = A_0 \left(\frac{1}{2}\right)^{30/T_{1/2}} = \frac{A_0}{16} \Rightarrow A_0 = 80 \text{ sec}^{-1}$
47. (d) $n_\alpha = \frac{A - A'}{4} = \frac{200 - 168}{4} = 8$
 $n_\beta = 2n_\alpha - Z + Z' = 2 \times 8 - 90 + 80 = 6$
48. (d) Similar to Q. 47
49. (b) $N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$. Hence fraction of atoms decayed
 $= 1 - \frac{N}{N_0} = 1 - \left(\frac{1}{2}\right)^{t/T_{1/2}} = 1 - \left(\frac{1}{2}\right)^{3 \times 60 / 60} = \frac{7}{8}$
In percentage it is $\frac{7}{8} \times 100 = 87.5\%$
50. (a) C-14 is carbon dating substance.
51. (b) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T} \Rightarrow \left(\frac{1}{16}\right) = \left(\frac{1}{2}\right)^{2/T} \Rightarrow \left(\frac{1}{2}\right)^4 = \left(\frac{1}{2}\right)^{2/T}$
 $\Rightarrow T = 0.5 \text{ hour} = 30 \text{ minutes}$.
52. (c) $\frac{dN}{dt} = -\lambda N \Rightarrow n = -\lambda N$ (Given $\frac{dN}{dt} = n$)
 $\therefore \lambda = -\frac{n}{N} \therefore \text{Half life} = \frac{0.693}{\lambda} = \frac{0.693}{\frac{n}{N}} = \frac{0.693 N}{n} \text{ sec}$
53. (a) ${}_{92}\text{X}^{235} \xrightarrow{\alpha} {}_{90}\text{X}^{231} \xrightarrow{-1e^0} {}_{91}\text{Y}^{231}$
54. (d) ${}_{7}\text{N}^{13} \rightarrow {}_{6}\text{C}^{13} + {}_{+1}\text{e}^0$
55. (c) $A = A_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow 100 = 1600 \left(\frac{1}{2}\right)^{8/T_{1/2}} \Rightarrow T_{1/2} = 2 \text{ sec}$
Again at $t = 6 \text{ sec}$, $A = 1600 \left(\frac{1}{2}\right)^{6/2} = 200 \text{ counts/sec}$
56. (d) ${}_{92}\text{U}^{238} \rightarrow {}_{92}\text{Th}^{234} + {}_2\text{He}^4$
57. (b)
58. (d)
59. (d)
60. (b) By using $N = N_0 e^{-\lambda t} \Rightarrow \frac{N_0}{2} = N_0 e^{-\lambda T_{1/2}} \Rightarrow 2 = e^{\lambda T_{1/2}}$
By taking log both the side
 $\log_e 2 = \lambda T_{1/2} \Rightarrow \lambda T_{1/2} = 0.693$
61. (a) Number of half lives in 20 min = $n = \frac{20}{5} = 4$
Fraction of material remains after four half lives = $\frac{1}{16}$
Hence fraction that decays = $1 - \frac{1}{16} = \frac{15}{16} = 93.75\%$
62. (d) In the given case, 12 days = 3 half lives Number of atoms left after 3 half lives.
 $= 6.4 \times 10^{10} \times \frac{1}{2^3} = 0.8 \times 10^{10}$
63. (a) Decay constant remains unchanged in a chemical reaction.
64. (d) $n_\alpha = \frac{A - A'}{4} = \frac{238 - 206}{4} = 8$
65. (b) ${}_Z\text{X}^A \xrightarrow{\alpha} {}_{Z-2}\text{Y}^{A-4} \xrightarrow{-2\beta} {}_Z\text{X}^{A-4}$
66. (a) Both the β -rays and the cathode rays are made up of electrons. γ -rays are EM waves, α -particles are doubly ionized helium atoms and protons and neutrons have approximately the same mass.
67. (b) ${}_{10}^{22}\text{Ne} \rightarrow {}_2^4\text{He} + {}_6^4\text{He} + {}_6^{14}\text{X}$; hence X is carbon.
68. (c) For 80 minutes, number of half lives of sample
 $A = n_A = \frac{80}{20} = 4$ and number of half lives of sample
 $B = n_B = \frac{80}{40} = 2$. Also by using $N = N_0 \left(\frac{1}{2}\right)^n$
 $\Rightarrow N \propto \frac{1}{2^n} \Rightarrow \frac{N_A}{N_B} = \frac{2^{n_B}}{2^{n_A}} = \frac{2^2}{2^4} = \frac{1}{4}$
69. (d) ${}_n\text{X}^m \xrightarrow{\alpha} {}_{n-2}\text{X}^{m-4} \xrightarrow{-\beta} {}_{n-1}\text{X}^{m-4}$
70. (c) Half-life $T_{1/2} = \frac{0.693}{\lambda} = \frac{0.693}{1.07 \times 10^{-4}} = 6476 \text{ years}$
71. (d) Number of nuclei decreases exponentially
 $N = N_0 e^{-\lambda t}$ and Rate of decay $\left(-\frac{dN}{dt}\right) = \lambda N$
Therefore, decay process lasts up to $t = \infty$.
Therefore, a given nucleus may decay at any time after $t = 0$.
72. (a) To become $\frac{1}{4}$ th, it requires time of two half lives
i.e., $t = 2(T_{1/2}) = 2 \times 5800 = 2 \times 58 \text{ centuries}$
73. (a) Carbon dating
74. (d) ${}_7\text{X}^{15} + {}_2\text{He}^4 \rightarrow {}_1\text{P}^1 + {}_8\text{Y}^{18}$
75. (c)
76. (d) ${}_{92}\text{U}^{238} \xrightarrow{\alpha} {}_{90}\text{X}^{234} \xrightarrow{-\beta^-} {}_{91}\text{Y}^{234}$
77. (d)

78. (c) After three half lives (*i.e.*, 30 days) it remains $\left(\frac{1}{2}\right)^3 = \frac{1}{8}$, so it will remain $\frac{1}{10}$ *th*, approximately in 33 days.
79. (a) ${}_{92}\text{U}^{238} \xrightarrow{\alpha} {}_{90}\text{Th}^{234} \xrightarrow{\beta} {}_{91}\text{Pa}^{234} \xrightarrow{E_{(-1)\beta^0}} {}_{92}\text{U}^{234}$
80. (d) By using $A = A_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow \frac{A}{A_0} = \left(\frac{1}{2}\right)^{9/3} = \frac{1}{8}$
81. (d) ${}_{48}\text{Cd}^{115} \xrightarrow{2(-1)\beta^0} {}_{50}\text{Sn}^{115}$
82. (b) In two half lives, the activity becomes one fourth.
83. (a) α decay decreases the mass number by 4 and atomic number by 2, β decay increases the atomic number by 1. Here atomic number of C is same as that of A.
84. (a) Number of half lives in two days four substance 1 and 2 respectively are $n_1 = \frac{2 \times 24}{12} = 4$ and $n_2 = \frac{2 \times 24}{1.6} = 3$
- By using $N = N_0 \left(\frac{1}{2}\right)^n \Rightarrow \frac{N_1}{N_2} = \frac{(N_0)_1}{(N_0)_2} \times \frac{\left(\frac{1}{2}\right)^{n_1}}{\left(\frac{1}{2}\right)^{n_2}}$
- $$= \frac{2}{1} \times \frac{\left(\frac{1}{2}\right)^4}{\left(\frac{1}{2}\right)^3} = \frac{1}{1}$$
85. (b)
86. (c) $\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{2.3} = 0.3$
87. (d) Number of α - particles emitted = $\frac{238 - 222}{4} = 4$
- This decreases atomic number to $90 - 4 \times 2 = 82$
- Since atomic number of ${}_{83}\text{Y}^{222}$ is 83, this is possible if one β - particle is emitted.
88. (d) Number of half lives in 150 years $n = \frac{150}{75} = 2$
- Fraction of the atom of decayed = $1 - \left(\frac{1}{2}\right)^n$
- $$= 1 - \left(\frac{1}{2}\right)^2 = \frac{3}{4} = 0.75 \Rightarrow \text{Percentage decay} = 75\%$$
89. (b) $A = A_0 e^{-\lambda t} \Rightarrow 975 = 9750 e^{-\lambda \times 5} \Rightarrow e^{5\lambda} = 10$
- $$\Rightarrow 5\lambda = \log_e 10 = 2.3026 \log_{10} 10 = 2.3026$$
- $$\Rightarrow \lambda = 0.461$$
90. (a) Mass number decreases by $8 \times 4 = 32$
91. (b) Atomic number decreases by $8 \times 2 - 5 = 11$
92. (d) $A = A_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow 5 \times 10^{-6} = 64 \times 10^{-5} \left(\frac{1}{2}\right)^{t/3}$
- $$\Rightarrow \frac{1}{128} = \left(\frac{1}{2}\right)^{t/3} \Rightarrow t = 21 \text{ days}$$
93. (c) Decayed fraction = $\frac{3}{4}$, so undecayed fraction = $\frac{1}{4}$
- Now $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \frac{1}{4} = \left(\frac{1}{2}\right)^n \Rightarrow n = 2$
- $$\Rightarrow t = n \times T_{1/2} = 2 \times 3.8 = 7.6 \text{ days}$$
94. (c) ${}_{84}\text{X}^{202} \xrightarrow{\alpha\text{-decay}} {}_{82}\text{Y}^{198} + {}_2\text{He}^4$ and ${}_{82}\text{Y}^{198} \xrightarrow{\beta\text{-decay}} {}_{83}\text{Z}^{198} + {}_{-1}\beta^0$
95. (a) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \frac{1}{8} = \left(\frac{1}{2}\right)^n \Rightarrow n = 3$
- Now $t = n \times T_{1/2} = 3 \times 3.8 = 11.4 \text{ days}$
96. (d) $n = \frac{72000}{24000} = 3$; Now $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n = \frac{1}{8}$
97. (d)
98. (a)
99. (b) $n_\alpha = \frac{A - A'}{4} = \frac{232 - 208}{4} = 6$
- $$n_\beta = 2n_\alpha - Z + Z' = 2 \times 6 - 90 + 82 = 4$$
100. (c)
101. (a) Number of half lives $n = \frac{5}{1} = 5$
- Now $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \frac{N}{N_0} = \left(\frac{1}{2}\right)^5 = \frac{1}{32}$
102. (d)
103. (b) Number of half lives $n = \frac{10}{5} = 2$, now $\frac{N}{N_0} = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$
- Fraction decayed = $1 - \frac{N}{N_0} = 1 - \frac{1}{4} = \frac{3}{4}$
- $$\Rightarrow \text{In percentage} = \frac{3}{4} \times 100 = 75\%$$
104. (b) Number of half lives $n = \frac{19}{3.8} = 5$; Now $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n$
- $$\Rightarrow \frac{N}{10.38} = \left(\frac{1}{2}\right)^5 \Rightarrow N = 10.38 \times \left(\frac{1}{2}\right)^5 = 0.32 \text{ gm}$$
105. (b) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n = \left(\frac{1}{2}\right)^5$
106. (d) $T_{1/2} = \frac{\log_e 2}{\lambda} = \frac{2.303 \log_{10} 2}{\lambda}$

1518 Atomic and Nuclear Physics

107. (a) ${}^A_Z X \rightarrow {}^{A-4}_{Z-2} Y + {}^4_2 He \rightarrow {}^{A-4}_{Z-3} Z + {}^0_1 B$
108. (d) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n, n = 2 \Rightarrow \frac{N}{N_0} = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$
 So disintegrated part = $1 - \frac{N}{N_0} = 1 - \frac{1}{4} = \frac{3}{4}$
109. (c) Number of half lives $n = \frac{10}{2.5} = 4$
 $\Rightarrow \frac{A}{A_0} = \frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow A = 1.6 \times \left(\frac{1}{2}\right)^4 = 0.1 \text{ curie}$
110. (b) By using $N = N_0 e^{-\lambda t}$ and $t = \tau = \frac{1}{\lambda}$
 Substance remains = $N = \frac{N_0}{e} = 0.37 N_0 \approx \frac{N_0}{3}$
 \therefore Substance disintegrated = $N_0 - \frac{N_0}{3} = \frac{2N_0}{3}$
111. (c) $\frac{3}{4}$ th active decay takes place in time
 $t = 2(T_{1/2}) \Rightarrow \frac{3}{4} = 2(T_{1/2}) \Rightarrow T_{1/2} = \frac{3}{8} \text{ sec}$
112. (c) By using $N = N_0 e^{-\lambda t}$ and average life time $t = \frac{1}{\lambda}$
 So $N = N_0 e^{-\lambda \times 1/\lambda} = N_0 e^{-1} \Rightarrow \frac{N}{N_0} = e^{-1} = \frac{1}{e}$
 Now disintegrated fraction = $1 - \frac{N}{N_0} = 1 - \frac{1}{e} = \frac{e-1}{e}$
113. (d) Complete reaction will be as follows
 ${}_{92}X^{235} \rightarrow {}_{82}Y^{207} + 7{}_2He^4 + 4{}_{-1}e^0$
 i.e., seven α - particles and four β - particles will be emitted.
114. (d) $n_\alpha = \frac{A - A'}{4} = \frac{235 - 207}{4} = 7$
 $n_\beta = (2n_\alpha - Z + Z') = (2 \times 7 - 92 + 82) = 4$
115. (c) During β -decay, a neutron is transformed into a proton and an electron.
116. (b) ${}_Z X^A \xrightarrow{\alpha} {}_{Z-2} X^{A-4}$
117. (a)
118. (a)
119. (d) After emitting β -particle (e) mass of nucleus doesn't change.
120. (a) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \frac{1}{16} = \left(\frac{1}{2}\right)^4 = \left(\frac{1}{2}\right)^n \Rightarrow n = 4$
 Also $n = \frac{t}{T_{1/2}} \Rightarrow T_{1/2} = \frac{40}{4} = 10 \text{ days}$
121. (c) As the γ - particle has no charge and mass.
122. (d) With emission of an α particle (${}_2He^4$) mass number decreases by 4 unit and atomic number decrease by 2 units and with emission of $2\beta^{-1}$ particle atomic number increases by 2 units. So Z will remain same and N will become $N - 4$.
123. (a) $N = N_0 \left(\frac{1}{2}\right)^n \Rightarrow \frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \frac{1}{100} = \left(\frac{1}{2}\right)^n \Rightarrow 2^n = 100$
n comes out in between 6 and 7.
124. (d) $N = N_0 \left(\frac{1}{2}\right)^n \Rightarrow N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$
 $\Rightarrow N = 1 \times \left(\frac{1}{2}\right)^{\frac{8.1}{2.7}} = \left(\frac{1}{2}\right)^3 = \frac{1}{8} \Rightarrow N = \frac{1}{8} \text{ mg} = 0.125 \text{ mg}$
125. (d) $N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T}} \Rightarrow \frac{N_0}{4} = N_0 \left(\frac{1}{2}\right)^{\frac{16}{T}} \Rightarrow T = 8 \text{ days.}$
126. (d) ${}_{92}U^{238} \rightarrow {}_2He^4 + {}_{90}X^{234} \rightarrow {}_{-1}e^0 + {}_{91}U^{234}$
 Hence, $A = 234, Z = 91$
127. (c) Mean life = $\frac{1}{\lambda} = 6.67 \times 10^8 \text{ sec.}$
128. (d) $\frac{dN}{dt} = -\lambda N \Rightarrow \left|\frac{dN}{dt}\right| = \frac{0.693}{T_{1/2}} \times N$
 $= \frac{0.693}{1.2 \times 10^7} \times 4 \times 10^{15} = 2.3 \times 10^8 \text{ atoms/sec}$
129. (c) Remaining material $N = \frac{N_0}{2^{t/T}}$
 $\Rightarrow N = \frac{10}{(2)^{20/15}} = \frac{10}{2.15} = 3.96 \text{ gm}$
 So decayed material = $10 - 3.96 = 6.04 \text{ gm}$
130. (b) Number of atoms remains undecayed $N = N_0 e^{-\lambda t}$
 Number of atoms decayed = $N_0(1 - e^{-\lambda t})$
 $= N_0 \left(1 - e^{-\lambda \times \frac{1}{\lambda}}\right) = N_0 \left(1 - \frac{1}{e}\right) = 0.63 N_0 = 63\% \text{ of } N_0.$
131. (d)
132. (b) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n \Rightarrow \left(\frac{1}{16}\right) = \left(\frac{1}{2}\right)^n \Rightarrow n = 4$
 also $n = \frac{t}{T_{1/2}} \Rightarrow T_{1/2} = \frac{120}{4} = 30 \text{ days}$
133. (c) $N = N_0 e^{-t/T_{1/2}} \Rightarrow \frac{1}{4} = e^{-t/10}$
 $\Rightarrow \left(\frac{1}{2}\right)^2 = \frac{1}{e^{t/10}} \Rightarrow t = 20 \text{ years}$
134. (a) $N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} = N_0 \left(\frac{1}{2}\right)^{\frac{15}{5}} = \frac{N_0}{8}$
135. (a)
136. (b) $A = A_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} \Rightarrow \frac{1}{64} = \left(\frac{1}{2}\right)^{\frac{60}{T_{1/2}}}$

$$\Rightarrow \left(\frac{1}{2}\right)^6 = \left(\frac{1}{2}\right)^{\frac{60}{T_{1/2}}} \Rightarrow T_{1/2} = 10 \text{ sec}$$

137. (a, c)

138. (b) $(Z=92)U^{(A=238)} \xrightarrow{(8\alpha, 6\beta)} Z'X^{A'}$

so $A' = A - 4n_\alpha = 238 - 4 \times 8 = 206$

and $Z' = n_\beta - 2n_\alpha + z = 6 - 2 \times 8 + 92 = 82$.

139. (c) $A = A_0 e^{-\lambda t} = A_0 e^{-t/\tau}$; where τ = mean life

So $A_1 = A_0 e^{-t_1/T} \Rightarrow A_0 = \frac{A_1}{e^{-t_1/T}} = A_1 e^{t_1/T}$

$\therefore A_2 = A_0 e^{-t_2/T} = (A_1 e^{t_1/T}) e^{-t_2/T} \Rightarrow A_2 = A_1 e^{(t_1 - t_2)/T}$

140. (c)

141. (d) $n_\alpha = \frac{228 - 212}{4} = 4$ and $n_\beta = 2 \times 4 - 90 + 83 = 1$

142. (c) In a gamma decay process. There is no change in either A or Z .

143. (a) The radioactivity of a sample decays to $\frac{1}{16}$ th of its initial value in four half lives.

144. (d) $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T} \Rightarrow \frac{1}{16} = \left(\frac{1}{2}\right)^{t/48}$
 $\Rightarrow \left(\frac{1}{2}\right)^4 = \left(\frac{1}{2}\right)^{t/48} \Rightarrow t = 192 \text{ hour.}$

145. (c) If λ is the decay constant of a radioactive substance than average life = $\frac{1}{\lambda}$

Also half life = $\frac{0.693}{\lambda} = 0.693 \times (\text{Average life})$

in single average life, more than 63% of radioactive nuclei decay

146. (b)

147. (d) $M = M_0 e^{-\lambda t}$; given $t = 2\left(\frac{1}{\lambda}\right)$

$\Rightarrow M = 10e^{-\lambda\left(\frac{2}{\lambda}\right)} = 10\left(\frac{1}{e}\right)^2 \Rightarrow M = 1.35 \text{ gm}$

148. (b)

149. (b) $\lambda = \frac{\log_e \frac{A_1}{A_2}}{t} = \frac{\log_e \frac{5000}{1250}}{5} = 0.4 \ln 2$

150. (c) $Z_{\text{Resuting nucleus}} = 92 - 8 \times 2 + 4 \times 1 - 2 \times 1 = 78$

151. (c) Radioactive nuclei that are injected into a patient collected at certain sites within its body, undergoing radioactive decay and emitting electromagnetic radiation. These radiation can than be recorded by a detector. This procedure provides an important diagnostic tool called radio tracer technique.

152. (a) By using $n_\alpha = \frac{A - A'}{4}$ and $n_\beta = 2n_\alpha - Z + Z'$

153. (b) $N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}}$

No of atoms at $t = 2hr$, $N_1 = 8 \times 10^{10} \left(\frac{1}{2}\right)^{\frac{2}{1}} = 2 \times 10^{10}$

No. of atoms at $t = 4hr$, $N_2 = 8 \times 10^{10} \left(\frac{1}{2}\right)^{\frac{4}{1}} = \frac{1}{2} \times 10^{10}$

\therefore No. of atoms decayed in given duration

$= \left(2 - \frac{1}{2}\right) \times 10^{10} = 1.5 \times 10^{10}$

154. (b)

155. (d) $A = A_0 \left(\frac{1}{2}\right)^n \Rightarrow 30 = 240 \left(\frac{1}{2}\right)^n \Rightarrow \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^n \Rightarrow n=3$

$\therefore \frac{t}{T_{1/2}} = 3 \Rightarrow T_{1/2} = \frac{t}{3} = \frac{1}{3} \text{ hr} = 20 \text{ min.}$

156. (b) $M = M_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} \Rightarrow 25 = 100 \left(\frac{1}{2}\right)^{\frac{t}{1600}} \Rightarrow t = 3200 \text{ years.}$

157. (c) Activity $R = R_0 e^{-\lambda t}$

$\frac{R_0}{3} = R_0 e^{-\lambda \times 9} \Rightarrow e^{-9\lambda} = \frac{1}{3} \dots(i)$

After further 9 years $R' = R e^{-\lambda t} = \frac{R_0}{3} \times e^{-\lambda \times 9} \dots(ii)$

From equation (i) and (ii) $R' = \frac{R_0}{9}$.

158. (c) To reduce one fourth it takes time $t = 2(T_{1/2}) = 2 \times 40 = 80 \text{ years.}$

Decay constant $\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{40} = 0.0173 \text{ years}$

159. (d) $M = M_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} = 20 \times \left(\frac{1}{2}\right)^{\frac{3.6}{3.6}} = 20 \times \left(\frac{1}{2}\right)^{10} = 0.019 \text{ mg}$

160. (c) $N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} \Rightarrow \frac{N}{N_0} = \left(\frac{1}{2}\right)^{\frac{30}{10}} = \frac{1}{8} = 0.125.$

161. (a) $\frac{7}{8}$ part decays i.e. remaining part is $\frac{1}{8}$

$N = N_0 \left(\frac{1}{2}\right)^{\frac{t}{T_{1/2}}} \Rightarrow \frac{1}{8} = \left(\frac{1}{2}\right)^{\frac{15}{T_{1/2}}} \Rightarrow T_{1/2} = 5 \text{ min.}$

162. (d) $\frac{A}{A_0} = \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow \frac{1}{8} = \left(\frac{1}{2}\right)^{t/8} \Rightarrow t = 24 \text{ years.}$

163. (c) After $\beta^+ ({}_{+1}e^0)$ emission atomic number decreases by one and mass number remain unchanged. γ -emission, there will be no change on mass number and atomic number.

164. (c) New mass number $A' = A - 4n_\alpha = 232 - 4 \times 6 = 208$

1520 Atomic and Nuclear Physics

atomic number $Z' = Z + n_{\beta} - 2n_{\alpha} = 90 + 4 - 2 \times 6 = 82$

165. (d)

166. (d) Using conservation of momentum $P_{daughter} = P_{\alpha}$

$$\Rightarrow \frac{E_d}{E_{\alpha}} = \frac{m_{\alpha}}{m_d} \Rightarrow E_d = \frac{E_{\alpha} \times m_{\alpha}}{m_d} = \frac{6.7 \times 4}{214} = 0.125 \text{ MeV}$$

167. (c)

168. (c)

169. (b) $N = N_0 \times \left(\frac{1}{2}\right)^{11400/5700} = N_0 \left(\frac{1}{2}\right)^2 = 0.25 N_0.$

170. (d) Mean life $(T) = 1/\lambda = 100 \text{ second}$

$$\text{Half-life} = \frac{0.693}{\lambda} = \frac{0.693 \times 100}{60} = 1.155 \text{ min.}$$

171. (b) By using $n_{\alpha} = \frac{A - A'}{4}$ and $n_{\beta} = 2n_{\alpha} - Z + Z'$

172. (c)

173. (b) $\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{77} = 9 \times 10^{-3} / \text{day}.$

174. (a) By using $n_{\alpha} = \frac{A - A'}{4} = \frac{232 - 204}{4} = 7.$

175. (c) $\therefore \frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T_{1/2}} \left(\frac{1}{2}\right)^{1/2} = \frac{1}{\sqrt{2}}.$

176. (d)

Critical Thinking Questions

1. (c) At closest distance of approach

Kinetic energy = Potential energy

$$\Rightarrow 5 \times 10^6 \times 1.6 \times 10^{-19} = \frac{1}{4\pi\epsilon_0} \times \frac{(ze)(2e)}{r}$$

For uranium $z = 92$, so $r = 5.3 \times 10^{-12} \text{ cm}$

2. (c) Speed of electron in n orbit of hydrogen atom $v = \frac{e^2}{2\epsilon_0 n h}$

In ground state $n = 1 \Rightarrow v = \frac{e^2}{2\epsilon_0 h}$

$$\Rightarrow \frac{v}{c} = \frac{e^2}{2\epsilon_0 ch} = \frac{(1.6 \times 10^{-19})^2}{2 \times 8.85 \times 10^{-12} \times 3 \times 10^8 \times 6.6 \times 10^{-34}}$$

$$= \frac{1}{137}.$$

3. (b) Recoil momentum = momentum of photon $= \frac{h}{\lambda}$

$$= hR \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) = \frac{hR \times 15}{16} = 6.8 \times 10^{-27} \text{ N} \times \text{sec}$$

4. (a) The average time that the atom spends in this excited state is equal to Δt , so by using $\Delta E \cdot \Delta t = \frac{h}{2\pi}$

$$\Rightarrow \text{Uncertainty in energy} = \frac{h/2\pi}{\Delta t}$$

$$= \frac{6.6 \times 10^{-34}}{2 \times 3.14 \times 10^{-8}} = 1.05 \times 10^{-26} \text{ J} = 6.56 \times 10^{-8} \text{ eV}$$

5. (a) After the removal of first electron remaining atom will be hydrogen like atom.

So energy required to remove second electron from the atom

$$E = 13.6 \times \frac{2^2}{1} = 54.4 \text{ eV}$$

$$\therefore \text{Total energy required} = 24.6 + 54.4 = 79 \text{ eV.}$$

6. (a) Electron after absorbing 10.2 eV energy goes to its first excited state ($n=2$) from ground state ($n=1$).

$$\therefore \text{Increase in momentum} = \frac{h}{2\pi}$$

$$= \frac{6.6 \times 10^{-34}}{6.28} = 1.05 \times 10^{-34} \text{ J-s.}$$

7. (a) Using $\Delta E \propto Z^2$ ($\because n_1$ and n_2 are same)

$$\Rightarrow \frac{hc}{\lambda} \propto Z^2 \Rightarrow \lambda Z^2 = \text{constant}$$

$$\Rightarrow \lambda_1 Z_1^2 = \lambda_2 Z_2^2 = \lambda_3 Z_3^2 = \lambda_4 Z_4^2$$

$$\Rightarrow \lambda_1 \times 1 = \lambda_2 \times 1^2 = \lambda_3 \times 2^2 = \lambda_4 \times 3^3$$

$$\Rightarrow \lambda_1 = \lambda_2 = 4\lambda_3 = 9\lambda_4.$$

8. (d) $mvr = \frac{h}{2\pi}$ (for first orbit)

$$\Rightarrow m\omega r^2 = \frac{h}{2\pi} \Rightarrow m \times 2\pi v \times r^2 = \frac{h}{2\pi} \Rightarrow v = \frac{h}{4\pi^2 m r^2}$$


$$= \frac{6.6 \times 10^{-34}}{4(3.14)^2 \times 9.1 \times 10^{-31} \times (0.53 \times 10^{-10})^2} = 6.5 \times 10^{15} \frac{\text{rev}}{\text{sec}}$$

9. (b) It will form a stationary wave

$$\lambda = 2l = 2 \times 10^{-9} \text{ m}$$

$$\Rightarrow \lambda = \frac{h}{\sqrt{2mE}}$$

$$\Rightarrow E = \frac{h^2}{2m\lambda^2} = 6 \times 10^{-20} \text{ J}$$



10. (d) Suppose closest distance is r , according to conservation of energy.

$$400 \times 10^3 \times 1.6 \times 10^{-19} = 9 \times 10^9 \frac{(ze)(2e)}{r}$$

$$\Rightarrow 6.4 \times 10^{-14} = \frac{9 \times 10^9 \times (82 \times 1.6 \times 10^{-19}) \times (2 \times 1.6 \times 10^{-19})}{r}$$

$$\Rightarrow r = 5.9 \times 10^{-13} \text{ m} = 0.59 \text{ pm.}$$

11. (a) Here radius of electron orbit $r \propto 1/m$ and energy $E \propto m$, where m is the mass of the electron.

Hence energy of hypothetical atom

$$E_0 = 2 \times (-13.6 \text{ eV}) = -27.2 \text{ eV} \text{ and radius } r_0 = \frac{a_0}{2}$$

12. (a) Electronic configuration of iodine is 2, 8, 18, 18, 7.

$$\text{Here } r_n = (0.053 \times 10^{-9} \text{ m}) \frac{n^2}{Z}$$

Here $n = 5$ and $Z = 53$, hence $r_n = 2.5 \times 10^{-11} \text{ m.}$

13. (a) $N \propto \left[\frac{1}{\sin^4 \theta/2} \right] \Rightarrow N_1 = 7 \times \frac{1}{(\sin 30^\circ)^4} = 112$

$$\text{and } N_2 = 7 \times \frac{1}{(\sin 60^\circ)^4} = 12.5.$$

14. (d) $E_n = -13.6 \frac{Z^2}{n^2} \text{ eV}$. Required energy for said transition

$$\Delta E = E_3 - E_1 = 13.6 Z^2 \left[\frac{1}{1^2} - \frac{1}{3^2} \right]$$

$$\Rightarrow \Delta E = 13.6 \times 3^2 \left[\frac{8}{9} \right] = 108.8 \text{ eV}$$

$$\Rightarrow \Delta E = 108.8 \times 1.6 \times 10^{-19} \text{ J}$$

$$\text{Now } \Delta E = \frac{hc}{\lambda} = 108.8 \times 1.6 \times 10^{-19}$$

$$\Rightarrow \lambda = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{108.8 \times 1.6 \times 10^{-19}} = 0.11374 \times 10^{-7} \text{ m} = 113.74 \text{ \AA}$$


15. (c) $\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$

$$\Rightarrow \frac{1}{970.6 \times 10^{-10}} = 1.097 \times 10^7 \left[\frac{1}{1^2} - \frac{1}{n_2^2} \right] \Rightarrow n_2 = 4$$

$$\therefore \text{Number of emission lines } N = \frac{n(n-1)}{2} = \frac{4 \times 3}{2} = 6$$

16. (d) Neutron velocity = v , mass = m

Deuteron contains 1 neutron and 1 proton, mass = $2m$



In elastic collision both momentum and $K.E.$ are conserved $p = p$.

$$mv = mv + m v \Rightarrow mv = mv + 2mv \quad \dots (i)$$

By conservation of kinetic energy

$$\frac{1}{2}mv^2 = \frac{1}{2}mv_1^2 + \frac{1}{2}(2m)v_2^2 \quad \dots (ii)$$

By solving (i) and (ii) we get

$$v_1 = \frac{m_1 - m_2}{m_1 + m_2} v + \frac{2m_2}{(m_1 + m_2)} v \Rightarrow v_1 = \frac{m_1 + 2m}{3m} = -\frac{v}{3}$$

$$K_i = \frac{1}{2}mv^2, \quad K_f = \frac{1}{2}mv_1^2 \Rightarrow \frac{K_i - K_f}{K_i} = 1 - \frac{v_1^2}{v^2}$$

$$= 1 - \frac{1}{9} = \frac{8}{9} \text{ (Fractional change in } K.E.)$$

17. (c) In hydrogen atom $E_n = -\frac{Rhc}{n^2}$

Also $E_n \propto m$; where m is the mass of the electron. Here the electron has been replaced by a particle whose mass is double of an electron. Therefore, for this hypothetical atom energy in

n orbit will be given by $E_n = -\frac{2Rhc}{n^2}$

The longest wavelength λ_{\max} (or minimum energy) photon will correspond to the transition of particle from $n = 3$ to $n = 2$

$$\Rightarrow \frac{hc}{\lambda_{\max}} = E_3 - E_2 = Rhc \left(\frac{1}{2^2} - \frac{1}{3^2} \right)$$

This gives $\lambda_{\max} = \frac{18}{5R}$.

18. (d) As the transition $n = 4$ and $n = 3$, results in UV radiation and infrared radiation involves smaller amounts of energy UV. So we require a transition involving initial values of n greater than 4 e.g. $5 \rightarrow 4$.

19. (c) $\frac{hc}{\lambda} = E = eV$

$$\Rightarrow \lambda = \frac{hc}{eV} = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{1.6 \times 10^{-19} \times 4.9} = 2525 \text{ \AA}$$

20. (d) Rydberg constant $R = \frac{\epsilon_0 n^2 h^2}{\pi m Z e^2}$

Velocity $v = \frac{Ze^2}{2\epsilon_0 nh}$ and energy $E = -\frac{mZ^2 e^4}{8\epsilon_0^2 n^2 h^2}$

Now, it is clear from above expressions $R, v \propto n$

21. (b) In second excited state $n = 3$

So $l_H = l_{Li} = 3 \left(\frac{h}{2\pi} \right)$

While $E \propto Z$ and $Z_+ = 1, Z_- = 3$

So $|E| = 9|E|$ or $|E_+| < |E_-|$

22. (c) Since the $^{133}_{55}\text{Cs}$ has larger size among the four atoms gives, thus the electrons present in the outermost orbit will be away from the nucleus and the electrostatic force experienced by electrons due to nucleus will be minimum. Therefore the energy required to liberate electron from outer will be minimum in the case of $^{133}_{55}\text{Cs}$.

23. (d)

24. (a) Potential energy $U = eV = eV_0 \ln \frac{r}{r_0}$

\therefore Force $F = -\frac{dU}{dr} = \frac{eV_0}{r}$.

\therefore The force will provide the necessary centripetal force.

Hence $\frac{mv^2}{r} = \frac{eV_0}{r} \Rightarrow v = \sqrt{\frac{eV_0}{m}}$ (i)

and $mvr = \frac{nh}{2\pi}$ (ii)

From equation (i) and(ii) $mr = \left(\frac{nh}{2\pi} \right) \sqrt{\frac{m}{eV_0}}$ or $r \propto n$

25. (d) $(r_m) = \left(\frac{m^2}{Z} \right) (0.53 \text{ \AA}) = (n \times 0.53 \text{ \AA}) \Rightarrow \frac{m^2}{Z} = n$

$m = 5$ for $^{100}\text{Fm}^{257}$ (the outermost shell)

and $z = 100 \Rightarrow n = \frac{(5)^2}{100} = \frac{1}{4}$

26. (d) Energy radiated $= 1.4 \text{ kW} / m^2$

$$= 1.4 \text{ kJ} / \text{sec } m^2 = \frac{1.4 \text{ kJ}}{86400 \text{ day } m^2} = \frac{1.4 \times 86400}{\text{day } m^2}$$

Total energy radiated/day

$$= \frac{4\pi \times (1.5 \times 10^{11})^2 \times 1.4 \times 86400 \text{ kJ}}{1 \text{ day}} = E$$

$\therefore E = mc^2 \Rightarrow m = \frac{E}{c^2}$

$$= \frac{4\pi(1.5 \times 10^{11})^2 \times 1.4 \times 86400}{(3 \times 10^8)^2} = 3.8 \times 10^{14} \text{ kg}.$$

27. (c) The equation is $O^{17} \rightarrow O^{n^1} + O^{16}$

\therefore Energy required = B.E. of O - B.E. of O
 $= 17 \times 7.75 - 16 \times 7.97 = 4.23 \text{ MeV}$

28. (c) $\Delta = mc^2 - m_0c^2 = \frac{m_0c^2}{\sqrt{1-(v^2/c^2)}} - m_0c^2$

$$= m_0c^2 \left(\frac{1}{\sqrt{1-(v^2/c^2)}} - 1 \right) = 0.511 \left(\frac{1}{\sqrt{0.75}} - 1 \right)$$

$$= 0.079 \text{ MeV}$$

29. (c,d) Due to mass defect (which is finally responsible for the binding energy of the nucleus), mass of a nucleus is always less than the sum of masses of its constituent particles $^{20}_{10}\text{Ne}$ is made up of 10 protons plus 10 neutrons. Therefore, mass of $^{20}_{10}\text{Ne}$ nucleus $M_1 < 10(m_p + m_n)$


Also heavier the nucleus, more is the mass defect thus $20(m_n + m_p) - M_2 > 10(m_p + m_n) - M_1$

or $10(m_p + m_n) > M_2 - M_1$

$\Rightarrow M_2 < M_1 + 10(m_p + m_n) \Rightarrow M_2 < M_1 + M_1$

$\Rightarrow M_2 < 2M_1$.

30. (a)



By conservation of momentum $m_1v_1 = m_2v_2$

$\Rightarrow \frac{v_1}{v_2} = \frac{8}{1} = \frac{m_2}{m_1}$ (i)

Also from $r \propto A^{1/3} \Rightarrow \frac{r_1}{r_2} = \left(\frac{A_1}{A_2} \right)^{1/3} = \left(\frac{1}{8} \right)^{1/3} = \frac{1}{2}$.

31. (a) Since nuclear density is constant hence mass \propto volume.

32. (c) Mass defect $= 3 \times 2.014 - 4.001 - 1.007 - 1.008$
 $= 0.026 \text{ amu} = 0.026 \times 931 \times 10^6 \times 1.6 \times 10^{-19} \text{ J}$
 $= 3.82 \times 10^{-12} \text{ J}$

Power of star = 10^W

$$\text{Number of deuterons used} = \frac{10^{16}}{\Delta M} = 0.26 \times 10^{28}$$

$$\text{Deuteron supply exhausts in } \frac{10^{40}}{0.26 \times 10^{28}} = 10^{12} \text{ s.}$$

33. (a) Since electron and positron annihilate

$$\lambda = \frac{hc}{E_{\text{Total}}} = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{(0.51 + 0.51) \times 10^6 \times 1.6 \times 10^{-19}}$$


$$= 1.21 \times 10^{-12} \text{ m} = 0.012 \text{ \AA}.$$

34. (a) Kinetic energy of the molecules of a gas at a temp. T is $\frac{3}{2}kT$

$$\therefore \text{To initiate the reaction } \frac{3}{2}kT = 7.7 \times 10^{-14} \text{ J}$$

$$\Rightarrow \frac{3}{2} \times 1.38 \times 10^{-23} T = 7.7 \times 10^{-14} \Rightarrow T = 3.7 \times 10^9 \text{ K.}$$

35. (b)



Q -value of the reaction is 5.5 eV

$$\text{i.e. } k_1 + k_2 = 5.5 \text{ MeV} \quad \dots(i)$$

By conservation of linear momentum

$$p_1 = p_2 \Rightarrow \sqrt{2(216)k_1} = \sqrt{2(4)k_2}$$

$$\Rightarrow k_1 = 54 k_2 \quad \dots(ii)$$

On solving equation (i) and (ii) we get $k_2 = 5.4 \text{ MeV}$.

36. (b) By the formula $N = N_0 e^{-\lambda t}$

$$\text{Given } \frac{N}{N_0} = \frac{1}{20} \text{ and } \lambda = \frac{0.6931}{3.8} \Rightarrow 20 = e^{\frac{0.6931 \times t}{3.8}}$$

Taking log of both sides

$$\text{or } \log 20 = \frac{0.6931 \times t}{3.8} \log_{10} e$$

$$\text{or } 1.3010 = \frac{0.6931 \times t \times 0.4343}{3.8} \Rightarrow t = 16.5 \text{ days.}$$

37. (b) $N = N_0 e^{-\lambda t}$

$$\therefore 0.9 N_0 = N_0 e^{-\lambda \times 5} \Rightarrow 5\lambda = \log_e \frac{1}{0.9} \quad \dots(i)$$

$$\text{and } x N_0 = N_0 e^{-\lambda \times 20} \Rightarrow 20\lambda = \log_e \left(\frac{1}{x} \right) \quad \dots(ii)$$

Dividing (i) by (ii), we get

$$\frac{1}{4} = \frac{\log_e(1/0.9)}{\log_e(1/x)} = \frac{\log_{10}(1/0.9)}{\log_{10}(1/x)} = \frac{\log_{10} 0.9}{\log_{10} x}$$

$$\Rightarrow \log_{10} x = 4 \log_{10} 0.9 \Rightarrow x = 0.658 = 65.8\%$$

38. (c) If in the rock there is no Y element, then the time taken by element X to reduce to $\frac{1}{8}$ th the initial value will be equal to

$$\frac{1}{8} = \left(\frac{1}{2} \right)^n \text{ or } n=3$$

Therefore, from the beginning three half life time is spent. Hence the age of the rock is

$$= 3 \times 1.37 \times 10^9 = 4.11 \times 10^9 \text{ years.}$$

39. (b) $\frac{N}{N_0} = \left(\frac{1}{2} \right)^n \Rightarrow \frac{1}{64} = \left(\frac{1}{2} \right)^6 = \left(\frac{1}{2} \right)^n \Rightarrow n = 6.$

After 6 half lives intensity emitted will be safe.

$$\therefore \text{Total time taken} = 6 \times 2 = 12 \text{ hrs.}$$

40. (a) $\frac{dN}{dt} = \lambda N; \lambda = \frac{0.6931}{t_{12}} = \frac{0.6931}{1620 \times 365 \times 24 \times 60 \times 60}$,

$$N = \frac{6.023 \times 10^{23}}{226}$$

$$\therefore \frac{dN}{dt} = \frac{0.6931 \times 6.023 \times 10^{23}}{1620 \times 365 \times 24 \times 60 \times 60 \times 226} = 3.61 \times 10^{10}$$

41. (a) $\lambda = \lambda_1 + \lambda_2 \Rightarrow \frac{1}{T} = \frac{1}{T_1} + \frac{1}{T_2}$

$$\therefore T = \frac{T_1 T_2}{T_1 + T_2} = \frac{810 \times 1620}{810 + 1620} = 540 \text{ years}$$

Hence $\frac{1}{4}$ th of material remain after 1080 years.

42. (b) Similar to Q. 40.

43. (c) $(T_{1/2})_x = (t_{\text{mean}})_y$

$$\Rightarrow \frac{0.693}{\lambda_x} = \frac{1}{\lambda_y} \Rightarrow \lambda_x = 0.693 \lambda_y \text{ or } \lambda_x < \lambda_y$$

Also rate of decay = λN

Initially number of atoms (N) of both are equal but since $\lambda_y > \lambda_x$, therefore, y will decay at a faster rate than x .

44. (c) $\lambda_\alpha = \frac{1}{1620} \text{ per year}$ and $\lambda_\beta = \frac{1}{405} \text{ per year}$ and it is given

$$\text{that the fraction of the remained activity } \frac{A}{A_0} = \frac{1}{4}$$

Total decay constant

$$\lambda = \lambda_\alpha + \lambda_\beta = \frac{1}{1620} + \frac{1}{405} = \frac{1}{324} \text{ per year}$$

$$\text{We know that } A = A_0 e^{-\lambda t} \Rightarrow t = \frac{1}{\lambda} \log_e \frac{A_0}{A}$$

$$\Rightarrow t = \frac{1}{\lambda} \log_e 4 = \frac{2}{\lambda} \log_e 2 = 324 \times 2 \times 0.693 = 449 \text{ years.}$$


45. (d) $n = \frac{24}{24 \times 138.6} = \frac{1}{138.6}$; Now $\frac{N}{N_0} = \left(\frac{1}{2} \right)^n = \left(\frac{1}{2} \right)^{1/138.6}$

$$\Rightarrow N = 10,000,000 \left(\frac{1}{2} \right)^{1/138.6} = 995011$$

So number of disintegration

$$= 1000000 - 995011 = 4989 \approx 5000.$$

46. (a)



Rest

According to conservation of momentum $4v = (A-4)v' \Rightarrow$

$$v' = \frac{4v}{A-4}.$$

47. (b) $\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{20} = 0.03465$

Now time of decay $t = \frac{2.303}{\lambda} \log \frac{N_0}{N}$
 $\Rightarrow t_1 = \frac{2.303}{0.03465} \log \frac{100}{67} = 11.6 \text{ min min}$

and $t_2 = \frac{2.303}{0.03465} \log \frac{100}{33} = 32 \text{ min}$

Thus time difference between points of time
 $= t_2 - t_1 = 32 - 11.6 = 20.4 \text{ min} \approx 20 \text{ min}$.

48. (d) $N_1 = N_0 e^{-10\lambda t}$ and $N_2 = N_0 e^{-9\lambda t}$
 $\Rightarrow \frac{N_1}{N_2} = \frac{1}{e} = e^{-1} = e^{-(10\lambda + 9\lambda)t} = e^{-9\lambda t} \Rightarrow t = \frac{1}{9\lambda}$.

49. (a) $N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}} \Rightarrow N_A = 10 \left(\frac{1}{2}\right)^{t/1}$ and $N_B = 1 \left(\frac{1}{2}\right)^{t/2}$

Given $N_A = N_B \Rightarrow 10 \left(\frac{1}{2}\right)^t = \left(\frac{1}{2}\right)^{t/2}$

$\Rightarrow 10 = \left(\frac{1}{2}\right)^{-t/2} \Rightarrow 10 = 2^{t/2}$. Taking log both the sides.

$\log_{10} 10 = \frac{t}{2} \log_{10} 2 \Rightarrow 1 = \frac{t}{2} \times 0.3010 \Rightarrow t = 6.62 \text{ years}$.

50. (b) Here $T_{1/2} = 20 \text{ minutes}$; we know $\frac{N}{N_0} = \left(\frac{1}{2}\right)^{t/T_{1/2}}$
 For 20% decay $\frac{N}{N_0} = \frac{80}{100} = \left(\frac{1}{2}\right)^{t_1/20}$ (i)

For 80% decay $\frac{N}{N_0} = \frac{20}{100} = \left(\frac{1}{2}\right)^{t_2/20}$ (ii)

Dividing (ii) by (i)

$\frac{1}{4} = \left(\frac{1}{2}\right)^{\frac{(t_2 - t_1)}{20}}$; on solving we get $t_2 - t_1 = 40 \text{ min}$.

51. (d) Here the activity of the radioactive sample reduces to half in 140 days. Therefore, the half life of the sample is 140 days. 280 days is its two half lives. So before two half lives its activity was ($2^2 \times$ present activity).

\therefore Initial activity = $2^2 \times 6000 = 24000 \text{ dps}$.

52. (a) Excitation energy $\Delta E = E_2 - E_1 = 13.6 Z^2 \left[\frac{1}{1^2} - \frac{1}{2^2} \right]$

$\Rightarrow 40.8 = 13.6 \times \frac{3}{4} \times Z^2 \Rightarrow Z = 2$.

Now required energy to remove the electron from ground state
 $= \frac{+13.6 Z^2}{(1)^2} = 13.6(Z)^2 = 54.4 \text{ eV}$.

53. (b) Rate of disintegration $\frac{dN}{dt} = 10^{17} \text{ s}^{-1}$

Half life $T_{1/2} = 1445 \text{ year}$

$= 1445 \times 365 \times 24 \times 60 \times 60 = 4.55 \times 10^8 \text{ sec}$

Now decay constant

$\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{4.55 \times 10^{10}} = 1.5 \times 10^{-11} \text{ per sec}$

The rate of disintegration

$\frac{dN}{dt} = \lambda \times N_0 \Rightarrow 10^{17} = 1.5 \times 10^{-11} \times N_0$

$\Rightarrow N_0 = 6.6 \times 10^8$.

54. (b) $P = \frac{nE}{t} \Rightarrow 300 \times 10^6 = \frac{n \times 170 \times 10^6 \times 1.6 \times 10^{-19}}{t}$

\therefore Number of atoms per sec $\frac{n}{t} = 1.102 \times 10^{19}$

Number of atoms per hour = $1.02 \times 10^8 \times 3600 = 3.97 \times 10^8$.

55. (a) According to kinetic interpretation of temperature

$K.E. = \left(\frac{1}{2} m v^2\right) = \frac{3}{2} kT$.

$\Rightarrow 10.2 \times 1.6 \times 10^{-19} = \frac{3}{2} \times (1.38 \times 10^{-23}) T$

$\Rightarrow T = 7.9 \times 10^4 K$

56. (a) R_0 = Initial activity = 1 micro curie = $3.7 \times 10^4 \text{ dps}$
 r = Activity in 1 cm of blood at $t = 5 \text{ hrs}$

$= \frac{296}{60} \text{ dps} = 4.93 \text{ dps}$

R = Activity of whole blood at time $t = 5 \text{ hr}$,

Total volume should be $V = \frac{R}{r} = \frac{R_0 e^{-\lambda t}}{r}$

$= \frac{3.7 \times 10^4 \times 0.7927}{4.93} = 5.94 \times 10^3 \text{ cm} = 5.94 \text{ Litre}$.

57. (b) Let ground state energy (in eV) be E_1
 Then from the given condition

$E_{2n} - E_1 = 204 \text{ eV}$ or $\frac{E_1}{4n^2} - E_1 = 204 \text{ eV}$

$\Rightarrow E_1 \left(\frac{1}{4n^2} - 1 \right) = 204 \text{ eV}$ (i)

and $E_{2n} - E_n = 40.8 \text{ eV}$

$\Rightarrow \frac{E_1}{4n^2} - \frac{E_1}{n^2} = E_1 \left(-\frac{3}{4n^2} \right) = 40.8 \text{ eV}$ (ii)

From equation (i) and (ii), $\frac{1 - \frac{1}{4n^2}}{\frac{3}{4n^2}} = 5 \Rightarrow n = 2$

58. (b) Here $\frac{N}{N_0} = \left(\frac{1}{2}\right)^n = \left(\frac{1}{2}\right)^{1/3}$

where n = Number of half lives = $\frac{1}{3}$

$\Rightarrow \frac{N}{N_0} = \frac{1}{1.26} \Rightarrow \frac{N_U}{N_{Pb} + N_U} = \frac{1}{1.26}$

$\Rightarrow N_{Pb} = 0.26 N_U \Rightarrow \frac{N_{Pb}}{N_U} = 0.26$

59. (b) For K_{α} X-ray line

$$\frac{1}{\lambda_{\alpha}} = R(Z-1)^2 \left[\frac{1}{1^2} - \frac{1}{2^2} \right] = \frac{3R}{4} (Z-1)^2$$
 On putting the given values

$$\frac{1}{0.76 \times 10^{-10}} = \frac{3}{4} \times 1.09 \times 10^7 (Z-1)^2$$

$$\Rightarrow (Z-1)^2 \approx 1600 \Rightarrow Z-1 = 40 \Rightarrow Z = 41$$

60. (a) Maximum energy is liberated for transition $E_n \rightarrow 1$ and minimum energy for $E_n \rightarrow E_{n-1}$
 Hence $\frac{E_1}{n^2} - E_1 = 52.224 \text{ eV}$ (i)
 and $\frac{E_1}{n^2} - \frac{E_1}{(n-1)^2} = 1.224 \text{ eV}$ (ii)
 Solving equations (i) and (ii) we get
 $E_1 = -54.4 \text{ eV}$ and $n = 2$
 Now $E_1 = -\frac{13.6 Z^2}{1^2} = -54.4 \text{ eV}$. Hence $Z = 2$

61. (a) Activity of substance that has 2000 disintegration/sec

$$= \frac{2000}{3.7 \times 10^{10}} = 0.054 \times 10^{-6} \text{ ci} = 0.054 \mu\text{ci}$$
 The number of radioactive nuclei having activity A

$$N = \frac{A}{\lambda} = \frac{2000 \times T_{1/2}}{\log_e 2}$$

$$= \frac{2000 \times 138.6 \times 24 \times 3600}{0.693} = 3.45 \times 10^{10}$$

62. (a) Maximum number of nuclei will be present when rate of decay = rate of formation $\Rightarrow \lambda N = \alpha \Rightarrow N = \frac{\alpha}{\lambda}$

63. (b) $r \propto A^{1/3} \Rightarrow \frac{r_1}{r_2} = \left(\frac{A_1}{A_2} \right)^{1/3}$

$$\Rightarrow \frac{3}{5} = \left(\frac{27}{A} \right)^{1/3} \Rightarrow \frac{27}{125} = \frac{27}{A} \Rightarrow A = 125$$
 Number of nuclei in atom $X = A - 52 = 125 - 52 = 73$.

64. (c) 1 week ≈ 7 days $\approx 7 \times 24 \text{ hrs} \approx 14$ half lives
 Number of atoms left = $\frac{N_0}{(2)^{14}}$, Activity = $N\lambda$
 \therefore Activity left is $\frac{1}{(2)^{14}}$ times the initial

$$\Rightarrow \frac{1}{(2)^{14}} \times 1 \text{ curie} = \frac{1}{16384} \times 1 \text{ curie} \approx 61 \times 10^{-6} \text{ curie}$$

$$\approx 60 \mu \text{ curie}$$

65. (a) $m_0 c^2 = 0.54 \text{ MeV}$ and K.E. = $mc^2 - m_0 c^2$

Also
$$m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{m_0}{\sqrt{1 - (0.8)^2}} = \frac{m_0}{0.6}$$

$$\therefore E = mc^2 = \frac{m_0}{0.6} c^2 = \frac{m_0 c}{0.6} = \frac{0.54 \text{ MeV}}{0.6} = 0.9 \text{ MeV}$$

$$\therefore \text{K.E.} = (0.9 - 0.54) = 0.36 \text{ MeV}$$

Graphical Questions

- (a) B.E. per nucleon is maximum for Fe^{56} . For further detail refer theory.
- (a) $\omega = 2\pi\nu = \frac{2\pi c}{\lambda} = 2\pi c \bar{\nu} \Rightarrow \omega \propto \bar{\nu}$.
- (c)
- (d) The total number of atoms neither remains constant (as in option (a)) nor can ever increase (as in option (b) and (c)). They will continuously decrease with time. Therefore option (d) is correct.
- (c) $N = N_0 e^{-\lambda t} \Rightarrow \frac{dN}{dt} = -N_0 \lambda e^{-\lambda t}$
i.e. Rate of decay $\left(\frac{dN}{dt} \right)$ varies exponentially with time (t).
- (d) Rate $R = -\frac{dN}{dt} = \lambda N_0 e^{-\lambda t} = \lambda N \Rightarrow \frac{R}{N} = \lambda$ (constant)
i.e. graph between $\frac{R}{N}$ and t , be a straight line parallel to the time axis.
- (b) Read time for 50 count rate, it gives half life period of 3 hrs, one small square gives 600 counts (10×60). The number of small squares between graph and time axis are approx 24
 Hence count rate = $24 \times 600 = 14400$
- (b) Number of atoms undecayed $N = N_0 e^{-\lambda t}$
 Number of atoms decayed = $N_0 - N = N_0(1 - e^{-\lambda t})$
 \Rightarrow Decayed fraction $f = \frac{N_0 - N}{N_0} = 1 - e^{-\lambda t}$
i.e. fraction will rise up to 1, following exponential path as shown in graph (B).
- (c) Energy is released in a process when total Binding energy (B.E.) of the nucleus is increased or we can say when total B.E. of products is more than the reactants. By calculation we can see that only in case of option (c), this happens.
 Given $W \rightarrow 2Y$
 B.E. of reactants = $120 \times 75 = 900 \text{ MeV}$
 and B.E. of products = $2 \times (60 \times 85) = 1020 \text{ MeV}$
i.e. B.E. of products > B.E. of reactants.
- (d) $N = N_0 e^{-\lambda t}$ and $A = A_0 e^{-\lambda t} = \lambda N_0 e^{-\lambda t}$

$$\therefore N_{\text{rem}} = N_0 - N = N_0 - N_0 e^{-\lambda t} \Rightarrow N_{\text{rem}} = N_0 \left(1 - e^{-\lambda t}\right)$$

This is equation of straight line with negative slope.

11. (d) Radius of n orbit $r_n \propto n^2$, graph between r_n and n is a

parabola. Also, $\frac{r_n}{r_1} = \left(\frac{n}{1}\right)^2 \Rightarrow \log_e \left(\frac{r_n}{r_1}\right) = 2 \log_e (n)$

Comparing this equation with $y = mx + c$,

Graph between $\log_e \left(\frac{r_n}{r_1}\right)$ and $\log_e (n)$ will be a straight line, passing from origin.

Similarly it can be proved that graph between $\log_e \left(\frac{f_n}{f_1}\right)$ and $\log n$ is not a straight line.

12. (d) By using $N = N_0 e^{-\lambda t}$ and $\frac{dN}{dt} = -\lambda N$.

It shows that N decreases exponentially with time.

13. (b) Activity $= -\frac{dN}{dt} = \lambda N = \lambda N_0 e^{-\lambda t}$

i.e., graph between activity and t , be exponential having negative slope.

14. (d) Activity $A = \lambda N_0 e^{-\lambda t} \Rightarrow \log_e A = \log_e \lambda N_0 + \log_e e^{-\lambda t}$

$$\Rightarrow \log_e A = \log_e C - \lambda t \quad (\text{Take } \lambda N_0 = C)$$

$$\Rightarrow \log_e A = -\lambda t + \log_e C$$

This is the equation of a straight line having negative slope ($= -\lambda$) and positive intercept on $\log A$ axis.

15. (c) Charge density is uniform inside and then falls rapidly near the surface of the nucleus.

16. (a) $R = R_0 A^{1/3}$; where $R_0 = 1.2 \times 10^{-15} \text{ m}$.

$$\Rightarrow \log R = \log R_0 + \frac{1}{3} \log_e A$$

This is the equation of a straight line with positive slope.

17. (b) $\left|\frac{dN}{dt}\right| = \lambda N \Rightarrow \left|\frac{dN}{dt}\right| \propto N$

18. (c) Number of atom decayed $N' = N_0(1 - e^{-\lambda t})$

N' will increase with time (t) exponentially.

19. (a) $A_n = \pi r_n^2 \Rightarrow \frac{A_n}{A_1} = \left(\frac{r_n}{r_1}\right)^2 = \left(\frac{n}{1}\right)^4 \quad (\because r_n \propto n^2)$

Taking log both the side $\log_e \frac{A_n}{A_1} = 4 \log_e (n)$

Comparing it with $y = mx + c$, graph (4) is correct.

2. (a) ${}^{90}_{38}\text{Sr}$ decays to ${}^{90}_{39}\text{Y}$ by the emission of β -rays. Sr gets absorbed in bones along with calcium.

Reason is also true. ${}^{90}\text{Sr} \xrightarrow{\beta} {}^{90}\text{Y}$ which emits β -rays of very high energy. Sr does not emit γ -rays. The damage is by the β -rays only.

3. (b) Neutron is about 0.1 more massive than proton. But the unique thing about the neutron is that while it is heavy, it has no charge (it is neutral). This lack of charge gives it the ability to penetrate matter without interacting as quickly as the beta particles or alpha particles.

4. (b) Bohr postulated that electrons in stationary orbits around the nucleus do not radiate.

This is the one of Bohr's postulate. According to this the moving electrons radiate only when they go from one orbit to the next lower orbit.

5. (c) Nuclear stability depends upon the ratio of neutron to proton. If the n/p ratio is more than the critical value, then a neutron gets converted into a proton forming a β^- particle in the process. $n \rightarrow p + e^-$

The β^- particle (e^-) is emitted from the nucleus in some radioactive transformation. So electrons do not exist in the nucleus but they result in some nuclear transformation.

6. (a) ${}_Z X^A \rightarrow 2({}_2\text{He}^4) + 2({}_{-1}e^0) + 2\gamma + {}_{z-2}X^{A-8}$

7. (a) Experimentally, it is found that the average radius of a nucleus is given by

$$R = R_0 A^{1/3} \text{ where } R_0 = 1.1 \times 10^{-15} \text{ m} = 1.1 \text{ fm}$$

and A = mass number

8. (b)

9. (b) Rutherford confirmed the repulsive force on α -particle due to nucleus varies with distance according to inverse square law and that the positive charges are concentrated at the centre and not distributed throughout the atom.

10. (a) In α -particle scattering experiment, Rutherford found a small number of α -particles which were scattered back through an angle approaching to 180° . This is possible only if the positive charges are concentrated at the centre or nucleus of the atom.

11. (e) According to classical electromagnetic theory, an accelerated charge continuously emits radiation. As electrons revolving in circular paths are constantly experiencing centripetal acceleration, hence they will be losing their energy continuously and the orbital radius will go on decreasing and form spiral and finally the electron will fall on the nucleus.

12. (c) According to postulates of Bohr's atom model, the electron revolve round the nucleus in fixed orbit of definite radii. As long as the electron is in a certain orbits it does not radiate any energy.

13. (b) Maximum number of photon is given by all the transitions possible $= 4C_2 = 6$

Minimum number of transition = 1, that is directly jump from 4 to 1.


14. (b) When the atom gets appropriate energy from outside, then this electron rises to some higher energy level. Now it can return either directly to the lower energy level or come to the lowest energy level after passing through other lower energy levels,

Assertion and Reason

1. (c) In fusion, lighter nuclei are used so, fusion is not possible with ${}^{35}\text{Cl}$. Also binding energy of ${}^{35}\text{Cl}$ is not too small.

hence all possible transitions take place in the source and many lines are seen in the spectrum.

15. (d) Emission transitions can take place between any higher energy level and any energy level below it while absorption transitions start from the lowest energy level only and may end at any higher energy level. Hence number of absorptions transitions between two given energy levels is always less than the number of emission transitions between same two levels.



16. (a) We know that an electron is a very light particle as compared to an α -particle. Hence electron cannot scatter the α -particle at large angles, according to law of conservation of momentum. On the other hand, mass of nucleus is comparable with the mass of α -particle, hence only the nucleus of atom is responsible for scattering of α -particles.
17. (c) All those elements which are heavier than lead are radioactive. This is because in the nuclei of heavy atoms, besides the nuclear attractive forces, repulsive forces between the protons are also effective and these forces reduce the stability of the nucleus. Hence, the nuclei of heavier elements are being converted into lighter and lighter elements by emission of radioactive radiation. When they are converted into lead, the emission is stopped because the nucleus of lead is stable (or lead is most stable elements in radioactive series).
18. (d) The penetrating power is maximum in case of gamma rays because gamma rays are an electromagnetic radiation of very small wavelength.
19. (b) β -particles, being emitted with very high speed compared to α -particles, pass very little time near the atoms of the medium. So the probability of the atoms being ionised is comparatively less. But due to this reason, their loss of energy is very slow and they can penetrate the medium through a sufficient depth.
20. (b) β -particles are emitted with very high velocity (up to $0.99c$). So, according to Einstein's theory of relativity, the mass of a β -particle is much higher compared to its rest mass (m_0). The velocity of electrons obtained by other means is very small compared to c (Velocity of light). So its mass remains nearly m_0 . But β -particle and electron both are similar particles.

21. (c) Radioactivity $= -\frac{dN}{dt} = \lambda N = \frac{0.693N}{T}$

$$= \frac{0.693 \times 10^8}{50} = \frac{0.693 \times 1.2 \times 10^8}{60} = 0.693 \times 2 \times 10^6.$$

Radioactivity is proportional to $1/T$, and not to T .


22. (c) Fragments produced in the fission of U^{235} are radioactive. When uranium undergoes fission, barium and krypton are not the only products. Over 100 different isotopes of more than 20 different elements have been detected among fission products. All of these atoms are, however, in the middle of the periodic table, with atomic numbers ranging from 34 to 58. Because the neutron-proton ratio needed for stability in this range is much smaller than that of the original uranium nucleus, the residual nuclei called fission fragments, always have too many neutrons

for stability. A few free neutrons are liberated during fission and the fission fragments undergo a series of beta decays (each of which increases Z by one and decreases N by one) until a stable nucleus is reached. During decay of the fission fragments, an average of 15 MeV of additional energy is liberated.


23. (b) Electron capture occurs more often than positron emission in heavy elements. This is because if positron emission is energetically allowed, electron capture is necessarily allowed, but the reverse is not true *i.e.* when electron capture is energetically allowed, positron emission is not necessarily allowed.
24. (e) The whole mass of the atom is concentrated at nucleus and $M_{\text{e}} < (\text{Sum of the masses of nucleons})$ because, when nucleons combine, some energy is wasted.

Atomic and Nuclear Physics

Self Evaluation Test -26

1. In Bohr model of hydrogen atom, the force on the electron depends on the principal quantum number as
- (a) $F \propto 1/n^3$ (b) $F \propto 1/n^4$
 (c) $F \propto 1/n^5$ (d) Does not depend on n
2. A nucleus X emits 9α -particles and $5p$ particle. The ratio of total protons and neutrons in the final nucleus is
- (a) $\frac{Z-13}{A-Z-23}$ (b) $\frac{(Z-18)}{(A-36)}$
 (c) $\frac{(Z-13)}{(A-36)}$ (d) $\frac{(Z-13)}{(A-Z-13)}$
3. If $t_{1/2}$ is the half life of a substance then t_{\dots} is the time in which substance
- (a) Decays $\frac{3}{4}$ th (b) Remains $\frac{3}{4}$ th
 (c) Decays $\frac{1}{2}$ (d) Remains $\frac{1}{2}$
4. The energy level diagram for an hydrogen like atom is shown in the figure. The radius of its first Bohr orbit is
- | | | |
|-----------|-------|--------------|
| 0 eV | _____ | $n = \infty$ |
| - 6.04 eV | _____ | $n = 3$ |
| - 13.6 eV | _____ | $n = 2$ |
| - 54.4 eV | _____ | $n = 1$ |
- (a) 0.265 Å (b) 0.53 Å
 (c) 0.132 Å (d) None of these
5. How much work must be done to pull apart the electron and the proton that make up the Hydrogen atom, if the atom is initially in the state with $n = 2$
- (a) $13.6 \times 1.6 \times 10^{-19} J$ (b) $3.4 \times 1.6 \times 10^{-19} J$
 (c) $1.51 \times 1.6 \times 10^{-19} J$ (d) 0
6. The nuclide ^{131}I is radioactive, with a half-life of 8.04 days. At noon on January 1, the activity of a certain sample is 60089. The activity at noon on January 24 will be
- (a) 75 Bq (b) Less than 75 Bq
 (c) More than 75 Bq (d) 150 Bq
7. U^{238} decays into Th^{234} by the emission of an α -particle. There follows a chain of further radioactive decays, either by α -decay or by β -decay. Eventually a stable nuclide is reached and after that,
- no further radioactive decay is possible. Which of the following stable nuclides is the end product of the U^{238} radioactive decay chain
- (a) Pb^{206} (b) Pb^{207}
 (c) Pb^{208} (d) Pb^{209}
8. If the mass of a radioactive sample is doubled, the activity of the sample and the disintegration constant of the sample are respectively
- (a) Increases, remains the same
 (b) Decreases, increases
 (c) Decreases, remains same
 (d) Increases, decreases
9. When a sample of solid lithium is placed in a flask of hydrogen gas then following reaction happened
- $${}^1_1H + {}^7_3Li \rightarrow {}^4_2He + {}^4_2He$$
- This statement is
- (a) True
 (b) False
 (c) May be true at a particular pressure
 (d) None of these
- 
10. Consider an initially pure M gm sample of X , an isotope that has a half life of T hour, what is its initial decay rate ($N_A =$ Avogadro No.)
- (a) $\frac{M N_A}{T}$ (b) $\frac{0.693 M N_A}{T}$
 (c) $\frac{0.693 M N_A}{AT}$ (d) $\frac{2.303 M N_A}{AT}$
11. At a given instant there are 25% undecayed radioactive nuclei in a sample. After 10 sec the number of undecayed nuclei reduces to 6.25%, the mean life of the nuclei is
- (a) 14.43 sec (b) 7.21 sec
 (c) 5 sec (d) 10 sec
12. Highly energetic electrons are bombarded on a target of an element containing 30 neutrons. The ratio of radii of nucleus to that of Helium nucleus is $14^{1/3}$. The atomic number of nucleus will be
- (a) 25 (b) 26
 (c) 56 (d) 30
13. The ratio of ionization energy of Bohr's hydrogen atom and Bohr's hydrogen like lithium atom is
- (a) 1 : 1 (b) 1 : 3

- (c) 1 : 9 (d) None of these
14. What is the angular momentum of an electron in Bohr's hydrogen atom whose energy is -0.544 eV .
- (a) $\frac{h}{\pi}$ (b) $\frac{2h}{\pi}$
 (c) $\frac{5h}{2\pi}$ (d) $\frac{7h}{2\pi}$
15. Consider a hypothetical annihilation of a stationary electron with a stationary positron. What is the wavelength of resulting radiation.
- (a) $\frac{h}{2m_0c}$ (b) $\frac{h}{m_0c}$
 (c) $\frac{2h}{m_0c}$ (d) $\frac{h}{m_0c^2}$
- (h = Planck's constant, c = speed of light, m_0 = rest mass)
16. Nuclear reactions are given as
- (i) $\square (n, p)_{15}P^{32} \square (p, \alpha)_8O^{16}$ (iii) ${}_7\square^4 (\quad p)$
 ${}_6C^{14}$
- missing particle or nuclide (in box \square) in these reactions are respectively
- (a) $S, F, {}_0n^1$ (b) $F, S, {}_0n^1$
 (c) $Be, F, {}_0n^1$ (d) None of these
17. In a sample of hydrogen like atoms all of which are in ground state, a photon beam containing photons of various energies is passed. In absorption spectrum, five dark lines, are observed. The number of bright lines in the emission spectrum will be (assume that all transitions takes place).
- (a) 5 (b) 10
 (c) 15 (d) None of these
18. A hydrogen atom emits a photon corresponding to an electron transition from $n = 5$ to $n = 1$. The recoil speed of hydrogen atom is almost (mass of proton $\approx 1.6 \times 10^{-27} \text{ kg}$).
- (a) 10 ms (b) $2 \times 10^{-4} \text{ ms}$
 (c) 4 ms (d) $8 \times 10^{-4} \text{ ms}$
19. Number of nuclei of a radioactive substance at time $t = 0$ are 1000 and 900 at time $t = 2 \text{ s}$. Then number of nuclei at time $t = 4 \text{ s}$ will be
- (a) 800 (b) 810
 (c) 790 (d) 700
20. The ratio between total acceleration of the electron in singly ionized helium atom and hydrogen atom (both in ground state) is
- (a) 1 (b) 8
 (c) 4 (d) 16
21. If the series limit of Lyman series for Hydrogen atom is equal to the series limit of Balmer series for a hydrogen like atom, then atomic number of this hydrogen like atom will be
- (a) 1 (b) 2
 (c) 3 (d) 4
22. Which sample contains greater number of nuclei :
- a $5.00\text{-}\mu\text{Ci}$ sample of ${}^{239}\text{Pu}$ (half-life 6560 y) or a $4.45\text{-}\mu\text{Ci}$ sample of ${}^{241}\text{Am}$ (half-life 7370 y)
- (a) ${}^{239}\text{Pu}$ (b) ${}^{241}\text{Am}$
 (c) Equal in both (d) None of these
23. The fission of ${}^{235}\text{U}$ can be triggered by the absorption of a slow neutrons by a nucleus. Similarly a slow proton can also be used. This statement is
- (a) Correct (b) Wrong
 (c) Information is insufficient (d) None of these
24. The radioactivity of a given sample of whisky due to tritium (half life 12.3 years) was found to be only 3% of that measured in a recently purchased bottle marked "7 years old". The sample must have been prepared about
- (a) 220 years back (b) 300 years back
 (c) 400 years back (d) 70 years back
25. The following diagram indicates the energy levels of a certain atom when the system moves from $4E$ level to E . A photon of wavelength λ_1 is emitted. The wavelength of photon produced during its transition from $\frac{7}{3}E$ level to E is λ_2 . The ratio $\frac{\lambda_1}{\lambda_2}$ will be



- (a) $\frac{9}{4}$
 (c) $\frac{3}{2}$

- (b) $\frac{4}{9}$
 (d) $\frac{7}{3}$

AS Answers and Solutions

(SET -26)

1. (b) $F \propto \frac{v^2}{r}$ also $v \propto \frac{1}{n}$ and $r \propto n^2 \Rightarrow F \propto \frac{1}{n^4}$
2. (a) ${}_Z X^A \xrightarrow{9\alpha} {}_{Z-18} X^{A-36} \xrightarrow{5\beta} {}_{Z-13} X^{A-36}$

Number of protons = ($Z = 13$)

Number of neutrons = $(A - 36) - (Z - 13) = (A - Z - 23)$

$$\therefore \frac{P}{N} = \frac{(Z - 13)}{(A - Z - 23)}$$

1530 Atomic and Nuclear Physics

3. (a) You must remember that $t_{1/2}$ is time in which substance decays half. Hence in $t_{1/2}$ time substance decays $\frac{3}{4}$ th.

4. (a) We know that $E_n = -13.6 \frac{Z^2}{n^2} eV$ and $r_n = 0.53 \frac{n^2}{Z} (\text{\AA})$

Here for $n=1$, $E_1 = -54.4 eV$

$$\text{Therefore } -54.4 = -13.6 \frac{Z^2}{1^2} \Rightarrow Z = 2$$

$$\text{Hence radius of first Bohr orbit } r = \frac{0.53(1)^2}{2} = 0.265 \text{ \AA}$$


5. (b) The electrostatic P.E. is zero when the electron and proton are far apart from each other. Work done in pulling electron and proton far away from each other

$$W = E_f - E_i = 0 - E_i = -\left(-\frac{13.6}{n^2} eV\right)$$

$$\Rightarrow W = \frac{13.6}{(2)^2} \times 1.6 \times 10^{-19} J = 3.4 \times 1.6 \times 10^{-19} J$$

6. (c) Number of days from January 1 to January 24 = 23 days.

$$\text{Number of half lives } n = \frac{23}{8.04} = 2.86 (< 3)$$



In three half lives activity becomes 75 Bq, but the given number of half lives are lesser than 3 so activity becomes greater than 75 Bq.

7. (a) $(4n+2)$ series starts from U^{238} and it's stable end product is Pb^{206} .
8. (a) Activity depends upon mass, but λ doesn't change.
9. (b) The given reaction is a nuclear reaction, which can take place only if a proton (a hydrogen nucleus) comes into contact with a lithium nucleus. If the hydrogen is in the atomic form, the interaction between its electron cloud and the electron cloud of a lithium atom keeps the two nuclei from getting close to each other. Even if isolated protons are used, they must be fired at the Li atom with enough kinetic energy to overcome the electric repulsion between the proton and Li atom.

10. (c) $N = N_0 e^{-\lambda t} \Rightarrow \left| \frac{dN}{dt} \right| = N_0 \lambda e^{-\lambda t}$

Initially at $t = 0$, $\left| \frac{dN}{dt} \right|_{t=0} = N_0 \lambda$

where N_0 = Initial number of undecayed atoms

$$= \frac{\text{Mass of the sample}}{\text{Mass of a single atom of X}} = \frac{M}{A/N_A} = \frac{MN_A}{A}$$

$$\therefore \left| \frac{dN}{dt} \right|_{t=0} = \frac{MN_A \lambda}{A} = \frac{0.693 MN_A}{AT}$$

11. (b) In 10 sec, number of nuclei has been reduced to one fourth (25% to 6.25%).

Therefore its half life is $T_{1/2} = 5 \text{ sec}$.

$$\therefore \text{Mean life } T = \frac{T_{1/2}}{0.693} = \frac{5}{0.693} = 7.21 \text{ sec}$$

12. (b) By using $R = R_0 A^{1/3} \Rightarrow \frac{R_1}{R_2} = \left(\frac{A_1}{A_2}\right)^{1/3}$

$$\Rightarrow \frac{R}{R_{He}} = \left(\frac{A}{4}\right)^{1/3} \Rightarrow (14)^{1/3} = \left(\frac{A}{4}\right)^{1/3}$$

$$\Rightarrow A = 56 \text{ so } Z = 56 - 30 = 26.$$

13. (c) Energy of an electron in ground state of an atom (Bohr's hydrogen like atom) is given as

$$E = -13.6 Z^2 eV \quad (Z = \text{atomic number of the atom})$$

$$\Rightarrow E_{He} = 13.6 Z$$

$$\Rightarrow \frac{(E_{ion})_H}{(E_{ion})_{Li}} = \left(\frac{Z_H}{Z_{Li}}\right)^2 = \left(\frac{1}{3}\right)^2 = \frac{1}{9}$$

14. (c) By using $E = -\frac{13.6}{n^2} eV$ (for H atom)

$$\Rightarrow -0.544 = -\frac{13.6}{n^2} \Rightarrow n = 25 \Rightarrow n = 5$$

$$\therefore \text{Angular momentum} = n \frac{h}{2\pi} = \frac{5h}{2\pi}$$

15. (b) From conservation of momentum, two identical photons must travel in opposite directions with equal magnitude of momentum and energy $\frac{hc}{\lambda}$

$$\text{from conservation of energy } \frac{hc}{\lambda} + \frac{hc}{\lambda} = m_0 c^2 + m_0 c^2$$

$$\Rightarrow \lambda = \frac{h}{m_0 c}$$

16. (a) (i) ${}_{16}S^{32} + {}_0n^1 \rightarrow {}_{15}P^{32} + {}_1H^1$
 (ii) ${}_9F^{19} + {}_1H^1 \rightarrow {}_2He^4 + {}_8O^{16}$
 (iii) ${}_7N^{14} + {}_0n^1 \rightarrow {}_6C^{14} + {}_1H^1$

17. (c) Number of lines in absorption spectrum = $(n-1)$

$$\Rightarrow 5 = n - 1 \Rightarrow n = 6$$

\therefore Number of bright lines in the emission spectrum

$$= \frac{n(n-1)}{2} = \frac{6(6-1)}{2} = 15.$$

18. (c) The Hydrogen atom before the transition was at rest. Therefore from conservation of momentum.

$$P_{H\text{-atom}} = P_{\text{photon}} = \frac{E_{\text{radiated}}}{c} = \frac{13.6 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) eV}{c}$$

$$1.6 \times 10^{-27} \times v = \frac{13.6 \left(\frac{1}{1^2} - \frac{1}{5^2} \right) \times 1.6 \times 10^{-19}}{3 \times 10^8}$$

$$\Rightarrow v = 4.352 \text{ m/s} \approx 4 \text{ m/sec}$$

19. (b) In 2 sec only 90% nuclei are left behind. Thus in next two second 90% of 900 or 810 nuclei will be left.

$$(4E - E) = \frac{hc}{\lambda_1} \Rightarrow \lambda_1 = \frac{hc}{3E} \quad \dots (i)$$

20. (b) Acceleration $a \propto \frac{v^2}{r}$

where $v \propto \frac{Z}{n}$ and $r \propto \frac{n^2}{Z} \Rightarrow a \propto \frac{Z^3}{n^4}$

Since both are in ground state i.e., $n = 1$

$$\text{so } a \propto Z \Rightarrow \frac{a_{He^+}}{a_H} = \left(\frac{Z_{He^+}}{Z_H} \right)^3 = \left(\frac{2}{1} \right)^3 = \frac{8}{1}$$

Transition from $\frac{7}{3}E$ to E

$$\left(\frac{7}{3}E - E \right) = \frac{hc}{\lambda_2} \Rightarrow \lambda_2 = \frac{3hc}{4E} \quad \dots (i)$$

From equation (i) and (ii) $\frac{\lambda_1}{\lambda_2} = \frac{4}{9}$

21. (b) By using $\frac{1}{\lambda} = RZ^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$

For Hydrogen atom $\frac{1}{(\lambda_{\min})_H} = R \left[\frac{1}{1^2} - \frac{1}{\infty} \right] = R$

$$\Rightarrow (\lambda_{\min})_H = \frac{1}{R} \quad \dots (i)$$

For hydrogen like atom $\left(\frac{1}{\lambda_{\min}} \right)_{\text{atom}} = RZ^2 \left(\frac{1}{2^2} - \frac{1}{\infty} \right)$

$$\Rightarrow (\lambda_{\min})_{\text{atom}} = \frac{4}{RZ^2} \quad \dots (ii)$$

From equation (i) and (ii) $\frac{1}{R} = \frac{4}{RZ^2} \Rightarrow Z = 2$.

22. (c) The activity $\left(-\frac{dN}{dt} \right) = \lambda N \Rightarrow N = \left(-\frac{dN}{dt} \right) \left(\frac{T_{1/2}}{\log_e 2} \right)$

Taking the ratio of this expression for ^{240}Pu to this same expression for ^{243}Am ,

$$\frac{N_{Pu}}{N_{Am}} = \frac{\left(-\frac{dN_{Pu}}{dt} \right) (T_{1/2})_{Pu}}{\left(-\frac{dN_{Am}}{dt} \right) (T_{1/2})_{Am}} = \frac{(5 \mu\text{Ci}) \times (6560\text{y})}{(4.45 \mu\text{Ci}) \times (7370\text{y})} = 1$$

i.e. the two samples contains equal number of nuclei.

23. (b) Because the neutron has no electric charge, it experience no electric repulsion from a U^+ nucleus. Hence a slow moving neutron can approach and enter a U^+ nucleus, thereby providing the excitation needed to trigger fission. By contrast a slow moving proton feels a strong repulsion from a U^+ nucleus. It never get's close to the nucleus, so it cannot trigger fission.
24. (d) After one half life period, the activity of Tritium becomes 50%.
 After 2 half life period 25%
 After 3 half life period 12.5%
 After 4 half life period 6.25%
 After 5 half life period 3.12% \approx 3%
 It is 5×12.5 years + 7 years i.e. approximately 70 years only .
25. (b) Transition from $4E$ to E