



Organic Chemistry is the study of compds. containing carbon.




Characteristic of C :-


1. Tetraivalent : C form 4 bond in excited state.




2. Tetrahedral : 4 bonds are directed towards corner of tetrahedron.



3. Tendency to form multiple bond :





4. Catenation : Self linking property of C to form long chain.




5. Hybridisation :

σ bond : It is formed by coaxial overlapping of atomic orbitals.



π -bond : formed by colateral / Sidewise overlapping of atomic orbitals.




Note :- ① σ bond is stronger than π -bond becoz ~~the~~ extent of overlapping is more in σ bond.

② π -bond is more reactive than σ -bond.


Str.	σ, π bond	hyb.	B.A.	Shape
$\begin{array}{c} \\ C \\ / \backslash \end{array}$	4, 0	sp^3	$109^\circ 28'$	tetrahedral
$>C=$	3, 1	sp^2	120°	trigonal planar
$-C\equiv$	2, 2	sp	180°	linear
$=C=$	2, 2	sp	180°	linear

S.No. = no. of σ bond + no. of lp.

CH_3^+



CH_3^-



E.N. $\propto$ % s-character

% s char	sp $\frac{1}{2} \times 100$ = 50%	sp^2 $\frac{1}{3} \times 100$ = 33.3%	sp^3 $\frac{1}{4} \times 100$ = 25%
----------	---	---	---

$$E.N. \rightarrow sp > sp^2 > sp^3$$

$3.25 \quad 2.75 \quad 2.5$






$$B.L. \propto \frac{1}{E.N.} \quad sp < sp^2 < sp^3$$

$$B.S. \propto E.N. \quad sp > sp^2 > sp^3$$


or

$$B.S. \propto \frac{1}{B.L.}$$


Ques Find hyb. of underlined C ?




Ques Find hybridisation of C-C single bond













Ques Find hyb. of C₂ & C₃ in following




Ques Find hyb. of Carbonium ion ?



Ques find no. of σ & π -bond in following?




13.



16σ, 4π

14.




14σ, 1π

15.



23σ, 2π




16.



10π, 2π



Ques Arrange following in ↑sing order of B.O.L?

①






B.O.L. α 1
E.N

②



b > c > a

③






γ > α > β

IIT

PAGE NO.: 7
DATE: / /

(4)




Type of Carbon and Hydrogen :-



1. Primary Carbon & Primary Hydrogen :-

$1^\circ \text{C} \rightarrow \text{C}$ which is attached to only 1° carbon.

$1^\circ \text{H} \rightarrow \text{H}$ which is attached to only 1°C .



$1^\circ \text{C} \rightarrow 4$





$1^\circ \text{H} \rightarrow 6$

2. Secondary Carbon & Secondary Hydrogen :-

$2^\circ \text{C} \rightarrow \text{C}$ which is attached to 2° carbon.


$2^\circ \text{H} \rightarrow \text{H}$ which is attached to 2°C .



3. Tertiary Carbon & Tertiary Hydrogen :-




$3^\circ \text{C} \rightarrow \text{C}$ which is attached to 3 carbon.

$3^\circ \text{H} \rightarrow \text{H}$ which is attached to 3°C .



4. Quaternary Carbon :-

$4^\circ\text{C} \rightarrow$ Carbon which is attached to 4 carbon atom



$$1^\circ\text{C} \rightarrow 7\text{C}$$

$$1^\circ\text{H} \rightarrow 19\text{H}$$

$$2^\circ\text{C} \rightarrow 3\text{C}$$

$$2^\circ\text{H} \rightarrow 5\text{H}$$





$$3^\circ\text{C} \rightarrow 3$$

$$3^\circ\text{H} \rightarrow 3$$


$$4^\circ\text{C} \rightarrow 1$$

Type of Alcohol :-


1. Primary Alcohol :- (1°OH) The alcoholic compd. in which OH gp. is attached to 1°C .



2. **Secondary Alcohol** : The compd in which OH gp. is attached to 2° C.




3. **Tertiary Alcohol** : The alcoholic compd. in which OH gp is attached to 3° C




Type of Amine :-


1. **Primary Amine** : N is attached to only 1 C atom.





2. **Secondary Amine** :- N is attached to 2 C atom.




3. **Tertiary Amine** :- ~~in which~~ N is attached to 3 C atom.



$$\text{NH}_4^+$$


ex



Classification :-

- ① Based on structure
- ② Based on Group
- ③ Based on Homologous

① Based on structure :-

Open chain / aliphatic / acyclic

Closed chain / cyclic

Saturated


Unsaturated

Homocyclic




Heterocyclic

C-C

+ alkanes





Paraffins




Olefinic bond

alkene



acetylinic bond



alkyne



alicyclic aromatic



Alicyclic



Aromatic
















Pyridine





Pyrrole

Ques which of the following is unsaturated compo






15.






Ques which of the following is unsaturated?

✓ Maleic acid



2. Succinic acid








A:- Acetic acid is not an unsaturated compd. ✓
R:- acetic acid do not have olefinic ~~comp~~ bond. (2)

A:- acetic acid is saturated compd. ✓ (2)
R: acetic acid have $-\text{C}-$ bond. ✓
 O


A: acetic acid have $-\text{C}-$ bond. (2)
 O

R: acetic acid is saturated compd.

Ques find no. of heteroatoms in following ?




4.




0

5.




2

6.




2

7.



0

(2) Classification based on Group :-



Hydrocarbon

hyd-radical

phy. property

B.P., M.P. &

density

etc

(G₁)

funcⁿ gp.

→ chemical property

- OH

- NH₂

- X

- COOH

- CH=O

Pentyl = amyl

Hydrocarbon Radical (R-) :-

Saturated :

Monovalent :-

alkane $\xrightarrow{-H}$ alkyl

$CH_4 \xrightarrow{-H} CH_3-$
Methane Methyl

$CH_3-CH_3 \xrightarrow{-H} CH_3-CH_2-$
ethane ethyl

$CH_3-CH_2-CH_3 \xrightarrow{-H} \begin{cases} CH_3-CH_2-CH_2- & n\text{-propyl} \\ CH_3-\underset{|}{CH}-CH_3 & iso\text{-propyl} \end{cases}$
Propane

Normal $\div$ • straight chain hydrocarbon
• no branching

eg $\begin{array}{c} CH_2-CH_3 \\ | \\ CH_3-CH_2 \end{array}$
n-butane


 $CH_3-CH_2-CH_2-CH_2-CH_2-CH_3$
n-hexane

$\begin{array}{c} CH_3 \\ | \\ CH_3-CH_2 \end{array}$
propane


n-pentane

$CH_3(CH_2)_5CH_3$
n-heptane








$CH_3-CH_2-CH_2-$

n-propyl

$CH_3-CH_2-CH_2-CH_2-$

n-butyl

Iso : When methyl gp. is attached at 2nd last C or 2nd C (in case of alkanes) is c/a iso-comp.




iso-butyl


t-butyl

isopropyl



Neo :- When 2 methyl gp. is attached to 2nd last C or 2nd C (in case of alkane) c/a neo compd.




neo-Pentane




neo-hexane




Exception






iso octane


PAGE NO.: 17
DATE: / /




n-pentyl



Sec. pentyl




iso-pentyl





neo pentyl

Bivalent:- 2 valency free
from same C-atom $\rightarrow$ alkylidene
from vicinal C-atom $\rightarrow$ alkylene
terminal C-atom $\rightarrow$ polymethylene




Methylidene
or

Methylene




ethylidene

ethylene











propylidene

propylene




isopropylidene





Aromatic ÷




benzene
or
phene




phenyl





phenylene





toluene





benzyl




benzal




benzo



o-tolyl





m-tolyl



p-tolyl

Concept of Homology :-

Those compounds which have same structural features as well as same general formula but differ by $-CH_2-$ groups/a homologs & series is c/a homologous series.




G.F.




alkane	C_nH_{2n+2}	$CH_4, CH_3-CH_3, CH_3-CH_2-CH_3$
alkene	C_nH_{2n}	$CH_2=CH_2, CH_3-CH=CH_2, CH_3-CH_2-CH=CH_2$
alkyne	C_nH_{2n-2}	$CH \equiv CH, CH_3-C \equiv CH$
Alcohol	$C_nH_{2n+2}O$	$CH_3-OH, CH_3-CH_2-OH, CH_3-CH_2-CH_2-OH$
ether	$C_nH_{2n+2}O$	$CH_3-O-CH_3, CH_3-O-CH_2-CH_3$
Halide	$C_nH_{2n+1}X$	
aldehyde or	$C_nH_{2n}O$	$H-\overset{\overset{O}{ }}{C}-H, CH_3-CH=O, CH_3-CH_2-CHO$
Ketone	$C_nH_{2n}O$	$CH_3-\overset{\overset{O}{ }}{C}-CH_3$
acid/ester	$C_nH_{2n}O_2$	$H-\overset{\overset{O}{ }}{C}-OH, CH_3-COOH$

Characteristics of Homologues :-


1. Have same general formula i.e. same method of preparation.
2. Have ~~a~~ same type of atom.
3. have same functional gp. hence similar chemical property.
4. Have different physical property.
5. 2 successive members differ by CH_2 -gp i.e. a weight of 14 unit.
















Ques Write lower & higher homologue of isopentyl alcohol.



Ques Which of the following is pair of Homologues?









Nomenclature :-

Common Name	Derived Name	IUPAC system
-------------	--------------	--------------

1. Common Name OR TRIVIAL NAMING :-

- based on source of origin.

Structure	Source	Common Name
CH_4	Marshy area	Marsh Gas
$\text{CH}_3 - \text{OH}$	distillation of wood	wood spirit
HCOOH	formica (Red Ant)	formic acid
$\text{CH}_3 - \text{COOH}$	acetum (Vinegar)	acetic acid
$\text{CH}_3 - \underset{\text{OH}}{\underset{ }{\text{CH}}} - \text{COOH}$	Milk	Lactic acid
$\text{C}_3\text{H}_7 - \text{COOH}$	Butter	butyric acid
$\text{HOOC} - \text{CH}_2 - \underset{\text{OH}}{\underset{ }{\text{CH}}} - \text{COOH}$	Apple	Malic acid
$\text{HOOC} - \underset{\text{OH}}{\underset{ }{\text{CH}}} - \underset{\text{OH}}{\underset{ }{\text{CH}}} - \text{COOH}$	tamarind tree	tartaric acid



Common Name functional Group

① Non-Terminating
 (with ^{out} C containing func gp)
 eg -OH, -NH₂, -O-, $\underset{\text{O}}{\underset{||}{\text{C}}}$ -


② terminating
 (C containing func gp)
 -COOH, -CHO

1. Non terminating Suffix


-OH	alcohol
-SH	thio alcohol
-X	Halide
-NH ₂	amine
-NH-	amine
-N- 	amine
-O-	ether
-S-	thio ether
$\underset{\text{O}}{\underset{ }{\text{C}}}$ -	ketone

Polyvalent func gp:-
 1. If same hydrocarbon gp is attached then di, tri, tetra etc used.
 2. If diff. hydrocarbon are attached then write them in alphabetical order


Note:- di, tri, tetra, primary, secondary, etc. are not considered in alphabetical order while terms iso, neo are considered in alphabetical order




dimethyl ether



isopropyl methyl ether




dimethyl ketone




ethyl methyl ketone




ethyl alcohol




isopropyl chloride




isobutyl amine




neopentyl alcohol




Vinyl chloride



allyl alcohol



dimethyl amine



ethyl isopropyl methyl amine

H Terminating fun^cgp :-

Prefix :-

Total no. of C

1C

2C

3C

4C

5C

Prefix

form

acet

propion

butyr $\rightarrow$ n

Valer

Iso

①

Cu

eq

3C + 1db

 $C-C=C$

acryl

4C + 1db

 $C-C=C-C$

croton

Suffix :-

P. fg

 $-COOH$ $-C(=O)-O-R$ $-C(=O)-X$ $-CHO$ $-C(=O)-O-C(=O)-$ $-CN$ $-NC$ $-C(=O)-NH_2$

Suffix

ic acid

alkyl - - - ate

yl halide


aldehyde

ic anhydride


o nitrile

o isonitrile


amide




formic acid




acetic acid




isobutyric acid




n-butyric acid




neo valeric acid




formaldehyde




acetaldehyde




propionaldehyde




isobutyraldehyde




isovaler aldehyde




acryl aldehyde




~~croton~~ acrylic acid




croton aldehyde




crotonic acid




methyl acetate




ethyl formate




isopropyl etho acetate




acetyl chloride




formyl chloride




propiononitrile





acetone isonitrile





formamide



acetamide



acetic anhydride



acetic propionic anhydride

DERIVED NAME :


selection of parent chain

③ Based on famous homolog


$4^\circ\text{C} > 3^\circ\text{C} > 2^\circ\text{C} > 1^\circ\text{C}$

alkane :- CH_4


methane



methyl methane




ethyl methyl methane




ethyl dimethyl methane

Ques write derived name of iso-octane?




isobutyl trimethyl methane

alkene :



ethylene

Eg > multiple bond
 $\equiv > =$



methyl ethylene




sym. dimethyl ethylene




unsym. dimethyl ethylene

alkyne :




acetylene



methyl acetylene



ethyl methyl acetylene




Vinyl acetylene

alcohol :



Carbinol



ethyl Carbinol




ethyl methyl carbinol




aldehyde :




acetaldehyde



methyl acetaldehyde



isopropyl methyl acetaldehyde



acid : $\text{CH}_3 - \text{COOH}$ acetic acid

$\begin{array}{c} \text{CH}_2 - \text{CH}_3 \\ | \\ \text{CH}_3 - \text{CH} - \text{COOH} \end{array}$ ethyl methyl acetic acid

$\text{Cl}_3 \text{C COOH}$

$\begin{array}{c} \text{Cl} \diagdown \\ \text{Cl} - \text{C} - \text{COOH} \\ \text{Cl} \diagup \end{array}$ trichloro acetic acid

Ketone : $\text{CH}_3 - \underset{\text{O}}{\overset{\parallel}{\text{C}}} - \text{CH}_3$ acetone

$\text{CH}_3 - \text{CH}_2 - \underset{\text{O}}{\overset{\parallel}{\text{C}}} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$

ethyl methyl acetone

IUPAC Naming

International Union of Pure & Applied Chemistry

Parts of IUPAC

1. Word Root
2. Prefix [Pri
 [Sec.
3. Suffix [Pri.
 [Sec.

Secondary Prefix 2°	Primary Prefix 1°	Word Root WR	Primary Suffix 1°	Secondary Suffix 2°
substituent is written in alphabetical order	used when compd is cyclic	represent total no. of C in Principal chain	tells whether compd. is saturated or unsaturated	tells about Principal func gp.
- Cl Chloro	"Cyclo"	1C - meth	C - C ane	- COOH
- NO ₂ Nitro		2C - eth	1C = C ene	- CHO
		3C - Prop	1C ≡ C yne	- OH
		⋮	2C = C diene	- NH ₂
		⋮	2C ≡ C diyne	
		10C - dec	C = C & C ≡ C enyne	
		11C - undec		
		12 - dodec		
		⋮		
		⋮		
		20 Eicos		

PRIORITY ORDER OF FUNCTIONAL GROUP :-

Functional Gp.	Prefix	Suffix
- (C)OOH (carboxylic acid)	x	oic acid
- COOH	Carboxy	carboxylic acid
- SO ₃ H (Sulphonic acid)	Sulpho	Sulphonic acid
$\begin{array}{c} \text{O} \\ \\ \text{-(C)} \end{array} \begin{array}{c} \text{O} \\ \\ \text{-(C)} \end{array} > \text{O} \text{ (anhydride)}$	x	oic anhydride

Functional Group	Prefix	Suffix
- (C)OOR (ester)	x	alkyl --- oate
- COOR	alkoxycarbonyl or carbalkony	alkyl --- carboxylate
- (C)OX (acid halide)	x	oyl halide
- COX	halo formyl	carbonyl halide
- (C)ONH ₂ (amide)	x	amide
- CONH ₂	carbamoyl	carboxamide
- (C)N (cyanide)	x	Nitrile
- CN	cyano	carbonitrile
- N $\equiv$ (C) (isocyanide)	x	isonitrile
- NC	isocyano/ carbonyl amine	carbonyl amine
- (C)HO (aldehyde)	oxo	al
- CHO	formyl	carbaldehyde
- (C)- (Ketone) O	keto/oxo	one
- OH (alcohol)	hydroxy	ol
- SH (thio alcohol)	mercapto	thiol
- NH ₂ (amine)	amino	amine

Substituents

- R	Prefix
- X	alkyl
- NH ₂	halo
- N $\equiv$ O	amino
- O - N = O	nitro
- N = O	nitrite
	nitroso

Substituents	Prefix
$-OCH_2CH_3$	ethoxy
$-CH_2-OH$	hydroxy methyl
$-CH_2-Cl$	chloro methyl
$-NH-CH_3$	methyl amino
$-S-$	thio
$-S-R$	alkyl thio
$CH_3-\overset{\overset{O}{\parallel}}{C}-O-$	acetoxy/ethanoxyloxy
$CH_3CH_2-\overset{\overset{O}{\parallel}}{C}-O-$	propanoxyloxy
$C_6H_5-\overset{\overset{O}{\parallel}}{C}-O-$	benzoxyloxy
$-OR$	alkoxy


Rules of IUPAC :-















1. Selection of Principal Carbon chain
2. Numbering of P.C.C.

1. Selection of Principal Carbon Chain (P.C.C)


P.C.C. will be that largest longest continuous chain of carbon which have




Pfg > Max^m no. of multiple bond > max. no. of C-atom








If symmetric molecule is +nt then priority is given to double & bond over triple bond












If 2 diff. P.C.C. have same no. of C in a molecule ~~the~~ then that P.C.C. will be selected as P.C.C. which have max^m no. of Sb (/ more branching)





Numbering of P.C.C.:-



- ① Lowest no. will given to
Pfg > Mb > Sb




② If more than one substituent are +nt then least locant no. Rule is followed.




3. If 2 or more substituent are +nt at same posⁿ from either side then no. starts from the side where alphabetically preferred substituent get lowest no.




4. If db & tb are at similar posⁿ from either side of PCC then priority is given to db






mb
LHS ← 1,5 3,4 ✓
RHS 1,5 3,4
tb




mb
LHS 2,5
RHS 1,4 ✓


Ques




2-Methyl Butane




3-Bromo - 2 Chloro Pentane




3,4 diethyl - 2,5 dimethyl
Hexane




2-chloro - 3,5 - diethyl - 6-methyl
heptane




2-Bromo - 4-Chloro - 3,3-diethyl
Pentane




5-Bromo-3-Chloro, 2,6-dimethyl
Octane




5-Chloro - ~~ene~~ Hex-2-ene




3 (1'-chloro Propyl) ~~ene~~ Pent-
1,4-diene



3-Bromo-4-Chloro hex-1,5 diene




4-ethyl-3-methyl-hept-1,6-di
ene




Pent-1-ene-4-yne


AI PMT-10




Hex-4-ene-1-yne




2-Ethyl-3-Methyl
Butane-1,3-diene




4-Ethenyl hept-1-ene-5-yne
or
4-ethenyl hept-1-en-5-yne



4-Bromo-3-chloro-Hex-1-ene
5-yne




3-ethenyl-Oct-1-ene-6-yne




4-Bromo-2-chloro-3-Ethenyl-
Hex-1-ene-5-yne

4-Bromo-2-chloro-3-Vinyl Hex-1-ene-
5-yne


-OH

Prefix
HydroxySuffix
ol


4-Bromo - 5-Chloro - 2-Hexanol




2-Propyl - But - 3-ene - 1-ol




3-Chloro - Hex - 3-ene - 2,5-diol



hept - 5-ene - 2,3,6-triol


6 Chloro - 4 Hydroxy Methyl hept -
2,5-diol

	Prefix	Suffix
- SH	mercapto	thiol




Hept - 6-ene-1-yne
- 3-thiol


	Prefix	Suffix
- NH ₂	amino	amine



2-Ethyl - Pent - 3-ene -
1-amine




N-ethyl - 2-methyl
Propanamine




N-isopropyl - N-methyl
ethene amine

	Prefix	Suffix
- CHO	oxo formyl	al carbaldehyde




2-Propyl But - 3-ene al




5-Bromo - 3-chloro - 4-formyl - 2,6-dimethyl
heptan-1,7-dial

PAGE NO.: 43

DATE: / /




3-Bromo - 2-ethenyl - 4-methyl
Pentan-1,5-dial




Note: When more than 2 carbons contain
same func gp are attached to main
chain then exclude all of them.

5-Bromo - 3-chloro hept - 3-ene - 2,4,6
- tricarbaldehyde




4-ethyl - 3-formyl methyl -
hexan-1,6-dial




Prefix
oxo / keto

Suffix
one



3-Bromo - Pent - 4-en - 2-one




5-Bromo - 3-chloro hexan - 2,4-dione

-CN


-(C)N
-CN

x
cyano


nitrile
carbonitrile



3-chloro-2-methyl-4-propyl-hex-5-ene nitrile




4-bromo-2-ethyl-5-methylhept-3-ene nitrile






PKg
LHS 1,3,4
RHS 1,2,4 ✓

4-bromo-1-chlorobutane-1,2,4-tricarbonitrile




5-bromo-2-chloro-3-cyano-1,6-dinitrile hexane


5-bromo-2-chloro-4-cyano-methylheptane-1,7-dinitrile






x
carbamoyl
amide
carboxamide



4-ethyl-2-methylhex-5-enamide



N-ethenyl-3,N-dimethyl Pentanamide




X


halo formyl

acyl halide




carbonyl halide



3-Chloro-Butanoyl Chloride



4-Bromo-Pent-2-enoyl Chloride




X


alkyl --- oate

alkoxy carbonyl


alkyl --- carboxylate



methyl - Propanoate





Ethyl - Methanoate




Propyl

Ethyl - 3-Methyl-Pent 2-ene oate





X

oic anhydride



Butanoic anhydride





ethanoic - Propanoic anhydride




sulpho

sulphonic acid

4-Bromo - 3-Methyl
Pentane sulphonic
acid.


X


oic acid




carboxy

carboxylic acid

4-Bromo - 2,5 - diethyl
hex - 5-ene - oic acid




4 (1-Chloro Ethyl) - 2, 3 - dimethyl
hex - 5 - ene - oic acid




2-bromo - 4 - carboxy hex - 2 - ene -
1, 6 - dioic acid

4 - Bromo - 1, 2, 4 - dicarboxy
dicarboxylic acid But -
3 - ene




5 - Bromo - 4 - carboxy methyl
2 - Methyl - Hept - 3 - ene - 1, 7 - dioic
acid


IUPAC Naming




methoxy methane



methoxy ethane




2 - ethoxy butane




butoxy ethene

Oxa - System

→ 0th C atom numbering Least locant no. given




2-oxa-butane




2,5-dioxa heptane

Cyclic ether




epoxy ethane


or oxirane (trivial name)



epoxy




1,2-epoxy propane




2,4-epoxy - 3,3,5-trimethyl hexane


IUPAC Naming of Polyfunctional group :-




3-amino - 4-hydroxy
Pentanal




3-amino - 2-Bromo - Pent-3-ene
oic acid




5-amino-4-hydroxy
hex-2-ene amide




2-Chloro-5-hydroxy heptan-
3-one




2-Bromo-4-hydroxy
Pent-2-ene yl chloride




ethyl-2-hydroxy-3-Propyl
Pent-4-enoate




2-amino-4-Bromo-3-hydroxy Pentan^{3-ol}
~~amine~~




6-amino hept-4-ene-3-ol




amino-methanoyl chloride




methanamide




methoxy methanoic acid




chloro methanoic acid




ethyl chloro methanoate



amino methanoic acid







methanoic acid




methyl amino methanoate

Note: In polyfunctional gp containing substance if C-containing func gp is not a substituent then its C is not considered in P.C.C. but in case of $-\text{CHO}$ & $-\overset{\text{O}}{\parallel}\text{C}-$ its C can be included in chain.





chloro formyl
3-methyl chloride - Propanoic acid3-amino - 2-hydroxy - 4-methyl
carbonyl - 3-butenoic acid


3-Bromo-3-Chloroformyl
Propanoic acid

5-Oxo-Pentanoic acid




5-Keto-Hexanamide

4-ethyl-methoxycarbonyl
5-keto Hex-2-ene methyl
Hex-2-ene oate~~5-Keto-3-Bromo Propyl
Hex-2-enoic acid~~
4-Bromo-3-methyl formyl hex-2-enoic
acidformyl methyl
4-Methyl formyl-5-hexenoyl
chloride4-Bromo-5-keto
Hexanal



3-amino-3-chloro-2-hydroxybutanoic acid
or
2-hydroxy-3-amino-3-chlorobutanoic acid




4-ethoxybutanoic acid

or

4-ethoxybutanoic acid


Complex Substituent :-



3-isopropylpent-4-enoic acid

or

3-(1-methylethyl)pent-4-enoic acid










bis-3,4-(1-chloroethyl)heptanoic acid



IUPAC name of Cyclic Compounds :-

Rule ① If an org. comp. consists of open chain & closed chain then that chain will be PCC which have -

Pfg > max^m no. of mb. > max^m no. of C-atom or longest chain > ring if open & closed chain have same no. of C-atom



Rule-2 If C-containing Pfg is directly attached to ring then it is consider as part of ring but its C is not included in P.C.C.




cyclo hexane
carbonitrile


cyclo hexane
carboxylic acid

Numbering as usual


- Note:- (1) ring prefix $\rightarrow$ cyclo
(2) word cyclo is considered in alphabetical order




3-cyclopentyl-2-hydroxy
Propanoic acid




3-amino-4-cyclopentyl
-Butan-2-ol




4 (2'-cyclobutenyl) - 3-hydroxy
- 2-butanone




3 (2'-butenyl) butanol




1-cyclobutyl But-1-ene



2-Butyl-cyclo Propene




1-Butyl-2-Chloro-cyclo Butane




1-Ethyl-4-Methyl cyclo hexane




4-Ethyl-1,1-dimethyl cyclohexane.




4-Ethyl-1,2-dimethyl cyclohexane




4-amino-3-chloro cyclo hexanol




2-Bromo-5-methyl cyclo Pentanol




3,5-dimethyl cyclo Penteneol




6-Ethyl-3-Methyl cyclo Hexeneamine




6-Bromo-5-Chloro-cyclo hex-3-eneol




6-bromo-3-chloro-cyclohex-1,4-diene-1-ol




5-mercapto-cyclohex-5-ene-1,3-diol




6-bromo-3-chloro-cyclohex-1,4-diene-1,5-diol




3-bromo-6-chloro-cyclohex-1,4-diene-1,4-diol




2-bromo-4-chloro-cyclopent-3-en-1-ol



4-amino-6-methyl-cyclohex-3-en-1-ol



1-carbonyl chloride-cyclohexane



Pt	mb	sb
1,4	1,4	3,3,6 ✓
1,4	1,4	3,6,6

PAGE NO.: 57
DATE: / /

4


CH₂CHO

2-cyclohexyl ethanal




COOH

3-amino-5-hydroxy-cyclohexane carboxylic acid




CHO

6-Bromo-cyclohex-3-ene-carbaldehyde




CN

3-amino-6-bromo-cyclohexene carbonitrile




COOH

3-ethoxy carbonyl-cyclohexane carboxylic acid



O

2-bromo-5-carbamoyl cyclohexane carbonyl halide



IUPAC Name of Aromatic




Benzene




toluene



Phenol




aniline




Benzoic acid




Benzaldehyde



Benzonitrile




anisole




Salicylic acid

2-hydroxy benzoic acid




Salicylaldehyde

2-hydroxy benzaldehyde




1, 2-dimethyl benzene
o-Xylene



m-Xylene



P-Xylene




1, 3, 5-trimethyl benzene
or
mesitylene


Rule: If aromatic compd is consist of open and closed chain then PCC will be

- ① which have P func gp.
- ② Open chain if it have more than 2 carbon

(in case when fun^c gp not +nt)




2-methyl Phenol
O-Cresol




2-ethyl - 4-methyl
benzoic acid




2-phenyl ethanoic acid




3-phenyl propanal



2-hydroxy methyl phenol
≡ (2-hydroxy phenyl)-methanol



Phenol



Alcoholic gp has more priority than phenolic gp.




methyl benzene



ethyl benzene




2-phenyl
propane




1-Chloro - 3 phenyl propane


If more than 2 fun^c gp are +nt




benzene-1,3-dicarboxylic acid



catechol
benzene-1,2-diol




2-ethanoyl oxy benzoic acid
2-acetoxy benzoic acid
acetyl salicylic acid




N-(4-hydroxy phenyl)
ethanamide


Paracetamol




IUPAC methyl-2-hydroxy Benzoate
common methyl salicylate
→ "oil of winter green"




2-Bromo-5-hydroxy phenyl cyano benzo nitrile




Benzene carbonyl chloride




6-Bromo-4-hydroxy
~~benzene~~ benzonitrile





2-Bromo-4-chloro-benzanol



3-amino-6-bromo-2-chloro
Benzo-nitrile.



2-Bromo-6-chloro-Benzaldehyde








Phosgene gas

poisonous gas


chloro-methyl anoyl chloride

CCl_4 tetrachloro methane
 $CHCl_3 \rightarrow$ chloroform / 1, 1, 1 - trichloro methane
 tear gas $\rightarrow CCl_3 \cdot NO_2$
 tri chloro nitro methane




Bicyclo Compounds

4 rings are fused




General formula $\rightarrow C_n H_{2n-2}$
 min C $\rightarrow 4$ C






Naming:


- Prefix $\rightarrow$ bicyclo
- no. of C-atoms in bridges are written in $\downarrow$ order in square bracket separated by full stop (.)
- word root $\rightarrow$ total no. of C-atom



bicyclo [4.2.0] Octane




bicyclo [2.2.1] heptane




bicyclo [2.2.2] Octane

If subs. is +nt then numbering start from bridged carbon & then we move from large ring to small ring



bicyclo [4.3.0] -7-Bromo
nonane




bicyclo [3.2.1]
6-bromo-
Oct-2-ene

Spiro - compounds :-

When 2 rings are fused at common carbon (4°C)

eg



$$G.F. \rightarrow C_n H_{2n-2}$$


min. no. of C $\rightarrow 5\text{C}$



Naming :-

• Prefix spiro

- no. of C atoms in ring are written in $\uparrow$ order
- word root, suffix.



Spiro [3.5] nonane




Spiro [2.2] Pentane




Spiro [2.4] heptane

* Sub is +nt


Numbering starts from adjacent carbon to common C & move through smaller ring to large ring



Spiro [3.4] - 6 - Bromo
Octane




Spiro [4.5] dec - 1, 6 -
diene



spiro [3.5] 2, 5 dione
nonane

Extra different org i.e. no common carbon



1-Chloro-4-Methyl benzene
4-Chloro-toluene