

## CHAPTER 2

### INVERSE TRIGONOMETRIC FUNCTIONS

#### VERY SHORT ANSWER TYPE QUESTIONS (1 MARK)

1. Write the principal value of

(i)  $\sin^{-1}(-\sqrt{3}/2)$

(ii)  $\cos^{-1}(\sqrt{3}/2)$ .

(iii)  $\tan^{-1}\left(-\frac{1}{\sqrt{3}}\right)$

(iv)  $\operatorname{cosec}^{-1}(-2)$ .

(v)  $\cot^{-1}\left(\frac{1}{\sqrt{3}}\right)$ .

(vi)  $\sec^{-1}(-2)$ .

(vii)  $\sin^{-1}\left(\frac{-\sqrt{3}}{2}\right) + \cos^{-1}\left(\frac{-1}{2}\right) + \tan^{-1}(-1/\sqrt{3})$

2. What is value of the following functions (using principal value).

(i)  $\tan^{-1}\left(\frac{1}{\sqrt{3}}\right) - \sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$ . (ii)  $\sin^{-1}\left(-\frac{1}{2}\right) - \cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$ .

(iii)  $\tan^{-1}(1) - \cot^{-1}(-1)$ . (iv)  $\operatorname{cosec}^{-1}(\sqrt{2}) + \sec^{-1}(\sqrt{2})$ .

(v)  $\tan^{-1}(1) + \cot^{-1}(1) + \sin^{-1}(1)$ .

(vi)  $\sin^{-1}\left(\sin \frac{4\pi}{5}\right)$ .

(vii)  $\tan^{-1}\left(\tan \frac{5\pi}{6}\right)$ .

(viii)  $\operatorname{cosec}^{-1}\left(\operatorname{cosec} \frac{3\pi}{4}\right)$ .

#### SHORT ANSWER TYPE QUESTIONS (4 MARKS)

3. Show that  $\tan^{-1}\left(\frac{\sqrt{1+\cos x} + \sqrt{1-\cos x}}{\sqrt{1+\cos x} - \sqrt{1-\cos x}}\right) = \frac{\pi}{4} + \frac{x}{2}$ .  $x \in [0, \pi]$

4. Prove

$$\tan^{-1}\left(\frac{\cos x}{1 - \sin x}\right) - \cot^{-1}\left(\sqrt{\frac{1 + \cos x}{1 - \cos x}}\right) = \frac{\pi}{4} \quad x \in (0, \pi/2).$$

5. Prove  $\tan^{-1}\left(\frac{x}{\sqrt{a^2 - x^2}}\right) = \sin^{-1}\frac{x}{a} = \cos^{-1}\left(\frac{\sqrt{a^2 - x^2}}{a}\right).$

6. Prove

$$\cot^{-1}\left[2 \tan\left(\cos^{-1}\frac{8}{17}\right)\right] + \tan^{-1}\left[2 \tan\left(\sin^{-1}\frac{8}{17}\right)\right] = \tan^{-1}\left(\frac{300}{161}\right).$$

7. Prove  $\tan^{-1}\left(\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}}\right) = \frac{\pi}{4} + \frac{1}{2}\cos^{-1}x^2.$

8. Solve  $\cot^{-1} 2x + \cot^{-1} 3x = \frac{\pi}{4}.$

9. Prove that  $\tan^{-1}\left(\frac{m}{n}\right) - \tan^{-1}\left(\frac{m-n}{m+n}\right) = \frac{\pi}{4}, m, n > 0$

10. Prove that  $\tan\left[\frac{1}{2}\sin^{-1}\left(\frac{2x}{1+x^2}\right) + \frac{1}{2}\cos^{-1}\left(\frac{1-y^2}{1+y^2}\right)\right] = \frac{x+y}{1-xy}$

11. Solve for  $x$ ,  $\cos^{-1}\left(\frac{x^2-1}{x^2+1}\right) + \frac{1}{2}\tan^{-1}\left(\frac{-2x}{1-x^2}\right) = \frac{2\pi}{3}$

12. Prove that  $\tan^{-1}\frac{1}{3} + \tan^{-1}\frac{1}{5} + \tan^{-1}\frac{1}{7} + \tan^{-1}\frac{1}{8} = \frac{\pi}{4}$

13. Solve for  $x$ ,  $\tan(\cos^{-1}x) = \sin(\tan^{-1}2); x > 0$

14. Prove that  $2\tan^{-1}\left(\frac{1}{5}\right) + \tan^{-1}\left(\frac{1}{4}\right) = \tan^{-1}\left(\frac{32}{43}\right)$

15. Evaluate  $\tan\left[\frac{1}{2}\cos^{-1}\left(\frac{3}{\sqrt{11}}\right)\right]$
16. Prove that  $\tan^{-1}\left(\frac{a\cos x - b\sin x}{b\cos x + a\sin x}\right) = \tan^{-1}\left(\frac{a}{b}\right) - x$
17. Prove that  $\cot\left\{\tan^{-1}x + \tan^{-1}\left(\frac{1}{x}\right)\right\} + \cos^{-1}(1 - 2x^2) + \cos^{-1}(2x^2 - 1) = \pi, \quad x > 0$
18. Prove that  $\tan^{-1}\left(\frac{a-b}{1+ab}\right) + \tan^{-1}\left(\frac{b-c}{1+bc}\right) + \tan^{-1}\left(\frac{c-a}{1+ca}\right) = 0$  where  $a, b, c > 0$
19. Solve for  $x$ ,  $2 \tan^{-1}(\cos x) = \tan^{-1}(2 \operatorname{cosec} x)$
20. Express  $\sin^{-1}(x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2})$  in simplest form.
21. If  $\tan^{-1}a + \tan^{-1}b + \tan^{-1}c = \pi$ , then prove that  $a + b + c = abc$
22. If  $\sin^{-1}x > \cos^{-1}x$ , then  $x$  belongs to which interval?

## ANSWERS

1. (i)  $-\frac{\pi}{3}$  (ii)  $\frac{\pi}{6}$  (iii)  $\frac{-\pi}{6}$  (iv)  $\frac{-\pi}{6}$   
 (v)  $\frac{\pi}{3}$  (vi)  $\frac{2\pi}{3}$  (vii)  $\frac{\pi}{6}$ .
2. (i) 0 (ii)  $\frac{-\pi}{3}$  (iii)  $-\frac{\pi}{2}$  (iv)  $\frac{\pi}{2}$   
 (v)  $\pi$  (vi)  $\frac{\pi}{5}$  (vii)  $\frac{-\pi}{6}$  (viii)  $\frac{\pi}{4}$ .

8. 1

11.  $\tan \frac{\pi}{12} = 2 - \sqrt{3}$

13.  $\frac{\sqrt{5}}{3}$

15.  $\sqrt{\frac{\sqrt{11}-3}{3+\sqrt{11}}}$

19.  $x = \frac{\pi}{4}.$

20  $\sin^{-1} x - \sin^{-1} \sqrt{x}.$

22.  $\left( \frac{1}{\sqrt{2}}, 1 \right]$

21. **Hint:** Let  $\tan^{-1} a = \alpha$   
 $\tan^{-1} b = \beta$   
 $\tan^{-1} c = \gamma$

then given,  $\alpha + \beta + \gamma = \pi$

$\therefore \alpha + \beta = \pi - \gamma$

take tangent on both sides,

$\tan (\alpha + \beta) = \tan (\pi - \gamma)$