

**Topic : Mathematical Tools**

**Type of Questions**

Single choice Objective ('-1' negative marking) Q.1 to Q.4

(3 marks, 3 min.)

**M.M., Min.**

**[12, 12]**

Subjective Questions ('-1' negative marking) Q.5 to Q. 6

(4 marks, 5 min.)

**[8, 10]**

Comprehension ('-1' negative marking) Q.7 to Q.9

(3 marks, 3 min.)

**[9, 9]**

1. If  $y = \sqrt{\sin \sqrt{x}}$ , then  $\frac{dy}{dx}$  is :

- (A)  $\frac{1}{4\sqrt{x}} \cdot \frac{\cos \sqrt{x}}{\sin \sqrt{x}}$  (B)  $\frac{1}{4\sqrt{x}} \cdot \sqrt{\tan \sqrt{x}} \sqrt{\cos \sqrt{x}}$  (C)  $\frac{1}{4\sqrt{x}} \sqrt{\frac{\cos \sqrt{x}}{\sin \sqrt{x}}}$  (D)  $\frac{1}{4\sqrt{x}} \sqrt{\cot \sqrt{x}} \cdot \sqrt{\cos \sqrt{x}}$

2. A particle moves along a straight line such that its displacement at any time  $t$  is given by :  
 $s = t^3 - 6t^2 + 3t + 4$  metres. The velocity when the acceleration is zero is :

- (A)  $3 \text{ ms}^{-1}$  (B)  $-12 \text{ ms}^{-1}$  (C)  $42 \text{ ms}^{-1}$  (D)  $-9 \text{ ms}^{-1}$

3. The area of region between  $y = \sin x$  and  $x$ -axis in the interval  $\left[0, \frac{\pi}{2}\right]$  will be :

- (A) 1 (B) 0 (C) 2 (D)  $\frac{1}{2}$

4. The value of  $\int_0^{\pi/2} \sin^2 x \, dx$  will be :

- (A) 1 (B) 0 (C)  $\frac{\pi}{4}$  (D)  $\frac{\pi}{2}$

**Evaluate : -**

5.  $\int_0^1 (3x^2 + 4) \, dx$

6.  $\int_0^{\pi/2} (\sin x + \cos x) \, dx$

**COMPREHENSION**

If  $a = (3t^2 + 2t + 1) \text{ m/s}^2$  is the expression according to which the acceleration of a particle varies moving along a straight line. Then -

7. The expression for instantaneous velocity at any time ' $t$ ' will be (if the particle was initially at rest) -  
(A)  $t^3 + 2t + 1$  (B)  $t^3 + t + 1$  (C)  $t^3 + t^2 + t$  (D)  $t^3 + t^2 + t + C$
8. The change in velocity after 3 seconds of its start is :  
(A)  $30 \text{ m/s}$  (B)  $39 \text{ m/s}$  (C)  $3 \text{ m/s}$  (D)  $20 \text{ m/s}$
9. Find displacement of the particle after 2 seconds of start -  
(A)  $26 \text{ m}$  (B)  $26/3 \text{ m}$  (C)  $30/7 \text{ m}$  (D)  $26/7 \text{ m}$

# Answers Key

## DPP NO. - 7

- |        |        |        |        |      |
|--------|--------|--------|--------|------|
| 1. (D) | 2. (D) | 3. (A) | 4. (C) | 5. 5 |
| 6. 2   | 7. (C) | 8. (B) | 9. (B) |      |

# Hint & Solutions

## DPP NO. - 7

$$1. \frac{d}{dx} \left[ (\sin \sqrt{x})^{1/2} \right] = \frac{1}{2} (\sin \sqrt{x})^{-1/2} \cdot [\cos \sqrt{x}] \cdot \frac{1}{2} (x)^{-1/2} \quad (\text{By power chain rule})$$

$$= \frac{1}{4\sqrt{x}} \cdot \frac{\cos \sqrt{x}}{\sqrt{\sin \sqrt{x}}} = \frac{1}{4\sqrt{x}} \cdot \sqrt{\cot \sqrt{x}} \cdot \sqrt{\cos \sqrt{x}}$$

$$2. v = \frac{ds}{dt} = 3t^2 - 12t + 3, \quad a = \frac{dv}{dt} = 6t - 12 = 0$$

$$\Rightarrow t = 2s$$

$$v_{t=2} = 3 \times 4 - 12 \times 2 + 3 = -9 \text{ m/s}$$

$$3. \int_0^{\pi/2} \sin x dx = [-\cos x]_0^{\pi/2} = 1.$$

$$4. \int_0^{\pi/2} \sin^2 x dx = \left[ \frac{x}{2} - \frac{\sin 2x}{4} + c \right]_0^{\pi/2} = \frac{\pi}{4}.$$

**Evaluate :**

$$5. \int_0^1 (3x^2 + 4) dx = \left[ x^3 \right]_0^1 + 4 \left[ x \right]_0^1 = 1 + 4 = 5$$

$$6. \int_0^{\pi/2} (\sin x + \cos x) dx = [-\cos x]_0^{\pi/2} + [\sin x]_0^{\pi/2} \\ = 1 - 0 + 1 - 0 = 2$$

7.  $a = 3t^2 + 2t + 1$

$$\int_0^v dv = \int_0^t (3t^2 + 2t + 1) dt \quad v = t^3 + t^2 + t$$

8.  $V(t = 0) = 0$

$$\begin{aligned} V_{t=3} &= (3)^3 + (3)^2 + 3 \\ &= 27 + 9 + 3 \\ &= 39 \end{aligned}$$

$$\Delta V = 39 - 0 = 39 \text{ m/s.}$$

9.  $\int_0^s dS = \int_0^2 (t^3 + t^2 + t) dt \quad S = \left[ \frac{t^4}{4} + \frac{t^3}{3} + \frac{t^2}{2} \right]_0^2 \quad S$

$$= 4 + \frac{8}{3} + 2S = \frac{12 + 8 + 6}{3} = \frac{26}{3}$$