


Chapter - 2

Polynomial

Exercise – 2.1

Q. 1 The graphs of $y = p(x)$ are given in Fig. 2.10 below, for some polynomials, $p(x)$. Find the number of zeroes of $p(x)$ in each case.



The number of zeroes for any graph is the number of values of x for which y is equal to zero. And y is equal to zero at the point where a graph cuts x axis.

So, to find the number of zeroes of a polynomial, watch the number of times it cuts the x axis.

(i) The number of zeroes is 0 as the graph does not cut the x -axis at any point.

(ii) The number of zeroes is 1 as the graph intersects the x -axis at only 1 point.

(iii) The number of zeroes is 3 as the graph intersects the x -axis at 3 points.

(iv) The number of zeroes is 2 as the graph intersects the x-axis at 2 points.

(v) The number of zeroes is 4 as the graph intersects the x-axis at 4 points.

(vi) The number of zeroes is 3 as the graph intersects the x-axis at 3 points.

Exercise – 2. 2

Q. 1 Find the zeroes of the following quadratic polynomials and verify the relationship between the zeroes and the coefficients.

(i) $x^2 - 2x - 8$

(ii) $4s^2 - 4s + 1$

(iii) $6x^2 - 3 - 7x$

(iv) $4u^2 + 8u$

(v) $t^2 - 15$

(vi) $3x^2 - x - 4$

Solution:

Zeroes of the polynomial are the values of the variable of the polynomial when the polynomial is put equal to zero.

Let $p(x)$ be a polynomial with any number of terms any number of degree. Now, zeroes of the polynomial will be the values of x at which $p(x) = 0$. If $p(x) = ax^2 + bx + c$ is a quadratic polynomial (highest power is equal to 2) and its roots are α and β , then

$$\text{Sum of the roots} = \alpha + \beta = -b/a$$

$$\text{Product of roots} = \alpha\beta = c/a$$

(i) $p(x) = x^2 - 2x - 8$

So, the zeroes will be the values of x at which $p(x) = 0$.

Therefore,

$$\Rightarrow x^2 - 4x + 2x - 8 = 0$$

(We will factorize 2 such that the product of the factors is equal to 8 and difference is equal to 2)

$$\Rightarrow x(x - 4) + 2(x - 4) = 0 = (x - 4)(x + 2)$$

The value of $x^2 - 2x - 8$ is zero when $x - 4 = 0$ or $x + 2 = 0$,

i.e, $x = 4$ or $x = -2$

Therefore, The zeroes of $x^2 - 2x - 8$ are 4 and -2 .

$$\text{Sum of zeroes} = 4 + (-2) = 2$$

$$= \frac{-(-2)}{1} = \frac{-(-\text{coefficient of } x)}{(\text{coefficient of } x^2)}$$

$$\text{Hence, it is verified that, sum of Zeros} = \frac{-(\text{coefficient of } x)}{\text{coefficient of } x^2}$$

$$\text{Product of zeroes} = 4 + (-2) = -8 = \frac{(-8)}{1} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

$$\text{Hence, it is verified, Product of zeroes} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

$$(ii) 4s^2 - 4s + 1$$

$$= (2s)^2 - 2(2s)1 + 1^2$$

As, we know $(a - b)^2 = a^2 - 2ab + b^2$, the above equation can be written as $= (2s - 1)^2$

The value of $4s^2 - 4s + 1$ is zero when $2s - 1 = 0$, when, $s = 1/2, 1/2$.

Therefore, the zeroes of $4s^2 - 4s + 1$ are $\frac{1}{2}$ and $\frac{1}{2}$.

$$\text{Sum of zeroes} = \frac{1}{2} + \frac{1}{2} = 1 = \frac{-(-4)}{4} = \frac{-(\text{coefficient of } s)}{\text{coefficient of } s^2}$$

Product of zeroes =

$$\frac{1}{2} \times \frac{1}{2} = \frac{1}{4} = \frac{\text{constant term}}{\text{coefficient of } s^2}$$

Hence Verified.

(iii) $6x^2 - 3 - 7x$

$$= 6x^2 - 7x - 3$$

(We will factorize 7 such that the product of the factors is equal to 18 and the difference is equal to - 7)
 $= 6x^2 + 2x - 9x - 3$
 $= 2x(3x + 1) - 3(3x + 1) = (3x + 1)(2x - 3)$

The value of $6x^2 - 3 - 7x$ is zero when $3x + 1 = 0$ or $2x - 3 = 0$,

i.e. $x = \frac{-1}{3}$ or $\frac{3}{2}$

Therefore, the zeroes of $6x^2 - 3 - 7x$ are $\frac{-1}{3}$ or $\frac{3}{2}$.

$$\text{Sum of zeroes} = \frac{-1}{3} + \frac{3}{2} = \frac{7}{6} = \frac{-(-7)}{6} = \frac{-(\text{coefficient of } x)}{\text{coefficient of } x^2}$$

$$\text{Product of zeroes} = \frac{-1}{3} \times \frac{3}{2} = \frac{-1}{2} = \frac{-3}{6} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Hence, verified.

(iv) $4u^2 + 8u$

$$= 4u^2 + 8u + 0$$

$$= 4u(u + 2)$$

The value of $4u^2 + 8u$ is zero when $4u = 0$ or $u + 2 = 0$,

i.e., $u = 0$ or $u = -2$

Therefore, the zeroes of $4u^2 + 8u$ are 0 and -2.

$$\text{Sum of zeroes} = 0 + (-2) = -2 = \frac{-(8)}{4} = \frac{-(\text{coefficient of } u)}{\text{coefficient of } u^2}$$

$$\text{Product of zeroes} = 0 + (-2) = 0 = \frac{0}{4} = \frac{-\text{constant term}}{\text{coefficient of } u^2}$$

$$(v) t^2 - 15$$

$$= t^2 - (\sqrt{15})^2$$

$$= (t - \sqrt{15})(t + \sqrt{15}) \quad [\text{As, } x^2 - y^2 = (x - y)(x + y)]$$

The value of $t^2 - 15$ is zero when $(t - \sqrt{15}) = 0$ or $(t + \sqrt{15}) = 0$,
i.e., when $t = \sqrt{15}$ or $t = -\sqrt{15}$

Therefore, the zeroes of $t^2 - 15$ are $\sqrt{15}$ and $-\sqrt{15}$.

$$\text{Sum of zeroes} = \sqrt{15} + (-\sqrt{15}) = 0 = \frac{-0}{1} = \frac{-(\text{coefficient of } t)}{\text{coefficient of } t^2}$$

$$\sqrt{15} + (-\sqrt{15}) = 0 = \frac{-0}{1} = \frac{-(\text{coefficient of } t)}{\text{coefficient of } t^2}$$

Product of zeroes =

$$(\sqrt{15})(-\sqrt{15}) = -15 = \frac{-15}{1} = \frac{-\text{constant term}}{\text{coefficient of } x^2}$$

Hence verified.

$$(vi) 3x^2 - x - 4$$

(We will factorize 1 in such a way that the product of factors is equal to 12 and the difference is equal to 1) $= 3x^2 - 4x + 3x - 4$
 $= x(3x - 4) + 1(3x - 4) = (3x - 4)(x + 1)$

The value of $3x^2 - x - 4$ is zero when $3x - 4 = 0$ or $x + 1 = 0$,

$$\text{when } x = \frac{4}{3} \text{ or } x = -1$$

Therefore, the zeroes of $3x^2 - x - 4$ are $\frac{4}{3}$ and -1

$$\text{Sum of zeroes} = \frac{4}{3} + (-1) = \frac{1}{3} = \frac{-(-1)}{3} = \frac{-\text{coefficient of } x}{\text{coefficient of } x^2}$$

$$\text{Product of zeroes} = \frac{4}{3}(-1) = \frac{-4}{3} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

Hence, verified.

Q. 2 Find a quadratic polynomial each with the given numbers as the sum and product of its zeroes respectively.

(i) $\frac{1}{4}, -1$

(ii) $\sqrt{2}, \frac{1}{3}$

(iii) $0, \sqrt{5}$

(iv) $1, 1$

(v) $-\frac{1}{4}, \frac{1}{4}$

(vi) $4, 1$

Solution: If α, β are roots of an equation, then the quadratic form of this equation can be given by $x^2 - (\alpha + \beta)x + \alpha\beta = 0$

(i) $\frac{1}{4}, -1$

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be, $ax^2 + bx + c$, then

$$\alpha + \beta = \frac{1}{4} = \frac{-b}{a}$$

$$\alpha\beta = -1 = \frac{-4}{4} = \frac{c}{a}$$

Let $a = 4$, then $b = -1$, $c = -4$

Therefore, the quadratic polynomial is $4x^2 - x - 4$.

(ii) $\sqrt{2}, \frac{1}{3}$

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be, $ax^2 + bx + c$, then

$$\alpha + \beta = \sqrt{2} = \frac{-b}{a}$$

and

If $a = 3$, then $b = -3\sqrt{2}$, and $c = 1$

Therefore, the quadratic polynomial is $3x^2 - 3\sqrt{2}x + 1$

(iii) $0, \sqrt{5}$

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be $ax^2 + bx + c$, then

$$\alpha + \beta = 0 = \frac{0}{1} = \frac{-b}{a}$$

$$\alpha\beta = \sqrt{5} = \frac{\sqrt{5}}{1} = \frac{c}{a}$$

If $a = 1$, then $b = 0$, $c = \sqrt{5}$

Therefore, the quadratic polynomial is $x^2 + \sqrt{5}$.

(iv) 1, 1

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be $ax^2 + bx + c$, then

$$\alpha + \beta = 1 = \frac{1}{1} = \frac{-b}{a}$$

$$\alpha\beta = 1 = \frac{1}{1} = \frac{c}{a}$$

If $a = 1$, then $b = -1$, $c = 1$

Therefore, the quadratic polynomial is $x^2 - x + 1$

(v) $-\frac{1}{4}, \frac{1}{4}$

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be $ax^2 + bx + c$, then

$$\alpha + \beta = 1 = \frac{-1}{4} = \frac{-b}{a}$$

$$\alpha\beta = \frac{1}{4} = \frac{c}{a}$$

If $a = 4$, then $b = 1$, $c = 1$

Therefore, the quadratic polynomial is $4x^2 + x + 1$.

(vi) 4, 1

we know that for a quadratic equation in the form $ax^2 + bx + c = 0$, and its zeros are α and β , then

sum of zeroes is $\alpha + \beta = \frac{-b}{a}$

and product of zeroes is $\alpha\beta = \frac{c}{a}$

Let the polynomial be $ax^2 + bx + c$, then

$$\alpha + \beta = 4 = \frac{4}{1} = \frac{-b}{a}$$

$$\alpha\beta = 1 = \frac{1}{1} = \frac{c}{a}$$

If $a = 1$, then $b = -4$, $c = 1$

Therefore, the quadratic polynomial is $x^2 - 4x + 1$.

Exercise 2.3

Divide the polynomial by the polynomial and find the quotient and remainder in each of the following:

(i) $p(x) = x^3 - 3x^2 + 5x - 3$, $g(x) = x^2 - 2$

(ii) $p(x) = x^4 - 3x^2 + 4x + 5$, $g(x) = x^2 + 1 - x$

(iii) $p(x) = x^4 - 5x + 6$, $g(x) = 2 - x^2$

(i) By long division method we have,

$$\begin{array}{r} x-3 \\ x^2 - 2 \overline{) x^3 - 3x^2 + 5x - 3} \\ \underline{x^3 - 2x} \\ - + \\ \underline{-3x^2 + 7x - 3} \\ -3x^2 + 6 \\ \underline{-3x^2 + 6} \\ 7x - 9 \end{array}$$

Quotient = $x - 3$

Remainder = $7x - 9$

(ii) By long division method we have,

$$\begin{array}{r}
 x^2 + x - 3 \\
 x^2 - x + 1 \sqrt{x^4 + 0x^3 - 3x^2 + 4x + 5} \\
 \underline{x^4 - x^3 + x^2} \\
 - + - \\
 x^3 - 4x^2 + 4x + 5 \\
 \underline{x^3 - x^2 + x} \\
 - + - \\
 -3x^2 + 3x + 5 \\
 -3x^2 + 3x - 3 \\
 + - + \\
 \hline
 8 \\
 \hline
 \end{array}$$

Quotient = $x^2 + x - 3$

Remainder = 8

(iii) By long division method we have,

$$\begin{array}{r}
 -x^2 - 2 \\
 -x^2 + 2 \sqrt{x^4 + 0x^2 - 5x + 6} \\
 \underline{x^4 - 2x^2} \\
 - + \\
 2x^2 - 5x + 6 \\
 2x^2 - 4 \\
 - + \\
 \hline
 -5x + 10 \\
 \hline
 \end{array}$$

Quotient = $-x^2 - 2$

Remainder = $-5x + 10$

Q. 2 Check whether the first polynomial is a factor of the second polynomial by dividing the second polynomial by the first polynomial:

(i) $t^2 - 3$, $2t^4 + 3t^3 - 2t^2 - 9t - 12$

(ii) $x^2 + 3x + 1$, $3x^4 + 5x^3 - 7x^2 + 2x + 2$

(iii) $x^3 - 3x + 1$, $x^5 - 4x^3 + x^2 + 3x + 1$

(i) $t^2 - 3 = t^2 + 0t - 3$

$$\begin{array}{r}
 2t^2 + 3t + 4 \\
 t^2 + 0t - 3 \sqrt{2t^4 + 3t^3 - 2t^2 - 9t - 12} \\
 \underline{2t^4 + 0t^3 - 6t^2} \\
 - - + \\
 3t^3 + 4t^2 - 9t - 12 \\
 \underline{3t^3 + 0t^2 - 9t} \\
 - - + \\
 4t^2 + 0t - 12 \\
 \underline{4t^2 + 0t - 12} \\
 - - + \\
 0
 \end{array}$$

Since the remainder is 0,

Hence, $t^2 - 3$ is a factor of $2t^4 + 3t^3 - 2t^2 - 9t - 12$.

(ii)

$$\begin{array}{r} 3x^2 - 4x + 2 \\ x^2+3x+1 \overline{\sqrt{3x^4 + 5x^3 - 7x^2 + 2x + 2}} \\ 3x^4 + 9x^3 + 3x^2 \\ \underline{- \quad - \quad -} \\ -4x^3 - 10x^2 + 2x + 2 \\ -4x^3 - 12x^2 - 4x \\ \underline{+ \quad + \quad +} \\ 2x^2 + 6x + 2 \\ 2x^2 + 6x + 2 \\ \underline{ 0} \\ \underline{ 0} \end{array}$$

Since the remainder is 0,

Hence, x^2+3x+1 is a factor of $3x^4+5x^3-7x^2+2x+2$.

(iii)

$$\begin{array}{r} x^2 - 1 \\ x^2 - 3x + 1 \overline{\sqrt{x^5 - 4x^3 + x^2 + 3x + 1}} \\ x^5 - 3x^2 + x^2 \\ \underline{- \quad + \quad -} \\ -x^3 + 3x + 1 \\ -x^3 + 3x - 1 \\ \underline{+ \quad - \quad +} \\ 2 \end{array}$$

Since the remainder $\neq 0$,

Hence, x^3-3x+1 is not a factor of $x^5-4x^3+ x^2+3x+1$.

Q. 3 Obtain all other zeroes of $3x^4 + 6x^3 - 2x^2 - 10x - 5$, if two of its zeroes are $\sqrt{\frac{5}{3}}$ and $-\sqrt{\frac{5}{3}}$.

$$p(x) = 3x^4 + 6x^3 - 2x^2 - 10x - 5$$

Since the two zeroes are $\sqrt{\frac{5}{3}}$ and $-\sqrt{\frac{5}{3}}$

$$\therefore \left(x - \sqrt{\frac{5}{3}}\right)\left(x + \sqrt{\frac{5}{3}}\right) = \left(x^2 - \frac{5}{3}\right) \text{ is a factor of } 3x^4 + 6x^3 - 2x^2 - 10x - 5.$$

Therefore, we divide the given polynomial by $x^2 - \frac{5}{3}$.

$$\begin{array}{r}
 3x^2 + 6x + 3 \\
 x^2 + 0x - \frac{5}{3} \quad \sqrt{3x^4 + 6x^3 - 2x^2 - 10x - 5} \\
 \underline{3x^4 + 0x^3 - 5x^2} \\
 - - + \\
 \underline{6x^3 + 3x^2 - 10x - 5} \\
 6x^3 + 0x^2 - 10x \\
 - - + \\
 \underline{3x^2 + 0x - 5} \\
 3x^2 + 0x - 5 \\
 - - + \\
 \underline{0}
 \end{array}$$

We know, $\text{Dividend} = (\text{Divisor} \times \text{quotient}) + \text{remainder}$

$$3x^4 + 6x^3 - 2x^2 - 10x - 5 = \left(x^2 - \frac{5}{3}\right)(3x^2 + 6x + 3)$$

$$3x^4 + 6x^3 - 2x^2 - 10x - 5 = \left(x^2 - \frac{5}{3}\right)(x^2 + 2x + 1)$$

$$\text{As } (a+b)^2 = a^2 + b^2 + 2ab$$

$$\text{So, } x^2 + 2x + 1 = (x+1)^2$$

$$3x^4 + 6x^3 - 2x^2 - 10x - 5 = 3\left(x^2 - \frac{5}{3}\right)(x+1)^2$$

Therefore, its zero is given by $x + 1 = 0$.

$$\Rightarrow x = -1, -1$$

Hence, the zeroes of the given polynomial are $\sqrt{\frac{5}{3}}$ and $-\sqrt{\frac{5}{3}}$ and $-1, -1$.

Q. 4 On dividing $x^3 - 3x^2 + x + 2$ by a polynomial $g(x)$ the quotient and remainder were $(x - 2)$ and $(-2x + 4)$, respectively. Find $g(x)$.

Solution: Given,

Polynomial, $p(x) = x^3 - 3x^2 + x + 2$ (dividend)

$$\text{Quotient} = (x - 2)$$

$$\text{Remainder} = (-2x + 4)$$

To find: divisor = $g(x)$ we know, Dividend = Divisor \times Quotient + Remainder

$$\Rightarrow x^3 - 3x^2 + x + 2 = g(x) \times (x - 2) + (-2x + 4)$$

$$\Rightarrow x^3 - 3x^2 + x + 2 + 2x - 4 = g(x)(x - 2)$$

$$\Rightarrow x^3 - 3x^2 + 3x - 2 = g(x)(x - 2)$$

$g(x)$ is the quotient when we divide $(x^3 - 3x^2 + 3x - 2)$ by $(x - 2)$

$$\begin{array}{r}
 x^2 - x + 1 \\
 x - 2 \overline{) x^3 - 3x^2 + 3x - 2} \\
 \underline{x^3 - 2x^2} \\
 -x^2 + 3x - 2 \\
 \underline{-x^2 + 2x} \\
 x - 2 \\
 \underline{x - 2} \\
 0
 \end{array}$$

$\therefore g(x) = (x^2 - x + 1)$

Q. 5 Give examples of polynomials and which satisfy the division algorithm and

(i) $\deg p(x) = \deg q(x)$

(ii) $\deg q(x) = \deg r(x)$

(iii) $\deg r(x) = 0$

Degree of a polynomial is the highest power of the variable in the polynomial. For example if $f(x) = x^3 - 2x^2 + 1$, then the degree of this polynomial will be 3.

(i) By division Algorithm : $p(x) = g(x) \times q(x) + r(x)$

It means when $P(x)$ is divided by $g(x)$ then quotient is $q(x)$ and remainder is $r(x)$. We need to start with $p(x) = q(x)$. This means that the degree of polynomial $p(x)$ and quotient $q(x)$ is same. This can only happen if the degree of $g(x) = 0$ i.e $p(x)$ is divided by a constant. Let $p(x) = x^2 + 1$ and $g(x) = 2$

$$\frac{p(x)}{g(x)} = \frac{x^2 + 1}{2}$$

The,

$$p(x) = g(x) \times \left(\frac{x^2+1}{2}\right)$$

Clearly, Degree of $p(x)$ = Degree of $q(x)$

2. Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x)$$

$$= 6x^2 + 2x + 2 = 3(3x^2 + x + 1)$$

$$= 6x^2 + 2x + 2$$

Thus, the division algorithm is satisfied.

(ii) Let us assume the division of $x^3 + x$ by x^2 ,

Here,

$$p(x) = x^3 + x$$

$$g(x) = x^2$$

$$q(x) = x \text{ and } r(x) = x$$

Clearly, the degree of $q(x)$ and $r(x)$ is the same i.e.,

Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x) \quad x^3 + x$$

$$= (x^2) \times x + x \quad x^3 + x = x^3 + x$$

Thus, the division algorithm is satisfied.

(iii) Degree of the remainder will be 0 when the remainder comes to a constant.

Let us assume the division of $x^3 + 1$ by x^2 .

Here,

$$p(x) = x^3 + 1 \quad g(x) = x^2$$

$$q(x) = x \text{ and } r(x) = 1$$

Clearly, the degree of $r(x)$ is 0. Checking for division algorithm,

$$p(x) = g(x) \times q(x) + r(x)x^3 + 1$$

$$= (x^2) \times x + 1 \quad x^3 + 1 = x^3 + 1$$

Thus, the division algorithm is satisfied.

Exercise 2.4

Q. 1 Verify that the numbers given alongside of the cubic polynomials below are their zeroes. Also verify the relationship between the zeroes and the coefficients in each case:

(i) $2x^3 + x^2 - 5x + 2; \frac{1}{2}, 1, -2$

(ii) $x^3 - 4x^2 + 5x - 2; 2, 1, 1$

Answer:

(i) $P(x) = 2x^3 + x^2 - 5x + 2$

Now for zeroes, putting the given values in x.

$$\begin{aligned} P(1/2) &= 2(1/2)^3 + (1/2)^2 - 5(1/2) + 2 \\ &= (1/4) + (1/4) - (5/2) + 2 = (1 + 1 - 10 + 8)/2 = 0/2 = 0 \end{aligned}$$

$$P(1) = 2 \times 1 + 1 - 5 \times 1 + 2 = 2 + 1 - 5 + 2 = 0$$

$$\begin{aligned} P(-2) &= 2 \times (-2)^3 + (-2)^2 - 5(-2) + 2 = (2 \times -8) + 4 + 10 + 2 = -16 + 16 \\ &= 0 \end{aligned}$$

Thus, $1/2$, 1 and -2 are zeroes of given polynomial.

Comparing given polynomial with $ax^3 + bx^2 + cx + d$ and Taking zeroes as α , β , and γ , we have

$$a = 2, b = 1, c = -5, d = 2 \text{ and } \alpha = \frac{1}{2}, \beta = 1, \gamma = -2$$

Now, We know the relation between zeroes and the coefficient of a standard cubic polynomial as

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

Substituting value, we have

$$\frac{1}{2} + 1 - 2 = \frac{1}{2}$$

$$\frac{1}{2} = \frac{1}{2}$$

Since, LHS = RHS (Relation Verified)

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$$

$$\left(\frac{1}{2} \times 1\right) + (1 \times -2) + \left(-2 \times \frac{1}{2}\right) = -\frac{5}{2}$$

$$\frac{1}{2} - 2 - 1 = -\frac{5}{2}$$

$$-\frac{5}{2} = -\frac{5}{2}$$

Since LHS = RHS, Relation verified.

$$\alpha\beta\gamma = -\frac{d}{a}$$

$$\left(\frac{1}{2} \times 1 \times -2\right) = -\frac{2}{2}$$

$$-\frac{2}{2} = -\frac{2}{2}$$

Since LHS = RHS, Relation verified.

Thus, all three relationships between zeroes and the coefficient is verified.

$$(ii) p(x) = x^3 - 4x^2 + 5x - 2$$

Now for zeroes, put the given value in x.

$$P(2) = 2^3 - 4(2)^2 + 5 \times 2 - 2 = 8 - 16 + 10 - 2 = 18 - 18 = 0$$

$$P(1) = 1^3 - 4(1)^2 + 5 \times 1 - 2 = 1 - 4 + 5 - 2 = 6 - 6 = 0$$

$$P(1) = 1^3 - 4(1)^2 + 5 \times 1 - 2 = 1 - 4 + 5 - 2 = 6 - 6 = 0$$

Thus, 2, 1, 1 are the zeroes of the given polynomial.

Now,

Comparing the given polynomial with $ax^3 + bx^2 + cx + d$, we get

$$a = 1, b = -4, c = 5, d = -2 \text{ and } \alpha = 2, \beta = 1, \gamma = 1$$

Now,

$$2 + 1 + 1 = \frac{4}{1}$$

$$4 = 4$$

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$$

$$(2 \times 1) + (1 \times 1) + (1 \times 2) = \frac{5}{1}$$

$$2 + 1 + 2 = 5$$

$$5 = 5$$

$$\alpha\beta\gamma = -\frac{d}{a}$$

$$2 \times 1 \times 1 = 2$$

$$2 = 2$$

Thus, all three relationships between zeroes and the coefficient is verified.

Q. 2 Find a cubic polynomial with the sum, sum of the product of its zeroes taken two at a time, and the product of its zeroes as 2, -7, -14 respectively.

Answer:

For a cubic polynomial equation, $ax^3 + bx^2 + cx + d$, and zeroes α , β and γ

we know that

$$\alpha + \beta + \gamma = \frac{-b}{a}$$

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$$

$$\alpha\beta\gamma = \frac{-d}{a}$$

Let the polynomial be $ax^3 + bx^2 + cx + d$, and zeroes α , β and γ .

A cubic polynomial with respect to its zeroes is given by, $x^3 - (\text{sum of zeroes}) x^2 + (\text{Sum of the product of roots taken two at a time}) x - \text{Product of Roots} = 0$

$$x^3 - (2) x^2 + (-7) x - (-14) = 0$$

$$x^3 - (2) x^2 + (-7) x + 14 = 0$$

Hence, the polynomial is $x^3 - 2x^2 - 7x + 14$.

Q. 3 If the zeroes of the polynomial $x^3 - 3x^2 + x + 1$ are $(a - b)$, a and $(a + b)$. Find a and b .

Answer:

Given

$$P(x) = x^3 - 3x^2 + x + 1$$

Zeroes are $= a - b$, $a + b$, a

Comparing the given polynomial with $mx^3 + nx^2 + px + q$, we get,

$$m = 1, n = -3, p = 1, q = 1$$

$$\text{Sum of zeroes} = a - b + a + a + b = -\frac{n}{m}$$

$$3a = -\frac{-3}{1} = 3$$

$$a = \frac{3}{3} = 1$$

The zeroes are $(1 - b)$, 1 and $(1 + b)$

$$\text{Product of zeroes} = (1 - b)(1 + b)$$

$$(1 - b)(1 + b) = -q/m$$

$$1 - b^2 = -\frac{1}{1} = -1$$

$$b^2 = 2$$

$$b = \pm\sqrt{2}$$

So,

We get, $a = 1$ and $b = \pm\sqrt{2}$

Q. 4 If two zeroes of the polynomial $x^4 - 6x^3 - 26x^2 + 138x - 35$ are $2 \pm \sqrt{3}$ find other zeroes.

Answer:

Given:

$2 + \sqrt{3}$ and $2 - \sqrt{3}$ are zeroes of given equation,

Therefore, $(x - 2 + \sqrt{3})(x - 2 - \sqrt{3})$ should be a factor of given equation.

$$\text{Also, } (x - 2 + \sqrt{3})(x - 2 - \sqrt{3}) = x^2 - 2x - \sqrt{3}x - 2x + 4 + 2\sqrt{3} + \sqrt{3}x - 2\sqrt{3} - 3$$

$$= x^2 - 4x + 1$$

To find other zeroes, we divide given equation by $x^2 - 4x + 1$

$$x^2 - 2x - 35$$

$$\begin{array}{r}
 x^2 - 4x + 1 \sqrt{x^4 - 6x^3 - 26x^2 + 138x - 35} \\
 \underline{x^4 - 4x^3 + x^2} \\
 -2x^3 - 27x^2 + 138x - 35 \\
 \underline{-2x^3 + 8x^2 - 2x} \\
 -35x^2 + 140x - 35 \\
 \underline{-35x^2 + 140x - 35} \\
 0
 \end{array}$$

We get ,

$$x^4 - 6x^3 - 26x^2 + 138x - 35 = (x^2 - 4x + 1)(x^2 - 2x - 35)$$

Now factorizing $x^2 - 2x - 35$ we get,

$$x^2 - 2x - 35 \text{ is also a factor of given polynomial and } x^2 - 2x - 35 = (x - 7)(x + 5)$$

The value of polynomial is also zero when,

$$x - 7 = 0$$

$$\text{or } x = 7$$

$$\text{And, } x + 5 = 0$$

$$\text{Or } x = -5$$

Hence, 7 and -5 are also zeroes of this polynomial.

Q. 5 If the polynomial $x^4 - 6x^3 + 16x^2 - 25x + 10$ is divided by another polynomial $x^2 - 2x + k$ the remainder comes out to be $x + a$, find k and a .

Answer:

To solve this question divide $x^4 - 6x^3 + 16x^2 - 25x + 10$ by $x^2 - 2x + k$ by long division method

Let us divide, by $x^4 - 6x^3 + 16x^2 - 25x + 10$ by $x^2 - 2x + k$

$$\begin{array}{r}
 x^2 - 4x + (8 - k) \\
 x^2 - 2x + k \overline{) x^4 - 6x^3 + 16x^2 - 25x + 10} \\
 \underline{x^4 - 2x^3 + kx^2} \\
 -4x^3 + (16 - k)x^2 - 25x + 10 \\
 \underline{-4x^3 + 8x^2 - 4kx} \\
 + - + \\
 \hline
 (8 - k)x^2 + (4k - 25)x + 10 \\
 \underline{(8 - k)x^2 + (16 - 2k)x + (8 - k)x} \\
 - + - \\
 \hline
 (2k - 9)x + (10 - 8k + k^2)
 \end{array}$$

So, remainder = $(2k - 9)x + (10 - 8k + k^2)$

But given remainder = $x + a \Rightarrow (2k - 9)x + (10 - 8k + k^2) = x + a$

Comparing coefficient of x , we have $2k - 9 = 1 \Rightarrow 2k = 10 \Rightarrow k = 5$ and Comparing constant term, $10 - 8k + k^2 = a$

$$\Rightarrow a = 10 - 8(5) + 5^2$$

$$\Rightarrow a = 10 - 40 + 25 \Rightarrow a = -5. \text{ So, the value of } k \text{ is } 5 \text{ and } a \text{ is } -5.$$