FRACTIONS

A fraction is a part of whole and is written as $\frac{a}{b}$, where 'a' and 'b' are integers and b \neq 0 Ex: $\frac{2}{3}$

- The upper trem 'a' is called as "numerator" and lower term 'b' is called the "denominator". Thus $Fraction = \frac{Numerator}{Demominator}$
- Value of fraction =1, if numerator = denominator
- ♦ Value of fraction =0, if numerator = 0; and denominator ≠ 0
- Value of fraction is not defined if denominator = 0
- Value of fraction remains unaltered if it is multiplied and divided by same number

KINDS OF FRACTIONS:



Improper fraction expressed as Mixed fraction.

Ex: $\frac{49}{12}$ can be written as $4\frac{5}{12}$ On dividing 49 by 12 we get 4 as quotient (Integer) as 5 as remainder (Numerator) and denominator is unaltered

Mixed fraction expressed as Improper fraction.

Ex: $5\frac{7}{8}$ can be written as $\frac{47}{8}$; Numerator = (5×8) + 7 = 40 + 7 = 47 and Denominator is unaltered.

FRACTIONS



Converting Unlike to like fractions:

- Find LCM of denominators of all given fractions
- Multiply each fraction's numerator and denominator with same number such that denominator is equal to the LCM

 $\frac{3}{4}, \frac{3}{5}, \frac{7}{8}, \frac{9}{16} \gg Lcm \ of \ 4, 5, 7, 8 \ is \ 80 \ \frac{3 \times 20}{4 \times 20}, \frac{3 \times 16}{5 \times 16}, \frac{7 \times 10}{8 \times 10}, \frac{9 \times 5}{16 \times 5} = \frac{60}{80}, \frac{48}{80}, \frac{70}{80}, \frac{45}{80}, \frac{10}{80}, \frac{10$

<u>Comparing Fractions:</u> Convert the fractions into like fractions and then the fraction with the greater numerator is greater.

Inserting Fraction between given fraction:

- Add numerator of the given fractions to get the required numerator
- Similarly add denominator to get the required denominator
 - $\frac{3}{4}, \frac{5}{6}, \gg \frac{3}{5}, \quad \frac{3+5}{4+6}, \frac{5}{6} \gg > \frac{3}{5}, \quad \frac{8}{10}, \frac{5}{6}$

Fundamental Operations on Fractions

Multiplication

- **Multiplication by an integer** Keep denominator unchanged, multiply numerator by integer and simplify Ex: $4 \times \frac{4}{12} = \frac{16}{12} = \frac{4}{3}$
- **Multiplication by fraction** Multiply numerators together and denominators separately together and simplify Ex: $\frac{3}{2} \times \frac{4}{12} = \frac{3 \times 4}{12 \times 2} = \frac{12}{24} = \frac{1}{2}$

FRACTIONS

Addition and Subtraction

• Fractions having same denominator $6\frac{2}{5}$ and $4\frac{4}{5}$

Convert into improper fraction $\frac{32}{5} \pm \frac{24}{5}$ Add / Sub the numerators of each fraction and denominator is retained. $\frac{32\pm24}{5}$ If the resultant is improper

fraction convert to Mixed fraction

• Fractions having different denominator $6\frac{2}{5}$ and $4\frac{4}{4}$

Convert into improper fraction $\frac{32}{5} \pm$

 $\frac{20}{4}$ Take L.C.M of numerators L.C.M

(5,4)=20 which is the denominator Multiply each fraction with LCM resultant are added/subtracted to numerator $\left(\frac{32}{5} \times 20\right) \pm \left(\frac{20}{4} \times 1\right)$

20 = (32×4)± (20×5) = 128±100

Division

- **Division in fraction** Multiply fraction by the reciprocal of the divisor and simplify Ex: $\frac{4}{12} \div \frac{4}{2} = \frac{4}{12} \times \frac{2}{4} = \frac{1}{6}$
- Division in decimal fraction Division by 10, 100,1000,.....shift decimal point to left as many digits as the number of zeros Ex: 43.4 ÷ 100 = 0.434
- Division of decimal fraction by an integer is done like ordinary division and the decimal point is placed to quotient such that its number of digits after decimal point is equal to digits after decimal point in the dividend
 Ex: 5.64 ÷ 2 = 2.82