18

OCCURRENCE AND EXTRACTION OF METALS

Metals and their alloys are extensively used in our day-to-day life. They are used for making machines, railways, motor vehicles, bridges, buildings, agricultural tools, aircrafts, ships etc. Therefore, production of a variety of metals in large quantities is necessary for the economic growth of a country. Only a few metals such as gold, silver, mercury etc. occur in free state in nature. Most of the other metals, however, occur in the earth's crust in the combined form, i.e., as compounds with different anions such as oxides, sulphides, halides etc. In view of this, the study of recovery of metals from their ores is very important. In this lesson, you shall learn about some of the processes of extraction of metals from their ores, called metallurgical processes.

Objectives

After reading this lesson, you will be able to :

- differentiate between minerals and ores;
- recall the occurrence of metals in native form and in combined form as oxides, sulphides, carbonates and chlorides;
- list the names and formulae of some common ores of Na, Al, Sn, Pb ,Ti, Fe, Cu, Ag and Zn;
- list the occurrence of minerals of different metals in India;
- list different steps involved in the extraction of metals;

Notes

Chemistry of Elements

MODULE-6

^{*} An alloy is a material consisting of two or more metals, or a metal and a non-metal. For example, brass is an alloy of copper and zinc; steel is an alloy of iron and carbon.

MODULE-6

Chemistry

Chemistry of Elements



list and explain various methods for concentration of or

- list and explain various methods for concentration of ores (gravity separation, magnetic separation, froth floatation and chemical method);
- explain different metallurgical operations : roasting, calcination and smelting with suitable examples;
- choose the reducing agent for a given ore;
- differentiate between flux and slag, and
- explain different methods for refining of metals : poling, liquation, distillation and electrolytic refining.

18.1 Occurrence of Metals

Metals occur in nature in free as well as combined form. Metals having low reactivity show little affinity for air, moisture, carbon dioxide or other non-metals present in nature. Such metals may remain in elemental or native (free) state in nature. Such metals are called "noble metals" as they show the least chemical reactivity. For example gold, silver, mercury and platinum occur in free state.

On the other hand, most of the metals are active and combine with air, moisture, carbon dioxide and non-metals like oxygen, sulphur, halogens, etc. to form their compounds, like oxides, sulphides, carbonates, halides and silicates. i.e., they occur in nature in a combined state.

A naturally occurring material in which a metal or its compound occurs is called a *mineral*. A mineral from which a metal can be extracted economically is called an *ore*.

An ore is that mineral in which a metal is present in appreciable quantities and from which the metal can be extracted economically.

The main active substances present in nature, expecially in the atmosphere are oxygen and carbon dioxide. In the earth's crust, sulphur and silicon are found in large quantities. Sea-water contains large quantities of chloride ions (obtained from dissolved sodium chloride). Most avtive metals are highly electropositive and therefore exist as ions. It is for this reason that most of the important *ores* of these metals occur as (i) oxides (ii) sulphides (iii) carbonates (iv) halides and (v) silicates. Some sulphide ores undergo oxidation by air to form sulphates. This explains the occurrence of sulphate ores.

Ores are invariably found in nature in contact with rocky materials. These rocky or earthy impurities accompanying the ores are termed as *gangue* or *matrix*.

Some important ores and the metals present in these ores are listed in Table 18.1

Type of Ore	Metals (Common Ores)
Native Metals	Gold (Au), silver (Ag)
Oxide ores	Iron (Haematite, Fe_2O_3); Aluminium (Bauxite, $Al_2O_3 \cdot 2H_2O$); Tin (Cassiterite, SnO_2); Copper (Cuprite, Cu_2O); Zinc (Zincite, ZnO); Titanium (Ilmenite, $FeTiO_3$, Rutile, TiO_2)
Sulphide ores	Zinc (Zinc blende, ZnS); Lead (Galena, PbS); Copper (Copper glance, Cu_2S); Silver (Silver glance or Argentite, Ag_2S); Iron (Iron pyrites, FeS_2)
Carbonate ores	Iron (Siferite, $FeCO_3$); Zinc (Calamine, $ZnCO_3$), Lead (Cerrusite, PbCO ₃)
Sulphate ores	Lead (Anglesite, PbSO ₄)
Halide ores	Silver (Horn silver, AgCl); Sodium (Common salt or Rock salt, NaCl); Aluminium (Cryolite, Na_3AlF_6)
Silicate ores	Zinc (Hemimorphite, 2ZnO.SiO ₂ .H ₂ O)

Table 18.1 Some Important Ores

MODULE-6

Chemistry of Elements



18.1.1 Mineral Wealth of India

India possesses large deposits of minerals of some important metals such as iron, manganese, aluminium, magnesium, chromium, thorium, uranium, titanium and lithium. They constitute one-quarter of the world's known mineral resources. Mineral fuels (like coal ,petroleum and natural gas) constitute more than 80% while metallic minerals constitute only about 10% of the total volume of mineral production in a year. In this section, we shall list some of the important minerals of a few common metals such as Fe, Cu, Ag, Zn, Ti, Al Sn, Pb and Na and their locations in India.

Important Ores of Some Common Elements

Iron	Iron ore reserves in the country are estimated at 1750 crore tonnes. Major iron ore mining is done in Goa, Madhya Pradesh, Bihar, Karnataka, Orissa and Maharashtra.
Aluminium	Its chief ore, bauxite is available in India in abundance. The annual production of bauxite is estimated to be more than 2 million tonnes. Important deposits of bauxite in the country occur in Bihar, Goa, Madhya Pradesh, Maharashtra, Tamil Nadu, Gujarat, Karnataka, Orissa, Uttar Pradesh, Andhra Pradesh, Jammu & Kashmir and Rajasthan.
Copper	India does not have rich copper ore deposits. The total reserves of the ore in the country are estimated around 60 crore tonnes. Major ore producing areas are Singhbum district (Bihar), Balaghat district (Madhya Pradesh) and Jhunjhunu district (Rajasthan).
Zinc and Lead	India has no significant deposits of lead ores. Lead and Zinc have been

located at Zawar mines near Udaipur (Rajasthan) and at Hazaribagh

3

MODULE-6



Chemistry of Elements

Chemistry

Tin

(Jharkhand), Sargipalli mines in Orissa and Bandalamottu lead project in Andhra Pradesh. Some reserves have been located in Gujarat and Sikkim. The total reserves are estimated at about 360 million tonnes in which lead content is estimated to be around 5 million tonnes and zinc to be around 16 million tonnes.

Deposits of tinstone (SnO_2) are found in Hazaribagh (Jharkhand) and Orissa.

- Silver India does not possess rich silver deposits. Gold from Kolar fields and Hutti gold fields (Karnataka) and lead-zinc ores of Zawar mines (Rajasthan) contain some silver.
- Titanium Ilmenite (FeTiO₃) is recovered from beach sands of Kerala and Tamil Nadu. The estimated reserves are around 100 to 150 million tonnes.


Tincal or Native borax (Na₂B₄O₇.10H₂O) is found in Ladakh and Sodium : Kashmir.

18.2. General Principles of Extraction of Metals

The process of extracting the metals from their ores and refining them is called *metallurgy*. The choice of the process depends upon the nature of the ore and the type of the metal. The metal content in the ore can vary depending upon the impurities present and chemical composition of the ore. Some common steps involved in the extraction of metals from their ores are :

- (i) Crushing and pulverization
- (ii) Concentration or dressing of the ore
- Calcination or roasting of the ore (iii)
- (iv) Reduction of metal oxides to free metal
- (v) Purification and refining of metal.

18.2.1. Crushing and Pulverization



The ore is generally obtained as big rock pieces. These big lumps of the ore are crushed to smaller pieces by using jawcrushers and grinders. It is easier to work with crushed ore. The big lumps of the ore are brought in between the plates of a crusher forming a jaw. One of the plates of the crusher is stationary while the other moves to and fro and the crushed pieces are collected below (Fig. 18.1).

Fig. 18.1 : Jaw Crusher

Crushed ore The crushed pieces of the ore are then pulverized (powdered) in a stamp mill shown in Fig. 18.2. The heavy stamp rises and falls on a hard die to powder the ore. The powdered ore is then taken out through a screen by a stream of water.

Pulverization can also be carried out in a ball mill. The crushed ore is taken in a steel cylinder containing iron balls. The cylinder is set into revolving motion. The striking balls pulverize the crushed ore into fine powder.




Fig. 18.2 : The Stamp mill

18.2.2 Concentration or Dressing of the Ore

Generally, the ores are found mixed with earthy impurities like sand, clay, lime stone etc. These unwanted impurities in the ore are called *gangue* or *matrix*.

The process of removal of gangue from powdered ore is called concentration or ore dressing.

There are several methods for concentrating the ores. The choice of method depends on the nature of the ore. Some important methods are :

(i) Gravity separation (Hydraulic washing) : In this method, the light (low specific gravity) earthy impurities are removed from the heavier metallic ore particles by washing with water. It is therefore, used for the concentration of heavier oxide ores, like haematite (Fe₂O₃) tinstone (SnO₂) and gold (Au). In this method, as shown in the Fig. 18.3 the powdered ore is agitated with water or washed with a strong current of water. The heavier ore settles down rapidly in the grooves and the lighter sandy and earthy materials (gangue particles) are washed away.




Fig. 18.3 : Gravity Separation (Hydraulic Washing)

MODULE-6

Chemistry of Elements



Chemistry of Elements



(ii) Magnetic separation method :

By this method, those ores can be concentrated which either contain impurities which are magnetic or are themselves magnetic in nature.

For example, the tin ore, tin stone (SnO_2) itself is non-magnetic but contains magnetic impurities such as iron tungstate (FeWO₄) and manganese tungstate (MnWO₄).





Fig. 18.4. : Magnetic Separation

The finely powdered ore is passed over a conveyer belt moving over two rollers, one of which is fitted with an electromagnet (Fig. 18.4). The magnetic material is attracted by the magnet and falls in a separate heap. In this way magnetic impurities are separated from non-magnetic material.

(iii) Froth floatation method :

This method is especially applied to sulphide ores, such as **galena** (PbS), **zinc blende** (ZnS), or **copper pyrites** (CuFeS₂). It is based on the different wetting properties of the surface of the ore and gangue particles. The sulphide ore particles are wetted



preferentially by oil and gangue particles by water. In this process, finely powdered ore is mixed with either pine oil or eucalyptus oil. It is then mixed with water. Air is blown through the mixture with a great force. Froth is produced in this process which carries the weted ore upwards with it. Impurities (gangue particles) are left in water and sink to the bottom from which these are drawn off (Fig. 18.5).

(iv) Chemical method :

In this method the ore is treated with a suitable chemical reagent

which dissolves the ore leaving behind insoluble impurities. The ore is then recovered from the solution by a suitable chemical method. This is applied for extraction of aluminium from bauxite $(Al_2O_3.2H_2O)$. Bauxite is contaminated with iron (III) oxide (Fe_2O_3) , titanium (IV) oxide (TiO_2) and silica (SiO_2) . These impurities are removed by digesting the powdered ore with aqueous solution of sodium hydroxide at 420 K

under pressure. Aluminium oxide dissolves in sodium hydroxide, whereas, iron (III) oxide, silica and titanium (IV) oxide remain insoluble and are removed by filtration.

$$\begin{array}{c} Al_2O_3 + 6NaOH & \longrightarrow 2Na_3AlO_3 + 3H_2O \\ & \text{Sodium aluminate} \end{array}$$

Sodium aluminate is diluted with water to obtain precipitate of aluminium hydroxide. It is filtered and ignited to obtain pure alumina.

 $Na_3AlO_3 + 3H_2O \longrightarrow Al(OH)_3 + 3NaOH$

 $2\text{Al}(\text{OH})_3 \xrightarrow{\Delta} \text{Al}_2\text{O}_3 + 3\text{H}_2\text{O}$

Intext Questions 18.1

1. Write the names of eight important metals. Give an example of one important ore of each metal.

.....

.....

What is the difference between an ore and a mineral?

3. Name some important methods of concentrating the ores.

-
- 4. Which method of concentration is applied in the following cases?

(i) Magnetic ores (ii) Sulphide ores (iii) Bauxite ore

5. Which metal is extracted from the mineral zinc blene ?

.....

.....

18.2.3 Calcination and Roasting of the Ore

The concentrated ore is converted into metal oxide by calcination or roasting.

(A) Calcination :

2.

Calcination involves heating of the concentrated ore in a limited supply of air so that it loses moisture, water of hydration and gaseous volatile substances. The ore is heated to a temperature so that it does not melt. Two examples of calcination are given below:

(i) Removal of water of hydration

 $Al_2O_3.2H_2O \longrightarrow Al_2O_3 + 2H_2O$

(ii) Expulsion of CO_2 from carbonate

 $ZnCO_3 \longrightarrow ZnO + 2CO_2$

Chemistry of Elements

MODULE-6



Chemistry of Elements



(B) Roasting :

Roasting is a process in which the concentrated ore is heated in a free supply of air at a temperature insufficient to melt it. The following changes take place during roasting :

(i) Drying of the ore.

(ii) Removal of the volatile impurities like arsenic, sulphur, phosphorus and organic matter.

 $4As + 3O_2 \longrightarrow 2As_2O_3(g)$ $S + O_2 \longrightarrow SO_2(g)$ $4P + 5O_2 \longrightarrow P_4O_{10}(g)$

(iii) Conversion of the sulphide ores into oxides

 $2PbS + 3O_2 \longrightarrow 2PbO + 2SO_2$ $2ZnS + 3O_2 \longrightarrow 2ZnO + 2SO_2$

Calcination and roasting are generally carried out in a reverberatory furnace or in a multiple hearth furnace.

18.2.4 Reduction of the Metal Oxides to Free Metal

This process is carried out after calcination or roasting of ores. In this process called *smelting*, the oxide ores are converted into the metallic state by reduction.

(A) Smelting :

Smelting is a process in which the oxide ore in molten state is reduced by carbon or other reducing agents to free metal.

(i) by using carbon as a reducing agent :

This method is used for the isolation of iron, tin and zinc metals from their respective oxides. The oxide ores are strongly heated with charcoal or coke. Reduction occurs by the action of carbon and/or carbon monoxide which is produced by the partial combustion of coke or charcoal.

$$Fe_{2}O_{3} + 3C \longrightarrow 2Fe + 3CO$$

$$Fe_{2}O_{3} + CO \longrightarrow 2FeO + CO_{2}$$

$$FeO + CO \longrightarrow Fe + CO_{2}$$

$$SnO_{2} + 2C \longrightarrow Sn + 2CO$$

$$ZnO + C \longrightarrow Zn + CO$$

Although the ore has been concentrated in an earlier step, it is still contaminated with some gangue material which is finally removed in the reduction process by the addition of *flux* during smelting.

Flux is a chemical substance which combines with gangue at higher temperatures to form easily fusible material called *slag* which is not soluble in the molten metal. Flux are of two types :

Basic Flux :

On heating, lime stone is converted into calcium oxide used as basic flux which combines with acidic impurities like silica in metallurgy of iron and forms fusible calcium silicate ($CaSiO_3$).

 $\begin{array}{ccc} CaCO_3 & \longrightarrow & CaO + CO_2(g) \\ \text{(Limestone)} & & \end{array}$

 $\begin{array}{ccc} CaO &+ & SiO_2 & \longrightarrow & CaSiO_3 \\ \textbf{(Basic flux)} & \textbf{(Acidic gangue)} & \textbf{(Slag)} \end{array}$

Acidic flux :

 ${\rm SiO}_2$ is used as acidic flux to remove basic impurity of FeO in metallurgy of Cu.

 $\begin{array}{ccc} SiO_2 &+ & FeO & \longrightarrow & FeSiO_3 \\ \textbf{(Acidic flux)} & \textbf{(Basic gangue)} & & \textbf{(Slag)} \end{array}$

The fusible slag, such as calcium silicate formed during smelting floats over the molten metal and is thus easily removed. Another advantage is that the slag provides a covering to the molten metal thus preventing it from getting oxidized by air.

(ii) Other reducing agents :

Oxide ores which cannot be reduced by carbon or metals which show affinity to carbon by forming metal carbides, are reduced by reducing agents like aluminium, sodium, magnesium or hydrogen. *Oxide* like chromium oxide (Cr_2O_3) or manganese oxide (Mn_3O_4) are reduced by aluminium powder is a highly exothermic reaction. This process is known as *Goldschmidt's Alumino-thermite reduction method*.

 $Cr_2O_3 + 2Al \longrightarrow 2Cr + Al_2O_3 + Heat$ $3Mn_3O_4 + 8Al \longrightarrow 9Mn + 4Al_2O_3 + Heat$

Heat is generated in the process due to the formation of Al_2O_3 which is a highly exothermic reaction.

Titanium is obtained by the reduction of $TiCl_4$ (produced by the action of carbon and chlorine on TiO_2) by Mg in an inert atmosphere of argon (Kroll process).

$$TiCl_4 + 2Mg \xrightarrow{Heat} Ti + 2MgCl_2$$

Titanium can also be obtained by the reduction of TiO_2 by sodium

$$TiO_2 + 4Na \rightarrow Ti + 2Na_2O$$

Tungsten and molybedenum can be obtained by the reduction of their oxides by hydrogen,

$$MoO_3 + 3H_2 \longrightarrow Mo + 3H_2O$$

MODULE - 6 Chemistry of Elements



Chemistry of Elements



(iii) Self-reduction :

This is applied to the sulphide ores of copper, mercury and lead. The ores are heated in air, a part of these sulphide ores is changed into the oxide or sulphate which then reacts with the remaining part of the sulphide ore to give the metal and sulphur dioxide. The reactions showing their extraction are given below :

(1)
$$\begin{array}{ccc} 2Cu_2S + 3O_2 & \longrightarrow & 2Cu_2O + 2SO_2 \\ Copper \ glance \end{array}$$

 $2Cu_2O + Cu_2S \longrightarrow 6Cu + SO_2$

Copper produced at this stage is called *Blister copper*. The evolution of sulphur dioxide produces blisters on the surface of solidified copper metal.

(2) $\begin{array}{cccc} 2HgS + 3O_2 & \longrightarrow & 2HgO + 2SO_2 \\ Cinnabar & & \\ 2HgO + HgS & \longrightarrow & 3Hg + SO_2 \\ \end{array}$ (3) $\begin{array}{cccc} 2PbS + 3O_2 & \longrightarrow & 2PbO + 2SO_2 \\ Galena & & \\ PbS + 2O_2 & \longrightarrow & PbSO_4 \\ PbS + 2PbO & \longrightarrow & 3Pb + SO_2 \\ PbS + PbSO_4 & \longrightarrow & 2Pb + 2SO_2 \end{array}$

(B) Reduction of concentrated ores by other methods:

Some metals cannot be obtained from their ores by using common reducing agents such as C, CO, H_2 etc. Other methods of reduction are used for such cases.

(i) Reduction by precipitation :

Noble metals like silver and gold are extracted from their concentrated ores by dissolving metal ions in the form of their soluble complexes. The metal ions are then regenerated by adding a suitable reagent. For example, concentrated argentite ore (Ag_2S) is treated with a dilute solution of sodium cyanide (NaCN) to form a soluble complex :

 $Ag_2S + 4NaCN \longrightarrow 2Na[Ag(CN)_2] + Na_2S$

This solution is decanted off and treated with zinc to precipitate silver,

 $2Na[Ag(CN)_2] + Zn \longrightarrow Na_2[Zn(CN)_4] + 2Ag \downarrow$

(ii) Electrolytic Reduction :

Active metals like sodium, potassium and aluminium etc., are extracted by the electrolysis of their fused (*molten*) salts. For example, sodium is obtained by the electrolysis of fused sodium chloride (Down's process). The reactions taking place in the electrolytic cell are :

NaCl \implies Na⁺ + Cl⁻

Na⁺ ions move towards the cathode and Cl⁻ ions move towards the anode. Following reactions take place at the electrodes : At the Cathode $Na^+ + e^- \longrightarrow Na$ (Reduction) (Negative electrode) $Cl^- \longrightarrow Cl + e^-$ At the Anode (Oxidation) (Positive electrode) $Cl + Cl \longrightarrow$ Cl_2 Aluminium is extracted from molten alumina (Al₂O₂) by electrolysis. The melting point of alumina is quite high (2323K) which is inconvenient for electrolysis. It dissolves in molten cryolite (Na_AIF_c) at around 1273 k. The reactions which take place in the cell are:. At the Cathode $Al^{3+} + 3e^{-} \longrightarrow Al$ (metal) At the Anode $C + 2O^{2-} \longrightarrow CO_2 + 4e^{-}$ **Intext Questions 18.2** Explain the following terms : 1. Calcination, Roasting, Smelting, Flux and Slag. 2. Which is the cheapest and most abundant reducing agent employed in the extraction of metals? Name the process which is used for converting sulphide ores into oxides. 3. 4. How are oxide ores reduced ? 5. Name few materials which are used as flux in metallurgical processes. 6. What happens to (i) Bauxite and (ii) Calamine ores during calcination?

18.2.5 Refining of Metals

Except in the electrolytic reduction method, metals produced by any other method are generally impure. The impurities may be in the form of (i) other metals (ii) unreduced oxide of the metal (iii) non-metals like carbon, silicon, phosphorus, sulphur etc. and (iv)

_

11

Chemistry of Elements

MODULE-6




Occurrence and Extraction of Metals


Chemistry of Elements




flux or slag. Crude metal may be refined by using one or more of the following methods :

(i) *Liquation :* Easily fusible metals like tin, lead etc. are refined by this process. In this method, the impure metal is poured on the sloping hearth of a reverberatory furnace (Fig.18.6) and heated slowly to a temperature little above the melting point of the metal. The pure metal drains out leaving behind infusible impurities .







For example, in the electrolytic refining of crude copper (blister copper), a large

piece of impure copper is made anode and a thin piece of pure copper is made the cathode. An acidified solution of copper sulphate is used as an electrolyte. On passing an electric current of low voltage through the solution copper (II) ions obtained from copper sulphate solution go to the cathode where they are reduced to the free copper metal and get deposited.

$$Cu^{2+} + 2e^- \longrightarrow Cu$$
 (at cathode)

An equivalent amount of the metal from the anode dissolves into the electrolyte as C_{11}^{2+} ions

$$Cu \longrightarrow Cu^{2+} + 2e^{-}$$
 (at anode)

As the process goes on, anode becomes thinner while the cathode becomes thicker. The impurities like silver, gold settle down at the bottom of the cell as *'anode mud'*.



Intext Questions 18.3

- 1. State the nature of materials used for constructing cathodes and anodes in the electrolytic cell for refining of copper. Write chemical equations for the reactions which take place.
- 2. Which method is used for the refining of metals that are easily fused ?
-
- 3. Which metals are refined by poling ?
- 4. Name a metal which is purified by distillation.

5. Name any three elements which are refined by electrolytic refining

What You Have Learnt

- Metals play an extremely useful role in day-to-day life.
- Most metals are found in nature in combined form. Only a few noble metals such as gold, silver, occur in nature in free state,
- The process of extraction of metals from their ores is called *metallurgical process*.
- A naturally occurring material in which a metal or its compound occurs is called a mineral. A mineral from which a metal can be extracted economically is called an ore. Thus all minerals do not serve the purpose of ores.

MODULE-6

Chemistry of Elements



Chemistry of Elements



- Most active metals are highly electropositive and exist as Mⁿ⁺ ions. Therefore, they are found in nature in association with some common anions, i.e., as salts like *oxides*, *sulphides*, *carbonates*, *halides*, *silicates* etc. Some sulphides slowly undergo oxidation by air to form *sulphates*.
- India possesses rich mineral wealth with abundance of iron, aluminium and some amount of copper, tin, lead, silver and gold.
- Various steps involved in the extraction of metal are :
 - (i) Crushing and pulverization
 - (ii) Concentration or dressing of the ore
 - (iii) Calcination or Raosting of the concentrated ore
 - (iv) Reduction of the oxides to free metal
- The metals thus obtained are then purified by employing some suitable method, viz. *liquation, poling, distillation or electrolytic refining.*

Terminal Exercise

- 1. Name the metal oxides that are not reduced to metallic state by heating with carbon. Which reducing agent is used for these ores ?
- 2. Which metal sulphide combines with its oxide to form metal? Give reactions.
- 3. Name four reducing agents other than carbon, used during smelting.
- 4. What is the difference between calcination and roasting ?
- 5. Give the name and formula of at least one ore, of the following metals :

(i) copper (ii) zinc (iii) iron (iv) tin

- 6. What happens when
 - (i) Calamine is calcined.
 - (ii) Zinc blende is roasted.
 - (iii) Silica is heated with lime stone.
- 7. Describe the froth floatation method for the concentration of sulphide ore.
- 8. Give chemical equations involved in the self reduction method for the extraction of copper.

Answers to Intext Questions

18.1

1. Sodium (Na), Aluminium (Al), Silver (Ag), Iron (Fe), Zinc (Zn), Lead (Pb), Gold (Au), Mercury (Hg).

Ores : rock salt, bauxite, argentite, haematite, zincite, galena, native gold, cinnabar, respectively.

2. *Mineral* is a naturally occurring material in which a metal or its compound occurs.

An *ore* is that mineral in which metal is present in large quantity and from which the metal can be extracted in pure and high quality, economically.

- 3. Gravity separation, magnetic separation, froth floatation and chemical method.
- 4. (i) Magnetic separation method
 - (ii) Froth floatation method
 - (iii) Chemical method
- 5. Zinc.

18.2

1. *Calcination* : it involves heating of the ore in a limited supply of air to a temperature that the ore does not melt.

Roasting : It involves heating of the ore in a free supply of air to a temperature that the ore does not melt.

Smelting : The extraction of metal in the fused state is termed smelting. The ore is reduced by carbon to the free metal.

Flux : it is a chemical substance which combines with gangue at higher temperatures to form easily fusible material called slag.

Slag is a fusible chemical compound formed by the reaction of flux with gangue. Slag is not soluble in the molten metal and is thus separated.

- 2. Carbon in the form of coke.
- 3. Roasting; $2ZnS + 3O_2 \longrightarrow 2ZnO + 2SO_2(g)$
- 4. Oxide ores are reduced to metal by heating them with coke.
- 5. Silica, borax and other non-metallic oxides are acidic fluxes. Lime stone (CaCO₃) is a basic flux.
- 6. (i) Al_2O_3 . $2H_2O \longrightarrow Al_2O_3 + 2H_2O$

(ii) $ZnCO_3 \longrightarrow ZnO + CO_2$

18.3

- 1. Cathode : Pure metalAt Cathode : $Cu^{2+} + 2e \rightarrow Cu$ Anode : Impure metalAt Anode : $Cu \rightarrow Cu^{2+} + 2e$
- 2. Liquation
- 3. Copper and tin
- 4. Zinc
- 5. Copper, Silver and Tin

MODULE - 6 Chemistry of Elements

