

TRIGONOMETRIC RATIOS & IDENTITIES [JEE ADVANCED PREVIOUS YEAR SOLVED PAPER]

JEE ADVANCED

Single Correct Answer Type

1. If $\tan \theta + \sin \theta = m$ and $\tan \theta - \sin \theta = n$, then

- a. $m^2 - n^2 = 4mn$ b. $m^2 + n^2 = 4mn$
c. $m^2 - n^2 = m^2 + n^2$ d. $m^2 - n^2 = 4\sqrt{mn}$

(IIT-JEE 1970)

2. If $\alpha + \beta + \gamma = 2\pi$, then

- a. $\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}$
b. $\tan \frac{\alpha}{2} \tan \frac{\beta}{2} + \tan \frac{\beta}{2} \tan \frac{\gamma}{2} + \tan \frac{\gamma}{2} \tan \frac{\alpha}{2} = 1$

- c. $\tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = -\tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}$
d. none of these (IIT-JEE 1979)
3. If $\tan \theta = -\frac{4}{3}$, then $\sin \theta$ is
a. $-\frac{4}{5}$ but not $\frac{4}{5}$ b. $-\frac{4}{5}$ or $\frac{4}{5}$
c. $\frac{4}{5}$ but not $-\frac{4}{5}$ d. none of these
(IIT-JEE 1979)
4. Given $A = \sin^2 \theta + \cos^4 \theta$, then for all real θ ,
a. $1 \leq A \leq 2$ b. $3/4 \leq A \leq 1$
c. $13/16 \leq A \leq 1$ d. $3/4 \leq A \leq 13/16$
(IIT-JEE 1980)
5. The value of $\left(1 + \cos \frac{\pi}{8}\right) \left(1 + \cos \frac{3\pi}{8}\right) \left(1 + \cos \frac{5\pi}{8}\right) \left(1 + \cos \frac{7\pi}{8}\right)$ is
a. $1/4$ b. $3/4$ c. $1/8$ d. $3/8$
(IIT-JEE 1984)
6. The value of the expression $\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$ is equal to
a. 2 b. $2 \sin 20^\circ / \sin 40^\circ$
c. 4 d. $4 \sin 20^\circ / \sin 40^\circ$
(IIT-JEE 1988)
7. Let $0 < x < \pi/4$, then $(\sec 2x - \tan 2x)$ equals
a. $\tan \left(x - \frac{\pi}{4}\right)$ b. $\tan \left(\frac{\pi}{4} - x\right)$
c. $\tan \left(x + \frac{\pi}{4}\right)$ d. $\tan^2 \left(x + \frac{\pi}{4}\right)$
(IIT-JEE 1994)
8. Let n be a positive integer such that $\sin \frac{\pi}{2n} + \cos \frac{\pi}{2n} = \frac{\sqrt{n}}{2}$. Then
a. $6 \leq n \leq 8$ b. $4 < n \leq 8$
c. $4 \leq n \leq 8$ d. $4 < n < 8$ (IIT-JEE 1994)
9. $3(\sin x - \cos x)^4 + 6(\sin x + \cos x)^2 + 4(\sin^6 x + \cos^6 x)$ is equal to
a. 11 b. 12 c. 13 d. 14
(IIT-JEE 1995)
10. $\sec^2 \theta = \frac{4xy}{(x+y)^2}$ is true if and only if
a. $x + y \neq 0$ b. $x = y, x \neq 0$
c. $x = y$ d. $x \neq 0, y \neq 0$
(IIT-JEE 1996)
11. Let $f(\theta) = \sin \theta (\sin \theta + \sin 3\theta)$. Then $f(\theta)$ is
a. ≥ 0 only when $\theta \geq 0$ b. ≤ 0 for all real θ
c. ≥ 0 for all real θ d. ≤ 0 only when $\theta \leq 0$
(IIT-JEE 2000)

12. If $\alpha + \beta = \pi/2$ and $\beta + \gamma = \alpha$, then $\tan \alpha$ equals
a. $2(\tan \beta + \tan \gamma)$ b. $\tan \beta + \tan \gamma$
c. $\tan \beta + 2 \tan \gamma$ d. $2 \tan \beta + \tan \gamma$
(IIT-JEE 2001)
13. The maximum value of $(\cos \alpha_1)(\cos \alpha_2) \cdots (\cos \alpha_n)$ under the restrictions $0 \leq \alpha_1, \alpha_2, \dots, \alpha_n \leq \pi/2$ and $(\cot \alpha_1)(\cot \alpha_2) \cdots (\cot \alpha_n) = 1$ is
a. $1/2^{n/2}$ b. $1/2^n$ c. $1/2n$ d. 1
(IIT-JEE 2001)
14. Given both θ and ϕ are acute angles and $\sin \theta = 1/2$, $\cos \phi = 1/3$, then the value of $\theta + \phi$ belongs to
a. $\left(\frac{\pi}{3}, \frac{\pi}{2}\right]$ b. $\left(\frac{\pi}{2}, \frac{2\pi}{3}\right]$
c. $\left(\frac{2\pi}{3}, \frac{5\pi}{6}\right]$ d. $\left(\frac{5\pi}{6}, \pi\right]$
(IIT-JEE 2004)
15. Let $\theta \in (0, \pi/4)$ and $t_1 = (\tan \theta)^{\tan \theta}$, $t_2 = (\tan \theta)^{\cot \theta}$, $t_3 = (\cot \theta)^{\tan \theta}$ and $t_4 = (\cot \theta)^{\cot \theta}$, then
a. $t_1 > t_2 > t_3 > t_4$ b. $t_4 > t_3 > t_1 > t_2$
c. $t_3 > t_1 > t_2 > t_4$ d. $t_2 > t_3 > t_1 > t_4$
(IIT-JEE 2006)
16. Let $P = \{\theta : \sin \theta - \cos \theta = \sqrt{2} \cos \theta\}$ and $Q = \{\theta : \sin \theta + \cos \theta = \sqrt{2} \sin \theta\}$ be two sets. Then
a. $P \subset Q$ and $Q - P \neq \emptyset$ b. $Q \not\subset P$
c. $P \not\subset Q$ d. $P = Q$ (IIT-JEE 2011)

Multiple Correct Answers Type

1. The expression $3 \left[\sin^4 \left(\frac{3}{2} \pi - \alpha \right) + \sin^4 (3\pi + \alpha) \right] - 2 \left[\sin^6 \left(\frac{1}{2} \pi + \alpha \right) + \sin^6 (5\pi - \alpha) \right]$ is equal to
a. 0 b. 1 c. 3 d. none of these
(IIT-JEE 1984)
2. For $0 < \phi \leq \pi/2$, if $x = \sum_{n=0}^{\infty} \cos^{2n} \phi$, $y = \sum_{n=0}^{\infty} \sin^{2n} \phi$, $z = \sum_{n=0}^{\infty} \cos^{2n} \phi \sin^{2n} \phi$, then
a. $xyz = xz + y$ b. $xyz = xy + z$
c. $xyz = x + y + z$ d. $xyz = yz + x$
(IIT-JEE 1992)
3. The minimum value of the expression $\sin \alpha + \sin \beta + \sin \gamma$, where α, β, γ are real numbers satisfying $\alpha + \beta + \gamma = \pi$ is
a. positive b. zero
c. negative d. -3 (IIT-JEE 1995)
4. Which of the following number(s) is/are rational?
a. $\sin 15^\circ$ b. $\cos 15^\circ$
c. $\sin 15^\circ \cos 15^\circ$ d. $\sin 15^\circ \cos 75^\circ$
(IIT-JEE 1998)

5. For a positive integer n , let
 $f_n(\theta) = (\tan \theta / 2)(1 + \sec \theta)(1 + \sec 2\theta)(1 + \sec 4\theta) \dots (1 + \sec 2^{n-1}\theta)$. Then
 a. $f_2(\pi/16) = 1$ b. $f_3(\pi/32) = 1$
 c. $f_4(\pi/64) = 1$ d. $f_5(\pi/128) = 1$
 (IIT-JEE 1999)

6. If $\frac{\sin^4 x}{2} + \frac{\cos^4 x}{3} = \frac{1}{5}$, then
 a. $\tan^2 x = \frac{2}{3}$ b. $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{1}{125}$
 c. $\tan^2 x = \frac{1}{3}$ d. $\frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{2}{125}$
 (IIT-JEE 2009)

7. Let $f: (-1, 1) \rightarrow \mathbb{R}$ be such that $f(\cos 4\theta) = \frac{2}{2 - \sec^2 \theta}$
 for $\theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$. Then the value(s) of $f\left(\frac{1}{3}\right)$
 is (are)
 a. $1 - \sqrt{\frac{3}{2}}$ b. $1 + \sqrt{\frac{3}{2}}$ c. $1 - \sqrt{\frac{2}{3}}$ d. $1 + \sqrt{\frac{2}{3}}$
 (IIT-JEE 2012)

Matching Column Type

1. Match List I with List II and select the correct answer using the code given below the lists :

List I	List II
(p) $\left(\frac{1}{y^2} \left(\frac{\cos(\tan^{-1} y) + y \sin(\tan^{-1} y)}{\cot(\sin^{-1} y) + \tan(\sin^{-1} y)} \right)^2 + y^4 \right)^{1/2}$ takes value	(1) $\frac{1}{2}\sqrt{\frac{5}{3}}$
(q) If $\cos x + \cos y + \cos z = 0 = \sin x + \sin y + \sin z$ then possible value of $\cos \frac{x-y}{2}$ is	(2) $\sqrt{2}$
(r) If $\cos \left(\frac{\pi}{4} - x \right) \cos 2x + \sin x \sin 2x \sec x$ $= \cos x \sin 2x \sec x + \cos \left(\frac{\pi}{4} + x \right) \cos 2x$ then possible value of $\sec x$ is	(3) $1/2$
(s) If $\cot \left(\sin^{-1} \sqrt{1-x^2} \right) = \sin \left(\tan^{-1} (x\sqrt{6}) \right)$, $x \neq 0$, then possible value of x is	(4) 1

Codes:

- (p) (q) (r) (s)
 a. (4) (3) (1) (2)
 b. (4) (3) (2) (1)
 c. (3) (4) (2) (1)
 d. (3) (4) (1) (2)

(IIT-JEE 2013)

Integer Answer Type

1. The positive integer value of $n > 3$ satisfying the equation

$$\frac{1}{\sin\left(\frac{\pi}{n}\right)} = \frac{1}{\sin\left(\frac{2\pi}{n}\right)} + \frac{1}{\sin\left(\frac{3\pi}{n}\right)} \text{ is } \quad (\text{IIT-JEE 2011})$$

2. The maximum value of the expression

$$\frac{1}{\sin^2 \theta + 3 \sin \theta \cos \theta + 5 \cos^2 \theta} \text{ is } \quad (\text{IIT-JEE 2010})$$

Fill in the Blanks Type

1. Suppose $\sin^3 x \sin 3x = \sum_{m=0}^n C_m \cos mx$ is an identity in x , where $C_0, C_1, \dots, C_n$ are constants, and

$C_n \neq 0$, then the value of n is _____. (IIT-JEE 1981)

2. The side of a triangle inscribed in a given circle subtends angles α, β , and γ at the center. The minimum value of the arithmetic mean of $\cos\left(\alpha + \frac{\pi}{2}\right)$, $\cos\left(\beta + \frac{\pi}{2}\right)$, and $\cos\left(\gamma + \frac{\pi}{2}\right)$ is equal to _____. (IIT-JEE 1987)

3. The value of

$$\sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \sin \frac{7\pi}{14} \sin \frac{9\pi}{14} \sin \frac{11\pi}{14} \sin \frac{13\pi}{14}$$

is equal to _____. (IIT-JEE 1991)

4. If $K = \sin(\pi/18) \sin(5\pi/18) \sin(7\pi/18)$, then the numerical value of K is _____. (IIT-JEE 1993)

5. If $A > 0, B > 0$ and $A + B = \pi/3$, then the maximum value of $\tan A \tan B$ is _____. (IIT-JEE 1993)

6. If $\cos(x-y), \cos x$ and $\cos(x+y)$ are in H.P., then $\cos x \sec\left(\frac{y}{2}\right) =$ _____. (IIT-JEE 1997)

True/False Type

1. If $\tan A = \frac{1 - \cos B}{\sin B}$, then $\tan 2A = \tan B$.
 (IIT-JEE 1983)

Subjective Type

1. If $\tan \alpha = \frac{m}{m+1}$ and $\tan \beta = \frac{1}{2m+1}$, find the possible values of $(\alpha + \beta)$. (IIT-JEE 1978)

2. a. Draw the graph of $y = \frac{1}{\sqrt{2}} (\sin x + \cos x)$ from

$$x = -\frac{\pi}{2} \text{ to } x = \frac{\pi}{2}. \quad (\text{IIT-JEE 1979})$$

- b. If $\cos(\alpha + \beta) = \frac{4}{5}$, $\sin(\alpha - \beta) = \frac{5}{13}$, and α, β lie between 0 and $\pi/4$, find $\tan 2\alpha$.

3. Prove that $5 \cos \theta + 3 \cos \left(\theta + \frac{\pi}{3} \right) + 3$ lies between -4 and 10 . (IIT-JEE 1979)
4. Given $\alpha + \beta - \gamma = \pi$, prove that $\sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma = 2 \sin \alpha \sin \beta \cos \gamma$. (IIT-JEE 1980)
5. For all θ in $[0, \pi/2]$ show that $\cos(\sin \theta) \geq \sin(\cos \theta)$. (IIT-JEE 1981)
6. Without using tables, prove that $(\sin 12^\circ)(\sin 48^\circ)(\sin 54^\circ) = 1/8$. (IIT-JEE 1980)
7. Show that $16 \cos \left(\frac{2\pi}{15} \right) \cos \left(\frac{4\pi}{15} \right) \cos \left(\frac{8\pi}{15} \right) \cos \left(\frac{16\pi}{15} \right) = 1$. (IIT-JEE 1983)

8. Prove that $\tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 8 \cot 8\alpha = \cot \alpha$. (IIT-JEE 1988)
9. ABC is a triangle such that $\sin(2A + B) = \sin(C - A) = -\sin(B + 2C) = 1/2$. If A, B and C are in A.P. determine the values of A, B , and C . (IIT-JEE 1990)
10. Show that the value of $\frac{\tan x}{\tan 3x}$, wherever defined, never lies between $\frac{1}{3}$ and 3 . (IIT-JEE 1992)
11. Prove that $\sum_{k=1}^{n-1} (n-k) \cos \frac{2k\pi}{n} = -\frac{n}{2}$, where $n \geq 3$ is an integer. (IIT-JEE 1997)
12. Find the range of values of t for which $2 \sin t = \frac{1-2x+5x^2}{3x^2-2x-1}$, $t \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$. (IIT-JEE 2005)

Answer Key

JEE Advanced

Single Correct Answer Type

- | | | | | |
|--------|--------|--------|--------|--------|
| 1. d. | 2. a. | 3. b. | 4. b. | 5. c. |
| 6. c. | 7. b. | 8. d. | 9. c. | 10. b. |
| 11. c. | 12. c. | 13. a. | 14. b. | 15. b. |
| 16. d. | | | | |

Multiple Correct Answers Type

- | | | | |
|-------------------|-----------|-----------|-------|
| 1. b. | 2. b., c. | 3. c. | 4. c. |
| 5. a., b., c., d. | 6. a., b. | 7. a., b. | |

Matching Column Type

1. b.

Integer Answer Type

1. 7 2. 2

Fill in the Blanks Type

1. $n = 6$ 2. $-\frac{\sqrt{3}}{2}$ 3. $1/64$ 4. $1/8$ 5. $1/3$
 6. $\pm\sqrt{2}$

True/False Type

1. True

Subjective Type

1. $n\pi + \frac{\pi}{4}$, $n \in \mathbb{Z}$
 2. b. $56/33$
 9. $A = 45^\circ$, $B = 60^\circ$, $C = 75^\circ$

Hints and Solutions

JEE Advanced

Single Correct Answer Type

1. d. From the given relations,

$$m + n = 2 \tan \theta, m - n = 2 \sin \theta.$$

$$\text{Thus, } m^2 - n^2 = 4 \tan \theta \sin \theta$$

$$\text{Also } \sqrt{mn} = 4\sqrt{\tan^2 \theta - \sin^2 \theta} = 4 \sin \theta \tan \theta$$

$$\text{From Eqs. (i) and (ii), we get } m^2 - n^2 = 4\sqrt{mn}.$$

2. a. $\alpha + \beta + \gamma = 2\pi$ or $\frac{\alpha}{2} + \frac{\beta}{2} + \frac{\gamma}{2} = \pi$

$$\therefore \tan\left(\frac{\alpha}{2} + \frac{\beta}{2}\right) = \tan\left(\pi - \frac{\gamma}{2}\right) = -\tan \frac{\gamma}{2}$$

$$\Rightarrow \frac{\tan \alpha/2 + \tan \beta/2}{1 - \tan \alpha/2 \tan \beta/2} = -\tan \gamma/2$$

$$\Rightarrow \tan \alpha/2 + \tan \beta/2 + \tan \gamma/2 = \tan \alpha/2 \tan \beta/2 \tan \gamma/2$$

3. b. $\tan \theta = \frac{-4}{3}$

$$\Rightarrow \theta \in \text{2nd quadrant or 4th quadrant}$$

$$\Rightarrow \sin \theta = \pm 4/5$$

$$\text{If } \theta \in \text{2nd quadrant, } \sin \theta = 4/5$$

$$\text{If } \theta \in \text{4th quadrant, } \sin \theta = -4/5$$

4. b. We have

$$A = \sin^2 \theta + \cos^4 \theta$$

$$= 1 - \cos^2 \theta + \cos^4 \theta$$

$$= 1 + (\cos^4 \theta - \cos^2 \theta)$$

$$= 1 + \left(\cos^2 \theta - \frac{1}{2}\right)^2 - \frac{1}{4}$$

$$= \frac{3}{4} + \left(\cos^2 \theta - \frac{1}{2}\right)^2 \geq \frac{3}{4}$$

$$\text{Now } 0 \leq \cos^2 \theta \leq 1$$

$$\Rightarrow -\frac{1}{2} \leq \cos^2 \theta - \frac{1}{2} \leq \frac{1}{2}$$

$$\Rightarrow 0 \leq \left(\cos^2 \theta - \frac{1}{2}\right)^2 \leq \frac{1}{4}$$

$$\Rightarrow \frac{3}{4} \leq \left(\cos^2 \theta - \frac{1}{2}\right)^2 + \frac{3}{4} \leq 1$$

$$\Rightarrow \frac{3}{4} \leq A \leq 1$$

Alternative Method:

$$\text{We have } A = \sin^2 \theta + \cos^4 \theta$$

$$= 1 - \cos^2 \theta + \cos^4 \theta$$

$$= 1 - \cos^2 \theta (1 - \cos^2 \theta)$$

$$= 1 - \cos^2 \theta \sin^2 \theta$$

$$= 1 - \frac{1}{4} \sin^2 2\theta$$

$$\text{Now } 0 \leq \frac{1}{4} \sin^2 2\theta \leq \frac{1}{4}$$

$$\Rightarrow -\frac{1}{4} \leq -\frac{1}{4} \sin^2 2\theta \leq 0$$

$$\Rightarrow \frac{3}{4} \leq 1 - \frac{1}{4} \sin^2 2\theta \leq 1$$

$$\text{Thus } \frac{3}{4} \leq A \leq 1$$

5. c. We have $\cos \frac{7\pi}{8} = \cos \left(\pi - \frac{\pi}{8}\right) = -\cos \frac{\pi}{8}$

(i)

(ii)

$$\text{and } \cos \frac{5\pi}{8} = \cos \left(\pi - \frac{3\pi}{8}\right) = -\cos \frac{3\pi}{8}$$

$$\therefore \text{L.H.S.} = \left(1 + \cos \frac{\pi}{8}\right) \left(1 + \cos \frac{3\pi}{8}\right) \left(1 - \cos \frac{3\pi}{8}\right) \left(1 - \cos \frac{\pi}{8}\right)$$

$$= \left(1 - \cos^2 \frac{\pi}{8}\right) \left(1 - \cos^2 \frac{3\pi}{8}\right)$$

$$= \sin^2 \frac{\pi}{8} \sin^2 \frac{3\pi}{8}$$

$$= \frac{1}{4} \left(2 \sin^2 \frac{\pi}{8}\right) \left(2 \sin^2 \frac{3\pi}{8}\right)$$

$$= \frac{1}{4} \left[\left(1 - \cos \frac{\pi}{4}\right) \left(1 - \cos \frac{3\pi}{4}\right) \right]$$

$$\left[\because 1 - \cos \theta = 2 \sin^2 \frac{\theta}{2} \right]$$

$$= \frac{1}{4} \left[\left(1 - \frac{1}{\sqrt{2}}\right) \left(1 + \frac{1}{\sqrt{2}}\right) \right] = \frac{1}{4} \left(1 - \frac{1}{2}\right) = \frac{1}{8}$$

6. c. The given expression is

$$\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$$

$$= \frac{\sqrt{3}}{\sin 20^\circ} - \frac{1}{\cos 20^\circ}$$

$$= \frac{\sqrt{3} \cos 20^\circ - \sin 20^\circ}{\sin 20^\circ \cos 20^\circ}$$

$$= 4 \left[\frac{\frac{\sqrt{3}}{2} \cos 20^\circ - \frac{1}{2} \sin 20^\circ}{2 \sin 20^\circ \cos 20^\circ} \right]$$

$$= 4 \left[\frac{\sin 60^\circ \cos 20^\circ - \cos 60^\circ \sin 20^\circ}{\sin 2 \times 20^\circ} \right]$$

$$= \frac{4 \sin(60^\circ - 20^\circ)}{\sin 40^\circ} = \frac{4 \sin 40^\circ}{\sin 40^\circ} = 4$$

$$7. \text{ b. } \sec 2x - \tan 2x = \frac{1 - \sin 2x}{\cos 2x} = \frac{1 - \cos 2\left(\frac{\pi}{4} - x\right)}{\sin 2\left(\frac{\pi}{4} - x\right)}$$

$$= \frac{2 \sin^2\left(\frac{\pi}{4} - x\right)}{2 \sin\left(\frac{\pi}{4} - x\right) \cos\left(\frac{\pi}{4} - x\right)} = \tan\left(\frac{\pi}{4} - x\right)$$

$$8. \text{ d. } \sin \frac{\pi}{2n} + \cos \frac{\pi}{2n} = \frac{\sqrt{n}}{2}$$

$$\text{or } \sin^2 \frac{\pi}{2n} + \cos^2 \frac{\pi}{2n} + 2 \sin \frac{\pi}{2n} \cos \frac{\pi}{2n} = \frac{n}{4}$$

$$\Rightarrow 1 + \sin \frac{\pi}{n} = \frac{n}{4} \quad \text{or} \quad \sin \frac{\pi}{n} = \frac{n-4}{4}$$

For $n = 2$, the given equation is not satisfied.

Considering that $n > 1$ and $n \neq 2$, $0 < \sin \frac{\pi}{n} < 1$

$$\Rightarrow 0 < \frac{n-4}{4} < 1 \quad \Rightarrow 4 < n < 8$$

$$\begin{aligned} 9. \text{ c. } & 3(\sin x - \cos x)^4 + 6(\sin x + \cos x)^2 + 4(\sin^6 x + \cos^6 x) \\ &= 3(1 - \sin 2x)^2 + 6(1 + \sin 2x) + 4[(\sin^2 x + \cos^2 x)^3 - 3 \sin^2 x \cos^2 x (\sin^2 x + \cos^2 x)] \\ &= 3(1 - 2 \sin 2x + \sin^2 2x) + (6 + 6 \sin 2x) \\ &\quad + 4\left[1 - \frac{3}{4} \sin^2 2x\right] \\ &= 13 + 3 \sin^2 2x - 3 \sin^2 2x = 13 \end{aligned}$$

$$10. \text{ b. } \text{Given, } \sec^2 \theta = \frac{4xy}{(x+y)^2}$$

$$\text{Now } \sec^2 \theta \geq 1 \Rightarrow \frac{4xy}{(x+y)^2} \geq 1$$

$$\text{or } (x+y)^2 \leq 4xy$$

$$\text{or } (x+y)^2 - 4xy \leq 0$$

$$\text{or } (x-y)^2 \leq 0$$

But for real values of x and y ,

$$(x-y)^2 \geq 0 \text{ or } (x-y)^2 = 0$$

$$\therefore x = y$$

$$\text{Also } x + y \neq 0 \Rightarrow x \neq 0, y \neq 0$$

$$\begin{aligned} 11. \text{ c. } f(\theta) &= \sin \theta (\sin \theta + \sin 3\theta) \\ &= (\sin \theta + 3 \sin \theta - 4 \sin^3 \theta) \sin \theta \\ &= (4 \sin \theta - 4 \sin^3 \theta) \sin \theta \\ &= 4 \sin^2 \theta (1 - \sin^2 \theta) \\ &= 4 \sin^2 \theta \cos^2 \theta \\ &= (2 \sin \theta \cos \theta)^2 = (\sin 2\theta)^2 \geq 0 \end{aligned}$$

which is true for all θ .

$$12. \text{ c. } \alpha + \beta = \frac{\pi}{2} \quad \text{or} \quad \alpha = \frac{\pi}{2} - \beta$$

$$\Rightarrow \tan \alpha = \cot \beta \quad \text{or} \quad \tan \alpha \tan \beta = 1$$

$$\text{Again, } \beta + \gamma = \alpha \text{ or } \gamma = \alpha - \beta$$

$$\Rightarrow \tan \gamma = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta} = \frac{\tan \alpha - \tan \beta}{2}$$

[using Eq. (i)]

$$\text{or } \tan \alpha = \tan \beta + 2 \tan \gamma$$

13. a. We are given that

$$(\cot \alpha_1)(\cot \alpha_2) \cdots (\cot \alpha_n) = 1$$

$$\Rightarrow (\cos \alpha_1)(\cos \alpha_2) \cdots (\cos \alpha_n) = (\sin \alpha_1)(\sin \alpha_2) \cdots (\sin \alpha_n) \quad (i)$$

$$\text{Let } y = (\cos \alpha_1)(\cos \alpha_2) \cdots (\cos \alpha_n)$$

(to be maximum)

Squaring both sides, we get

$$\begin{aligned} y^2 &= (\cos^2 \alpha_1)(\cos^2 \alpha_2) \cdots (\cos^2 \alpha_n) \\ &= \cos \alpha_1 \sin \alpha_1 \cos \alpha_2 \sin \alpha_2 \cdots \cos \alpha_n \sin \alpha_n \end{aligned}$$

[using Eq. (i)]

$$= \frac{1}{2^n} [\sin 2\alpha_1 \sin 2\alpha_2 \cdots \sin 2\alpha_n]$$

As $0 \leq \alpha_1, \alpha_2, \dots, \alpha_n \leq \pi/2$, we have

$$0 \leq 2\alpha_1, 2\alpha_2, \dots, 2\alpha_n \leq \pi$$

$$\Rightarrow 0 \leq \sin 2\alpha_1, \sin 2\alpha_2, \dots, \sin 2\alpha_n \leq 1$$

$$\therefore y^2 \leq \frac{1}{2^n} \times 1 \quad \text{or} \quad y \leq \frac{1}{2^{n/2}}$$

Therefore, the maximum value of y is $1/2^{n/2}$.

14. b. Given that $\sin \theta = 1/2$ and $\cos \phi = 1/3$, and θ and ϕ are acute angles. Therefore,

$$\theta = \pi/6 \text{ and } 0 < \frac{1}{3} < \frac{1}{2}$$

$$\text{or } \cos \pi/2 < \cos \phi < \cos \pi/3 \text{ or } \pi/3 < \phi < \pi/2$$

$$\therefore \frac{\pi}{3} + \frac{\pi}{6} < \theta + \phi < \frac{\pi}{2} + \frac{\pi}{6}$$

$$\text{or } \frac{\pi}{2} < \theta + \phi < \frac{2\pi}{3} \Rightarrow \theta + \phi \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$$

$$15. \text{ b. } \theta \in \left(0, \frac{\pi}{4}\right) \Rightarrow \tan \theta < 1 \text{ and } \cot \theta > 1.$$

Let $\tan \theta = 1 - x$ and $\cot \theta = 1 + y$, where $x, y > 0$ and are very small, then

$$t_1 = (1-x)^{1-x}, t_2 = (1-x)^{1+y}, t_3 = (1+y)^{1-x}, t_4 = (1+y)^{1+y}$$

Clearly, $t_4 > t_3$ and $t_1 > t_2$. Also $t_3 > t_1$.

Thus, $t_4 > t_3 > t_1 > t_2$.

$$16. \text{ d. } \text{In set } P, \sin \theta = (\sqrt{2} + 1) \cos \theta$$

$$\text{or } \tan \theta = \sqrt{2} + 1$$

$$\text{In set } Q, (\sqrt{2} - 1) \sin \theta = \cos \theta$$

$$\text{or } \tan \theta = \frac{1}{\sqrt{2} - 1} = \sqrt{2} + 1 \Rightarrow P = Q$$

Multiple Correct Answers Type

$$1. \text{ b. } 3 \left[\sin^4 \left(\frac{3}{2} \pi - \alpha \right) + \sin^4 (3\pi + \alpha) \right] - 2 \left[\sin^6 \left(\frac{1}{2} \pi + \alpha \right) + \sin^6 (5\pi - \alpha) \right] \quad (i)$$

$$\begin{aligned}
&= 3(\cos^4 \alpha + \sin^4 \alpha) - 2(\cos^6 \alpha + \sin^6 \alpha) \\
&= 3(1 - 2 \sin^2 \alpha \cos^2 \alpha) - 2[(\sin^2 \alpha + \cos^2 \alpha)^3 \\
&\quad - 3 \sin^2 \alpha \cos^2 \alpha (\sin^2 \alpha + \cos^2 \alpha)] \\
&= 3(1 - 2 \sin^2 \alpha \cos^2 \alpha) - 2[1 - 3 \sin^2 \alpha \cos^2 \alpha] \\
&= 1
\end{aligned}$$

2. b., c.

All are infinite geometric progression with common ratio < 1

$$\begin{aligned}
x &= \frac{1}{1 - \cos^2 \phi} = \frac{1}{\sin^2 \phi}, \quad y = \frac{1}{1 - \sin^2 \phi} = \frac{1}{\cos^2 \phi}, \\
z &= \frac{1}{1 - \cos^2 \phi \sin^2 \phi}
\end{aligned}$$

$$\begin{aligned}
\text{Now, } xy + z &= \frac{1}{\sin^2 \phi \cos^2 \phi} + \frac{1}{1 - \sin^2 \phi \cos^2 \phi} \\
&= \frac{1}{\sin^2 \phi \cos^2 \phi (1 - \sin^2 \phi \cos^2 \phi)}
\end{aligned}$$

$$\text{or } xy + z = xyz \quad (i)$$

$$\text{Clearly, } x + y = \frac{\sin^2 \phi + \cos^2 \phi}{\sin^2 \phi \cos^2 \phi} = xy$$

$$\therefore x + y + z = xyz \quad [\text{using Eq. (i)}]$$

3. c. For $\alpha = -\pi/2$, $\beta = -\pi/2$ and $\gamma = 2\pi$

$$\sin \alpha + \sin \beta + \sin \gamma = -2$$

Hence, the minimum value of the expression is negative.

4. c. We know that $\sin 15^\circ = \frac{\sqrt{3}-1}{2\sqrt{2}}$ (irrational)

$$\cos 15^\circ = \frac{\sqrt{3}+1}{2\sqrt{2}} \text{ (irrational)}$$

$$\begin{aligned}
\sin 15^\circ \cos 15^\circ &= \frac{1}{2} (2 \sin 15^\circ \cos 15^\circ) \\
&= \frac{1}{2} \sin 30^\circ = \frac{1}{4} \text{ (rational)}
\end{aligned}$$

$$\begin{aligned}
\sin 15^\circ \cos 75^\circ &= \sin 15^\circ \cos (90^\circ - 15^\circ) \\
&= \sin 15^\circ \sin 15^\circ = \sin^2 15^\circ \\
&= \frac{1}{2} (1 + \cos 30^\circ) \\
&= \frac{1}{2} \left(1 - \frac{\sqrt{3}}{2} \right) \text{ (irrational)}
\end{aligned}$$

5. a., b., c., d.

$$\begin{aligned}
f_n(\theta) &= \frac{\sin(\theta/2)}{\cos(\theta/2)} \times \left[\frac{2 \cos^2(\theta/2)}{\cos \theta} \frac{2 \cos^2 \theta}{\cos 2\theta} \frac{2 \cos^2 2\theta}{\cos 4\theta} \dots \right] \\
&= \frac{\sin \theta}{\cos \theta} \left[\frac{2 \cos^2 \theta}{\cos 2\theta} \frac{2 \cos^2 2\theta}{\cos 4\theta} \dots \right] \\
&= \frac{\sin 2\theta}{\cos 2\theta} \left[\frac{2 \cos^2 2\theta}{\cos 4\theta} \dots \right] = \tan 2^n \theta
\end{aligned}$$

$$\therefore f_2\left(\frac{\pi}{16}\right) = \tan 4 \cdot \frac{\pi}{16} = \tan \frac{\pi}{4} = 1$$

Similarly, $f_3(\pi/32)$, $f_4(\pi/64)$, and $f_5(\pi/128)$ are found to be $\tan \frac{\pi}{4} = 1$.

6. a., b.

$$\begin{aligned}
\frac{\sin^4 x}{2} + \frac{\cos^4 x}{3} &= \frac{1}{5} \\
3 \sin^4 x + 2(1 - \sin^2 x)^2 &= \frac{6}{5} \\
\Rightarrow 25 \sin^4 x - 20 \sin^2 x + 4 &= 0 \\
\Rightarrow \sin^2 x &= \frac{2}{5} \therefore \cos^2 x = \frac{3}{5} \\
\therefore \tan^2 x &= \frac{2}{3}
\end{aligned}$$

$$\text{and } \frac{\sin^8 x}{8} + \frac{\cos^8 x}{27} = \frac{1}{125}$$

7. a., b.

$$\text{For } \theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$$

$$\text{Let } \cos 4\theta = \frac{1}{3}$$

$$\Rightarrow \cos 2\theta = \pm \sqrt{\frac{1 + \cos 4\theta}{2}} = \pm \sqrt{\frac{2}{3}}$$

$$f\left(\frac{1}{3}\right) = \frac{2}{2 - \sec^2 \theta} = \frac{2 \cos^2 \theta}{2 \cos^2 \theta - 1} = 1 + \frac{1}{\cos 2\theta}$$

$$\Rightarrow f\left(\frac{1}{3}\right) = 1 - \sqrt{\frac{3}{2}} \text{ or } 1 + \sqrt{\frac{3}{2}}$$

Matching Column Type

1. b.

(q) $\cos x + \cos y + \cos z = 0$ and $\sin x + \sin y + \sin z = 0$

$$\cos x + \cos y = -\cos z \quad (1)$$

$$\text{and } \sin x + \sin y = -\sin z \quad (2)$$

Squaring and adding we get,

$$1 + 1 + 2(\cos x \cos y + \sin x \sin y) = 1$$

$$\Rightarrow 2 + 2 \cos(x - y) = 1$$

$$\Rightarrow 2 \cos(x - y) = -1$$

$$\Rightarrow \cos(x - y) = -\frac{1}{2}$$

$$\Rightarrow 2 \cos^2\left(\frac{x - y}{2}\right) - 1 = -\frac{1}{2}$$

$$\Rightarrow \cos\left(\frac{x - y}{2}\right) = \frac{1}{2}$$

$$\begin{aligned}
(r) \quad \cos\left(\frac{\pi}{4} - x\right) \cos 2x + \sin x \sin 2x \sec x \\
&= \cos x \sin 2x \sec x + \cos\left(\frac{\pi}{4} + x\right) \cos 2x \\
&\Rightarrow \left[\cos\left(\frac{\pi}{4} - x\right) - \cos\left(\frac{\pi}{4} + x\right)\right] \cos 2x \\
&= (\cos x \sin 2x \sec x - \sin x \sin 2x \sec x)
\end{aligned}$$

$$\Rightarrow \frac{2}{\sqrt{2}} \sin x \cos 2x = (\cos x - \sin x) \sin 2x \sec x$$

$$\Rightarrow \sqrt{2} \sin x \cos 2x = (\cos x - \sin x) 2 \sin x$$

$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{1}{\cos x + \sin x}$$

$$\Rightarrow x = \frac{\pi}{4} \Rightarrow \sec x = \sec \frac{\pi}{4} = \sqrt{2}$$

Note: Solutions of the remaining parts are given in their respective chapters.

Integer Answer Type

1. (7)

$$\frac{1}{\sin \frac{\pi}{n}} - \frac{1}{\sin \frac{3\pi}{n}} = \frac{1}{\sin \frac{2\pi}{n}}$$

$$\text{or } \frac{\sin \frac{3\pi}{n} - \sin \frac{\pi}{n}}{\sin \frac{\pi}{n} \sin \frac{3\pi}{n}} = \frac{1}{\sin \frac{2\pi}{n}}$$

$$\text{or } \frac{\left(2 \sin \frac{\pi}{n} \cos \frac{2\pi}{n}\right) \sin \frac{2\pi}{n}}{\sin \frac{\pi}{n} \sin \frac{3\pi}{n}} = 1$$

$$\text{or } \sin \frac{4\pi}{n} = \sin \frac{3\pi}{n} \Rightarrow \frac{4\pi}{n} + \frac{3\pi}{n} = \pi \Rightarrow n = 7.$$

2. (2) $\frac{1}{4 \cos^2 \theta + 1 + \frac{3}{2} \sin 2\theta}$

$$= \frac{1}{2[1 + \cos 2\theta] + 1 + \frac{3}{2} \sin 2\theta}$$

$$= \frac{1}{2 \cos 2\theta + \frac{3}{2} \sin 2\theta + 3}$$

Now

$$-\sqrt{2^2 + \left(\frac{3}{2}\right)^2} \leq 2 \cos 2\theta + \frac{3}{2} \sin 2\theta \leq \sqrt{2^2 + \left(\frac{3}{2}\right)^2}$$

$$\text{or } -\frac{5}{2} \leq 2 \cos 2\theta + \frac{3}{2} \sin 2\theta \leq \frac{5}{2}$$

$$\Rightarrow \frac{1}{2} \leq 2 \cos 2\theta + \frac{3}{2} \sin 2\theta + 3 \leq \frac{11}{2}$$

$$\Rightarrow \frac{2}{11} \leq \frac{1}{2 \cos 2\theta + \frac{3}{2} \sin 2\theta + 3} \leq 2$$

Hence, the maximum value is 2.

Fill in the Blanks Type

1. According to the given question, we have to express L.H.S. in the form

$$C_0 + C_1 \cos x + C_2 \cos 2x + \dots + C_n \cos nx$$

Now, $\sin^3 x \sin 3x$

$$= \frac{3 \sin x - \sin 3x}{4} \sin 3x$$

$$= \frac{3 \sin x \sin 3x - \sin^2 3x}{4}$$

$$= \frac{3(\cos 2x - \cos 4x) - (1 - \cos 6x)}{8}$$

Hence, $n = 6$.

2. We know that A.M. $\geq$ G.M.

It implies that the minimum value of A.M. is obtained when A.M. = G.M.

Therefore, the quantities whose A.M. is being taken are equal. Thus,

$$\cos \left(\alpha + \frac{\pi}{2} \right) = \cos \left(\beta + \frac{\pi}{2} \right) = \cos \left(\gamma + \frac{\pi}{2} \right)$$

$$\text{or } \sin \alpha = \sin \beta = \sin \gamma$$


$$\text{Also, } \alpha + \beta + \gamma = 360^\circ$$

$$\text{or } \alpha = \beta = \gamma = 120^\circ = \frac{2\pi}{3}$$

$\therefore$ Minimum value of A.M.

$$\frac{\cos \left(\frac{2\pi}{3} + \frac{\pi}{2} \right) + \cos \left(\frac{2\pi}{3} + \frac{\pi}{2} \right) + \cos \left(\frac{2\pi}{3} + \frac{\pi}{2} \right)}{3}$$

$$= \frac{-3 \sin \frac{2\pi}{3}}{3} = -\frac{\sqrt{3}}{2}$$



3. $\sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \sin \frac{7\pi}{14} \sin \frac{9\pi}{14} \sin \frac{11\pi}{14} \sin \frac{13\pi}{14}$

$$= \sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \sin \frac{\pi}{2} \sin \left(\pi - \frac{5\pi}{14} \right)$$

$$\times \sin \left(\pi - \frac{3\pi}{14} \right) \sin \left(\pi - \frac{\pi}{14} \right)$$

$$= \sin^2 \frac{\pi}{14} \sin^2 \frac{3\pi}{14} \sin^2 \frac{5\pi}{14}$$

$$= \left(\sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \right)^2$$

$$= \left[\cos \left(\frac{\pi}{2} - \frac{\pi}{14} \right) \cos \left(\frac{\pi}{2} - \frac{3\pi}{14} \right) \cos \left(\frac{\pi}{2} - \frac{5\pi}{14} \right) \right]^2$$

$$= \left[\cos \frac{3\pi}{7} \cos \frac{2\pi}{7} \cos \frac{\pi}{7} \right]^2$$

$$= \left[\frac{1}{2 \sin \pi/7} \left\{ 2 \cos \frac{\pi}{7} \sin \frac{\pi}{7} \cos \frac{2\pi}{7} \cos \frac{3\pi}{7} \right\} \right]^2$$

$$\begin{aligned}
&= \left[\frac{1}{2^2 \sin \pi/7} \left\{ 2 \sin \frac{2\pi}{7} \cos \frac{2\pi}{7} \cos \left(\pi - \frac{4\pi}{7} \right) \right\} \right]^2 \\
&= \left[\frac{1}{2^3 \sin \pi/7} \left(2 \sin \frac{4\pi}{7} \cos \frac{4\pi}{7} \right) \right]^2 \\
&= \left(\frac{1}{8 \sin \pi/7} \sin \frac{8\pi}{7} \right)^2 \\
&= \left(\frac{\sin(\pi + \pi/7)}{8 \sin \pi/7} \right)^2 \\
&= \left(\frac{-\sin \pi/7}{8 \sin \pi/7} \right)^2 = \left(\frac{1}{8} \right)^2 = \frac{1}{64}
\end{aligned}$$

$$\begin{aligned}
4. \quad k &= \sin \frac{\pi}{18} \sin \frac{5\pi}{18} \sin \frac{7\pi}{18} \\
&= \cos \left(\frac{\pi}{2} - \frac{\pi}{18} \right) \cos \left(\frac{\pi}{2} - \frac{5\pi}{18} \right) \cos \left(\frac{\pi}{2} - \frac{7\pi}{18} \right) \\
&= \cos \frac{\pi}{9} \cos \frac{2\pi}{9} \cos \frac{4\pi}{9} \\
&= \frac{1}{2^3 \sin \frac{\pi}{9}} \sin \frac{8\pi}{9} = \frac{1}{8 \sin \frac{\pi}{9}} \sin \frac{\pi}{9} = \frac{1}{8} \\
&\quad \left[\because \sin \frac{8\pi}{9} = \sin \left(\pi - \frac{\pi}{9} \right) = \sin \frac{\pi}{9} \right]
\end{aligned}$$

Alternative solution:

$$\begin{aligned}
k &= \sin \frac{\pi}{18} \sin \frac{5\pi}{18} \sin \frac{7\pi}{18} \\
&= \sin 10^\circ \sin 50^\circ \sin 70^\circ \\
&= \sin 10^\circ \sin(60^\circ - 10^\circ) \sin(60^\circ + 10^\circ) \\
&= \frac{1}{4} \sin(3 \times 10^\circ) \\
&= \frac{\sin 30^\circ}{4} \\
&= \frac{1}{8}
\end{aligned}$$

$$\begin{aligned}
5. \quad A + B &= \pi/3 \Rightarrow \tan(A + B) = \sqrt{3} \\
\Rightarrow \frac{\tan A + \tan B}{1 - \tan A \tan B} &= \sqrt{3} \\
\text{or } \frac{\tan A + y}{1 - y} &= \sqrt{3} \quad [\text{where } y = \tan A \tan B] \\
\text{or } \tan^2 A + \sqrt{3}(y - 1) \tan A + y &= 0 \\
\text{For real value of } \tan A, \\
D &\geq 0 \\
\therefore 3(y - 1)^2 - 4y &\geq 0 \\
\text{or } 3y^2 - 10y + 3 &\geq 0 \\
\text{or } (y - 3) \left(y - \frac{1}{3} \right) &\geq 0 \\
\text{or } y &\leq \frac{1}{3} \text{ or } y \geq 3
\end{aligned}$$

But $A, B > 0$ and $A + B = \pi/3 \Rightarrow A, B < \pi/3$
 $\Rightarrow \tan A \tan B < 3$
 Therefore, $y \leq 1/3$, i.e., the maximum value of y is $1/3$.

$$\begin{aligned}
6. \quad \text{We have } \frac{2}{\cos x} &= \frac{1}{\cos(x-y)} + \frac{1}{\cos(x+y)} \\
&= \frac{2 \cos x \cos y}{\cos^2 x - \sin^2 y} \\
\text{or } \cos^2 x - \sin^2 y &= \cos^2 x \cos y \\
\text{or } \cos^2 x (1 - \cos y) &= \sin^2 y \\
\text{or } \cos^2 x \cdot 2 \sin^2 \frac{y}{2} &= 4 \sin^2 \frac{y}{2} \cos^2 \frac{y}{2} \\
\text{or } \cos^2 x &= 2 \cos^2 \frac{y}{2} \\
\text{or } \cos^2 x \sec^2 \frac{y}{2} &= 2 \\
\text{or } \cos x \sec \frac{y}{2} &= \pm \sqrt{2}
\end{aligned}$$

True/False Type

$$\begin{aligned}
1. \quad \text{True. } \tan A &= \frac{1 - \cos B}{\sin B} = \frac{2 \sin^2 \frac{B}{2}}{2 \sin \frac{B}{2} \cos \frac{B}{2}} = \tan \frac{B}{2} \\
\text{Hence, } \tan 2A &= \frac{2 \tan A}{1 - \tan^2 A} = \frac{2 \tan \frac{B}{2}}{1 - \tan^2 \frac{B}{2}} = \tan B
\end{aligned}$$

Therefore, the statement is true.

Subjective Type

$$\begin{aligned}
1. \quad \text{We have } \tan(\alpha + \beta) &= \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \\
&= \frac{\frac{m}{m+1} + \frac{1}{2m+1}}{1 - \frac{m}{m+1} \times \frac{1}{2m+1}} \\
&= \frac{2m^2 + 2m + 1}{2m^2 + 2m + 1} = 1
\end{aligned}$$

$$\Rightarrow \alpha + \beta = n\pi + \pi/4, \text{ where } n \in \mathbb{Z}.$$


$$\begin{aligned}
2. \quad \text{a. To draw the graph of } y &= \frac{1}{\sqrt{2}} (\sin x + \cos x) \text{ from } x = -\frac{\pi}{2} \\
\text{to } x &= \frac{\pi}{2}
\end{aligned}$$

$$y = \frac{1}{\sqrt{2}} (\sin x + \cos x) = \sin \left(x + \frac{\pi}{4} \right)$$

To draw the graph of $y = \sin \left(x + \frac{\pi}{4} \right)$ we first draw the graph of $y = \sin x$ and then shift the graph $\frac{\pi}{4}$ to the left along x -axis.

$$\text{Also for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}, \text{ we have } -\frac{\pi}{4} \leq x + \frac{\pi}{4} \leq \frac{3\pi}{4}$$

Thus graph of $y = \sin\left(x + \frac{\pi}{4}\right)$ is to be considered for $-\frac{\pi}{4} \leq x \leq \frac{3\pi}{4}$



b. We have

$$\begin{aligned}\cos(\alpha + \beta) &= \frac{4}{5} \text{ and } \sin(\alpha - \beta) = \frac{5}{13} \\ \Rightarrow \tan(\alpha + \beta) &= \frac{3}{4} \text{ and } \tan(\alpha - \beta) = \frac{5}{12} \\ \tan 2\alpha &= \tan[(\alpha + \beta) + (\alpha - \beta)] \\ &= \frac{\tan(\alpha - \beta) + \tan(\alpha + \beta)}{1 - \tan(\alpha - \beta)\tan(\alpha + \beta)} \\ &= \frac{\frac{5}{12} + \frac{3}{4}}{1 - \left(\frac{5}{12}\right)\left(\frac{3}{4}\right)} = \frac{56}{33}\end{aligned}$$

3. We have

$$\begin{aligned}5 \cos \theta + 3 \cos\left(\theta + \frac{\pi}{3}\right) + 3 \\ = 5 \cos \theta + 3 \cos \theta \cos \frac{\pi}{3} - 3 \sin \theta \sin \frac{\pi}{3} + 3 \\ = \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta + 3 \\ \text{Now, } -\sqrt{\left(\frac{13}{2}\right)^2 + \left(\frac{3\sqrt{3}}{2}\right)^2} \leq \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta \\ \leq \sqrt{\left(\frac{13}{2}\right)^2 + \left(\frac{3\sqrt{3}}{2}\right)^2} \\ \Rightarrow -7 \leq \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta \leq 7 \\ \Rightarrow -4 \leq \frac{13}{2} \cos \theta - \frac{3\sqrt{3}}{2} \sin \theta + 3 \leq 10\end{aligned}$$

4. Given $\alpha + \beta - \gamma = \pi$ and we want to prove that

$$\begin{aligned}\sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma &= 2 \sin \alpha \sin \beta \cos \gamma \\ \text{L.H.S.} &= \sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma \\ &= \sin^2 \alpha + \sin(\beta + \gamma) \sin(\beta - \gamma) \\ &= \sin^2 \alpha + \sin(\beta + \gamma) \sin(\pi - \alpha) \quad (\because \alpha + \beta - \gamma = \pi) \\ &= \sin^2 \alpha + \sin(\beta + \gamma) \sin \alpha \\ &= \sin \alpha [\sin \alpha + \sin(\beta + \gamma)] \\ &= \sin \alpha [\sin(\pi - (\beta - \gamma)) + \sin(\beta + \gamma)] \\ &= \sin \alpha [\sin(\beta - \gamma) + \sin(\beta + \gamma)] \\ &= \sin \alpha [2 \sin \beta \cos \gamma] \\ &= 2 \sin \alpha \sin \beta \cos \gamma \\ &= \text{R.H.S.}\end{aligned}$$

5. We have,

$$\begin{aligned}\cos \theta + \sin \theta &= \sqrt{2} \left[\frac{1}{\sqrt{2}} \cos \theta + \frac{1}{\sqrt{2}} \sin \theta \right] \\ &= \sqrt{2} \sin(\pi/4 + \theta)\end{aligned}$$

$$\therefore \cos \theta + \sin \theta \leq \sqrt{2} < \pi/2$$

$$\therefore \cos \theta + \sin \theta < \pi/2$$

$$\text{or } \cos \theta < \pi/2 - \sin \theta \quad (i)$$

As $\theta \in [0, \pi/2]$ in which $\sin \theta$ increases. Taking sin on both the sides of Eq. (i), we get

$$\sin(\cos \theta) < \sin(\pi/2 - \sin \theta)$$

$$\text{or } \sin(\cos \theta) < \cos(\sin \theta)$$

$$\text{or } \cos(\sin \theta) > \sin(\cos \theta) \quad (ii)$$

6. L.H.S. = $\sin 12^\circ \sin 48^\circ \sin 54^\circ$

$$= \frac{1}{2} [2 \sin 12^\circ \cos 42^\circ] \sin 54^\circ$$

$$= \frac{1}{2} [\sin 54^\circ - \sin 30^\circ] \sin 54^\circ$$

$$= \frac{1}{2} \left[\sin^2 54^\circ - \frac{1}{2} \sin 54^\circ \right]$$

$$= \frac{1}{4} [2 \sin^2 54^\circ - \sin 54^\circ]$$

$$= \frac{1}{4} \left[2 \left(\frac{1+\sqrt{5}}{4} \right)^2 - \left(\frac{1+\sqrt{5}}{4} \right) \right]$$

$$= \frac{1}{4} \left[2 \left(\frac{1+5+2\sqrt{5}}{16} \right) - \left(\frac{1+\sqrt{5}}{4} \right) \right]$$

$$= \frac{1}{4} \times \frac{1}{8} [6 + 2\sqrt{5} - 2 - 2\sqrt{5}]$$

$$= \frac{1}{32} \times 4 = \frac{1}{8} = \text{R.H.S.}$$

7. We know that

$$\cos A \cos 2A \cos 4A \cdots \cos 2^n A$$

$$= \frac{1}{2^{n+1} \sin A} \sin(2^{n+1} A)$$

$$\therefore 16 \cos \frac{2\pi}{15} \cos 2 \left(\frac{2\pi}{15} \right) \cos 2^2 \left(\frac{2\pi}{15} \right)$$

$$\times \cos 2^3 \left(\frac{2\pi}{15} \right)$$

$$= 16 \frac{\sin(2^4 A)}{2^4 \sin A}$$

(where $A = 2\pi/15$)

$$= 16 \frac{\sin(32\pi/15)}{16 \sin 2\pi/15} = \frac{\sin(32\pi/15)}{\sin(2\pi + 2\pi/15)}$$

$$= \frac{\sin(32\pi/15)}{\sin(32\pi/15)} = 1$$

8. We know that

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$$

$$\text{or } \frac{1 - \tan^2 \alpha}{\tan \alpha} = 2 \cot 2\alpha$$

$$\text{or } \cot \alpha - \tan \alpha = 2 \cot 2\alpha \quad (i)$$

Now we have to prove

$$\tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 8 \cot 8\alpha = \cot \alpha$$

$$\text{L.H.S.} = \tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 4 (2 \cot 8\alpha)$$

$$= \tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 4 (\cot 4\alpha - \tan 4\alpha) \quad [\text{using Eq. (i)}]$$

$$= \tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 4 \cot 4\alpha - 4 \tan 4\alpha$$

$$= \tan \alpha + 2 \tan 2\alpha + 2 (2 \cot 4\alpha)$$

$$= \tan \alpha + 2 \tan 2\alpha + 2 (\cot 2\alpha - \tan 2\alpha)$$

$$[\text{using Eq. (i)}]$$

$$= \tan \alpha + 2 \cot 2\alpha$$

$$= \tan \alpha + (\cot \alpha - \tan \alpha) \quad [\text{using Eq. (i)}]$$

$$= \cot \alpha = \text{R.H.S.}$$

9. Given that in $\triangle ABC$, A , B , and C are in A.P. Therefore,

$$A + B = 2B$$

$$\text{Also } A + B + C = 180^\circ$$

$$\Rightarrow B + 2B = 180^\circ \quad \text{or } B = 60^\circ$$

Also given that

$$\sin(2A + B) = \sin(C - A) = -\sin(B + 2C) = \frac{1}{2}$$

$$\Rightarrow \sin(2A + 60^\circ) = \sin(C - A) = -\sin(60^\circ + 2C) = \frac{1}{2} \quad (i)$$

From Eq. (i), we have

$$\sin(2A + 60^\circ) = \frac{1}{2}$$

$$\Rightarrow 2A + 60^\circ = 150^\circ$$

$$\text{or } 2A = 90^\circ$$

$$\text{or } A = 45^\circ$$

$$\Rightarrow C = \pi - A - B = 75^\circ$$

$$10. \text{ Let } y = \frac{\tan x}{\tan 3x}$$

$$= \frac{\tan x (1 - 3 \tan^2 x)}{3 \tan x - \tan^3 x}$$


$$= \frac{1 - 3 \tan^2 x}{3 - \tan^2 x}$$

$$\text{or } 3y - (\tan^2 x)y = 1 - 3 \tan^2 x$$

$$\text{or } (y - 3) \tan^2 x = 3y - 1$$

$$\text{or } \tan^2 x = \frac{3y - 1}{y - 3}$$

$$\text{or } \frac{3y - 1}{y - 3} \geq 0 \quad (\text{L.H.S. is a perfect square})$$



From the sign scheme $y < \frac{1}{3}$ or $y \geq 3$

Thus, y never lies between $1/3$ and 3 .

$$11. S = \sum_{k=1}^{n-1} (n-k) \cos \frac{2k\pi}{n}$$

$$= (n-1) \cos \frac{2\pi}{n} + (n-2) \cos 2 \frac{2\pi}{n} + \dots$$

$$+ 1 \cos (n-1) \frac{2\pi}{n} \quad (i)$$

We know that $\cos \theta = \cos (2\pi - \theta)$. Replacing each angle θ by $2\pi - \theta$ in Eq. (i), we get

$$S = (n-1) \cos (n-1) \frac{2\pi}{n} + (n-2) \cos (n-2) \frac{2\pi}{n} + \dots + 1 \cos \frac{2\pi}{n} \quad [\text{using Eq. (i)}] \quad (ii)$$

Adding terms having the same angle and taking n common, we have

$$2S = n \left[\cos \frac{2\pi}{n} + \cos \frac{4\pi}{n} + \cos \frac{6\pi}{n} + \dots + \cos (n-1) \frac{2\pi}{n} \right]$$

$$= n \left[\frac{\sin(n-1) \frac{\pi}{n} \cos \frac{2\pi}{n} + (n-1) \frac{2\pi}{n}}{\sin \frac{\pi}{n}} \right]$$

$$= n \left[\frac{\sin \left(\pi - \frac{\pi}{n} \right)}{\sin \frac{\pi}{n}} \cdot \cos \pi \right] = -n$$

$$\therefore S = -n/2$$

12. Given that

$$2 \sin t = \frac{1 - 2x + 5x^2}{3x^2 - 2x - 1}, t \in [-\pi/2, \pi/2]$$

This can be written as

$$(6 \sin t - 5)x^2 + 2(1 - 2 \sin t)x - (1 + 2 \sin t) = 0$$

For the given equation to hold, x should be a real number; therefore, the above equation should have real roots, i.e., $D \geq 0$. Thus,

$$4(1 - 2 \sin t)^2 + 4(6 \sin t - 5)(1 + 2 \sin t) \geq 0$$

$$\text{or } 16 \sin^2 t - 8 \sin t - 4 \geq 0$$

$$\text{or } (4 \sin^2 t - 2 \sin t - 1) \geq 0$$

$$\text{or } 4 \left(\sin t - \frac{\sqrt{5}+1}{4} \right) \left(\sin t + \frac{\sqrt{5}-1}{4} \right) \geq 0$$

$$\text{or } \sin t \leq -\left(\frac{\sqrt{5}-1}{4} \right) \quad \text{or } \sin t \geq \frac{\sqrt{5}+1}{4}$$

$$\text{or } \sin t \leq \sin(-\pi/10) \text{ or } \sin t \geq \sin(3\pi/10)$$

$$\text{or } t \leq -\pi/10 \text{ or } t \geq 3\pi/10$$

(Note that $\sin x$ is an increasing function from $-\pi/2$ to $\pi/2$.)

Therefore, the range of t is $[-\pi/2, -\pi/10] \cup [3\pi/10, \pi/2]$.