CBSE Board Class IX Mathematics Sample Paper 9

Time: 3 hrs

Total Marks: 80

General Instructions:

- **1.** All questions are **compulsory**.
- The question paper consists of 30 questions divided into four sections A, B, C, and D.
 Section A comprises of 6 questions of 1 mark each, Section B comprises of 6 questions of 2 marks each, Section C comprises of 10 questions of 3 marks each and Section D comprises of 8 questions of 4 marks each.
- **3.** Use of calculator is **not** permitted.

Section A (Questions 1 to 6 carry 1 mark each)

1. Rationalise the denominator of
$$\frac{1}{3+\sqrt{2}}$$
.

OR

Multiply $5\sqrt{11}$ and $3\sqrt{11}$.

- 2. Is point (2, 1) lie on a line whose equation is 2x + y = 5?
- 3. In \triangle ABC, m \angle A = x, m \angle B = 2x, m \angle C = 3x. Find the value of m \angle C.
- 4. Point (-2, -5) will lie in which Quadrant?

OR

Write the ordinate of every point on the x-axis.

- 5. If the range of the data is 28 and number of classes is 7, then find the class size of the data?
- 6. O is a center of a Circle and $OR \perp PQ$, distance of a chord PQ of a circle from the center is 12 cm and the length of the chord is 10 cm, what is the length of a radius?

Section B

(Questions 7 to 12 carry 2 marks each)

- 7. Express $0.\overline{975}$ in the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$.
- 8. Factorise: $7\sqrt{2}x^2 10x 4\sqrt{2}$
- 9. The perpendicular distance of a point from the x-axis is 2 units and the perpendicular distance from the y-axis is 5 units. Write the coordinates of such a point if it lies in one of the following quadrants:
 - (i) I Quadrant (ii) II Quadrant (iii) III Quadrant (iv) IV Quadrant
- 10. In the figure, $\angle AOC$ and $\angle BOC$ form a linear pair. If a b = 80°, then find the values of a and b.



OR

The angles of a triangle are in the ratio 4 : 5 : 6. Find the greatest angle.

- 11. The inner diameter of a circular well is 3.5 m. It is 10 m deep. Find
 - i. Its inner curved surface area,
 - ii. The cost of plastering this curved surface at the rate of Rs 40 per m².

OR

Find the volume, total surface area of a cube whose edges measures 20 cm.

12. Find the value of a and b if y = 1 and x = 2 is solution of linear equation ax + by = 3 and 3a - 2b = 1.

Section C

(Questions 13 to 22 carry 3 marks each)

13. Simplify:

$$\frac{\left(25\right)^{\frac{3}{2}} \times \left(343\right)^{\frac{3}{5}}}{16^{\frac{5}{4}} \times 8^{\frac{4}{3}} \times 7^{\frac{3}{5}}}$$

- 14. If the polynomials $x^2 5x 3a$ and $ax^2 5x 7$ leave the same remainder when they are divided by (x 1), then what is the value of a?
- 15. Find the value of $x^3 8y^3 36xy 216$ when x = 2y + 6.
 - 16. In the figure, PQ is a line segment and O is the mid-point of PQ. R and S are on the same side of PQ such that $\angle PQS = \angle QPR$ and $\angle POS = \angle QOR$. Prove that:



- (i) $\Delta PQR \cong \Delta QOS$
- (ii) PR = QS
- 17. Show that the line segments joining the mid points of the opposite sides of a quadrilateral bisect each other.

OR

Show that if the diagonals of a quadrilateral bisect each other at right angles, then it is a rhombus.

18. A company selected 2400 families at random and surveyed them to determine relationship between income level and the number of television sets at home. The information gathered is listed in the table below:

Monthly income	Television per family			
in Rs.	0	1	2	Above 2
Less than 7,000	10	160	25	0
7,000 - 10,000	0	305	27	2
10,000-13,000	1	535	29	1
13,000-16,000	2	469	59	25
16,000 or more	1	579	82	88

If one family is chosen at random find the probability of choosing

- i. A family whose income is 16,000 or more and has more than 2 TV sets
- ii. A family whose income is less than 7,000 and has 2 TV's
- iii. A family whose income is between 10,000 and 13,000 and has 1 TV.

OR

A die is thrown once. Find the probability of getting

- i. A prime number
- ii. A number lying between 2 and 6
- iii. An odd number.
- 19. In the figure, PQ is the diameter of the circle and XY is chord equal to the radius of the circle. PX and QY when extended intersect at point E. Prove that $m \angle PEQ = 60^{\circ}$



OR

Two circles intersect at two points B and C. Through B, two line segments ABD and PBQ are drawn to intersect the circles at A, D and P, Q respectively. Prove that $\angle ACP = \angle QCD$.



20. In the given figure, E is the mid-point of side AD of trapezium ABCD with AB || CD, EF || AB. A line through E parallel to AB meets BC in F. Show that F is the mid-point of BC.



21. Two unbiased dice are tossed 50 times. The sum of integers obtained on the dice is noted below.

Sum	2	3	4	5	6	7	8	9	10	11	12
Frequency	3	9	8	8	4	5	1	3	7	2	0

Find the probability that:

- i. The sum of integers is more than 9.
- ii. The sum of integers is exactly 7.
- iii. The sum of integers is less than 6.
- 22. A wooden article was made by scooping out a hemisphere from each end of a solid cylinder. If the height of the cylinder is 10 cm and its base is 7 cm, find the total surface area of the article.

OR

A sphere, a cylinder and a cone have the same radius. Find the ratio of their curved surface areas.

(SECTION - D)

(Questions 23 to 30 carry 4 marks each)

23. Simplify: $\frac{16 \times 2^{n+1} - 4 \times 2^{n}}{16 \times 2^{n+2} - 2 \times 2^{n+2}}$

OR

Simplify:
$$\frac{3\sqrt{2} - 2\sqrt{3}}{3\sqrt{2} + 2\sqrt{3}} + \frac{\sqrt{12}}{\sqrt{3} - \sqrt{2}}$$

24. Find $x^3 + y^3$ when $x = \frac{1}{3 - 2\sqrt{2}}$ and $y = \frac{1}{3 + 2\sqrt{2}}$.

25.

- (i) Multiply $9x^2 + 25y^2 + 15xy + 12x 20y + 16$ by 3x 5y 4 using suitable identities.
- (ii) Factorise: $a^2 + b^2 2(ab ac + bc)$.
- 26. In the figure, PQRS is a square and SRT is an equilateral triangle. Prove that:



a) $\angle PST = \angle QRT$

b) PT = QT

27. The cost of painting the complete outside surface of a closed cylindrical oil tank at 60 paise per sq dm is Rs. 237.60. The height of the tank is 6 times the radius of the base of the tank. Find its volume corrected to two decimal places.

OR

A dome of a building is in the form of a hemisphere. From inside, it was whitewashed at the cost of Rs. 498.96. If the cost of white-washing is Rs. 2.00 per square metre, find the Inner surface area of the dome Volume of air inside the dome.

28. AC and BD are chords of a circle which bisect each other. Prove that (i) AC and BD are diameters; (ii) ABCD is a rectangle.



The lengths of two parallel chords of a circle are 6 cm and 8 cm. If the smaller chord is at distance 4 cm from the centre, what is the distance of the other chord from the centre?

- 29. Construct $\triangle ABC$ in which m $\angle B = 60^{\circ}$, m $\angle C = 45^{\circ}$ and the perimeter of the triangle is 11 cm.
- 30. The bus fare in a city is as follows: For the first kilometre, the fare is Rs. 8 and for the subsequent distance it is Rs. 5 per kilometre. Taking the distance covered as x km and total fares as Rs. y, write a linear equation for this information and draw its graph.

CBSE Board Class IX Mathematics Sample Paper 9 – Solution

Time: 3 hrs

Total Marks: 80

Section A

1. Here,

$$\frac{1}{3+\sqrt{2}} = \frac{1}{3+\sqrt{2}} \times \frac{3-\sqrt{2}}{3-\sqrt{2}} = \frac{3-\sqrt{2}}{9-2} = \frac{3-\sqrt{2}}{7}$$
OR
$$5\sqrt{11} \times 3\sqrt{11} = 5 \times 3\sqrt{11} \times \sqrt{11} = 15 \times 11 = 165$$

- 2. Substitute x = 2 and y = 1 in the equation 2x + y = 5, we get L.H.S. = 2(2) + 1 = 5 = R.H.S.
 Since L.H.S. = R.H.S.
 ∴ Point (2, 1) lies on a line 2x + y = 5.
- 3. In \triangle ABC,

m ∠A + m ∠B + m ∠C = 180° (sum of the angles of a triangle is 180°)

 $\therefore x + 2x + 3x = 180^{\circ}$

 $\therefore 6x = 180^{\circ}$

∴ x = 30°

Since, m $\angle C = 3x^{\circ}$

- \therefore m \angle C = 3x = 3 \times 30° = 90°
- 4. Here x = -2 and y = -5

Since both are negative,

 \therefore Point (-2, -5) will lie in 3rd Quadrant.

(Since, both the coordinates of any point in the third quadrant are negative)

OR

The point on the x axis is of the form (a, 0). Hence, the ordinate of any point x-axis is always zero.

5. Here, range = 28 and number of classes = 7

$$\therefore \text{ Class Size} = \frac{\text{Range}}{\text{Number of classes}} = \frac{28}{7} = 4$$

- \therefore Class size is the data is 4.
- 6. Given, PQ = 10 cm and OR = 12 cm Since, OR \perp PQ

$$\therefore$$
 PR = $\frac{1}{2}$ PQ

(Perpendicular from the centre of a circle to a chord bisects the chord) Since, OR \perp PQ

- $\therefore \Delta POR$ is a right angled triangle.
- : By Pythagoras theorem,

$$PO^2 = OR^2 + PR^2$$

: PO =
$$\sqrt{(12)^2 + (5)^2} = \sqrt{144 + 25} = \sqrt{169} = 13$$

∴ Radius = 13 cm

Section B

7. Let $x = 0.\overline{975} = 0.975975975$ (1)

On multiplying both sides of equation (1) by 1000, 1000x = 975.975975(2) On subtracting equation (1) from equation (2),

999x = 975

$$\Rightarrow x = \frac{975}{999} = \frac{325}{333}$$

8.
$$7\sqrt{2}x^2 - 10x - 4\sqrt{2}$$

= $7\sqrt{2}x^2 - 14x + 4x - 4\sqrt{2}$
= $7\sqrt{2}(x - \sqrt{2}) + 4(x - \sqrt{2})$
= $(7\sqrt{2} + 4)(x - \sqrt{2})$



9. (i) I quadrant: (5, 2)
(ii) II quadrant: (-5, 2)
(iii) III quadrant: (-5, -2)
(iv) IV quadrant: (5, -2)

10. $a + b = 180^{\circ}$ (Linear pair)....(i) $a - b = 80^{\circ}$ (given)....(ii) Adding (i) and (ii) $2a = 260^{\circ}$ $\Rightarrow a = 130^{\circ}$ $\Rightarrow b = 180^{\circ} - 130^{\circ} = 50^{\circ}$

OR

Let the angles of a triangle are 4x, 5x and 6x.

$$4x + 5x + 6x = 180^{\circ}$$

 $15x = 180^{\circ}$
 $x = 12^{\circ}$
The gradient engls in a triangle is 12 \times (= 72)

The greatest angle in a triangle is $12 \times 6 = 72^{\circ}$

11. Inner radius (r) of circular well =
$$\left(\frac{3.5}{2}\right)$$
 m = 1.75 m

Depth (h) of circular well = 10 m

- i. Inner curved surface area = $2\pi rh = \left(2 \times \frac{22}{7} \times 1.75 \times 10\right) m^2 = (44 \times 0.25 \times 10) m^2$ Inner curved surface area = $2\pi rh = 110 m^2$
- ii. Cost of plastering $1 \text{ m}^2 = \text{Rs. } 40$ Cost of plastering $110 \text{ m}^2 = \text{Rs. } (110 \times 40) = \text{Rs.} 4400.$

OR

Edge of a cube is 20 cm.

Volume of a cube = $a^3 = 20^3 = 8000 \text{ cm}^3$

Total surface area of a cube = $6a^2 = 6 \times 20^2 = 2400 \text{ cm}^2$

12. Substituting the values of x and y in the equation, we get

a(2) + b(1) = 3 $2a + b = 3 \quad \dots(1)$ Multiplying equation (1) by 2, we get $4a + 2b = 6 \quad \dots(2)$ Also, $3a - 2b = 1 \quad \dots(3)$ Adding equations (2) and (3) 7a = 7 a = 1Now, substituting a = 1 in equation (1) we get the value of b = 1.

Section C

13.
$$\frac{(25)^{\frac{3}{2}} \times (343)^{\frac{3}{5}}}{16^{\frac{5}{4}} \times 8^{\frac{4}{3}} \times 7^{\frac{3}{5}}} = \frac{(5^2)^{\frac{3}{2}} \times (7^3)^{\frac{3}{5}}}{(2^4)^{\frac{5}{4}} \times (2^3)^{\frac{4}{3}} \times 7^{\frac{3}{5}}} = \frac{5^3 \times 7^{\frac{9}{5}}}{2^5 \times 2^4 \times 7^{\frac{3}{5}}} = \frac{5^3 \times 7^{\frac{9}{5}}}{2^9 \times 7^{\frac{3}{5}}} = \frac{5^3 \times 7^{\frac{9}{5}}}{2^9 \times 7^{\frac{5}{5}}} = \frac{5^3 \times 7^{\frac{9}{5}}}{2^9}$$

14. Let $p(x) = x^2 - 5x - 3a$ and $q(x) = ax^2 - 5x - 7$.

According to remainder theorem, when the polynomial p(x) is divided by a linear polynomial (x - a), the remainder obtained is p(a).

Remainder obtained when p(x) is divided by x - 1 = p(1)

Remainder obtained when q(x) is divided by x - 1 = q(1)

It is given that p(1) = q(1).

 \Rightarrow (1)² - 5(1) - 3a = a(1)² - 5(1) - 7

 \Rightarrow 1 – 5 – 3a = a – 5 – 7

 $\Rightarrow -4 - 3a = a - 12$ $\Rightarrow 8 = 4a$ $\Rightarrow a = 2$ Thus, the value of a is 2.

15. Given: x = 2y + 6 or x - 2y - 6 = 0We know that if a + b + c = 0, then $a^3 + b^3 + c^3 = 3xyz$ Therefore, we have: $(x)^3 + (-2y)^3 + (-6)^3 = 3x(-2y)(-6)$ Or, $x^3 - 8y^3 - 36xy - 216 = 0$





By ASA congruence rule,

 $\Delta PQR\cong \Delta QOS$

$$\Rightarrow$$
 PR = QS (By CPCT)

17. Let ABCD be a quadrilateral. P, Q, R, and S are the mid points of AB, BC, CD and DA respectively.



Join PQ, QR, RS and SP. Join AC.

In ΔDAC , SR || AC

And SR = $\frac{1}{2}$ AC

(Mid-point theorem)

In \triangle BAC, PQ || AC

And PQ = $\frac{1}{2}$ AC

Clearly, PQ || SR and PQ = SR

In quadrilateral PQRS, one pair of opposite sides is equal and parallel to each other and hence it is a parallelogram.

Now, PR and SQ are the diagonals of PQRS and hence PR and SQ bisect each other.

OR

Let ABCD be a quadrilateral, whose diagonals AC and BD bisect each other at right angle i.e. OA = OC, OB = OD and $m \angle AOB = m \angle BOC = m \angle COD = m \angle AOD = 90^{\circ}$ To prove that ABCD is a rhombus, we need to prove that ABCD is a parallelogram and all sides of ABCD are equal.



Since opposite sides of quadrilateral ABCD are equal, so, we can say that ABCD is a parallelogram. Since all sides of a parallelogram ABCD are equal, so we can say that ABCD is a rhombus.

- 18. i. Total number of families surveyed = 2400 No. of families with income more than 16,000 and having more than 2 TV's = 88 ∴ Required probability = $\frac{88}{2400} = \frac{11}{300}$
 - ii. Total number of families surveyed = 2400Number of families with income less than 7,000 and having 2 TV sets = 25

$$\therefore$$
 Required probability $=\frac{25}{2400}=\frac{1}{96}$

- iii. Total number of families surveyed = 2400
 - Number of families with income between 10,000 and 13,000 having 1 TV set = 535

$$\therefore \text{ Required probability} = \frac{535}{2400} = \frac{107}{480}$$

OR

A dice is rolled once. Find i. Sample space $S = \{1, 2, 3, 4, 5, 6\}$ Number of outcomes n(S) = 6Favourable outcomes for prime number $E = \{2, 3, 5\}$ Number of favourable outcomes n(E) = 3Probability $P(E) = \frac{3}{6} = \frac{1}{2}$. ii. Sample space S = {1, 2, 3, 4, 5, 6} Number of outcomes n(S) = 6Favourable outcomes for number between $2 \& 6 E = \{3, 4, 5\}$ Number of favourable outcomes n(E) = 3Probability P(E)= $\frac{3}{6}=\frac{1}{2}$. iii. Sample space S = {1, 2, 3, 4, 5, 6} Number of outcomes n(S) = 6Favourable outcomes for odd number $E = \{1, 3, 5\}$ Number of favourable outcomes n(E) = 3Probability $P(E) = \frac{3}{6} = \frac{1}{2}$.

19. PQ is the diameter of the circle, chord XY = r (radius of circle)PX and QY extended intersect at a point E.



To prove m \angle PEQ = 60°

XY = OX = OY [radii of a circle]

 $\Rightarrow \Delta XOY$ is an equilateral triangle

 \therefore m \angle XOY = 60°

 $\Rightarrow \Delta XQY = 30^{\circ}$ [Inscribed angle is half of the central angle]

m∠PXQ = 90° [Angle in a semi circle]

 $m \angle QXE = 180^{\circ} - m \angle PXQ = 90^{\circ}$ [Linear pair]

In ΔXEQ,

 $m \angle XEQ = 180^{\circ} - (m \angle EXQ + \angle EQX)$ (Angle sum property)

 $m \angle XEQ = 180^{\circ} - (90^{\circ} + 30^{\circ}) = 60^{\circ}$

 \Rightarrow m∠PEQ = 60°

OR

Given, two circles intersect at two points B and C.

Through B, two line segments ABD and PBQ are drawn to intersect the circles at A, D and P, Q respectively.

To Prove: $\angle ACP = \angle QCD$.

Construction: Join chords AP and DQ



Proof: Consider chord AP, \angle PBA = \angle ACP (angles in the same segment) ... (1) Consider chord DQ, \angle DBQ = \angle QCD (angles in the same segment) ... (2) ABD and PBQ are line segments intersecting at B. $\therefore \angle$ PBA = \angle DBQ (vertically opposite angles) ... (3) From equations (1), (2), and (3), we obtain \angle ACP = \angle QCD

20. Given: ABCD is a trapezium. E is the mid-point of AD and AB || CD, EF || AB.To prove: F is the mid-point of BC



Construction: Join AC to intersect EF at point G.

Proof: EF || DC [Given]

∴ EG ∥ DC

Since E is the mid-point of AD.

:. G is the mid-point of AC. [By converse of midpoint theorem]

In $\triangle ABC$, FG || AB

G is the mid-point of AC

- \therefore F is the mid-point of BC.
- 21. Total number of trials = 50
 - i. Number of trials when the sum of integers is more than 9 = 7 + 2 + 0 = 9
 - : P(the sum of integers is more than 9) = $\frac{9}{50}$ = 0.18
 - ii. Number of trials when the sum of integers is exactly 7 = 5

 $\therefore P(\text{the sum of integers is exactly 7}) = \frac{5}{50} = \frac{1}{10} = 0.10$

iii. Number of trails when the sum of integers is less than 6 = 3 + 9 + 8 + 8 = 28

 $\therefore P(\text{the sum of integers is less than } 6) = \frac{28}{50} = \frac{14}{25} = 0.56$

22. Base radius, r = $\frac{7}{2}$ = 3.5 cm

Height of the cylinder, h = 10 cm

Curved surface area of cylinder = $2\pi rh = 2 \times \pi \times 3.5 \times 10 = 70\pi$ Curved surface area of two hemisphere = $2 \times 2\pi r^2 = 4 \times \pi \times 3.5^2 = 49\pi$ Total surface area = $70\pi + 49\pi = 374$ cm²

OR

Let r be the common radius of a sphere, a cone and cylinder. Height of sphere = diameter = 2r Then, the height of the cone=height of cylinder = height of sphere = 2r let l be the slant height of cone= $\sqrt{r^2 + h^2} = \sqrt{r^2 + (2r)^2} = r\sqrt{5}$ $S_1 = curved$ surface area of sphere= $4\pi r^2$ $S_2 = curved$ surface area of cylinder= $2\pi rh=2\pi r \times 2r=4\pi r^2$ $S_3 = curved$ surface area of cone= $\pi rl=\pi r \times r\sqrt{5} = \sqrt{5}\pi r^2$ Ratio of curved surface area as $\therefore S_1 : S_2 : S_3 = 4\pi r^2 : 4\pi r^2 : \sqrt{5}\pi r^2 = 4 : 4 : \sqrt{5}$

Section D

23.
$$\frac{16 \times 2^{n+1} - 4 \times 2^{n}}{16 \times 2^{n+2} - 2 \times 2^{n+2}} = \frac{2^{4} \times 2^{n+1} - 2^{2} \times 2^{n}}{2^{4} \times 2^{n+2} - 2 \times 2^{n+2}}$$
$$= \frac{2^{n+5} - 2^{n+2}}{2^{n+6} - 2^{n+3}}$$
$$= \frac{2^{n+5} - 2^{n+2}}{2 \cdot 2^{n+5} - 2 \cdot 2^{n+2}}$$
$$= \frac{\left(2^{n+5} - 2^{n+2}\right)}{2\left(2^{n+5} - 2^{n+2}\right)}$$
$$= \frac{1}{2}$$

We have,

$$\frac{3\sqrt{2} - 2\sqrt{3}}{3\sqrt{2} + 2\sqrt{3}} + \frac{\sqrt{12}}{\sqrt{3} - \sqrt{2}}$$

$$= \frac{3\sqrt{2} - 2\sqrt{3}}{3\sqrt{2} + 2\sqrt{3}} \times \frac{3\sqrt{2} - 2\sqrt{3}}{3\sqrt{2} - 2\sqrt{3}} + \frac{\sqrt{12}}{\sqrt{3} - \sqrt{2}} \times \frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} + \sqrt{2}}$$

$$= \frac{(3\sqrt{2} - 2\sqrt{3})^2}{(3\sqrt{2} + 2\sqrt{3})(3\sqrt{2} - 2\sqrt{3})} + \frac{\sqrt{12}(\sqrt{3} + \sqrt{2})}{(\sqrt{3} - \sqrt{2})(\sqrt{3} + \sqrt{2})}$$

$$= \frac{(3\sqrt{2})^2 + (2\sqrt{3})^2 - 2 \times 3\sqrt{2} \times 2\sqrt{3}}{(3\sqrt{2})^2 - (2\sqrt{3})^2} + \frac{2\sqrt{3}(\sqrt{3} + \sqrt{2})}{(\sqrt{3})^2 - (\sqrt{2})^2}$$

$$= \frac{18 + 12 - 12\sqrt{6}}{18 - 12} + \frac{6 + 2\sqrt{6}}{3 - 2}$$

$$= \frac{30 - 12\sqrt{6}}{6} + 6 + 2\sqrt{6}$$

$$= \frac{6(5 - 2\sqrt{6})}{6} + 6 + 2\sqrt{6}$$

$$= 11$$

$$\therefore \qquad \frac{3\sqrt{2} - 2\sqrt{3}}{3\sqrt{2} + 2\sqrt{3}} + \frac{\sqrt{12}}{\sqrt{3} - \sqrt{2}} = 11$$

24.

$$x = \frac{1}{3 - 2\sqrt{2}} = \frac{1}{3 - 2\sqrt{2}} \times \frac{3 + 2\sqrt{2}}{3 + 2\sqrt{2}} = \frac{3 + 2\sqrt{2}}{9 - 8} = 3 + 2\sqrt{2}$$
$$y = \frac{1}{3 + 2\sqrt{2}} = \frac{1}{3 + 2\sqrt{2}} \times \frac{3 - 2\sqrt{2}}{3 - 2\sqrt{2}} = \frac{3 - 2\sqrt{2}}{9 - 8} = 3 - 2\sqrt{2}$$

$$\Rightarrow x + y = 3 + 2\sqrt{2} + 3 - 2\sqrt{2} = 6$$
$$xy = \left(3 + 2\sqrt{2}\right)\left(3 - 2\sqrt{2}\right) = 9 - 8 = 1$$

$$x^{3} + y^{3} = (x + y)^{3} - 3xy(x + y)$$
$$= 6^{3} - 3.1.6$$
$$= 216 - 18$$
$$= 198$$

25. (i)
$$(3x - 5y - 4)(9x^2 + 25y^2 + 15xy + 12x - 20y + 16)$$

 $= (3x + (-5y) + (-4))[(3x)^2 + (-5y)^2 + (-4)^2 - (3x)(-5y) - (-5y)(-4) - (3x)(-4)]$
 $= (3x)^3 + (-5y)^3 + (-4)^3 - 3(3x)(-5y)(-4)$
 $[(a + b + c) (a^2 + b^2 + c^2 - ab - bc - ca) = a^3 + b^3 + c^3 - 3abc]$
 $= 27x^3 - 125y^3 - 64 - 180 xy$

(ii)
$$a^2 + b^2 - 2(ab - ac + bc)$$

= $a^2 + b^2 - 2ab + 2ac - 2bc$
= $(a^2 + b^2 - 2ab) + 2c(a - b)$
= $(a - b)^2 + 2c(a - b)$
= $(a - b)(a - b + 2c)$

26. \Box PQRS is a square. \therefore PQ = QR = RS = SP(i)

Also $\angle RSP = \angle SRQ = \angle RQP = \angle SPQ = 90^{\circ}$ (ii)

Also \triangle TSR is equilateral. TS = TR = SR.....(iii)

Also \angle STR = \angle TSR = \angle TRS = 60° TR = QR....from (i) and (ii)

Also $\angle TSP = \angle RSP + \angle TSR$ $\angle TSP = 90^\circ + 60^\circ = 150^\circ$

Similarly $\angle TRQ = 150^{\circ}$

In \triangle TSP and \triangle TRQ, PS = QR.....(:: by (i)) \angle TSP = \angle TRQ.....(:: Both 150°) TS = TR.....(:: by (iii)) $\therefore \triangle$ TSP $\cong \triangle$ TRQ(by SAS criterion) \therefore PT = QT(c.p.c.t)

27.

Let 'r' dm be the radius of the base and 'h' dm be the height of the cylindrical tank. Then, h = 6r (given)

Total surface area = $2\pi r (r + h) = 2\pi r (r + 6r) = 14\pi r^2$

$$\Rightarrow \text{Cost of painting} = \text{Rs. } 14\pi r^2 \times \frac{60}{100} = \text{Rs. } \frac{42}{5}\pi r^2$$

It is given that the cost of painting is Rs. 237.60

$$\therefore \frac{42}{5}\pi r^2 = 237.60$$
$$\Rightarrow \frac{42}{5} \times \frac{22}{7} \times r^2 = 237.60$$
$$\Rightarrow r^2 = 237.60 \times \frac{5}{42} \times \frac{7}{22} = 9 \Rightarrow r = 3 \text{ dm}$$



∴ h = 6r = 18 dm

Volume of the cylinder = $\pi r^2 h$

$$\Rightarrow \text{Volume of the cylinder} = (\pi \times 3 \times 3 \times 18) \text{ dm}^3 = \left(\frac{22}{7} \times 9 \times 18\right) \text{ dm}^2 = 509.14 \text{ dm}^2$$

OR

Cost of white washing 1 m^2 area = Rs. 2

$$\therefore$$
 C.S.A. of the inner side of the dome = $\left(\frac{498.96}{2}\right)$ m² = 249.48 m²

Let inner radius of the hemispherical dome be r. C.S.A. of the inner side of the dome = 249.48 m² $2\pi r^2$ = 249.48 m²

$$\Rightarrow 2 \times \frac{22}{7} \times r^2 = 249.48 \text{ m}^2$$
$$\Rightarrow r^2 = \left(\frac{249.48 \times 7}{2 \times 22}\right) \text{ m}^2 = 39.69 \text{ m}^2$$
$$\Rightarrow r = 6.3 \text{ m}$$

Volume of air inside the dome = Volume of the hemispherical dome = $\frac{2}{3}\pi r^3$

Volume of air inside the dome = $\left[\frac{2}{3} \times \frac{22}{7} \times (6.3)^3\right] \text{m}^3$

Volume of air inside the dome = 523.908 m³

Thus, the volume of air inside the dome is approximately 523.9 m³.

28. In $\triangle AOB$ and $\triangle COD$,

OA = OC	(given)
OB = OD	(given)



$\angle AOB = \angle COD$	(vertically opposite angles)				
$\Delta AOB \cong \Delta COD$	(SAS congruence rule)				
AB = CD	(by CPCT)				
Similarly, we can prove $\triangle AOD$	$\Omega \cong \Delta COB$				
$\therefore AD = CB$	(by CPCT)				
Since in \Box ABCD opposite sides are equal in length. Hence, ABCD is a parallelogram.					
Also, $\angle A = \angle C$	(Opposite angles of a parallelogram are equal)				
But $m \angle A + m \angle C = 180^{\circ}$	(ABCD is a cyclic quadrilateral)				
\Rightarrow m \angle A + m \angle A = 180°					
\Rightarrow 2m \angle A = 180°					
\Rightarrow m \angle A = 90°					
As ABCD is a parallelogram ar	nd one of its interior angles is 90°, so it is a rectangle.				

 $\angle A$ is the angle subtended by chord BD. And as $m \angle A = 90^\circ$, so BD should be the diameter of the circle. Similarly AC is the diameter of the circle.

OR

Let AB and CD be two parallel chords in a circle centered at O. Join OB and OD Distance of smaller chord AB from centre of the circle = 4 cm.



OD OB 5 cm radii of same circle $ND = \frac{CD}{2} = \frac{8}{2} = 4 \text{ cm}$ $ON^{2} + ND^{2} = OD^{2}$ $ON^{2} + (4)^{2} = (5)^{2}$ $ON^{2} = 25 - 16 = 9$ ON = 3So, the distance of the bigger chord from the centre is 3 cm.

- 29. Steps of Construction:
 - 1. Draw a line segment PQ = 11 cm (= AB + BC + CA).
 - 2. At P construct an angle of 60° and at Q an angle of 45°.
 - 3. Bisect these angles. Let the bisectors of these angles intersect at point A.
 - 4. Draw perpendicular bisectors DE of AP to intersect PQ at B and FG of AQ to intersect PQ at C.
 - 5. Join AB and AC

 Δ ABC is the required triangle.



30. Given,

Bus fare for the first kilometer = Rs. 8 Bus fare for the remaining distance = Rs. 5 Total distance covered = x Total fare = y Since the fare for first kilometer = Rs. 8 According to given condition,

Fare for (x - 1) kilometer = 5(x - 1)Therefore, the total fare = 5(x - 1) + 8y = 5(x - 1) + 8 $\Rightarrow y = 5x - 5 + 8$ $\Rightarrow y = 5x + 3$

Therefore, y = 5x + 3 is the required linear equation.

Now the equation is

y = 5x + 3 - (1)

Put the value x = 0 in equation (1)

 $y = 5 \times 0 + 3$

y = 0 + 3 = 3

The solution is (0, 3).

Putting the value x = 1 in equation (1) we get,

 $y = 5 \times 1 + 3$

y = 5 + 3 = 8.

The solution is (1, 8).

Putting the value x = 2 in equation (1)

 $y = 5 \times 2 + 3$

y = 10 + 3 = 13.

The solution is (2, 13).

X	0	1	2
У	3	8	13

Now plot the points (0, 3), (1, 8), and (2, 13) and a draw line passing through these points.

