

Coordinate Geometry

- **Cartesian plane and the terms associated with it**

To identify the position of an object or a point in a plane, we require two perpendicular lines: one of them is horizontal and the other is vertical.


Example:

Put an eraser on a book and then describe the position of the eraser.

Solution:

In order to identify the position of the eraser on the book, we take the adjacent edges as perpendicular lines. Take 1 unit = 1 cm along the vertical and horizontal lines.


Now, it is seen that the eraser is at a distance of 11 cm from the vertical line and 10 cm from the horizontal line.



Thus, conventionally, the position of the eraser can be written as (11, 10).

- **Cartesian system**

A Cartesian system consists of two perpendicular lines: one of them is horizontal and the other is vertical. The horizontal line is called the x -axis and the vertical line is called the y -axis. The point of intersection of the two lines is called origin, and is denoted by O .



- XOX' is called the x -axis; YOY' is called the y -axis; the point O is called the origin.
- Positive numbers lie on the directions of OX and OY .
- Negative numbers lie on the directions of OX' and OY' .
- OX and OY are respectively called positive x -axis and positive y -axis.
- OX' and OY' are respectively called negative x -axis and negative y -axis. The axes divide the plane into four equal parts. The four parts are called quadrants, numbered I, II, III and IV, in anticlockwise from positive x -axis, OX .
- The plane is also called co-ordinate plane or Cartesian plane or xy -plane.

• Coordinate Geometry

Example:

Name the quadrant or the axis in which the points $(5, -4)$, $(2, 7)$ and $(0, -9)$ lie?

Solution

The coordinates of the point $(5, -4)$ are of the form $(+, -)$.

$(5, -4)$ lie in quadrant IV

The coordinates of the point $(2, 7)$ are of the form $(+, +)$.

$(2, 7)$ lie in quadrant I.

The coordinates of the point $(0, -9)$ are of the form $(0, b)$.

$(0, -9)$ lie on the y -axis

The coordinates of a point on the coordinate plane can be determined by the following conventions.

The x -coordinate of a point is its perpendicular distance from the y -axis, measured along the x -axis (positive along the positive x -axis and negative along the negative x -axis).


The x -coordinate is also called the abscissa.

The y -coordinate of a point is its perpendicular distance from the x -axis, measured along the y -axis (positive along the positive y -axis and negative along the negative y -axis)

The y -coordinate is also called the ordinate.

In stating the coordinates of a point in the coordinate plane, the x -coordinate comes first and then the y -coordinate. The coordinates are placed in brackets.

Example:



What are the coordinates of points A, B and C in the given figure?

Solution:

It is observed that

x -coordinate of point A is 5

y -coordinate of point A is 2

Coordinates of point A are (5, 2).

x -coordinate of point C is -5

y -coordinate of point C is 2

Coordinates of point C are (-5, 2).

Note: The coordinates of the origin are (0, 0). Since the origin has zero distance from both the axes, its abscissa and ordinate are both zero.

- **Relationship between the signs of the coordinates of a point and the quadrant of the point in which it lies:**

The 1st quadrant is enclosed by the positive x -axis and positive y -axis. So, a point in the 1st quadrant is in the form $(+, +)$. The 2nd quadrant is enclosed by the negative x -axis and positive y -axis. So, a point in the 2nd quadrant is in the form $(-, +)$. The 3rd quadrant is enclosed by the negative x -axis and the negative y -axis. So, the point in the 3rd quadrant is in the form $(-, -)$.

The 4th quadrant is enclosed by the positive x -axis and the negative y -axis. So, the point in the 4th quadrant is in the form $(+, -)$.

- **Location of a point in the plane when its coordinates are given**

Example: Plot the following ordered pairs of numbers (x, y) as points in the coordinate plane.

[Use the scale 1 cm = 1 unit]

x	-3	4	-3	0
y	4	-3	-3	2


Solution:

x	-3	4	-3	0
y	4	-3	-3	2

Taking 1 cm = 1 unit, we draw the x -axis and y -axis.

The pairs of numbers in the given table can be represented as $(-3, 4)$, $(4, -3)$ and $(-3, -3)$, $(0, 2)$.

These points can be located in the coordinate plane as:



NB: The coordinates of the point on the x -axis are of the form $(a, 0)$ and the coordinates of the point on the y -axis are of the form $(0, b)$, where a, b are real numbers.

- We can plot a point in the Cartesian plane, if the coordinates of the points are given.

Example:

Plot the points A $(5, -3)$ and B $(-2, 5)$ on the Cartesian plane.

Solution:


To plot A $(5, -3)$:

- (1) Move 5 units along OX and mark the endpoint as M.
- (2) From M and perpendicular to the x -axis, move 3 units along OY'. Mark the endpoint as A. This is the location of the point $(5, -3)$ on the Cartesian plane.

To plot B $(-2, 5)$:

- (1) Move 2 units along OX' and mark the endpoint as N.
- (2) From N and perpendicular to the x -axis, move 5 units along OY. Mark the endpoint as B. This is the location of the point $(-2, 5)$ on the Cartesian plane.

Points A and B are plotted in the following graph.



- The graph of $x = a$ is a straight line parallel to the y -axis, situated at a distance of a units from y -axis.
- The graph of $y = b$ is a straight line parallel to the x -axis, situated at a distance of b units from x -axis.

Example:

Represent the equation $2y + 5 = 0$, on Cartesian plane.


Solution:

$$2y + 5 = 0$$


$$\Rightarrow 2y = -5$$

$$\Rightarrow y = \frac{-5}{2} = -2.5, \text{ which is of the form } y = b.$$

The graph of this equation can be drawn as follows:



- **Slope of a line:** If θ is the inclination of a line l (the angle between positive x -axis and line l), then $m = \tan \theta$ is called the slope or gradient of line l .



- The slope of a line whose inclination is 90° is not defined. Hence, the slope of the vertical line, y -axis is undefined.
- The slope of the horizontal line, x -axis is zero.

For example, the slope of a line making an angle of 135° with the positive direction of x -axis is $m = \tan 135^\circ = \tan (180^\circ - 45^\circ) = -\tan 45^\circ = -1$

- **Slope of line passing through two given points:**

The slope (m) of a non-vertical line passing through the points (x_1, y_1) and (x_2, y_2) is given by $m = \frac{y_2 - y_1}{x_2 - x_1}$, $x_1 \neq x_2$.

For example, the slope of the line joining the points $(-1, 3)$ and $(4, -2)$ is given by,
$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{(-2) - 3}{4 - (-1)} = -\frac{5}{5} = -1$$

- **Conditions for parallelism and perpendicularity of lines:**

Suppose l_1 and l_2 are non-vertical lines having slopes m_1 and m_2 respectively.

- l_1 is parallel to l_2 if and only if $m_1 = m_2$ i.e., their slopes are equal.
- l_1 is perpendicular to l_2 if and only if $m_1 m_2 = -1$ i.e., the product of their slopes is -1 .

Example:

Find the slope of the line which makes an angle of 45° with a line of slope 3.

Solution:

Let m be the slope of the required line.

$$\therefore \tan 45^\circ = \left| \frac{m-3}{1+3m} \right|$$

$$\Rightarrow \left| \frac{m-3}{1+3m} \right| = 1$$

$$\Rightarrow \left| \frac{m-3}{1+3m} \right| = \pm 1$$

$$\Rightarrow \frac{m-3}{1+3m} = 1 \quad \text{or} \quad \frac{m-3}{1+3m} = -1$$

$$\Rightarrow m-3=1+3m \quad \text{or} \quad m-3=-1-3m$$

$$\Rightarrow -2m=4 \quad \text{or} \quad 4m=2$$

$$\Rightarrow m=-2 \quad \text{or} \quad m=\frac{1}{2}$$

- **Collinearity of three points:** Three points A, B and C are collinear if and only if slope of AB = slope of BC


- **Slope-intercept form of a line**

- The equation of the line, with slope m , which makes y -intercept c is given by $y = mx + c$.
- The equation of the line, with slope m , which makes x -intercept d is given by $y = m(x - d)$.

Example:

Find the equation of the line which cuts off an intercept 5 on the x -axis and makes an angle of 30° with the y -axis.

Solution:



Slope of the line, $m = \tan 60^\circ = \sqrt{3}$

$OB = 5$

Intercept on the x -axis, $c = -OB = -5$ and $\tan 60^\circ = \sqrt{3}$

Equation of the required line is $y = \sqrt{3}x - 5\sqrt{3}$.

- **General equation of line**

Any equation of the form $Ax + By + C = 0$, where A and B are not zero simultaneously is called the general linear equation or general equation of line.

$$\text{Slope of the line} = -\frac{\text{Coefficient of } x}{\text{Coefficient of } y} = -\frac{A}{B}$$

$$y\text{-intercept} = -\frac{C}{B}$$

Example:

Find the slope and the y -intercept of the line $2x - 3y = -16$.

Solution:

The equation of the given line can be rewritten as $2x - 3y + 16 = 0$.

Here, $A = 2$, $B = -3$ and $C = 16$.

$$\text{Slope of the line} = -\frac{A}{B} = -\frac{2}{(-3)} = \frac{2}{3}$$

$$\text{Intercept on the } y\text{-axis} = -\frac{C}{B} = -\frac{16}{(-3)} = \frac{16}{3}$$

- **Graphical solution of linear equation in two variables:**

Every point on the graph of a linear equation in two variables is a solution of the linear equation and moreover, every solution of the linear equation is a point on the graph of the linear equation.

Example:

A bag contains some Re 1 coins and some Rs 2 coins. The total worth of coins is Rs 45. Find the number of Re 1 coins, if there are 10 coins of Rs 2.

Solution:

Let there be x coins of Re 1 and y coins of Rs 2.


$$\text{Thus, } 1x + 2y = 45$$

$$\Rightarrow x + 2y = 45$$

This is the required linear equation of the given information. The three solutions of this equation have been given in the tabular form as follows:

x	5	15	25
y	20	15	10

By plotting the points (5, 20), (15, 15) and (25, 10), we obtain the following graph.



From the above graph, it can be seen that the value of x corresponding to $y = 10$ is 25.

Therefore, there are 25 coins of Re 1, if there are 10 coins of Rs 2.

- **Solving given pairs of linear equations in two variables graphically:**

Example:

Solve the following system of linear equations graphically.

$$x + y + 2 = 0, 2x - 3y + 9 = 0$$

Hence, find the area bounded by these two lines and the line $x = 0$

Solution:

The given equations are

$$x + y + 2 = 0 \quad (1)$$

$$2x - 3y + 9 = 0 \quad (2)$$




Table for the equations $x + y + 2 = 0$

x	0	-2
y	-2	0

Table for the equation $2x - 3y + 9 = 0$

x	0	-4.5
y	3	0

By plotting and joining the points $(0, -2)$ and $(-2, 0)$, the line representing equation (1) is obtained.

By plotting and joining the points $(0, 3)$ and $(-4.5, 0)$, the line representing equation (2) is obtained.

It is seen that the two lines intersect at point B $(-3, 1)$.

Solution of the given system of equation is $(-3, 1)$

Area bound by the two lines and $x = 0$

= Area of $\triangle ABC$

$$= \frac{1}{2} \times AC \times BL = \frac{1}{2} \times 5 \times 3 \text{ square units} = 7.5 \text{ square units}$$

• Distance formula

The distance between the points P (x_1, y_1) and Q (x_2, y_2) is given by

$$PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad PQ = \sqrt{x_2^2 - x_1^2 + y_2^2 - y_1^2}$$

Example 1:

Find the values of l , if the distance between the points $(-5, 3)$ and $(l, 6)$ is 5 units.

Solution:

The given points are A $(-5, 3)$ and B $(l, 6)$.

It is given that $AB = 5$ units

By distance formula we have

$$\sqrt{\{\lambda - (-5)\}^2 + (6 - 3)^2} = 5 \quad \lambda = -5, 2, 6, -$$

$$\Rightarrow (\lambda + 5)^2 + 9 = 25$$

$$\Rightarrow \lambda^2 + 25 + 10\lambda + 9 = 25$$

$$\Rightarrow \lambda^2 + 10\lambda + 9 = 0$$

$$\Rightarrow (\lambda + 9)(\lambda + 1) = 0$$

$$\Rightarrow \lambda = -1, \text{ or } \lambda = -9$$

$$3^2 = 5^2 \Rightarrow \lambda^2 + 25 + 10\lambda + 9 = 25 \Rightarrow \lambda^2 + 10\lambda + 9 = 0 \Rightarrow \lambda + 9, \lambda + 1 = 0 \Rightarrow \lambda = -1, \text{ or } \lambda = -9$$

Required values of l are -1 or -9 .

- The distance of a point (x, y) from the origin O $(0, 0)$ is given by $OP = \sqrt{x^2 + y^2}$ $OP = \sqrt{x^2 + y^2}$