CHAPTER 18 Polygons and Quadrilaterals

18-1. Parallelograms


A **parallelogram** (\square) is a quadrilateral with two pairs of parallel opposite sides.

A **rhombus** is a parallelogram with four sides of equal measure. The diagonals of a rhombus are perpendicular to each other, and each diagonal of a rhombus bisects a pair of opposite angles. In rhombus *ABCD*, *AB* = *BC* = *CD* = *DA*, *AC* \perp *BD*, $m \angle 1 = m \angle 2 = m \angle 5 = m \angle 6$, and $m \angle 3 = m \angle 4 = m \angle 7 = m \angle 8$.

In $\square ABCD$, $\overline{AB} \parallel \overline{CD}$ and $\overline{BC} \parallel \overline{AD}$.

Properties of Parallelogram

Opposite sides are congruent. Opposite angles are congruent. Consecutive angles are supplementary. The diagonals bisect each other. $\overline{AB} \cong \overline{CD}$ and $\overline{BC} \cong \overline{AD}$ $\angle BAD \cong \angle BCD$ and $\angle ABC \cong \angle ADC$ $m \angle ABC + m \angle BAD = 180$ and $m \angle ADC + m \angle BCD = 180$ AE = CE and BE = DE



E


b

R

Theorem

The area of a parallelogram equals the product of a base and the height to that base. $A = b \cdot h$ The area of a rhombus is half the product of the lengths of its diagonals $(d_1 \text{ and } d_2)$. $A = \frac{1}{2}d_1 \cdot d_2$


Solution \Box x+11=3x-5 $16=2x \Rightarrow x=8$ y+6=2y-4y=10



Opposite sides of \square are \cong .

The diagonals of \square bisect each other.


Example 2 \Box Find the area of parallelogram *PQRS* shown at the right.




$$PT = \frac{1}{2}PQ = \frac{1}{2}(10) = 5$$

$$QT = \sqrt{3}PT = \sqrt{3}(5) = 5\sqrt{3}$$

Area of $PQRS = b \cdot h = 12 \cdot 5\sqrt{3} = 60\sqrt{3}$



Exercise - Parallelograms



18-2. Rectangles, Squares, and Trapezoids

A **rectangle** is a quadrilateral with four right angles. The diagonals of a rectangle are congruent and bisect each other. The diagonals divide the rectangle into four triangles of equal area. In rectangle *ABCD*, AE = BE = CE = DE. Area of $\triangle ABE =$ Area of $\triangle BCE =$ Area of $\triangle CDE =$ Area of $\triangle DAE$

If a quadrilateral is both a rhombus and a rectangle, it is a **square**. A square has four right angles and four congruent sides. The diagonals of a square are congruent and bisect each other.

In square ABCD, AB = BC = CD = DA, $\overline{AB} \perp \overline{BC} \perp \overline{CD} \perp \overline{DA}$, and AE = CE = BE = DE.

A trapezoid is a quadrilateral with exactly one pair of parallel sides.

The **midsegment** of a trapezoid is parallel to the bases and the length of the midsegment is the average of the lengths of the bases. Trapezoid

ABCD with median
$$\overline{MN}$$
, $\overline{AD} \parallel \overline{MN} \parallel \overline{BC}$ and $MN = \frac{1}{2}(b_1 + b_2)$.

If the legs of a trapezoid are congruent, the trapezoid is an **isosceles trapezoid**. The diagonals of an isosceles trapezoid are congruent. Each pair of base angles of an isosceles trapezoid is congruent. For isosceles trapezoid *ABCD* at the right,


AC = BD, $m \angle BAD = m \angle CDA$, and $m \angle ABC = m \angle BCD$.


Theorems - Areas of Rectangle, Square, and Trapezoid

The area of a rectangle is the product of its base and height. The area of a square is the square of the length of a side.

a.

The area of a trapezoid is half the product of its height and sum of the bases.





Solution \Box a. The quadrilateral is a rectangle.

$$h = \frac{1}{2}(16) = 8, \ b = h \cdot \sqrt{3} = 8\sqrt{3}$$
$$A = h \cdot h = 8\sqrt{3} \cdot 8 = 64\sqrt{3}$$


Use the $30^{\circ}-60^{\circ}-90^{\circ} \Delta$ ratio. Area formula for rectangle


b. The quadrilateral is a trapezoid.

$$h \cdot \sqrt{2} = 6\sqrt{2} \implies h = 6$$

 $A = \frac{1}{2}h(b_1 + b_2) = \frac{1}{2}(6)(8 + 12) = 60$


Use the $45^{\circ}-45^{\circ}-90^{\circ} \Delta$ ratio. Area formula for trapezoid












18-3. Regular Polygons

A regular polygon is a convex polygon with all sides congruent and all angles congruent. A polygon is **inscribed in a circle** and the circle is **circumscribed about the polygon** where each vertex of the polygon lies on the circle. The radius of a regular polygon is the distance from the center to a vertex of the polygon. A central angle of a regular polygon is an angle formed by two radii drawn to consecutive vertices. The apothem of a regular polygon is the distance from the center to a side.



Theorems - Angles and Areas of Regular Polygons


The sum of the measures of the interior angles of an *n*-sided polygon is (n-2)180.

The measure of each interior angle of a regular *n*-sided polygon is $\frac{(n-2)180}{n}$.

The sum of the measures of the exterior angles of any polygon is 360.

 $A = \frac{1}{2}ap$ The area of a regular polygon is half the product of the apothem a, and the perimeter p.

Regular Polygons Inscribed in Circles



Solution


$$m \angle PQR = 360 \div 6 = 60$$

$$m \angle PQS = \frac{1}{2}m \angle PQR = \frac{1}{2}(60) = 30$$

$$PS = \frac{1}{2}PR = \frac{1}{2}(4) = 2$$


$$a = \sqrt{3} \cdot PS = 2\sqrt{3}$$

$$A = \frac{1}{2}ap = \frac{1}{2}(2\sqrt{3})(24) = 24\sqrt{3}$$




 $30^{\circ}-60^{\circ}-90^{\circ}$ triangle ratio is used.

$$A = \frac{1}{2}ap$$




Chapter 18 Practice Test



What is the area of the isosceles trapezoid above?


- A) 238
- B) 252
- C) 276
- D) 308



What is the area of rhombus ABCD above?

- A) $20\sqrt{2}$
- B) 25√2
- C) $50\sqrt{2}$
- D) 100√2


3



In the figure above, \overline{EO} is the midsegment of trapezoid *TRAP* and \overline{RP} intersect \overline{EO} at point Z. If RA = 15 and EO = 18, what is the length of \overline{EZ} ? 4

A rectangle has a length that is 6 meters more than twice its width. What is the perimeter of the rectangle if the area of the rectangle is 1,620 square meters?

5




The figure above shows an equilateral triangle with sides of length a and three squares with sides of length a. If the area of the equilateral triangle is $25\sqrt{3}$, what is the sum of the areas of the three squares?

- A) 210
- B) 240
- C) 270
- D) 300


6

The perimeter of a rectangle is 5x and its length is $\frac{3}{2}x$. If the area of the rectangle is 294, what is the value of x?




In the figure above, what is the area of the region *ABCD* ?

- A) $22\sqrt{3} + 30$
- B) $22\sqrt{3} + 36$
- C) $22\sqrt{3} + 42$
- D) $22\sqrt{3} + 48$



In the figure above, *ABCD* is a rectangle and *BCFE* is a square. If AB = 40, BC = 16, and $m \angle AGD = 45$, what is the area of the shaded region?


- A) 240
- B) 248
- C) 256
- D) 264



The figure above shows parallelogram *ABCD*. Which of the following equations represents the area of parallelogram *ABCD*?

- A) $12\cos x^\circ \times 9\sin x^\circ$
- B) $12 \times 9 \tan x^{\circ}$
- C) $12 \times 9 \cos x^{\circ}$
- D) $12 \times 9 \sin x^{\circ}$

10



In the figure above, *ABCDEF* is a regular hexagon with side lengths of 4. *PQRS* is a rectangle. What is the area of the shaded region?


- A) $8\sqrt{3}$
- B) 9√3
- C) $10\sqrt{3}$
- D) $12\sqrt{3}$

Answer Key					
Section 18-1					
1. 4 6. 24	2. 6 7. 240	3.112	4.68	5.70	
Section 18-2					
1.25	2.7.5	3.27	4. C	5. D	
Section 18-3					
1. 120 6. 36	2.5 7.B	3. 15	4. D	5.108	
Chapter 18 Practice Test					
1. C 6. 14	2. B 7. A	3. 10.5 8. C	4. 174 9. D	5. D 10. A	

Answers and Explanations

Section 18-1

1. 4



 $PT = \frac{1}{2}PR$ Diagonals of \Box bisect each
other. $x + 2y = \frac{1}{2}(32) = 16$ SubstitutionST = TQDiagonals of \Box bisect each
other.8x - y = 26Substitution2(8x - y) = 2(26)Multiply each side by 2.16x - 2y = 52Simplify.Add x + 2y = 16 and 16x - 2y = 52.

16x - 2y = 52 $+ \underbrace{x + 2y = 16}_{17x = 68}$ x = 4

2. 6

Substitute 4 for x into the equation x + 2y = 16. 4 + 2y = 16

$$2y = 12$$
$$y = 6$$

3. 112

$m \angle 3 = m \angle 1$	If $\overline{PQ} \parallel \overline{RS}$, Alternate
	Interior $\angle s$ are \cong .
$a^2 - 7 = 6a$	Substitution
$a^2 - 6a - 7 = 0$	Make one side 0.
(a-7)(a+1) = 0	Factor.
a = 7 or $a = -1$	

Discard a = -1, because the measure of angles in parallelogram are positive. $m \angle 1 = 6a = 6(7) = 42$

$$m \angle 2 = 10a = 10(7) = 70$$

 $m \angle PQR = m \angle 1 + m \angle 2$
 $= 42 + 70$
 $= 112$

4. 68

Since $\overline{PQ} \parallel \overline{RS}$, consecutive interior angles are supplementary. Thus, $m \angle PQR + m \angle QRS = 180$.

$$112 + m \angle QRS = 180 \qquad m \angle PQR = 112$$
$$m \angle QRS = 68$$

5. 70

```
m \angle QTR = m \angle PRS + m \angle 3 Exterior Angle Theorem

m \angle 3 = m \angle 1 = 42


m \angle PRS = 4a Given

= 4(7) = 28 a = 7

m \angle QTR = 28 + 42 Substitution

= 70
```

6. 24




$$CE2 + DE2 = CD2$$

$$42 + DE2 = 52$$

$$DE2 = 9$$


$$DE = 3$$

Area of
$$ABCD = \frac{1}{2}AC \cdot BD = \frac{1}{2}(8)(6) = 24$$



Section 18-2


1. 25



Let
$$AD = CD = s$$
.
 $AD^2 + CD^2 = (5\sqrt{2})^2$ Pythagorean Theorem
 $s^2 + s^2 = 50$
 $2s^2 = 50$
 $s^2 = 25$

Area of $ABCD = s^2 = 25$.

2. 7.5




3. 27

Area of rectangle $ABCD = 12 \times 9 = 108$. In a rectangle, diagonals divide the rectangle into four triangles of equal area. Therefore,

Area of
$$\triangle CED = \frac{1}{4}$$
 the area of rectangle *ABCD*
= $\frac{1}{4}(108) = 27$.

4. C




Draw \overline{QT} , which is perpendicular to \overline{PS} , to make triangle PQT, a 30°-60°-90° triangle. In a 30°-60°-90° triangle, the hypotenuse is twice as long as the shorter leg. Therefore,

$$QT = \frac{1}{2}PQ = \frac{1}{2}(8) = 4$$
.

Area of trapezoid $PQRS = \frac{1}{2}(PS + QR) \cdot QT$

$$=\frac{1}{2}(10+3)\cdot 4=26$$

5. D



$$PR^{2} + PQ^{2} = QR^{2}$$
 Pythagorean Theorem

$$PR^{2} + 9^{2} = 15^{2}$$
 Substitution

$$PR^{2} = 15^{2} - 9^{2} = 144$$

$$PR = \sqrt{144} = 12$$

$$12^{2} + RS^{2} = 13^{2}$$
 Pythagorean Theorem

$$RS^{2} = 13^{2} - 12^{2} = 25$$


$$RS = \sqrt{25} = 5$$

Area of trapezoid *PQRS*

$$= \frac{1}{2}(PQ + RS) \cdot PR = \frac{1}{2}(9 + 5) \cdot 12$$

Section 18-3

1. 120



$$m \angle AOB = m \angle BOC = m \angle AOC = \frac{1}{3}(360) = 120$$

2. 5

$$m \angle COD = \frac{1}{2}m \angle AOC = \frac{1}{2}(120) = 60$$

Since triangle *COD* is a $30^{\circ}-60^{\circ}-90^{\circ}$ triangle, the hypotenuse is twice as long as the shorter leg.

Therefore, $OD = \frac{1}{2}CO = \frac{1}{2}(10) = 5$.

3. 15

In a circle all radii are equal in measure. Therefore, AO = BO = CO = 10. BD = BO + OD = 10 + 5 = 15

4. D

In a 30°-60°-90° triangle, the longer leg is $\sqrt{3}$ times as long as the shorter leg. Therefore, $CD = \sqrt{3}OD = 5\sqrt{3}$


$$CD = \sqrt{3}OD = 5\sqrt{3}$$

$$AC = 2CD = 10\sqrt{3}$$

Area of $\triangle ABC$

$$= \frac{1}{2}(AC)(BD) = \frac{1}{2}(10\sqrt{3})(15) = 75\sqrt{3}$$

5. 108



The measure of each interior angle of a regular *n*-sided polygon is $\frac{(n-2)180}{n}$. Therefore,

$$x = \frac{(5-2)180}{5} = 108$$

6. 36

$$m \angle PQR = \frac{360}{5} = 72$$
$$m \angle RQS = \frac{1}{2}m \angle PQR = \frac{1}{2}(72) = 36$$

7. B

In triangle RQS, QR is the hypotenuse and QS is adjacent to $\angle RQS$. Therefore the cosine ratio can be used to find the value of a.

$$\cos \angle RQS = \frac{\text{adjacent to } \angle RQS}{\text{hypotenuse}} = \frac{a}{6}$$

Chapter 18 Practice Test

1. C

$$A \xrightarrow{B} 18 C$$

$$A \xrightarrow{13} 12 F$$

$$E F$$

AE² + BE² = AB²AE² + 12² = 13²


 $AE^2 = 13^2 - 12^2 = 25$ $AE = \sqrt{25} = 5$

Also DF = 5.

AD = AE + EF + DF = 5 + 18 + 5 = 28Area of trapezoid $= \frac{1}{2}(AD + BC) \cdot BF$

$$=\frac{1}{2}(28+18)\cdot 12=276$$





Draw \overline{BF} perpendicular to \overline{AD} to form a 45°-45°-90° triangle.

In a $45^{\circ}-45^{\circ}-90^{\circ}$ triangle, the hypotenuse is $\sqrt{2}$ times as long as a leg. Therefore, $\sqrt{2}BF = AB$.

$$\sqrt{2BF} = 10$$
Substitution

$$BF = \frac{10}{\sqrt{2}} = \frac{10 \cdot \sqrt{2}}{\sqrt{2} \cdot \sqrt{2}} = \frac{10\sqrt{2}}{2} = 5\sqrt{2}$$
Area of rhombus ABCD

$$= \frac{1}{2}AD \cdot BF = \frac{1}{2}(10)(5\sqrt{2}) = 25\sqrt{2}$$

3. 10.5

The length of the midsegment of a trapezoid is the average of the lengths of the bases. Therefore,

$$EO = \frac{1}{2}(TP + RA) .$$

$$18 = \frac{1}{2}(TP + 15)$$
 Substitution

$$2 \times 18 = 2 \times \frac{1}{2}(TP + 15)$$

$$36 = TP + 15$$

$$21 = TP$$

In ΔTRP , $EZ = \frac{1}{2}TP = \frac{1}{2}(21) = 10.5$.

4. 174

Let w = the width of the rectangle in meters, then 2w+6= the length of the rectangle in meters.

Area of rectangle = length \times width

 $=(2w+6)\times w=2w^2+6w$.

Since the area of the rectangle is 1,620 square meters, you can set up the following equation. $2w^2 + 6w = 1620$

$$2w^{2} + 6w - 1620$$

 $2w^{2} + 6w - 1620 = 0$ Make one side 0.
 $2(w^{2} + 3w - 810) = 0$ Common factor is 2.

Use the quadratic formula to solve the equation, $w^2 + 3w - 810 = 0$.

$$w = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$= \frac{-3 \pm \sqrt{3^2 - 4(1)(-810)}}{2(1)}$$
$$= \frac{-3 \pm \sqrt{3249}}{2} = \frac{-3 \pm 57}{2}$$

Since the width is positive, $w = \frac{-3+57}{2} = 27$. The length is 2w+6 = 2(27)+6 = 60. The perimeter of the rectangle is 2(length + width) = 2(60 + 27) = 174

5. D

Area of an equilateral triangle with side length of $a = \frac{\sqrt{3}}{4}a^2$. Since the area of the equilateral triangle is given as $25\sqrt{3}$, you can set up the following equation.

$$\frac{\sqrt{3}}{4}a^2 = 25\sqrt{3}$$
$$a^2 = 25\sqrt{3} \cdot \frac{4}{\sqrt{3}} = 100$$

The area of each square is a^2 , or 100, so the sum of the areas of the three squares is 3×100 , or 300.

6. 14

Let w = the width of the rectangle. The perimeter of the rectangle is given as 5x. Perimeter of rectangle = 2(length + width)

$$5x = 2(\frac{3}{2}x + w)$$

$$5x = 3x + 2w$$

$$2x = 2w$$

$$x = w$$


Area of rectangle = length × width = 294

$$\frac{3}{2}x \cdot x = 294$$

2
$$x^{2} = 294 \cdot \frac{2}{3} = 196$$

 $x = \sqrt{196} = 14$

7. A



$$AC^{2} = AB^{2} + BC^{2}$$
 Pythagorean Theorem

$$AC^{2} = 11^{2} + (4\sqrt{3})^{2}$$
 Substitution

$$AC^{2} = 121 + 48 = 169$$

$$AC = \sqrt{169} = 13$$

$$AC^{2} = AD^{2} + CD^{2}$$
 Pythagorean Theorem

$$169 = 12^{2} + CD^{2}$$
 Substitution


$$25 = CD^2$$

5 = CD

The area of region *ABCD* is the sum of the area of $\triangle ABC$ and the area of $\triangle ADC$. Area of the region *ABCD*

$$=\frac{1}{2}(11)(4\sqrt{3}) + \frac{1}{2}(12)(5)$$
$$= 22\sqrt{3} + 30$$

8. C



Since
$$BCFE$$
 is a square,
 $BC = BE = CF = EF = 16$.
 $AE = AB - BE$
 $= 40 - 16 = 24$

Triangle AGD is a 45° - 45° - 90° triangle.


In a 45° - 45° - 90° triangle, the length of the two legs are equal in measure. Therefore,

$$AD = DG = 16$$
.
 $FG = DC - DG - CF$
 $= 40 - 16 - 16 = 8$

Area of the shaded region

$$= \frac{1}{2}(AE + FG) \cdot EF$$
$$= \frac{1}{2}(24 + 8) \cdot 16 = 256$$

9. D




С

Draw \overline{BE} perpendicular to \overline{AD} . In $\triangle ABE$, $\sin x^{\circ} = \frac{BE}{\Omega}$.

Therefore, $BE = 9 \sin x^{\circ}$. Area of parallelogram *ABCD* $= AD \times BE = 12 \times 9 \sin x^{\circ}$ 10. A

-



Draw the diagonals of a regular hexagon, \overline{AD} , \overline{BE} , and \overline{CF} .

BE = BO + OE = 8 and QR = BE = 8

Since *ABCDEF* is a regular hexagon, the diagonals intersect at the center of the hexagon. Let the point of intersection be O. The diagonals divide the hexagon into 6 equilateral triangles with side lengths of 4. Area of each equilateral triangle

with side lengths of 4 is
$$\frac{\sqrt{3}}{4}(4)^2 = 4\sqrt{3}$$
.

Draw \overline{OT} perpendicular to \overline{PS} .

Triangle AOT is a $30^{\circ}-60^{\circ}-90^{\circ}$ triangle.

Therefore,
$$AT = \frac{1}{2}AO = \frac{1}{2}(4) = 2$$
 and
 $OT = \sqrt{3}AT = 2\sqrt{3}$.

In rectangle *PQRS*, $RS = 2OT = 2(2\sqrt{3}) = 4\sqrt{3}$. Area of rectangle *PQRS* = $QR \times RS$

 $= 8 \times 4\sqrt{3} = 32\sqrt{3}$. Area of regular hexagon *ABCDEF* $= 6 \times \text{area of the equilateral triangle}$ $= 6 \times 4\sqrt{3} = 24\sqrt{3}$ Area of shaded region

= area of rectangle - area of hexagon

$$= 32\sqrt{3} - 24\sqrt{3} = 8\sqrt{3} .$$