

Trigonometrical Equations and Inequations

CONTENTS

2.1	Introduction
2.2	General Solution of Standard Trigonometrical Equations
2.3	General Solution of Some Particular Equations
2.4	General Solution of Square of Trigonometrical Equation
2.5	Solution in the case of Two Equations are given (Simultaneously Solving Equations)
2.6	General Solution of the form $a \cos \theta + b \sin \theta = c$
2.7	Some Particular Equations
2.8	Method for Finding Principal Value
2.9	Some Important Points to be taken in Case of while Solving Trigonometrical Equations
2.10	Miscellaneous Examples
2.11	Periodic Functions

Assignment (Basic & Advance Level)

Answer Sheet



The Greek Mathematician Euclid

The notion of equation is indeed a very old one. The babylonians knew of quadratic equations some 4000 years ago. The Greek Mathematician Euclid (300 B.C.) and Aryabhatta (476 A.D.) gives several Quadratic equations while solving Quadratic and Trigonometrical equations. It was Sridhara an Indian Mathematician around 900 A.D. who was the first to give an algebraic solution of the general equation, and an important treatment by factoring is found in Harriot's work in approximately 1613 A.D., Both these results are very helpful in solving Trigonometrical equations.

Trigonometrical Equations and Inequations

2.1 Introduction

An equation involving one or more trigonometrical ratio of an unknown angle is called a trigonometrical equation *i.e.*, $\sin x + \cos 2x = 1$, $(1 - \tan \theta)(1 + \sin 2\theta) = 1 + \tan \theta$; $\left| \sec\left(\theta + \frac{\pi}{4}\right) \right| = 2$ etc.

A trigonometric equation is different from a trigonometrical identities. An identity is satisfied for every value of the unknown angle *e.g.*, $\cos^2 x = 1 - \sin^2 x$ is true $\forall x \in R$ while a trigonometric equation is satisfied for some particular values of the unknown angle.

(1) **Roots of trigonometrical equation** : The value of unknown angle (a variable quantity) which satisfies the given equation is called the root of an equation *e.g.*, $\cos \theta = \frac{1}{2}$, the root is $\theta = 60^\circ$ or $\theta = 300^\circ$ because the equation is satisfied if we put $\theta = 60^\circ$ or $\theta = 300^\circ$.

(2) **Solution of trigonometrical equations** : A value of the unknown angle which satisfies the trigonometrical equation is called its solution.

Since all trigonometrical ratios are periodic in nature, generally a trigonometrical equation has more than one solution or an infinite number of solutions. There are basically three types of solutions:


- (i) **Particular solution** : A specific value of unknown angle satisfying the equation.
- (ii) **Principal solution** : Smallest numerical value of the unknown angle satisfying the equation (Numerically smallest particular solution.)
- (iii) **General solution** : Complete set of values of the unknown angle satisfying the equation. It contains all particular solutions as well as principal solutions.

When we have two numerically equal smallest unknown angles, preference is given to the positive value in writing the principal solution. *e.g.*, $\sec \theta = \frac{2}{\sqrt{3}}$ has

$\frac{\pi}{6}, -\frac{\pi}{6}, \frac{11\pi}{6}, -\frac{11\pi}{6}, \frac{23\pi}{6}, -\frac{23\pi}{6}$ etc.

As its particular solutions out of these, the numerically smallest are $\frac{\pi}{6}$ and $-\frac{\pi}{6}$ but the principal solution is taken as

$\theta = \frac{\pi}{6}$ to write the general solution we notice that the position on



P or P' can be obtained by rotation of OP or OP' around O through a complete angle (2π) any number of times and in any direction (clockwise or anticlockwise)

$\therefore$ The general solution is $\theta = 2k\pi \pm \frac{\pi}{6}, k \in \mathbb{Z}$.

2.2 General Solution of Standard Trigonometrical Equations

(1) **General solution of the equation $\sin \theta = \sin \alpha$:** If $\sin \theta = \sin \alpha$ or $\sin \theta - \sin \alpha = 0$

$$\text{or, } 2 \sin\left(\frac{\theta - \alpha}{2}\right) \cos\left(\frac{\theta + \alpha}{2}\right) = 0 \Rightarrow \sin\left(\frac{\theta - \alpha}{2}\right) = 0 \text{ or } \cos\left(\frac{\theta + \alpha}{2}\right) = 0$$

$$\text{or, } \frac{\theta - \alpha}{2} = m\pi; m \in I \text{ or } \frac{\theta + \alpha}{2} = (2m + 1)\frac{\pi}{2}; m \in I$$

$$\Rightarrow \theta = 2m\pi + \alpha; m \in I \text{ or } \theta = (2m + 1)\pi - \alpha; m \in I$$

$$\Rightarrow \theta = (\text{any even multiple of } \pi) + \alpha \text{ or } \theta = (\text{any odd multiple of } \pi) - \alpha$$

$$\boxed{\theta = n\pi + (-1)^n \alpha; n \in I}$$

Note: $\square$ The equation $\operatorname{cosec} \theta = \operatorname{cosec} \alpha$ is equivalent to $\sin \theta = \sin \alpha$. So these two equations having the same general solution.

(2) **General solution of the equation $\cos \theta = \cos \alpha$:** If $\cos \theta = \cos \alpha \Rightarrow \cos \theta - \cos \alpha = 0 \Rightarrow -2 \sin\left(\frac{\theta + \alpha}{2}\right) \sin\left(\frac{\theta - \alpha}{2}\right) = 0 \Rightarrow \sin\left(\frac{\theta + \alpha}{2}\right) = 0$ or $\sin\left(\frac{\theta - \alpha}{2}\right) = 0, \Rightarrow \frac{\theta + \alpha}{2} = n\pi; n \in I$ or $\frac{\theta - \alpha}{2} = n\pi; n \in I$
 $\Rightarrow \theta = 2n\pi - \alpha; n \in I$ or $\theta = 2n\pi + \alpha; n \in I$. for the general solution of $\cos \theta = \cos \alpha$, combine these two result which gives $\boxed{\theta = 2n\pi \pm \alpha; n \in I}$

Note: $\square$ The equation $\sec \theta = \sec \alpha$ is equivalent to $\cos \theta = \cos \alpha$, so the general solution of these two equations are same.

(3) **General solution of the equation $\tan \theta = \tan \alpha$:** If $\tan \theta = \tan \alpha \Rightarrow \frac{\sin \theta}{\cos \theta} = \frac{\sin \alpha}{\cos \alpha}$
 $\Rightarrow \sin \theta \cos \alpha - \cos \theta \sin \alpha = 0 \Rightarrow \sin(\theta - \alpha) = 0 \Rightarrow \theta - \alpha = n\pi; n \in I$ $\boxed{\theta = n\pi + \alpha; n \in I}$

Note: $\square$ The equation $\cot \theta = \cot \alpha$ is equivalent to $\tan \theta = \tan \alpha$ so these two equations having the same general solution.

2.3 General Solution of Some Particular Equations

$$(1) \sin \theta = 0 \Rightarrow \theta = n\pi, \cos \theta = 0 \Rightarrow \theta = (2n + 1)\frac{\pi}{2} \text{ or } n\pi + \frac{\pi}{2}, \tan \theta = 0 \Rightarrow \theta = n\pi$$

$$(2) \sin \theta = 1 \Rightarrow \theta = (4n + 1)\frac{\pi}{2} \text{ or } 2n\pi + \frac{\pi}{2}, \cos \theta = 1 \Rightarrow \theta = 2n\pi, \tan \theta = 1 \Rightarrow \theta = (4n + 1)\frac{\pi}{4} \text{ or } n\pi + \frac{\pi}{4}$$

$$(3) \sin \theta = -1 \Rightarrow \theta = (4n + 3)\frac{\pi}{2} \text{ or } 2n\pi + \frac{3\pi}{2}, \cos \theta = -1 \Rightarrow \theta = (2n + 1)\pi,$$

$$\tan \theta = -1 \Rightarrow \theta = (4n - 1)\frac{\pi}{4} \text{ or } n\pi - \frac{\pi}{4}$$

$$(4) \tan \theta = \text{not defined} \Rightarrow \theta = (2n + 1)\frac{\pi}{2}, \cot \theta = \text{not defined} \Rightarrow \theta = n\pi$$

48 Trigonometrical Equations and

$$\operatorname{cosec} \theta = \text{not defined} \Rightarrow \theta = n\pi, \quad \sec \theta = \text{not defined} \Rightarrow \theta = (2n+1)\frac{\pi}{2}.$$

Important Tips

☞ For equations involving two multiple angles, use multiple and sub-multiple angle formulas, if necessary.

☞ For equations involving more than two multiple angles (i) Apply $C \pm D$ formula to combine the two. (ii) Choose such pairs of multiple angle so that after applying the above formulae we get a common factor in the equation.

Example: 1 If $\sin \theta = \frac{\sqrt{3}}{2}$, then the general value of θ is [MP PET 1988]

- (a) $2n\pi \pm \frac{\pi}{6}$ (b) $2n\pi \pm \frac{\pi}{3}$ (c) $n\pi + (-1)^n \frac{\pi}{3}$ (d) $n\pi + (-1)^n \frac{\pi}{6}$

Solution: (c) $\sin \theta = \frac{\sqrt{3}}{2} \Rightarrow \sin \theta = \sin \frac{\pi}{3} \Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{3}.$

Example: 2 The general solution of $\tan 3x = 1$ is [Karnataka CET 1991]

- (a) $n\pi + \frac{\pi}{4}$ (b) $\frac{n\pi}{3} + \frac{\pi}{12}$ (c) $n\pi$ (d) $n\pi \pm \frac{\pi}{4}$

Solution: (b) $\tan 3x = \tan \frac{\pi}{4} \Rightarrow 3x = n\pi + \frac{\pi}{4} \Rightarrow x = \frac{n\pi}{3} + \frac{\pi}{12}.$

Example: 3 If $\sin 3\theta = \sin \theta$, then the general value of θ is

- (a) $2n\pi, (2n+1)\frac{\pi}{3}$ (b) $n\pi, (2n+1)\frac{\pi}{4}$ (c) $n\pi, (2n+1)\frac{\pi}{3}$ (d) None of these

Solution: (b) $\sin 3\theta = \sin \theta$ or $3\theta = m\pi + (-1)^m \theta$

For (m) even i.e., $m = 2n$ then $\theta = \frac{2n\pi}{2} = n\pi$

And for (m) odd, i.e., $m = (2n+1)$ then $\theta = (2n+1)\frac{\pi}{4}.$

Example: 4 The general solution of $2\sin^2 \theta - 3\sin \theta - 2 = 0$ is [Roorkee 1993]

- (a) $n\pi + (-1)^n \frac{\pi}{2}$ (b) $n\pi + (-1)^n \frac{\pi}{6}$ (c) $n\pi + (-1)^n \frac{7\pi}{6}$ (d) $n\pi - (-1)^n \frac{\pi}{6}$

Solution: (d) $2\sin^2 \theta - 3\sin \theta - 2 = 0 \Rightarrow 2\sin^2 \theta - 4\sin \theta + \sin \theta - 2 = 0 \Rightarrow 2\sin \theta(\sin \theta - 2) + (\sin \theta - 2) = 0$
 $(2\sin \theta + 1)(\sin \theta - 2) = 0$

$\sin \theta = +2$ (which is impossible) $\Rightarrow \therefore \sin \theta = -\frac{1}{2} \Rightarrow \sin \theta = \sin(-\pi/6) \Rightarrow \theta = n\pi - (-1)^n \pi/6.$

Example: 5 The number of solutions of the equation $3\sin^2 x - 7\sin x + 2 = 0$ in the interval $[0, 5\pi]$ is [MP PET 2001; IIT 1998]

- (a) 0 (b) 5 (c) 6 (d) 10

Solution: (c) $3\sin^2 x - 7\sin x + 2 = 0 \Rightarrow 3\sin^2 x - 6\sin x - \sin x + 2 = 0 \Rightarrow (3\sin x - 1)(\sin x - 2) = 0,$

But $\sin x \neq 2$ so $\sin x = \frac{1}{3}$. Hence from 0 to $2\pi = 2$ solution's (one in 1st quadrant and other in 2nd quadrant),

from 2π to $4\pi = 2$ solution's and 4π to $5\pi = 2$ solution's. So total number of solutions = 6.

Example: 6 Number of solutions of the equation $\tan x + \sec x = 2\cos x$, lying in the interval $[0, 2\pi]$ is

[AIEEE 2002; MP PET 2000; IIT 1993]

- (a) 0 (b) 1 (c) 2 (d) 3

Solution: (c) $\frac{\sin x}{\cos x} + \frac{1}{\cos x} = 2 \cos x \Rightarrow \sin x + 1 = 2 \cos^2 x \Rightarrow 2 \sin^2 x + \sin x - 1 = 0 \Rightarrow [2 \sin x - 1][\sin x + 1] = 0$

So, $\sin x = -1$ or $\sin x = \frac{1}{2} \Rightarrow x = \frac{3\pi}{2}$ or $x = \frac{\pi}{6}, \frac{5\pi}{6}$ but $\frac{3\pi}{2}$ does not satisfy the equation, So total number of solutions = 2.

Example: 7 If $\tan \theta + \tan 2\theta + \sqrt{3} \tan \theta \tan 2\theta = \sqrt{3}$, then [UPSEAT 2001]

- (a) $\theta = \frac{(6n+1)\pi}{18}, \forall n \in I$ (b) $\theta = \frac{(6n+1)\pi}{9}, \forall n \in I$ (c) $\theta = \frac{(3n+1)\pi}{9}, \forall n \in I$ (d) None of these

Solution: (c) $\tan \theta + \tan 2\theta + \sqrt{3} \tan \theta \tan 2\theta = \sqrt{3} \Rightarrow \tan \theta + \tan 2\theta = \sqrt{3}(1 - \tan \theta \tan 2\theta)$
 $\Rightarrow \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \tan 2\theta} = \sqrt{3} \Rightarrow \tan 3\theta = \tan\left(\frac{\pi}{3}\right) \Rightarrow 3\theta = n\pi + \frac{\pi}{3} \Rightarrow \theta = \frac{n\pi}{3} + \frac{\pi}{9} = (3n+1)\frac{\pi}{9}.$

Example: 8 The solution of the equation $\cos^2 x - 2 \cos x = 4 \sin x - \sin 2x, (0 \leq x \leq \pi)$ is [DCE 2001]

- (a) $\pi - \cot^{-1}\left(\frac{1}{2}\right)$ (b) $\pi - \tan^{-1}(2)$ (c) $\pi + \tan^{-1}\left(-\frac{1}{2}\right)$ (d) None of these

Solution: (c) Given equation is $\cos^2 x - 2 \cos x = 4 \sin x - \sin 2x$
 $\Rightarrow \cos^2 x - 2 \cos x = 4 \sin x - 2 \sin x \cos x \Rightarrow \cos x(\cos x - 2) = 2 \sin x(2 - \cos x) \Rightarrow (\cos x - 2)(\cos x + 2 \sin x) = 0$
 $\Rightarrow \cos x + 2 \sin x = 0 \quad (\because \cos x \neq 2) \Rightarrow \tan x = -\frac{1}{2} \Rightarrow x = n\pi + \tan^{-1}\left(-\frac{1}{2}\right), n \in I$
 As $0 \leq x \leq \pi$, therefore $x = \pi + \tan^{-1}\left(-\frac{1}{2}\right).$

Example: 9 The solution of the equation $\begin{vmatrix} \cos \theta & \sin \theta & \cos \theta \\ -\sin \theta & \cos \theta & \sin \theta \\ -\cos \theta & -\sin \theta & \cos \theta \end{vmatrix} = 0$, is

- (a) $\theta = n\pi$ (b) $\theta = 2n\pi \pm \frac{\pi}{2}$ (c) $\theta = n\pi \pm (-1)^n \frac{\pi}{4}$ (d) $\theta = 2n\pi \pm \frac{\pi}{4}$

Solution: (b) After solving the determinant $2 \cos \theta = 0 \Rightarrow \theta = 2n\pi \pm \frac{\pi}{2}.$

Example: 10 The general value of θ in the equation $2\sqrt{3} \cos \theta = \tan \theta$, is

- (a) $2n\pi \pm \frac{\pi}{6}$ (b) $2n\pi \pm \frac{\pi}{4}$ (c) $n\pi + (-1)^n \frac{\pi}{3}$ (d) $n\pi + (-1)^n \frac{\pi}{4}$

Solution: (c) $2\sqrt{3} \cos^2 \theta = \sin \theta \Rightarrow 2\sqrt{3} \sin^2 \theta + \sin \theta - 2\sqrt{3} = 0$
 $\Rightarrow \sin \theta = \frac{-1 \pm 7}{4\sqrt{3}} \Rightarrow \sin \theta = \frac{-8}{4\sqrt{3}}$ (impossible) or $\sin \theta = \frac{6}{4\sqrt{3}} = \frac{\sqrt{3}}{2} \Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{3}.$

Example: 11 The general value of θ is obtained from the equation $\cos 2\theta = \sin \alpha$ is [MP PET 1996]

- (a) $2\theta = \frac{\pi}{2} - \alpha$ (b) $\theta = 2n\pi \pm \left(\frac{\pi}{2} - \alpha\right)$ (c) $\theta = \frac{n\pi + (-1)^n \alpha}{2}$ (d) $\theta = n\pi \pm \left(\frac{\pi}{4} - \frac{\alpha}{2}\right)$

50 Trigonometrical Equations and

Solution: (d) $\cos 2\theta = \sin \alpha \Rightarrow \cos 2\theta = \cos\left(\frac{\pi}{2} - \alpha\right)$

$$\therefore 2\theta = 2n\pi \pm \left(\frac{\pi}{2} - \alpha\right) \Rightarrow \theta = n\pi \pm \left(\frac{\pi}{4} - \frac{\alpha}{2}\right)$$

Example: 12 If $\begin{vmatrix} \cos(A+B) & -\sin(A+B) & \cos 2B \\ \sin A & \cos A & \sin B \\ -\cos A & \sin A & \cos B \end{vmatrix} = 0$, then $B =$

[EAMCET 2003]

(a) $(2n+1)\frac{\pi}{2}$ (b) $n\pi$ (c) $(2n+1)\pi$ (d) $2n\pi$

Solution: (a) On expanding the determinant $\cos^2(A+B) + \sin^2(A+B) + \cos 2B = 0$

$$1 + \cos 2B = 0 \text{ or } \cos 2B = \cos \pi \text{ or } 2B = 2n\pi + \pi \text{ or } B = (2n+1)\frac{\pi}{2}.$$

Example: 13 If $\cos \theta + \cos 2\theta + \cos 3\theta = 0$, then the general value of θ is

(a) $\theta = 2m\pi \pm \frac{2\pi}{3}$ (b) $\theta = 2m\pi \pm \frac{\pi}{4}$ (c) $\theta = m\pi + (-1)^m \frac{2\pi}{3}$ (d) $\theta = m\pi + (-1)^m \frac{\pi}{3}$

Solution: (a) $\Rightarrow \cos \theta + \cos 2\theta + \cos 3\theta = 0 \Rightarrow (\cos \theta + \cos 3\theta) + \cos 2\theta = 0$

$$\Rightarrow 2\cos 2\theta \cdot \cos \theta + \cos 2\theta = 0 \Rightarrow \cos 2\theta(2\cos \theta + 1) = 0$$

$$\Rightarrow \cos 2\theta = 0 = \cos \frac{\pi}{2} \Rightarrow 2\theta = 2m\pi \pm \pi/2 \Rightarrow \theta = m\pi \pm \frac{\pi}{4} \text{ or } \cos \theta = -\frac{1}{2} = \cos \frac{2\pi}{3} \Rightarrow \theta = 2m\pi \pm \frac{2\pi}{3}.$$

Example: 14 $\sin 6\theta + \sin 4\theta + \sin 2\theta = 0$, then θ equal to

(a) $\frac{n\pi}{4}$ or $n\pi \pm \frac{\pi}{3}$ (b) $\frac{n\pi}{4}$ or $n\pi \pm \frac{\pi}{6}$ (c) $\frac{n\pi}{4}$ or $2n\pi \pm \frac{\pi}{6}$ (d) None of these

Solution: (a) $(\sin 6\theta + \sin 2\theta) + \sin 4\theta = 0 \Rightarrow 2\sin 4\theta \cos 2\theta + \sin 4\theta = 0 \Rightarrow \sin 4\theta(2\cos 2\theta + 1) = 0 \Rightarrow \sin 4\theta = 0 \text{ or } 2\cos 2\theta + 1 = 0$

$$\sin 4\theta = \sin 0 \quad \text{or} \quad \cos 2\theta = \cos \frac{2\pi}{3}$$

$$4\theta = n\pi \quad \text{or} \quad 2\theta = 2n\pi \pm \frac{2\pi}{3}$$

$$\theta = \frac{n\pi}{4} \quad \text{or} \quad \theta = n\pi \pm \frac{\pi}{3}.$$

2.4 General Solution of Square of Trigonometrical Equations

(1) **General solution of $\sin^2 \theta = \sin^2 \alpha$:** If $\sin^2 \theta = \sin^2 \alpha$ or, $2\sin^2 \theta = 2\sin^2 \alpha$ (Both the sides multiply by 2) or, $1 - \cos 2\theta = 1 - \cos 2\alpha$ or, $\cos 2\theta = \cos 2\alpha$, $2\theta = 2n\pi \pm 2\alpha$; $n \in I$, $\boxed{\theta = n\pi \pm \alpha; n \in I}$

(2) **General solution of $\cos^2 \theta = \cos^2 \alpha$:** If $\cos^2 \theta = \cos^2 \alpha$ or, $2\cos^2 \theta = 2\cos^2 \alpha$ (multiply both the side by 2) or, $1 + \cos 2\theta = 1 + \cos 2\alpha$ or, $2\theta = 2n\pi \pm 2\alpha$; $\boxed{\theta = n\pi \pm \alpha; n \in I}$

(3) **General solution of $\tan^2 \theta = \tan^2 \alpha$:** If $\tan^2 \theta = \tan^2 \alpha$ or, $\frac{\tan^2 \theta}{1} = \frac{\tan^2 \alpha}{1}$

Using componendo and dividendo rule, $\frac{\tan^2 \theta + 1}{\tan^2 \theta - 1} = \frac{\tan^2 \alpha + 1}{\tan^2 \alpha - 1}$

$$\text{or } \frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} = \frac{1 + \tan^2 \alpha}{1 - \tan^2 \alpha} \quad \text{or } \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - \tan^2 \alpha}{1 + \tan^2 \alpha} \quad \text{or } \cos 2\theta = \cos 2\alpha, \quad \boxed{\theta = n\pi \pm \alpha; n \in I}$$

Example: 15 General value of θ satisfying the equation $\tan^2 \theta + \sec 2\theta = 1$ is [IIT 1996]

- (a) $m\pi, n\pi + \frac{\pi}{3}$ (b) $m\pi, n\pi \pm \frac{\pi}{3}$ (c) $m\pi, n\pi \pm \frac{\pi}{6}$ (d) None of these

Solution: (b) $\tan^2 \theta + \sec 2\theta = 1 \Rightarrow \tan^2 \theta + \frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} = 1 \Rightarrow \tan^2 \theta - \tan^4 \theta + 1 + \tan^2 \theta = 1 - \tan^2 \theta$

$$\tan^4 \theta - 3\tan^2 \theta = 0 \Rightarrow \tan^2 \theta(\tan^2 \theta - 3) = 0 \Rightarrow \tan^2 \theta = 0 \text{ and } \tan^2 \theta = 3$$

$$\tan^2 \theta = \tan^2 0 \text{ and } \tan^2 \theta = \tan^2 \frac{\pi}{3} \Rightarrow \theta = m\pi \text{ and } \theta = n\pi \pm \frac{\pi}{3}.$$

Example: 16 If $\sec^2 \theta = \frac{4}{3}$, then the general value of θ is [MP PET 1988]

- (a) $2n\pi \pm \frac{\pi}{6}$ (b) $n\pi \pm \frac{\pi}{6}$ (c) $2n\pi \pm \frac{\pi}{3}$ (d) $n\pi \pm \frac{\pi}{3}$

Solution: (b) $\sec^2 \theta = \frac{4}{3} \Rightarrow \cos^2 \theta = \frac{3}{4} = \left(\frac{\sqrt{3}}{2}\right)^2 \Rightarrow \cos^2 \theta = \cos^2 \frac{\pi}{6}$

$$\therefore \theta = n\pi \pm \frac{\pi}{6}.$$

Example: 17 If $2\tan^2 \theta = \sec^2 \theta$, then the general value of θ is [MP PET 1989]

- (a) $n\pi + \frac{\pi}{4}$ (b) $n\pi - \frac{\pi}{4}$ (c) $n\pi \pm \frac{\pi}{4}$ (d) $2n\pi \pm \frac{\pi}{4}$

Solution: (c) $2\tan^2 \theta = 1 + \tan^2 \theta \Rightarrow \tan^2 \theta = 1 \Rightarrow \tan^2 \theta = \tan^2 \frac{\pi}{4}$

$$\therefore \theta = n\pi \pm \frac{\pi}{4}.$$

Example: 18 If $\sin^2 \theta = \frac{1}{4}$, then the most general value of θ is [MP PET 1984, 90; UPSEAT 1973]

- (a) $2n\pi \pm (-1)^n \frac{\pi}{6}$ (b) $\frac{n\pi}{2} \pm (-1)^n \frac{\pi}{6}$ (c) $n\pi \pm \frac{\pi}{6}$ (d) $2n\pi \pm \frac{\pi}{6}$

Solution: (c) $\sin^2 \theta = \frac{1}{4} = \left(\frac{1}{2}\right)^2 \Rightarrow \sin^2 \theta = \sin^2 \frac{\pi}{6}$

$$\therefore \theta = n\pi \pm \frac{\pi}{6}.$$

Example: 19 If $\frac{1 - \cos 2\theta}{1 + \cos 2\theta} = 3$, then the general value of θ is

52 Trigonometrical Equations and

(a) $2n\pi \pm \frac{\pi}{6}$

(b) $n\pi \pm \frac{\pi}{6}$

(c) $2n\pi + \frac{\pi}{3}$

(d) $n\pi \pm \frac{\pi}{3}$

Solution: (d) $\frac{1 - \cos 2\theta}{1 + \cos 2\theta} = 3 \Rightarrow \frac{2 \sin^2 \theta}{2 \cos^2 \theta} = 3 \Rightarrow \tan^2 \theta = 3 = (\sqrt{3})^2 \Rightarrow \tan^2 \theta = \tan^2 \frac{\pi}{3}$

$$\theta = n\pi \pm \frac{\pi}{3}.$$

2.5 Solutions in the Case of Two Equations are given (Simultaneously Solving Equation).

We may divide the problem into two categories. (1) Two equations in one 'unknown' satisfied simultaneously. (2) Two equations in two 'unknowns' satisfied simultaneously.

(1) **Two equations is one 'unknown'** : Two equations are given and we have to find the values of variables θ which may satisfy with the given equations.

(i) $\cos \theta = \cos \alpha$ and $\sin \theta = \sin \alpha$, so the common solution is $\boxed{\theta = 2n\pi + \alpha, n \in I}$

(ii) $\sin \theta = \sin \alpha$ and $\tan \theta = \tan \alpha$, so the common solution is $\boxed{\theta = 2n\pi + \alpha, n \in I}$

(iii) $\cos \theta = \cos \alpha$ and $\tan \theta = \tan \alpha$, so the common solution is $\boxed{\theta = 2n\pi + \alpha, n \in I}$

Example: 20 The most general value of θ satisfying the equation $\tan \theta = -1$ and $\cos \theta = \frac{1}{\sqrt{2}}$ is

[MP PET 2003; UPSEAT 2002, 1982; Roorkee

1990]

(a) $n\pi + \frac{7\pi}{4}$

(b) $n\pi + (-1)^n \frac{7\pi}{4}$

(c) $2n\pi + \frac{7\pi}{4}$

(d) None of these

Solution: (c) $\tan \theta = -1 = \tan\left(2\pi - \frac{\pi}{4}\right)$ and $\cos \theta = \frac{1}{\sqrt{2}} = \cos\left(2\pi - \frac{\pi}{4}\right)$

Hence, general value is $2n\pi + \left(2\pi - \frac{\pi}{4}\right) = 2n\pi + \frac{7\pi}{4}$.

Example: 21 The most general value of θ which will satisfy both the equations $\sin \theta = \frac{-1}{2}$ and $\tan \theta = \frac{1}{\sqrt{3}}$ is

[MNR 1980; MP PET 1989; DCE 1995]

(a) $n\pi + (-1)^n \frac{\pi}{6}$

(b) $n\pi + \frac{\pi}{6}$

(c) $2n\pi \pm \frac{\pi}{6}$

(d) None of these

Solution: (d) $\sin \theta = \frac{-1}{2} = \sin\left(\frac{-\pi}{6}\right) = \sin\left(\pi + \frac{\pi}{6}\right)$

$\tan \theta = \frac{1}{\sqrt{3}} = \tan\left(\frac{\pi}{6}\right) = \tan\left(\pi + \frac{\pi}{6}\right) \Rightarrow \theta = \left(\pi + \frac{\pi}{6}\right)$

Hence, general value of θ is $2n\pi + \frac{7\pi}{6}$.

(2) **System of equations (Two equations in two unknowns)** : Let $f(\theta, \phi) = 0, g(\theta, \phi) = 0$ be the system of two equations in two unknowns.

Step (i) : Eliminate any one variable, say ϕ . Let $\theta = \alpha$ be one solution.

Step (ii) : Then consider the system $f(\alpha, \phi) = 0, g(\alpha, \phi) = 0$ and use the method of two equations in one variable.

Note : □ It is preferable to solve the system of equations quadrant wise.

Example: 22 If $\left(\frac{\sin \theta}{\sin \phi}\right)^2 = \frac{\tan \theta}{\tan \phi} = 3$, then the value of θ and ϕ are

(a) $\theta = n\pi \pm \frac{\pi}{3}, \phi = n\pi \pm \frac{\pi}{6}$

(b) $\theta = n\pi - \frac{\pi}{3}, \phi = n\pi - \frac{\pi}{6}$

(c) $\theta = n\pi \pm \frac{\pi}{2}, \phi = n\pi + \frac{\pi}{3}$

(d) None of these

Solution: (a) $\left(\frac{\sin \theta}{\sin \phi}\right)^2 = \frac{\tan \theta}{\tan \phi} \Rightarrow \sin \theta \cos \theta = \sin \phi \cos \phi \Rightarrow \sin 2\theta = \sin 2\phi$

$$2\theta = \pi - 2\phi \Rightarrow \theta = \frac{\pi}{2} - \phi$$

But, $\frac{\tan \theta}{\tan \phi} = 3 \Rightarrow \frac{\tan \theta}{\cot \theta} = 3 \Rightarrow \tan^2 \theta = 3 \Rightarrow \theta = n\pi \pm \frac{\pi}{3}$, so that $\phi = n\pi \pm \frac{\pi}{6}$

Trick: Check with the options for $n=0, n=1$.

Example: 23 Solve the system of equations $\sec \theta = \sqrt{2} \sec \phi, \tan \theta = \sqrt{3} \tan \phi$

Solution: Usually students proceed this type of problems in the following way:

Squaring and subtracting, we get $\sec^2 \theta - \tan^2 \theta = 2 \sec^2 \phi - 3 \tan^2 \phi$,

i.e., $2 \tan^2 \phi + 2 - 3 \tan^2 \phi = 1$ or $\tan^2 \phi = 1$ or $\phi = n\pi \pm \frac{\pi}{4}$ (i)

Also we have $\sec^2 \phi - \tan^2 \phi = \frac{\sec^2 \theta}{2} - \frac{\tan^2 \theta}{3}$

which gives $6 = 3 \sec^2 \theta - 2 \tan^2 \theta$ or $\tan^2 \theta = 3$, and so $\theta = m\pi \pm \frac{\pi}{3}$.

Thus solution of this system is $\theta = m\pi \pm \frac{\pi}{3}$ and $\phi = n\pi \pm \frac{\pi}{4}, m, n \in I$ (ii)

Now see the fallacies: $\theta = \frac{\pi}{3}$ and $\phi = -\frac{\pi}{4}$ (from the solution) give $\sec\left(\frac{\pi}{3}\right) = \sqrt{2} \sec\left(-\frac{\pi}{4}\right)$

i.e., $2 = 2$, but $\tan\left(\frac{\pi}{3}\right) = \sqrt{3} \tan\left(-\frac{\pi}{4}\right)$ give $\sqrt{3} = -\sqrt{3}$.

Thus solution given in (ii) consists many extraneous (absurd) solutions. The simple reason for this is quite obvious. (ii) consists of solutions of following four systems:

$\sec \theta = \sqrt{2} \sec \phi, \tan \theta = \sqrt{3} \tan \phi$ (iii)

$\sec \theta = \sqrt{2} \sec \phi, \tan \theta = -\sqrt{3} \tan \phi$ (iv)

$\sec \theta = -\sqrt{2} \sec \phi, \tan \theta = \sqrt{3} \tan \phi$ (v)

and $\sec \theta = -\sqrt{2} \sec \phi, \tan \theta = -\sqrt{3} \tan \phi$ (vi)

While we have to find the values which satisfy (iii). Therefore, we have to verify the solutions and should retain only the valid ones.

Alternative Method : A better method for such type of equations is following:

The given system is $\sec \theta = \sqrt{2} \sec \phi$ (vii)

$\tan \theta = \sqrt{3} \tan \phi$ (viii)

54 Trigonometrical Equations and

$$(vii)^2 - (viii)^2 \text{ gives } \tan^2 \phi = 1 \quad \therefore \phi = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}.$$

Case 1 : $\phi = \frac{\pi}{4}$, the system reduces to $\sec \theta = 2$, $\tan \theta = \sqrt{3}$, so $\theta = \frac{\pi}{3}$.

$$\therefore \theta = 2n\pi + \frac{\pi}{3}, \phi = 2m\pi + \frac{\pi}{4}. \quad \dots\dots\dots(\text{ix})$$

Case 2 : $\phi = \frac{3\pi}{4}$, then we have $\sec \theta = -2$, $\tan \theta = -\sqrt{3}$, so $\theta = \frac{2\pi}{3}$.

$$\text{Thus general solution is } \theta = 2n\pi + \frac{2\pi}{3}, \phi = 2m\pi + \frac{3\pi}{4} \quad \dots\dots\dots(\text{x})$$

Case 3 : $\phi = \frac{5\pi}{4}$ (or can be taken as $-\frac{3\pi}{4}$)

Then $\sec \theta = -2$, $\tan \theta = \sqrt{3}$, $\therefore \theta \in Q_3$, so $\theta = \frac{4\pi}{3}$. Thus $\theta = 2n\pi + \frac{4\pi}{3}$, $\phi = 2m\pi + \frac{5\pi}{4}$ or $\theta = 2n\pi - \frac{2\pi}{3}$,

$$\phi = 2m\pi - \frac{3\pi}{4} \quad \dots\dots\dots(\text{xi})$$

Case 4: $\phi = \frac{7\pi}{4}$ (or $-\frac{\pi}{4}$).

Then $\sec \theta = 2$, $\tan \theta = -\sqrt{3}$, so $\theta = -\frac{\pi}{3}$ and so $\theta = 2n\pi - \frac{\pi}{3}$, $\phi = 2m\pi - \frac{\pi}{4}$. $\dots\dots\dots(\text{xii})$

Hence, the required solutions are given as $(\theta, \phi) = \left(2n\pi + \frac{\pi}{3}, 2m\pi + \frac{\pi}{4}\right); \left(2n\pi + \frac{2\pi}{3}, 2m\pi + \frac{3\pi}{4}\right),$
 $\left(2n\pi - \frac{2\pi}{3}, 2m\pi - \frac{3\pi}{4}\right), \left(2n\pi - \frac{\pi}{3}, 2m\pi - \frac{\pi}{4}\right).$

Note : $\square$ Do not write the solution as $\theta = 2n\pi + \frac{\pi}{3}, 2n\pi + \frac{2\pi}{3}, \dots\dots\dots, \phi = 2m\pi + \frac{\pi}{4}, \dots\dots\dots$.

2.6 General Solution of the form $a \cos \theta + b \sin \theta = c$

In $a \cos \theta + b \sin \theta = c$, put $a = r \cos \alpha$ and $b = r \sin \alpha$ where $r = \sqrt{a^2 + b^2}$ and $|c| \leq \sqrt{a^2 + b^2}$

$$\text{Then, } r(\cos \alpha \cos \theta + \sin \alpha \sin \theta) = c \Rightarrow \cos(\theta - \alpha) = \frac{c}{\sqrt{a^2 + b^2}} = \cos \beta \text{ (say)} \quad \dots\dots\dots(\text{i})$$

$\Rightarrow \theta - \alpha = 2n\pi \pm \beta \Rightarrow \theta = 2n\pi \pm \beta + \alpha$, where $\tan \alpha = \frac{b}{a}$, is the general solution

Alternatively, putting $a = r \sin \alpha$ and $b = r \cos \alpha$ where $r = \sqrt{a^2 + b^2} \Rightarrow \sin(\theta + \alpha) = \frac{c}{\sqrt{a^2 + b^2}} = \sin \gamma$

(say)

$\Rightarrow \theta + \alpha = n\pi + (-1)^n \gamma \Rightarrow \theta = n\pi + (-1)^n \gamma - \alpha$, where $\tan \alpha = \frac{a}{b}$, is the general solution.

Note : $\square$ $(-\sqrt{a^2 + b^2}) \leq a \cos \theta + b \sin \theta \leq (\sqrt{a^2 + b^2})$

$\square$ The general solution of $a \cos x + b \sin x = c$ is $x = 2n\pi + \tan^{-1}\left(\frac{b}{a}\right) \pm \cos^{-1}\left(\frac{c}{\sqrt{a^2 + b^2}}\right).$

Example: 24 The number of integral values of k , for which the equation $7 \cos x + 5 \sin x = 2k + 1$ has a solution is [IIT Screening]

(a) 4

(b) 8

(c) 10

(d) 12

Solution: (b) $-\sqrt{7^2 + 5^2} \leq (7 \cos x + 5 \sin x) \leq \sqrt{7^2 + 5^2}$

So, for solution $-\sqrt{74} \leq (2k+1) \leq \sqrt{74}$ or $-8.6 \leq 2k+1 \leq 8.6$ or $-9.6 \leq 2k \leq 7.6$ or $-4.8 \leq k \leq 3.8$.

So, integral values of k are $-4, -3, -2, -1, 0, 1, 2, 3$ (eight values)

Example: 25 If $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$, then general value of θ is

[MP PET 2002, 1991; UPSEAT 1999]

- (a) $n\pi + (-1)^n \frac{\pi}{4}$ (b) $(-1)^n \frac{\pi}{4} - \frac{\pi}{3}$ (c) $n\pi + \frac{\pi}{4} - \frac{\pi}{3}$ (d) $n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{3}$

Solution: (d) $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2} \Rightarrow \frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta = \frac{1}{\sqrt{2}}$

$$\Rightarrow \sin \frac{\pi}{3} \cos \theta + \cos \frac{\pi}{3} \sin \theta = \frac{1}{\sqrt{2}} \Rightarrow \sin \left(\theta + \frac{\pi}{3} \right) = \sin \frac{\pi}{4} \Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{3}.$$

Example: 26 If $\sin \theta + \cos \theta = 1$, then the general value of θ is

[Karnataka CET 2002; DCE 2000; MNR 1987; IIT 1981]

- (a) $2n\pi$ (b) $n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{4}$ (c) $2n\pi + \frac{\pi}{2}$ (d) None of these

Solution: $\sin \theta + \cos \theta = 1 \Rightarrow \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta = \frac{1}{\sqrt{2}} \quad \left[\text{Dividing by } \sqrt{1^2 + 1^2} = \sqrt{2} \right]$

$$\Rightarrow \sin \left(\theta + \frac{\pi}{4} \right) = \frac{1}{\sqrt{2}} = \sin \frac{\pi}{4} \Rightarrow \theta + \frac{\pi}{4} = n\pi + (-1)^n \frac{\pi}{4} \Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{4}.$$

Example: 27 The equation $\sqrt{3} \sin x + \cos x = 4$ has

[EAMCET 2001]

- (a) Only one solution (b) Two solutions
(c) Infinitely many solutions (d) No solution

Solution: (d) Given equation is $\sqrt{3} \sin x + \cos x = 4$ which is of the form $a \sin x + b \cos x = c$ with $a = \sqrt{3}, b = 1, c = 4$

Here $a^2 + b^2 = 3 + 1 = 4 < c^2$, Therefore the given equation has no solution.

Example: 28 The general solution of the equation $(\sqrt{3} - 1) \sin \theta + (\sqrt{3} + 1) \cos \theta = 2$ is

[Roorkee 1992]

- (a) $2n\pi \pm \frac{\pi}{4} + \frac{\pi}{12}$ (b) $n\pi + (-1)^n \frac{\pi}{4} + \frac{\pi}{12}$ (c) $2n\pi \pm \frac{\pi}{4} - \frac{\pi}{12}$ (d) $n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{12}$

Solution: (a) $(\sqrt{3} - 1) \sin \theta + (\sqrt{3} + 1) \cos \theta = 2$

Divided by $\sqrt{(\sqrt{3} - 1)^2 + (\sqrt{3} + 1)^2} = 2\sqrt{2}$ in both sides,

$$\text{We get, } \frac{(\sqrt{3} - 1)}{2\sqrt{2}} \sin \theta + \frac{(\sqrt{3} + 1)}{2\sqrt{2}} \cos \theta = \frac{2}{2\sqrt{2}}$$

$$\sin \theta \sin 15^\circ + \cos \theta \cos 15^\circ = \frac{1}{\sqrt{2}} \Rightarrow \sin \theta \cdot \sin \frac{\pi}{12} + \cos \theta \cdot \cos \frac{\pi}{12} = \cos \frac{\pi}{4}$$

$$\cos \left(\theta - \frac{\pi}{12} \right) = \cos \frac{\pi}{4} \Rightarrow \theta - \frac{\pi}{12} = 2n\pi \pm \frac{\pi}{4} \Rightarrow \theta = 2n\pi \pm \frac{\pi}{4} + \frac{\pi}{12}.$$

2.7 Some Particular Equations

(1) Equation of the form $a_0 \sin^n x + a_1 \sin^{n-1} x \cos x + a_2 \sin^{n-2} x \cos^2 x + \dots + a_n \cos^n x = 0$: Here $a_0, a_1, \dots, a_n$ are real numbers and the sum of the exponents in $\sin x$ and $\cos x$ in each term is equal to n , are said to be homogeneous with respect to $\sin x$ and $\cos x$. For $\cos x \neq 0$, above equation can be written as, $a_0 \tan^n x + a_1 \tan^{n-1} x + \dots + a_n = 0$.

56 Trigonometrical Equations and

Example: 29 The solution of equation $5 \sin^2 x - 7 \sin x \cos x + 16 \cos^2 x = 4$ is

(a) $x = n\pi + \tan^{-1} 3$ or $x = n\pi + \tan^{-1} 4$

(b) $x = n\pi + \frac{\pi}{6}$ or $x = n\pi + \frac{\pi}{4}$

(c) $x = n\pi$ or $x = n\pi + \frac{\pi}{4}$

(d) None of these

Solution: (a) To solve this kind of equation; we use the fundamental formula trigonometrical identity, $\sin^2 x + \cos^2 x = 1$

writing the equation in the form, $5 \sin^2 x - 7 \sin x \cos x + 16 \cos^2 x = 4 (\sin^2 x + \cos^2 x)$

$$\Rightarrow \sin^2 x - 7 \sin x \cos x + 12 \cos^2 x = 0$$

Dividing by $\cos^2 x$ on both sides we get, $\tan^2 x - 7 \tan x + 12 = 0$

Now it can be factorized as; $(\tan x - 3)(\tan x - 4) = 0 \Rightarrow \tan x = 3, 4$

i.e., $\tan x = \tan(\tan^{-1} 3)$ or $\tan x = \tan(\tan^{-1} 4) \Rightarrow x = n\pi + \tan^{-1} 3$ or $x = n\pi + \tan^{-1} 4$.

(2) **A trigonometric equation of the form $R(\sin kx, \cos nx, \tan mx, \cot lx) = 0$** : Here R is a rational function of the indicated arguments and (k, l, m, n are natural numbers) can be reduced to a rational equation with respect to the arguments $\sin x, \cos x, \tan x$, and $\cot x$ by means of the formulae for trigonometric functions of the sum of angles (in particular, the formulas for double and triple angles) and then reduce equation of the given form to a rational equation with respect to the unknown, $t = \tan \frac{x}{2}$ by means of the formulas,

$$\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}, \cot x = \frac{1 - \tan^2 \frac{x}{2}}{2 \tan \frac{x}{2}}$$

Example: 30 If $(\cos x - \sin x) \left(2 \tan x + \frac{1}{\cos x} \right) + 2 = 0$ then $x =$

(a) $2n\pi \pm \frac{\pi}{3}$

(b) $n\pi \pm \frac{\pi}{3}$

(c) $2n\pi \pm \frac{\pi}{6}$

(d) None of these

Solution: (a) Let $t = \tan \frac{x}{2}$, and using the formula. We get, $\left\{ \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} - \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \right\} \left\{ \frac{4 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} + \frac{1 + \tan^2 \frac{x}{2}}{1 - \tan^2 \frac{x}{2}} \right\} + 2 = 0$

$$\left(\frac{1 - t^2}{1 + t^2} - \frac{2t}{1 + t^2} \right) \left(\frac{4t}{1 - t^2} + \frac{1 + t^2}{1 - t^2} \right) + 2 = 0 \Rightarrow \frac{3t^4 + 6t^3 + 8t^2 - 2t - 3}{(t^2 + 1)(1 - t^2)} = 0$$

Its roots are; $t_1 = \frac{1}{\sqrt{3}}$ and $t_2 = -\frac{1}{\sqrt{3}}$.

Thus the solution of the equation reduces to that of two elementary equations,

$$\tan \frac{x}{2} = \frac{1}{\sqrt{3}}, \tan \frac{x}{2} = -\frac{1}{\sqrt{3}} \Rightarrow \frac{x}{2} = n\pi \pm \frac{\pi}{6} \Rightarrow x = 2n\pi \pm \frac{\pi}{3}, \text{ is required solution.}$$

(3) **Equation of the form $R(\sin x + \cos x, \sin x \cos x) = 0$** : where R is rational function of the arguments in brackets, Put $\sin x + \cos x = t$ (i) and use the following identity:

$$(\sin x + \cos x)^2 = \sin^2 x + \cos^2 x + 2 \sin x \cos x = 1 + 2 \sin x \cos x \Rightarrow \sin x \cos x = \frac{t^2 - 1}{2} \quad \text{.....(ii)}$$

Taking (i) and (ii) into account, we can reduce given equation into; $R\left(t, \frac{t^2 - 1}{2}\right) = 0$.

Similarly, by the substitution $(\sin x - \cos x) = t$, we can reduce the equation of the form; $R(\sin x - \cos x, \sin x \cos x) = 0$ to an equation; $R\left(t, \frac{1 - t^2}{2}\right) = 0$.

Example: 31 If $\sin x + \cos x - 2\sqrt{2} \sin x \cos x = 0$ then the general solution of x is

- (a) $x = 2n\pi + \frac{\pi}{4}$ (b) $x = n\pi + (-1)^n \frac{\pi}{6} - \frac{\pi}{4}$ (c) Both (a) and (b) (d) None of these

Solution: (c) Let $(\sin x + \cos x) = t$ and using the equation $\sin x \cos x = \frac{t^2 - 1}{2}$, we get $t - 2\sqrt{2}\left(\frac{t^2 - 1}{2}\right) = 0 \Rightarrow \sqrt{2}t^2 - t - \sqrt{2} = 0$

The numbers $t_1 = \sqrt{2}, t_2 = -\frac{1}{\sqrt{2}}$ are roots of this quadratic equation.

Thus the solution of the given equation reduces to the solution of two trigonometrical equation;

$$\sin x + \cos x = \sqrt{2} \text{ or } \sin x + \cos x = -\frac{1}{\sqrt{2}}$$

$$\text{or } \frac{1}{\sqrt{2}} \sin x + \frac{1}{\sqrt{2}} \cos x = 1 \text{ or } \frac{1}{\sqrt{2}} \sin x + \frac{1}{\sqrt{2}} \cos x = -\frac{1}{2}$$

$$\text{or } \sin x \cos \frac{\pi}{4} + \sin \frac{\pi}{4} \cos x = 1 \text{ or } \sin x \cos \frac{\pi}{4} + \sin \frac{\pi}{4} \cos x = -\frac{1}{2}$$

$$\Rightarrow \sin\left(x + \frac{\pi}{4}\right) = 1 \text{ or } \sin\left(x + \frac{\pi}{4}\right) = -\frac{1}{2} \Rightarrow x + \frac{\pi}{4} = (4n+1)\frac{\pi}{2} \text{ or } x + \frac{\pi}{4} = n\pi + (-1)^n \cdot \left(\frac{\pi}{6}\right)$$

$$\Rightarrow x = 2n\pi + \frac{\pi}{4} \text{ or } x = n\pi + (-1)^n \frac{\pi}{6} - \left(\frac{\pi}{4}\right).$$

Example: 32 If $\sin^{10} x + \cos^{10} x = \frac{29}{16} \cos^4 2x$. then $x =$

- (a) $x = \frac{n\pi}{4}$ (b) $x = \frac{n\pi}{4} + \frac{\pi}{8}$ (c) $x = \frac{n\pi}{4} + \frac{\pi}{3}$ (d) None of these

Solution: (b) Using half-angle formulae we can represent the given equation in the form,

$$\left(\frac{1 - \cos 2x}{2}\right)^5 + \left(\frac{1 + \cos 2x}{2}\right)^5 = \frac{29}{16} \cos^4 2x$$

$$\text{Put } \cos 2x = t, \left(\frac{1-t}{2}\right)^5 + \left(\frac{1+t}{2}\right)^5 = \frac{29}{16} t^4 \Rightarrow 24t^4 - 10t^2 - 1 = 0 \text{ whose only real root is, } t^2 = \frac{1}{2}.$$

$$\therefore \cos^2 2x = \frac{1}{2} \Rightarrow 1 + \cos 4x = 1 \Rightarrow \cos 4x = 0 \Rightarrow 4x = (2n+1)\frac{\pi}{2} \Rightarrow x = \frac{n\pi}{4} + \frac{\pi}{8}; n \in I$$


58 Trigonometrical Equations and

Note: □ Some trigonometric equations can sometimes be simplified by lowering their degrees. If the exponent of the sines and cosines occurring into an equation are even, the lowering of the degree can be done by half angle formulas as in above example.

2.8 Method for Finding Principal Value

Suppose we have to find the principal value of θ satisfying the equation $\sin \theta = -\frac{1}{2}$.

Since $\sin \theta$ is negative, θ will be in 3rd or 4th quadrant. We can approach 3rd or 4th quadrant from two directions. If we take anticlockwise direction the numerical value of the angle will be greater than π . If we approach it in clockwise direction the angle will be numerically less than π . For principal value, we have to take numerically smallest angle. So for principal value



(1) If the angle is in 1st or 2nd quadrant we must select anticlockwise direction and if the angle is in 3rd or 4th quadrant, we must select clockwise direction.

(2) Principal value is never numerically greater than π .

(3) Principal value always lies in the first circle (i.e., in first rotation). On the above criteria, θ will be $-\frac{\pi}{6}$ or $-\frac{5\pi}{6}$. Among these two $-\frac{\pi}{6}$ has the least numerical value. Hence $-\frac{\pi}{6}$ is the principal value of θ satisfying the equation $\sin \theta = -\frac{1}{2}$.

From the above discussion, the method for finding principal value can be summed up as follows :

- (i) First draw a trigonometrical circle and mark the quadrant, in which the angle may lie.
- (ii) Select anticlockwise direction for 1st and 2nd quadrants and select clockwise direction for 3rd and 4th quadrants.
- (iii) Find the angle in the first rotation.
- (iv) Select the numerically least angle. The angle thus found will be principal value.
- (v) In case, two angles one with positive sign and the other with negative sign qualify for the numerically least angle, then it is the convention to select the angle with positive sign as principal value.

Example: 33 If $\cos \theta + \sqrt{3} \sin \theta = 2$, then $\theta =$ (only principal value)


- (a) $\frac{\pi}{3}$ (b) $\frac{2\pi}{3}$ (c) $\frac{4\pi}{3}$ (d) $\frac{5\pi}{3}$

Solution: (a) $\frac{1}{2} \cos \theta + \frac{\sqrt{3}}{2} \sin \theta = \frac{2}{2} \Rightarrow \cos \theta \cos \frac{\pi}{3} + \sin \theta \sin \frac{\pi}{3} = 1 \Rightarrow \cos \left(\theta - \frac{\pi}{3} \right) = 1 = \cos 0^\circ \Rightarrow \theta - \frac{\pi}{3} = 0^\circ \Rightarrow \theta = \frac{\pi}{3}$.

Example: 34 Principal value of $\tan \theta = -1$ is

- (a) $-\frac{\pi}{4}$ (b) $\frac{\pi}{4}$ (c) $\frac{3\pi}{4}$

Solution: (a) $\because \tan \theta$ is negative.



$\therefore \theta$ will lie in 2nd or 4th quadrant. For 2nd quadrant we will select anticlockwise and for 4th quadrant, we will select clockwise direction.

In the first circle two values $\frac{-\pi}{4}$ and $\frac{3\pi}{4}$ are obtained.

Among these two, $\frac{-\pi}{4}$ is numerically least angle. Hence principal value is $\frac{-\pi}{4}$.

Important Tips

Any trigonometric equation can be solved without using any formula. Find all angles in $[0, 2\pi]$ which satisfy the equation and then add $2n\pi$ to each.

For example: Consider the equation $\sin \theta = \frac{1}{2}$, then $\theta = \frac{\pi}{6}, \frac{5\pi}{6}$. Hence required solutions are $\theta = 2n\pi + \frac{\pi}{6}, 2n\pi + \frac{5\pi}{6}$.

2.9 Important Points to be Taken in Case of While Solving Trigonometrical Equations

(1) Check the validity of the given equation, e.g., $2\sin \theta - \cos \theta = 4$ can never be true for any θ as the value $(2\sin \theta - \cos \theta)$ can never exceeds $\sqrt{2^2 + (-1)^2} = \sqrt{5}$. So there is no solution to this equation.

(2) Equation involving $\sec \theta$ or $\tan \theta$ can never have a solution of the form $(2n+1)\frac{\pi}{2}$

Similarly, equations involving $\operatorname{cosec} \theta$ or $\cot \theta$ can never have a solution of the form $\theta = n\pi$. The corresponding functions are undefined at these values of θ .

(3) If while solving an equation we have to square it, then the roots found after squaring must be checked whether they satisfy the original equation or not, e.g., Let $x = 3$. Squaring, we get $x^2 = 9 \therefore x = 3$ and -3 but $x = -3$ does not satisfy the original equation $x = 3$. e.g., $\sin x + \cos x = 1$

Square both sides, we get $1 + \sin 2x = 1 \therefore \sin 2x = 0$

$$\therefore 2x = n\pi \quad \text{or} \quad x = \frac{n\pi}{2}, n \in I$$

$$\therefore \text{Roots are } \dots, \frac{-3\pi}{2}, \frac{-2\pi}{2}, \frac{-\pi}{2}, 0, \frac{\pi}{2}, \frac{2\pi}{2}, \frac{3\pi}{2}, \dots$$

We find that 0 and $\pi/2$ are roots but π and $3\pi/2$ do not satisfy the given equation as it leads to $-1 = 1$

Similarly 0 and $\frac{-3\pi}{2}$ are roots but $-\frac{\pi}{2}$ and $-\pi$ are not roots as it will lead to $-1 = 1$.

As stated above, because of squaring we are solving the equations $\sin x + \cos x = 1$ and $\sin x + \cos x = -1$ both. The rejected roots are for $\sin x + \cos x = -1$.

(4) Do not cancel common factors involving the unknown angle on L.H.S. and R.H.S. because it may delete some solutions. e.g., In the equation $\sin \theta(2\cos \theta - 1) = \sin \theta \cos^2 \theta$ if we cancel $\sin \theta$ on

60 Trigonometrical Equations and

both sides we get $\cos^2 \theta - 2 \cos \theta + 1 = 0 \Rightarrow (\cos \theta - 1)^2 = 0 \Rightarrow \cos \theta = 1 \Rightarrow \theta = 2n\pi$. But $\theta = n\pi$ also satisfies the equation because it makes $\sin \theta = 0$. So, the complete solution is $\theta = n\pi, n \in \mathbb{Z}$.

(5) Any value of x which makes both R.H.S. and L.H.S. equal will be a root but the value of x for which $\infty = \infty$ will not be a solution as it is an indeterminate form.

Hence, $\cos x \neq 0$ for those equations which involve $\tan x$ and $\sec x$ whereas $\sin x \neq 0$ for those which involve $\cot x$ and $\operatorname{cosec} x$.

Also exponential function is always +ve and $\log_a x$ is defined if $x > 0$, $x \neq 0$ and $a > 0, a \neq 1$. $\sqrt{f(x)} = +ve$ always and not $\pm i.e.$ $\sqrt{(\tan^2 x)} = \tan x$ and not $\pm \tan x$.

(6) Denominator terms of the equation if present should never become zero at any stage while solving for any value of θ contained in the answer.

(7) Sometimes the equation has some limitations also e.g., $\cot^2 \theta + \operatorname{cosec}^2 \theta = 1$ can be true only if $\cot^2 \theta = 0$ and $\operatorname{cosec}^2 \theta = 1$ simultaneously as $\operatorname{cosec}^2 \theta \geq 1$. Hence the solution is $\theta = (2n+1)\pi/2$.

(8) If $xy = xz$ then $x(y-z) = 0 \Rightarrow$ either $x = 0$ or $y = z$ or both. But $\frac{y}{x} = \frac{z}{x} \Rightarrow y = z$ only and not $x = 0$, as it will make $\infty = \infty$. Similarly if $ay = az$, then it will also imply $y = z$ only as $a \neq 0$ being a constant.

Similarly $x+y = x+z \Rightarrow y = z$ and $x-y = x-z \Rightarrow y = z$. Here we do not take $x = 0$ as in the above because x is an additive factor and not multiplicative factor.

When $\cos \theta = 0$, then $\sin \theta = 1$ or -1 . We have to verify which value of $\sin \theta$ is to be chosen which satisfies the equation. $\cos \theta = 0 \Rightarrow \theta = \left(n + \frac{1}{2}\right)\pi$.

(9) Student are advised to check whether all the roots obtained by them, satisfy the equation and lie in the domain of the variable of the given equation.

2.10 Miscellaneous Examples

Example: 35 The equation $3 \sin^2 x + 10 \cos x - 6 = 0$ is satisfied if

(a) $x = n\pi \pm \cos^{-1}\left(\frac{1}{3}\right)$ (b) $x = 2n\pi \pm \cos^{-1}\left(\frac{1}{3}\right)$ (c) $x = n\pi \pm \cos^{-1}\left(\frac{1}{6}\right)$ (d) $x = 2n\pi \pm \cos^{-1}\left(\frac{1}{6}\right)$

Solution: (b) $3 \sin^2 x + 10 \cos x - 6 = 0 \Rightarrow 3(1 - \cos^2 x) + 10 \cos x - 6 = 0$

on solving, $(\cos x - 3)(3 \cos x - 1) = 0$. Either $\cos x = 3$ (which is not possible) or $\cos x = \frac{1}{3} \Rightarrow$

$$x = 2n\pi \pm \cos^{-1}\left(\frac{1}{3}\right).$$

Example: 36 If the solutions for θ of $\cos p\theta + \cos q\theta = 0, p > 0, q > 0$ are in A.P., then the numerically smallest common difference of A.P. is

[Kerala (Engg.) 2001]

(a) $\frac{\pi}{p+q}$ (b) $\frac{2\pi}{p+q}$ (c) $\frac{\pi}{2(p+q)}$ (d) $\frac{1}{p+q}$

Solution: (b) Given, $\cos p\theta = -\cos q\theta = \cos(\pi + q\theta) \Rightarrow p\theta = 2n\pi \pm (\pi + q\theta), n \in I \Rightarrow \theta = \frac{(2n+1)\pi}{p-q}$ or $\frac{(2n-1)\pi}{p+q}, n \in I$. Both the solutions form an A.P. $\theta = \frac{(2n+1)\pi}{p-q}$ gives us an A.P. with common difference $= \frac{2\pi}{p-q}$ and $\theta = \frac{(2n-1)\pi}{p+q}$ gives us an A.P. with common difference $= \frac{2\pi}{p+q}$. Certainly, $\frac{2\pi}{p+q} < \left| \frac{2\pi}{p-q} \right|$.

Example: 37 The set of values of x for which the expression $\frac{\tan 3x - \tan 2x}{1 + \tan 3x \tan 2x} = 1$ is [MP PET 1992; UPSEAT 1993, 2002]

- (a) ϕ (b) $\frac{\pi}{4}$
 (c) $\left\{ n\pi + \frac{\pi}{4} : n = 1, 2, 3, \dots \right\}$ (d) $\left\{ 2n\pi + \frac{\pi}{4} : n = 1, 2, 3, \dots \right\}$

Solution: (a) $\tan(3x - 2x) = \tan x = 1 \Rightarrow x = n\pi + \frac{\pi}{4}$. But this value does not satisfy the given equation.

Example: 38 If $\cos \theta + \cos 7\theta + \cos 3\theta + \cos 5\theta = 0$, then $\theta =$ [ISM Dhanbad 1972]

- (a) $\frac{n\pi}{4}$ (b) $\frac{n\pi}{2}$ (c) $\frac{n\pi}{8}$ (d) None of these

Solution: (c) Combining θ and $7\theta, 3\theta$ and 5θ , we get $2 \cos 4\theta (\cos 3\theta + \cos \theta) = 0$

$$\therefore 4 \cos 4\theta \cdot \cos 2\theta \cdot \cos \theta = 0 \Rightarrow 4 \frac{1}{2^3 \sin \theta} (\sin 2^3 \theta) = 0 \Rightarrow \sin 8\theta = 0. \text{ Hence } \theta = \frac{n\pi}{8}$$

Example: 39 If $\tan(\cot x) = \cot(\tan x)$, then $\sin 2x$ equal to

- (a) $(2n+1)\frac{\pi}{4}$ (b) $\frac{4}{(2n+1)\pi}$ (c) $4\pi(2n+1)$ (d) None of these

Solution: (b) $\tan(\cot x) = \cot(\tan x) \Rightarrow \tan(\cot x) = \tan\left(\frac{\pi}{2} - \tan x\right)$

$$\cot x = n\pi + \frac{\pi}{2} - \tan x$$

$$\tan x + \cot x = \frac{(2n+1)\pi}{2} \Rightarrow \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{(2n+1)\pi}{2}$$

$$\frac{1}{\sin x \cdot \cos x} = \frac{(2n+1)\pi}{2} \Rightarrow \sin 2x = \frac{4}{(2n+1)\pi}.$$

Example: 40 The sum of all solutions of the equation $\cos x \cdot \cos\left(\frac{\pi}{3} + x\right) \cdot \cos\left(\frac{\pi}{3} - x\right) = \frac{1}{4}, x \in [0, 6\pi]$ is

- (a) 15π (b) 30π (c) $\frac{110\pi}{3}$ (d) None of these

Solution: (b) Here, $\cos x \left(\frac{1}{4} \cos^2 x - \frac{3}{4} \sin^2 x \right) = \frac{1}{4}$ or $\frac{\cos x}{4} (4 \cos^2 x - 3) = \frac{1}{4}$ or $\cos 3x = 1$

$$\Rightarrow 3x = 2n\pi \Rightarrow x = \frac{2n\pi}{3}, \text{ where } n = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9; \therefore \text{The required sum} = \frac{2\pi}{3} \sum_{n=0}^9 n = 30\pi.$$

Example: 41 The equation $\sin x + \sin y + \sin z = -3$ for $0 \leq x \leq 2\pi, 0 \leq y \leq 2\pi, 0 \leq z \leq 2\pi$, has [Orissa JEE 2003]

- (a) One solution (b) Two sets of solution

62 Trigonometrical Equations and

(c) Four sets of solution

(d)

No solution

Solution: (a) Given, $\sin x + \sin y + \sin z = -3$ is satisfied only when $x = y = z = \frac{3\pi}{2}$; for $x, y, z, \in [0, 2\pi]$.

Example: 42 The solution set of $(5 + 4 \cos \theta)(2 \cos \theta + 1) = 0$ in the interval $[0, 2\pi]$ is

[EAMCET 2003]

(a) $\left\{\frac{\pi}{3}, \frac{2\pi}{3}\right\}$

(b) $\left\{\frac{\pi}{3}, \pi\right\}$

(c) $\left\{\frac{2\pi}{3}, \frac{4\pi}{3}\right\}$

(d) $\left\{\frac{2\pi}{3}, \frac{5\pi}{3}\right\}$

Solution: (c) $(5 + 4 \cos \theta)(2 \cos \theta + 1) = 0$

$\cos \theta = \frac{-5}{4}$ which is not possible

$\therefore 2 \cos \theta + 1 = 0$ or $\cos \theta = -\frac{1}{2} \Rightarrow \theta = \frac{2\pi}{3}, \frac{4\pi}{3}$. Solution set is $\left\{\frac{2\pi}{3}, \frac{4\pi}{3}\right\} \in [0, 2\pi]$

Example: 43 The equation $3 \cos x + 4 \sin x = 6$ has

[Orissa JEE 2002]

(a) Finite solution

(b) Infinite solution

(c) One solution

(d) No solution

Solution: (d) $3 \cos x + 4 \sin x = 6$

$\Rightarrow \frac{3}{5} \cos x + \frac{4}{5} \sin x = \frac{6}{5} \Rightarrow \cos(x - \theta) = \frac{6}{5}$ $\left[\therefore \theta = \cos^{-1}\left(\frac{3}{5}\right) \right]$

So that equation has no solution.

Example: 44 The equation $\sin x + \cos x = 2$ has

[EAMCET 1986; MP PET 1998]

(a) One solution

(b) Two solution

(c) Infinite number of solution

(d)

No solution

Solution: (d) No solution as $|\sin x| \leq 1, |\cos x| \leq 1$ and both of them do not attain their maximum value for the same angle.

Trick: Maximum value of $\sin x + \cos x = \sqrt{1^2 + 1^2} = \sqrt{2}$. Hence there is no x satisfying this equation.

Example: 45 If $2 \cos x < \sqrt{3}$ and $x \in [-\pi, \pi]$ then the solution set for x is

(a) $\left[-\pi, \frac{-\pi}{6}\right) \cup \left(\frac{\pi}{6}, \pi\right]$

(b) $\frac{-\pi}{6}, \frac{\pi}{6}$

(c) $\left[-\pi, \frac{-\pi}{6}\right] \cup \left[\frac{\pi}{6}, \pi\right]$


(d) None of these

Solution: (a) Here, $\cos x < \frac{\sqrt{3}}{2}$. The value scheme for this is shown below.

From the figure,

$-\pi \leq x < \frac{-\pi}{6}$ or $\frac{\pi}{6} < x \leq \pi$

$\therefore x \in \left[-\pi, \frac{-\pi}{6}\right) \cup \left(\frac{\pi}{6}, \pi\right]$



Example: 46 The number of pairs (x, y) satisfying the equations $\sin x + \sin y = \sin(x + y)$ and $|x| + |y| = 1$ is

(a) 2

(b) 4

(c) 6

(d) ∞

Solution: (c) The first equation can be written as, $2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y) = 2 \sin \frac{1}{2}(x+y) \cos \frac{1}{2}(x+y)$

$$\therefore \text{Either } \sin \frac{1}{2}(x+y) = 0 \text{ or } \sin \frac{1}{2}x = 0 \text{ or } \sin \frac{1}{2}y = 0$$

$\Rightarrow x+y=0$, or $x=0$ or $y=0$. As $|x| + |y| = 1$, therefore when $x+y=0$, we have to reject $x+y=1$, or $x+y=-1$ and solve it with $x-y=1$ or $x-y=-1$ which gives $\left(\frac{1}{2}, \frac{-1}{2}\right)$ or $\left(\frac{-1}{2}, \frac{1}{2}\right)$ as the possible solution. Again solving with $x=0$, we get $(0, \pm 1)$ and solving with $y=0$, we get $(\pm 1, 0)$ as the other solution. Thus we have six pairs of solution for x and y .

Example: 47 If $\cos \theta = -\frac{1}{2}$ and $0^\circ < \theta < 360^\circ$, then the values of θ are

- (a) 120° and 300° (b) 60° and 120° (c) 120° and 240° (d) 60° and 240°

Solution: (c) Given, $\cos \theta = -\frac{1}{2}$ and $0^\circ < \theta < 360^\circ$. We know that $\cos 60^\circ = \frac{1}{2}$ and $\cos(180^\circ - 60^\circ) = -\cos 60^\circ = -\frac{1}{2}$ or $\cos 120^\circ = -\frac{1}{2}$. Similarly $\cos(180^\circ + 60^\circ) = -\cos 60^\circ = -\frac{1}{2}$ or $\cos 240^\circ = -\frac{1}{2}$.

Therefore $\theta = 120^\circ$ and 240° .

Example: 48 If $\tan(\pi \cos \theta) = \cot(\pi \sin \theta)$, then the value of $\cos\left(\theta - \frac{\pi}{4}\right) =$

- (a) $\frac{1}{2\sqrt{2}}$ (b) $\frac{1}{\sqrt{2}}$ (c) $\frac{1}{3\sqrt{2}}$ (d) $\frac{1}{4\sqrt{2}}$

Solution: (a) $\tan(\pi \cos \theta) = \tan\left(\frac{\pi}{2} - \pi \sin \theta\right)$

$$\therefore \sin \theta + \cos \theta = \frac{1}{2} \Rightarrow \cos\left(\theta - \frac{\pi}{4}\right) = \frac{1}{2\sqrt{2}}.$$

Example: 49 The only value of x for which $2^{\sin x} + 2^{\cos x} > 2^{1-(1/\sqrt{2})}$ hold is

- (a) $\frac{5\pi}{4}$ (b) $\frac{3\pi}{4}$ (c) $\frac{\pi}{2}$ (d) All values of x .

Solution: (a) Since A.M. $\geq$ G.M.

$$\frac{1}{2}(2^{\sin x} + 2^{\cos x}) \geq \sqrt{2^{\sin x} \cdot 2^{\cos x}} \Rightarrow 2^{\sin x} + 2^{\cos x} \geq 2 \cdot 2^{\frac{\sin x + \cos x}{2}} \Rightarrow 2^{\sin x} + 2^{\cos x} \geq 2^{1 + \frac{\sin x + \cos x}{2}}$$

And, we know that $\sin x + \cos x \geq -\sqrt{2}$

$$\therefore 2^{\sin x} + 2^{\cos x} > 2^{1-(1/\sqrt{2})} \text{ for } x = \frac{5\pi}{4}.$$

2.11 Periodic Functions

A function $f(x)$ is called periodic function if there exists a least positive real number T such that $f(x+T) = f(x)$. T is called the period (or fundamental period) of function $f(x)$. Obviously, if T is the period of $f(x)$, then $f(x) = f(x+T) = f(x+2T) = f(x+3T) = \dots$

(i) If $f_1(x)$ and $f_2(x)$ are two periodic functions of x having the same period T , then the function $af_1(x) + bf_2(x)$ where a and b are any numbers, is also a periodic function having the same period T .

64 Trigonometrical Equations and

(ii) If T is the period of the periodic function $f(x)$, then the function $f(ax + b)$, where $a(> 0)$ and b are any numbers is also a periodic function with period equal to T/a .

(iii) If T_1 and T_2 are the periods of periodic functions $f_1(x)$ and $f_2(x)$ respectively, then the function $af_1(x) + bf_2(x)$, where a and b are any numbers is also periodic and its period is T which is the L.C.M. of T_1 and T_2 i.e. T is the least positive number which is divisible by T_1 and T_2 .

All trigonometric functions are periodic. The period of trigonometric function $\sin x, \cos x, \sec x$ and $\operatorname{cosec} x$ is 2π because $\sin(x + 2\pi) = \sin x, \cos(x + 2\pi) = \cos x$ etc.

The period of $\tan x$ and $\cot x$ is π because $\tan(x + \pi) = \tan x$ and $\cot(x + \pi) = \cot x$

The period of the function which are of the type: $\sin ax, \cos(ax + b); b \cos ax$ is $\frac{2\pi}{a}$

The period of $\tan ax$ and $\cot ax$ is $\frac{\pi}{|a|}$. Here $|a|$ is taken so as the value of the period is positive real number.

Some functions with their periods

Function	Period
$\sin(ax + b), \cos(ax + b), \sec(ax + b), \operatorname{cosec}(ax + b)$	$2\pi / a$
$\tan(ax + b), \cot(ax + b)$	π / a
$ \sin(ax + b) , \cos(ax + b) , \sec(ax + b) , \operatorname{cosec}(ax + b) $	π / a
$ \tan(ax + b) , \cot(ax + b) $	π / a

Example: 50 Period of $\sin^2 x$ is

[UPSEAT 2002; AIEEE 2002]

- (a) π (b) 2π (c) $\frac{\pi}{2}$ (d) None of these

Solution: (a) $\sin^2 x = \frac{1 - \cos 2x}{2} \Rightarrow \text{Period} = \frac{2\pi}{2} = \pi.$

Example: 51 The period of the function $y = \sin 2x$ is

[Kerala(Engg.) 2003]

- (a) 2π (b) π (c) $\frac{\pi}{2}$ (d) 4π

Solution: (a) Period of $\sin(ax + b) = \frac{2\pi}{|a|}$

$\therefore$ Period of $\sin 2x = \frac{2\pi}{|2|} = \pi.$

Example: 52 The period of the function $f(\theta) = \sin \frac{\theta}{3} + \cos \frac{\theta}{2}$ is

[EAMCET 2001]

- (a) 3π (b) 6π (c) 9π (d) 12π

Solution: (d) Period of $\sin\left(\frac{\theta}{3}\right) = 6\pi$ and period of $\cos\left(\frac{\theta}{2}\right) = 4\pi$

L.C.M. of 6π and $4\pi = 12\pi$

Example: 53 The function $f(x) = \sin \frac{\pi x}{2} + 2 \cos \frac{\pi x}{3} - \tan \frac{\pi x}{4}$ is periodic with period [Rajasthan PET 2001]

- (a) 6 (b) 3 (c) 4 (d) 12

Solution: (d) Period of $\sin \frac{\pi x}{2} = \frac{2\pi}{\pi/2} = 4$

Period of $\cos \frac{\pi x}{3} = \frac{2\pi}{\pi/3} = 6$ and period of $\tan \frac{\pi x}{4} = \frac{\pi}{\pi/4} = 4$

$\therefore$ Period of $f(x)$ = L.C.M. of (4, 6, 4) = 12.

Example: 54 If the period of the function $f(x) = \sin\left(\frac{x}{n}\right)$ is 4π , then n is equal to [Pb. CET 2000]

- (a) 1 (b) 4 (c) 8 (d) 2

Solution: (d) $\sin\left(\frac{x}{n}\right) = \sin\left(2\pi + \frac{x}{n}\right) = \sin\left[\frac{1}{n}(2n\pi + x)\right] \Rightarrow$ Period of the function $\sin\left(\frac{x}{n}\right)$ is $2n\pi$. $\Rightarrow 2n\pi = 4\pi \Rightarrow n = 2$.

Example: 55 The period of $\sin^4 x + \cos^4 x$ is

- (a) $\frac{\pi}{2}$ (b) π (c) 2π (d) $\frac{3\pi}{2}$

Solution: (a) $(\sin^2 x)^2 + (\cos^2 x)^2 = 1 - 2 \sin^2 x \cos^2 x = 1 - \frac{1}{4} 2 \sin^2 2x = 1 - \left(\frac{1 - \cos 4x}{4}\right) = \frac{3}{4} + \frac{1}{4} \cos 4x$

Therefore period is $\frac{2\pi}{4} = \frac{\pi}{2}$.

Trick: $f(x) = \sin^4 x + \cos^4 x \Rightarrow f\left(\frac{\pi}{2} + x\right) = \sin^4\left(\frac{\pi}{2} + x\right) + \cos^4\left(\frac{\pi}{2} + x\right) \Rightarrow f\left(\frac{\pi}{2} + x\right) = \cos^4 x + \sin^4 x = f(x)$

Hence the period is $\frac{\pi}{2}$.

Example: 56 Period of $\sin \theta - \sqrt{3} \cos \theta$ is [MP PET 1990]

- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$ (c) π (d) 2π

Solution: (d) $\sin \theta - \sqrt{3} \cos \theta = 2 \left(\frac{1}{2} \sin \theta - \frac{\sqrt{3}}{2} \cos \theta \right) = 2 \sin \left(\theta - \frac{\pi}{3} \right)$

Hence period = 2π .

Example: 57 Period of $|\sin 2x|$ is [MP PET 1989]

- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$ (c) π (d) 2π

Solution: (b) Period of $\sin 2x = \pi$ and period of $|\sin 2x| = \frac{\pi}{2}$.



Assignment

General Solution of a Trigonometric Equations

Basic Level

- If $\cos p\theta = \cos q\theta$, $p \neq q$, then [MP PET 1995]

(a) $\theta = 2n\pi$ (b) $\theta = \frac{2n\pi}{p \pm q}$ (c) $\theta = \frac{n\pi}{p + q}$ (d) None of these
- General solution of $\tan 5\theta = \cot 2\theta$ is [Karnataka CET 2000]

(a) $\theta = \frac{n\pi}{7} + \frac{\pi}{14}$ (b) $\theta = \frac{n\pi}{7} + \frac{\pi}{5}$ (c) $\theta = \frac{n\pi}{7} + \frac{\pi}{2}$ (d) $\theta = \frac{n\pi}{7} + \frac{\pi}{3}, n \in \mathbb{Z}$
- If $\tan 2\theta \tan \theta = 1$, then $\theta =$

(a) $\frac{\pi}{3}$ (b) $(6n \pm 1)\frac{\pi}{6}$ (c) $(4n \pm 1)\frac{\pi}{6}$ (d) None of these
- If $\tan \theta + \cot \theta = 2$, then $\theta =$

(a) $n\pi$ (b) $n\pi + \frac{\pi}{4}$ (c) $n\pi \pm \frac{\pi}{4}$ (d) $n\pi \pm \frac{\pi}{3}$
- If $\cot \theta + \tan \theta = 2\operatorname{cosec} \theta$, then general value of θ is [Roorkee 1971]

(a) $n\pi \pm \frac{\pi}{3}$ (b) $n\pi \pm \frac{\pi}{6}$ (c) $2n\pi \pm \frac{\pi}{3}$ (d) $2n\pi \pm \frac{\pi}{6}$
- If $1 + \cot \theta = \operatorname{cosec} \theta$, then the general value of θ is [Roorkee 1981]

(a) $n\pi + \frac{\pi}{2}$ (b) $2n\pi - \frac{\pi}{2}$ (c) $2n\pi + \frac{\pi}{2}$ (d) None of these
- The value of $\cos y \cos\left(\frac{\pi}{2} - x\right) - \cos\left(\frac{\pi}{2} - y\right) \cos x + \sin y \cos\left(\frac{\pi}{2} - x\right) + \cos x \sin\left(\frac{\pi}{2} - y\right)$ is zero if [Kerala (Engg.) 1993]

(a) $x = 0$ (b) $y = 0$ (c) $x = y$ (d) $x = n\pi - \frac{\pi}{4} + y (n \in \mathbb{I})$
- If $\sin^2 \theta - 2 \cos \theta + \frac{1}{4} = 0$, then the general value of θ is [MP PET 1984]

(a) $n\pi \pm \frac{\pi}{3}$ (b) $2n\pi \pm \frac{\pi}{3}$ (c) $2n\pi \pm \frac{\pi}{6}$ (d) $n\pi \pm \frac{\pi}{6}$
- The general value of θ satisfying $\sin^2 \theta + \sin \theta = 2$ is [AMU 1996, 99]

(a) $n\pi + (-1)^n \frac{\pi}{6}$ (b) $2n\pi + \frac{\pi}{4}$ (c) $n\pi + (-1)^n \frac{\pi}{2}$ (d) $n\pi + (-1)^n \frac{\pi}{3}$

66 Trigonometrical Equations and

Advance Level

10. If $\sec 4\theta - \sec 2\theta = 2$, then the general value of θ is [IIT 1963]
 (a) $(2n+1)\frac{\pi}{4}$ (b) $(2n+1)\frac{\pi}{10}$ (c) $n\pi + \frac{\pi}{2}$ or $\frac{n\pi}{5} + \frac{\pi}{10}$ (d) None of these
11. If $\sin\left(\frac{\pi}{4}\cot\theta\right) = \cos\left(\frac{\pi}{4}\tan\theta\right)$, then $\theta =$ [Karnataka CET 1988]
 (a) $n\pi + \frac{\pi}{4}$ (b) $2n\pi \pm \frac{\pi}{4}$ (c) $n\pi - \frac{\pi}{4}$ (d) $2n\pi \pm \frac{\pi}{6}$
12. The general solution of $\sin x - 3\sin 2x + \sin 3x = \cos x - 3\cos 2x + \cos 3x$ is [IIT 1989]
 (a) $n\pi + \frac{\pi}{8}$ (b) $\frac{n\pi}{2} + \frac{\pi}{8}$ (c) $(-1)^n \frac{n\pi}{2} + \frac{\pi}{8}$ (d) $2n\pi + \cos^{-1} \frac{3}{2}$
13. The solution of the equation $\sec \theta - \operatorname{cosec} \theta = \frac{4}{3}$ is [Roorkee 1994]
 (a) $\frac{1}{2}\left[n\pi + (-1)^n \sin^{-1}\left(\frac{3}{4}\right)\right]$ (b) $n\pi + (-1)^n \sin^{-1}\left(\frac{3}{4}\right)$ (c) $\frac{n\pi}{2} + (-1)^n \sin^{-1}\left(\frac{3}{4}\right)$ (d) None of these
14. If $4\sin^4 x + \cos^4 x = 1$, then x equal to [Roorkee 1989]
 (a) $n\pi$ (b) $n\pi \pm \sin^{-1} \frac{2}{5}$ (c) $n\pi + \frac{\pi}{6}$ (d) None of these
15. If $2(\sin x - \cos 2x) + \sin 2x(1 + 2\sin x) - 2\cos x = 0$, then $x =$ [Karnataka CET 2002]
 (a) $(2n \pm 1)\pi$ (b) $(2n \pm 1)\frac{\pi}{6}$ (c) $(n \pm 1)\frac{\pi}{4}$ (d) $(n \pm 1)\frac{\pi}{6}$
16. Expression $2^{\sin\theta} + 2^{-\cos\theta}$ is minimum when $\theta = \dots\dots\dots$ and its minimum value is $\dots\dots\dots$
 (a) $2n\pi + \frac{\pi}{4}, n \in I; 2$ (b) $2n\pi + \frac{7\pi}{4}, n \in I; 2^{1-\left(\frac{1}{\sqrt{2}}\right)}$ (c) $n\pi \pm \frac{\pi}{4}, n \in I; 2^{1-\left(\frac{1}{\sqrt{2}}\right)}$ (d) None of these
17. The general solution of the equation $2^{\cos 2x} + 1 = 3 \cdot 2^{-\sin^2 x}$ is
 (a) $n\pi$ (b) $n\pi + \pi$ (c) $n\pi - \pi$ (d) None of these
18. If $\theta = \tan^{-1}(2\tan^2 \theta) - \frac{1}{2}\sin^{-1}\left(\frac{3\sin 2\theta}{5+4\cos 2\theta}\right)$ then the general value of θ [Roorkee 1997; WBJEE 2000]
 (a) $n\pi$ (b) $n\pi + \frac{\pi}{4}$ (c) $n\pi + \tan^{-1}(-2)$ (d) All of these
19. If the expression $\frac{\sin \frac{x}{2} + \cos \frac{x}{2} - i \tan x}{1 + 2i \sin \frac{x}{2}}$ is real, then x is equal to
 (a) $2n\pi + \tan^{-1} K, K \in R, n \in Z$ (b) $2n\pi + 2 \tan^{-1} K$, where $K \in (0, 1), n \in Z$
 (b) $2n\pi + 2 \tan^{-1} K$, where $K \in (1, 2), n \in Z$ (d) $2n\pi + 2 \tan^{-1} K, K \in (2, 3), n \in Z$
20. If $\frac{1}{6}\sin x, \cos x, \tan x$ are in G.P., then x is equal to
 (a) $n\pi \pm \frac{\pi}{3}, n \in Z$ (b) $2n\pi \pm \frac{\pi}{3}, n \in Z$ (c) $n\pi + (-1)^n \frac{\pi}{3}, n \in Z$ (d) None of these

Trigonometrical Equations and Inequations

- 21.** If $32 \tan^8 \theta = 2 \cos^2 \alpha - 3 \cos \alpha$ and $3 \cos 2\theta = 1$, then the general value of α is [Roorkee 1996]
- (a) $2n\pi \pm \frac{\pi}{3}$ (b) $2n\pi \pm \cos^{-1} 2$ (c) $2n\pi \pm \frac{2\pi}{3}$ (d) None of these
- 22.** If $\max_{\theta \in R} \{5 \sin \theta + 3 \sin(\theta - \alpha)\} = 7$, then the set of possible values of α is
- (a) $\left\{x \mid x = 2n\pi \pm \frac{\pi}{3}, n \in Z\right\}$ (b) $\left\{x \mid x = 2n\pi \pm \frac{2\pi}{3}, n \in Z\right\}$ (c) $\left[\frac{\pi}{3}, \frac{2\pi}{3}\right]$ (d) None of these
- 23.** If $|\cos x|^{\sin^2 x - \frac{3}{2} \sin x + \frac{1}{2}} = 1$, then possible values of x are
- (a) $n\pi$ or $n\pi + (-1)^n \frac{\pi}{6}, n \in I$ (b) $n\pi$ or $2n\pi + \frac{\pi}{2}$ or $n\pi + (-1)^n \frac{\pi}{6}, n \in I$
- (c) $n\pi + (-1)^n \frac{\pi}{6}, n \in I$ (d) None of these
- 24.** The general solution of $\cos^{50} x - \sin^{50} x = 1$ is
- (a) $n\pi$ (b) $2n\pi$ (c) $n\pi + \frac{\pi}{2}$ (d) $2n\pi + \frac{\pi}{2}$
- 25.** Let $[x]$ = the greatest integer less than or equal to x and let $f(x) = \sin x + \cos x$. Then the most general solution of $f(x) = \left[f\left(\frac{\pi}{10}\right)\right]$ are
- (a) $2n\pi + \frac{\pi}{2}, n \in Z$ (b) $n\pi, n \in Z$ (c) $2n\pi, n \in Z$ (d) None of these
- 26.** The most general values of x for which $\sin x + \cos x = \min_{a \in R} \{1, a^2 - 4a + 6\}$ are given by
- (a) $2n\pi$ (b) $2n\pi + \frac{\pi}{2}$ (c) $n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{4}$ (d) None of these

Number of Solutions and Solutions in the Interval of a Trigonometric Equation

Basic Level

- 27.** The number of values of θ in $[0, 2\pi]$ satisfying the equation $2 \sin^2 \theta = 4 + 3 \cos \theta$ are [MP PET 1989]
- (a) 0 (b) 1 (c) 2 (d) 3
- 28.** The equation $3 \cos x + 4 \sin x = 6$ has....solution [Orissa JEE 2002]
- (a) Finite (b) Infinite (c) One (d) No
- 29.** The solution of the equation $\cos^2 \theta + \sin \theta + 1 = 0$, lies in the interval
- (a) $\left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$ (b) $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$ (c) $\left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$ (d) $\left(\frac{5\pi}{4}, \frac{7\pi}{4}\right)$
- 30.** The solution set of the system of equations $x + y = \frac{2\pi}{3}$, $\cos x + \cos y = \frac{3}{2}$, where x and y are real in
- (a) A finite non-empty set (b) Null set (c) Infinite (d)
- 31.** The equation $2 \cos^2\left(\frac{x}{2}\right) \sin^2 x = x^2 + \frac{1}{x^2}, 0 \leq x \leq \frac{\pi}{2}$ has [Kurukshetra CEE 1995]
- (a) No solution (b) One real solution (c) More than one real solution (d) None of these
- 32.** The smallest positive root of the equation $\tan x - x = 0$ lies on

68 Trigonometrical Equations and

- (a) $\left(0, \frac{\pi}{2}\right)$ (b) $\left(\frac{\pi}{2}, \pi\right)$ (c) $\left(\pi, \frac{3\pi}{2}\right)$ (d) $\left(\frac{3\pi}{2}, 2\pi\right)$

Advance Level

33. If a is any real number, then the number of roots of $\cot x - \tan x = a$ in the first quadrant are
 (a) 2 (b) 0 (c) 1 (d) None of these
34. The number of solutions of the equation $\sin\left(\frac{\pi x}{2\sqrt{3}}\right) = x^2 - 2\sqrt{3}x + 4$
 (a) Form an empty set (b) 1 (c) 2 (d) > 2
35. The number of all possible triplets (a_1, a_2, a_3) such that $a_1 + a_2 \cos 2x + a_3 \sin^2 x = 0$ for all x is
 (a) Zero (b) 1 (c) 2 (d) Infinite
36. The equation $(\cos p - 1)x^2 - (\cos p)x + \sin p = 0$, where x is a variable, has real roots. Then the interval of p may be any one of the followings [IIT 1990]
 (a) $(0, 2\pi)$ (b) $(-\pi, 0)$ (c) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (d) $(0, \pi)$
37. Let n be an odd integer if $\sin n\theta = \sum_{r=0}^n b_r \sin^r \theta$, for every value of θ , then [IIT 1998]
 (a) $b_0 = 1, b_1 = 3$ (b) $b_0 = 0, b_1 = 4$ (c) $b_0 = 0, b_1 = n$ (d) $b_0 = 0, b_1 = n^2 - n + 3$

Solution of Simultaneously Solving Equation

Basic Level

38. The most general value of θ which will satisfy both the equations $\sin \theta = -\frac{1}{2}$ and $\tan \theta = \frac{1}{\sqrt{3}}$ is [UPSEAT 1980; MP PET 1980]
 (a) $n\pi + (-1)^n \frac{\pi}{6}$ (b) $n\pi + \frac{\pi}{6}$ (c) $2n\pi \pm \frac{\pi}{6}$ (d) None of these
39. The most general value of θ satisfying the equations $\sin \theta = \sin \alpha$ and $\cos \theta = \cos \alpha$ is [DCE 1999; IIT 1971; Karnataka CET 1990]
 (a) $2n\pi + \alpha$ (b) $2n\pi - \alpha$ (c) $n\pi + \alpha$ (d) $n\pi - \alpha$
40. If $\sin A = \sin B, \cos A = \cos B$, then the value of A in terms of B is
 (a) $n\pi + B$ (b) $n\pi + (-1)^n B$ (c) $2n\pi + B$ (d) $2n\pi - B$
41. If $\cos \theta = -\frac{1}{\sqrt{2}}$ and $\tan \theta = 1$, then the general value of θ is
 (a) $2n\pi + \frac{\pi}{4}$ (b) $(2n+1)\pi + \frac{\pi}{4}$ (c) $n\pi + \frac{\pi}{4}$ (d) $n\pi \pm \frac{\pi}{4}$
42. If $\sin 2x + \sin 4x = 2 \sin 3x$, then $x =$ [EAMCET 1989]
 (a) $\frac{n\pi}{3}$ (b) $n\pi + \frac{\pi}{3}$ (c) $2n\pi \pm \frac{\pi}{3}$ (d) None of these
43. If $\tan \theta = \sqrt{3}$ and $\operatorname{cosec} \theta = \frac{-2}{\sqrt{3}}$, then the most general value of θ satisfying both the equations is

- (a) $2n\pi + \frac{\pi}{3}$ (b) $2n\pi + \frac{2\pi}{3}$ (c) $2n\pi + \frac{4\pi}{3}$ (d) None of these
44. If $\sin \theta = \frac{1}{2}$ and $\cos \theta = \frac{\sqrt{3}}{2}$, then the general value of θ which satisfies both the equations is
 (a) $2n\pi + \frac{\pi}{3}$ (b) $2n\pi + \frac{\pi}{4}$ (c) $2n\pi + \frac{\pi}{6}$ (d) None of these
45. General solution of $(1 + 2 \sin \theta)^2 + (\sqrt{3} \tan \theta - 1)^2 = 0$ is
 (a) $n\pi + \frac{\pi}{6}$ (b) $2n\pi + \frac{11\pi}{6}$ (c) $2n\pi + \frac{7\pi}{6}$ (d) None of these
46. If $3 \sin^2 \theta + 2 \sin^2 \phi = 1$ and $3 \sin 2\theta = 2 \sin 2\phi$, $0 < \theta < \frac{\pi}{2}$ and $0 < \phi < \frac{\pi}{2}$, then the value of $\theta + 2\phi$ is
 (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{4}$ (c) 0 (d) None of these

Principal Value and Particular Values of a Trigonometrical Equations

Basic Level

47. The smallest value of θ satisfying $\sqrt{3}(\cot \theta + \tan \theta) = 4$ is [EAMCET 1996]
 (a) $\frac{2\pi}{3}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{6}$ (d) $\frac{\pi}{12}$
48. If $\sin \theta = \sqrt{3} \cos \theta$, $-\pi < \theta < 0$, then $\theta =$ [MP PET 1992]
 (a) $-\frac{5\pi}{6}$ (b) $-\frac{4\pi}{6}$ (c) $\frac{4\pi}{6}$ (d) $\frac{5\pi}{6}$
49. If $\tan \theta = -\frac{1}{\sqrt{3}}$, $\sin \theta = \frac{1}{2}$ and $\cos \theta = -\frac{\sqrt{3}}{2}$, then the principal value of θ will be [MP PET 1984]
 (a) $\frac{\pi}{6}$ (b) $\frac{5\pi}{6}$ (c) $\frac{7\pi}{6}$ (d) $-\frac{\pi}{6}$
50. The smallest positive angle satisfying the equation $\sin^2 \theta - 2 \cos \theta + \frac{1}{4} = 0$ is
 (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$
51. If $\sin 5x + \sin 3x + \sin x = 0$, then the value of x other than 0 lying between $0 \leq x \leq \frac{\pi}{2}$ is [MNR 1985]
 (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{12}$ (c) $\frac{\pi}{3}$ (d) $\frac{\pi}{4}$
52. The value of x satisfying the equation $\sin x + \frac{1}{\sin x} = \frac{7}{2\sqrt{3}}$ is given by
 (a) 10° (b) 30° (c) 45° (d) 60°
53. If $\sin 2\theta = \cos 3\theta$ and θ is an acute angle, then $\sin \theta$ is equal to
 (a) $\frac{\sqrt{5}-1}{4}$ (b) $\frac{-\sqrt{5}-1}{4}$ (c) 0 (d) None of these
54. The values of θ satisfying $\sin 7\theta = \sin 4\theta - \sin \theta$ and $0 < \theta < \frac{\pi}{2}$ are

70 Trigonometrical Equations and

- (a) $\frac{\pi}{9}, \frac{\pi}{4}$ (b) $\frac{\pi}{3}, \frac{\pi}{9}$ (c) $\frac{\pi}{6}, \frac{\pi}{9}$ (d) $\frac{\pi}{3}, \frac{\pi}{4}$
55. If $(2 \cos x - 1)(3 + 2 \cos x) = 0$, $0 \leq x \leq 2\pi$, then $x =$ [UPSEAT 1988, 2000]
 (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{3}, \frac{5\pi}{3}$ (c) $\frac{\pi}{2}, \frac{5\pi}{3}, \cos^{-1}\left(-\frac{3}{2}\right)$ (d) $\frac{5\pi}{3}$
56. If $2 \cos^2 x + 3 \sin x - 3 = 0$, $0 \leq x \leq 180^\circ$, then $x =$ [MP PET 1986]
 (a) $30^\circ, 90^\circ, 150^\circ$ (b) $60^\circ, 120^\circ, 180^\circ$ (c) $0^\circ, 30^\circ, 150^\circ$ (d) $45^\circ, 90^\circ, 135^\circ$
57. If $r \sin \theta = 3$, $r = 4(1 + \sin \theta)$, $0 \leq \theta \leq 2\pi$, then $\theta =$ [Roorkee 1974]
 (a) $\frac{\pi}{6}, \frac{\pi}{3}$ (b) $\frac{\pi}{6}, \frac{5\pi}{6}$ (c) $\frac{\pi}{3}, \frac{\pi}{4}$ (d) $\frac{\pi}{2}, \pi$
58. If $\tan(\theta + x)\tan(\theta - x) = 1$ for all x , then value of θ must be
 (a) 0° (b) 30° (c) 45° (d) 60°
59. If $2 \sin^2 \theta = 3 \cos \theta$, where $0 \leq \theta \leq 2\pi$, then $\theta =$ [IIT 1963]
 (a) $\frac{\pi}{6}, \frac{7\pi}{6}$ (b) $\frac{\pi}{3}, \frac{5\pi}{3}$ (c) $\frac{\pi}{3}, \frac{7\pi}{3}$ (d) None of these
60. $\cot \theta = \sin 2\theta$ ($\theta \neq n\pi$, n is integer), if $\theta =$ [Karnataka CET 1993; BIT Ranchi 1991]
 (a) 45° and 60° (b) 45° and 90° (c) 45° only (d) 90° only
61. If $\cos \theta = \frac{-1}{2}$ and $0 < \theta < 360^\circ$, then the values of θ are [Karnataka CET 2001]
 (a) 120° and 300° (b) 60° and 120° (c) 120° and 240° (d) 60° and 240°
62. If $\tan(\pi \cos \theta) = \cot(\pi \sin \theta)$, then $\sin\left(\theta + \frac{\pi}{4}\right)$ equals [AMU 1999]
 (a) $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) $\frac{1}{2\sqrt{2}}$ (d) $\frac{\sqrt{3}}{2}$
63. If $2 \sec(2\alpha) = \tan \beta + \cot \beta$, then one of the values of $(\alpha + \beta)$ is
 (a) π (b) $n\pi - \frac{\pi}{4}$ (c) $\frac{\pi}{4}$ (d) None of these

Advance Level

64. If $3 \sin 2\theta = 2 \sin 3\theta$ and $0 < \theta < \pi$, then value of $\sin \theta$ is [T.S. Rajendra 1992]
 (a) $\frac{\sqrt{2}}{3}$ (b) $\frac{\sqrt{3}}{\sqrt{5}}$ (c) $\frac{\sqrt{15}}{4}$ (d) $\frac{\sqrt{2}}{\sqrt{5}}$
65. $2 \sin^2 x + \sin^2 2x = 2$, $-\pi < x < \pi$, then $x =$ [ISM Dhanbad 1989]
 (a) $\pm \frac{\pi}{6}$ (b) $\pm \frac{\pi}{4}$ (c) $\frac{3\pi}{4}$ (d) None of these
66. If $5 \cos 2\theta + 2 \cos^2 \frac{\theta}{2} + 1 = 0$, $-\pi < \theta < \pi$, then $\theta =$ [Roorkee 1984]

Trigonometrical Equations and Inequations

- (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{3}, \cos^{-1} \frac{3}{5}$ (c) $\cos^{-1} \frac{3}{5}$ (d) None of these
67. If $\sec x \cos 5x + 1 = 0$, where $0 < x < 2\pi$, then $x =$ [Roorkee 1978; IIT 1963]
 (a) $\frac{\pi}{5}, \frac{\pi}{4}$ (b) $\frac{\pi}{5}$ (c) $\frac{\pi}{4}$ (d) None of these
68. The equation $3^{\sin 2x + 2 \cos^2 x} + 3^{1 - \sin 2x + 2 \sin^2 x} = 28$ is satisfied for the values of x given by
 (a) $\cos x = 0$ (b) $\tan x = -1$ (c) $\tan x = 1$ (d) Both (a) and (b)
69. If $e^{((\sin^2 x + \sin^4 x + \sin^6 x + \dots \infty) \log_e 2)}$ satisfies the equation $x^2 - 9x + 8 = 0$, then the value of $\frac{\cos x}{\cos x + \sin x}$, $0 < x < \frac{\pi}{2}$ is [IIT 1991]
 (a) $\frac{1}{2}(\sqrt{3} + 1)$ (b) $\frac{1}{2}(\sqrt{3} - 1)$ (c) $\frac{1}{2}$ (d) 0
70. The value of θ lying between $\theta = 0$ and $\theta = \frac{\pi}{2}$ and satisfying the equation $\begin{vmatrix} 1 + \cos^2 \theta & \sin^2 \theta & 4 \sin 4\theta \\ \cos^2 \theta & 1 + \sin^2 \theta & 4 \sin 4\theta \\ \cos^2 \theta & \sin^2 \theta & 1 + 4 \sin 4\theta \end{vmatrix} = 0$ is [MNR 1991]
 (a) $\frac{11\pi}{24}$ (b) $\frac{7\pi}{24}$ (c) $\frac{5\pi}{24}$ (d) $\frac{\pi}{24}$
71. The smallest positive value of x and y , satisfying $x - y = \frac{\pi}{4}$ and $\cot x + \cot y = 2$ are [Roorkee 2000]
 (a) $x = \frac{\pi}{6}, y = \frac{5\pi}{2}$ (b) $x = \frac{5\pi}{12}, y = \frac{\pi}{6}$ (c) $x = \frac{\pi}{3}, y = \frac{7\pi}{12}$ (d) None of these
72. If $(1 - \tan \theta)(1 + \tan \theta) \sec^2 \theta + 2^{\tan^2 \theta} = 0$ then in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, the value of θ is [IIT 1996]
 (a) $\frac{\pi}{4}$ (b) $-\frac{\pi}{4}$ (c) $\frac{\pi}{3}$ (d) $-\frac{\pi}{3}$

Periodic Functions

Basic Level

73. Period of $\sin \theta \cos \theta$ is
 (a) $\frac{\pi}{2}$ (b) π (c) 2π (d) None of these
74. Period of $\tan 3\theta$ is
 (a) 2π (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{3}$ (d) π
75. Period of $\frac{\sin \theta + \sin 2\theta}{\cos \theta + \cos 2\theta}$ is
 (a) 2π (b) π (c) $\frac{2\pi}{3}$ (d) $\frac{\pi}{3}$
76. Period of $\cos(7x - 5)$ is
 (a) $\frac{2\pi - 5}{7}$ (b) $2\pi - 5$ (c) $\frac{2\pi}{7}$ (d) $\frac{\pi}{7}$
77. Period of $\sin \theta + \cos \theta$ is

72 Trigonometrical Equations and

- (a) π (b) 2π (c) $\frac{\pi}{4}$ (d) None of these
78. Period of $\sin \frac{x}{2} - \cos \frac{x}{3}$ is
 (a) 2π (b) 4π (c) 8π (d) 12π
79. Period of $\cot 3x - \cos(4x + 3)$ is
 (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{4}$ (c) π (d) 2π
80. Period of $|2 \sin 3\theta + 4 \cos 3\theta|$ is
 (a) $\frac{2\pi}{3}$ (b) π (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{3}$
81. The period of the function $\sin\left(\frac{\pi x}{2}\right) + \cos\left(\frac{\pi x}{2}\right)$ is [EAMCET 1990]
 (a) 4 (b) 6 (c) 12 (d) 24
82. The period of the function $|\sin \pi x|$ is [AMU 1999]
 (a) π^2 (b) 2π (c) 2 (d) 1
83. The period of $f(x) = \sin\left(\frac{\pi x}{n-1}\right) + \cos\left(\frac{\pi x}{n}\right), n \in \mathbb{Z}, n > 2$ is
 (a) $2\pi(n-1)$ (b) $4\pi(n-1)$ (c) $2\pi(n-1)$ (d) None of these
84. The period of $a \sin x + b \cos x$ is
 (a) $\frac{\pi}{2}$ (b) π (c) 2π (d) None of these
85. The period of $\sin 8x - \sqrt{3} \cos 8x$ is
 (a) π (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{4}$ (d) 2π
86. The period of $118 \sin 2x - 143 \cot 4x$ is
 (a) 2π (b) π (c) 4π (d) None of these

Advance Level

87. The period of $\sin x \cdot \cos\left(\frac{\pi}{4} - x\right)$ is
 (a) π (b) 2π (c) $\frac{\pi}{2}$ (d) None of these
88. The period of $\frac{7 \sin x + 5 \cos x}{7 \sin 2x + 11 \cos x}$ is
 (a) 2π (b) π (c) $\frac{\pi}{2}$ (d) None of these
89. The period of the function $f(x) = 3 \sin(2x + 1)$ in radians is

- (a) 2π (b) π (c) $\frac{\pi}{2}$ (d) $-\pi$

90. The function f given by $f(x) = \sin\left(\frac{\pi x}{2}\right) + 2 \cos\left(\frac{\pi x}{3}\right) - \tan\left(\frac{\pi x}{4}\right)$ is periodic with period [EAMCET 1992]

- (a) 6 (b) 3 (c) 4 (d) 12

Trigonometrical Inequations

Basic Level

91. The solution set of inequality $\cos^2 \theta < \frac{1}{2}$ is

- (a) $\left\{ \theta : (8n+1)\frac{\pi}{4} < \theta < (8n+3)\frac{\pi}{4}, n \in \mathbb{Z} \right\}$ (b) $\left\{ \theta : (8n-3)\frac{\pi}{4} < \theta < (8n-1)\frac{\pi}{4}, n \in \mathbb{Z} \right\}$
 (c) $\left\{ \theta : (4n+1)\frac{\pi}{4} < \theta < (4n+3)\frac{\pi}{4}, n \in \mathbb{Z} \right\}$ (d) None of these

92. Let $2 \sin^2 x + 3 \sin x - 2 > 0$ and $x^2 - x - 2 < 0$ (x measured in radians) then x lies in the interval

- (a) $\left(\frac{\pi}{6}, \frac{5\pi}{6}\right)$ (b) $\left(-1, \frac{5\pi}{6}\right)$ (c) $(-1, 2)$ (d) $\left(\frac{\pi}{6}, 2\right)$

93. The number of values of $x \in [0, 4\pi]$ satisfying $|\sqrt{3} \cos x - \sin x| \geq 2$ is

- (a) 2 (b) 0 (c) 4 (d) 8

94. If $4 \sin^2 x - 8 \sin x + 3 \leq 0$, $0 \leq x \leq 2\pi$, then the solution set for x is

- (a) $\left[0, \frac{\pi}{6}\right]$ (b) $\left[0, \frac{5\pi}{6}\right]$ (c) $\left[\frac{5\pi}{6}, 2\pi\right]$ (d) $\left[\frac{\pi}{6}, \frac{5\pi}{6}\right]$

95. The number of solutions of $|\cos x| = \sin x$, $0 \leq x \leq 4\pi$, is

- (a) 8 (b) 4 (c) 2 (d) None of these

96. If $\cos x - \sin x \geq 1$ and $0 \leq x \leq 2\pi$ then the solution set for x is


- (a) $\left[0, \frac{\pi}{4}\right] \cup \left[\frac{7\pi}{4}, 2\pi\right]$ (b) $\left[\frac{3\pi}{2}, \frac{7\pi}{4}\right] \cup \{0\}$ (c) $\left[\frac{3\pi}{2}, 2\pi\right] \cup \{0\}$ (d) None of these

97. If $|\tan x| \leq 1$ and $x \in [-\pi, \pi]$ then the solution for x is

- (a) $\left[-\pi, -\frac{3\pi}{4}\right] \cup \left[-\frac{\pi}{4}, \frac{\pi}{4}\right] \cup \left[\frac{3\pi}{4}, \pi\right]$ (b) $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right] \cup \left[\frac{3\pi}{4}, \pi\right]$
 (c) $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$ (d) None of these

98. The set of values of x for which $\sin x \cdot \cos^3 x > \cos x \cdot \sin^3 x$, $0 \leq x \leq 2\pi$ is

- (a) $(0, \pi)$ (b) $\left(0, \frac{\pi}{4}\right)$ (c) $\left(\frac{\pi}{4}, \pi\right)$ (d) None of these



Trigonometrical Equations and

Assignment (Basic & Advance Level)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
b	a	b	b	c	c	d	b	c	c	a	b	a	a	a	b	a	d	b	b
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
c	a	a	a	d	c	a	d	d	b	a	a	c	b	d	d	c	d	a	c
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
b	a	c	c	c	a	c	b	b	b	c	d	a	a	b	a	b	c	b	b
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
c	c	c	c	b	d	c	d	b	a,b	b	c,d	b	c	c	c	b	d	c	d
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98		
a	d	c	c	c	b	a	a	b	d	c	d	c	d	b	c	a	b		