

Sample Question Paper - 30
Mathematics-Standard (041)
Class- X, Session: 2021-22
TERM II

Time Allowed : 2 hours

Maximum Marks : 40

General Instructions :


- The question paper consists of 14 questions divided into 3 sections A, B, C.
- All questions are compulsory.
- Section A comprises of 6 questions of 2 marks each. Internal choice has been provided in two questions.
- Section B comprises of 4 questions of 3 marks each. Internal choice has been provided in one question.
- Section C comprises of 4 questions of 4 marks each. An internal choice has been provided in one question. It contains two case study based questions.

SECTION - A

- Solve the following quadratic equation for x :
 $4x^2 - 4a^2x + (a^4 - b^4) = 0$
- How many two-digit numbers are divisible by 3?
- If O is the centre of a circle, PQ is a chord and the tangent PR at P makes an angle of 50° with PQ , then find $\angle POQ$.

OR

In the given figure, PA and PB are tangents to the circle from an external point P . CD is another tangent touching the circle at Q . If $PA = 12$ cm, $QC = QD = 3$ cm, then find $PC + PD$.



- Data of 'missed catches' for the 40 matches played by a player is as follows :

Number of missed catches in a match	0-3	3-6	6-9	9-12	12-15
Number of matches	15	16	3	4	2

Calculate the mean number of catches missed by him.

- A toy is in the form of a cone mounted on a hemisphere. The diameter of the base of the cone and that of hemisphere is 6 cm and the height of cone is 4 cm. Calculate the surface area of the toy. [Take $\pi = 3.14$]

OR

A toy is in the shape of a cone mounted on a hemisphere of same base radius. If the volume of the toy is 231 cm^3 and its diameter is 7 cm, then find the height of the toy.

$$\left[\text{Use } \pi = \frac{22}{7} \right]$$

6. The mean of a set of numbers is \bar{x} . If each number is multiplied by k , then find the mean of the new set.

SECTION - B

7. Find that non-zero value of k , for which the quadratic equation $kx^2 + 1 - 2(k - 1)x + x^2 = 0$ has equal roots. Hence, find the roots of the equation.

OR

Find two consecutive positive integers, the sum of whose squares is 61.

8. If the sum of first 7 terms of an A.P. is 49 and that of its first 17 terms is 289, find the sum of first n terms of the A.P.
9. A 7 m long flagstaff is fixed on the top of a tower standing on the horizontal plane. From a point on the ground, the angles of elevation of the top and bottom of the flagstaff are 60° and 45° respectively. Find the height of the tower correct to one place of decimal. (Use $\sqrt{3} = 1.73$)
10. Draw a circle of radius 2.4 cm. Take a point P on it. Without using the centre of the circle, draw a tangent to the circle at point P .

SECTION - C

11. As observed from the top of a 100 m high light house from the sea-level, the angles of depression of two ships are 30° and 45° . If one ship is exactly behind the other on the same side of the light house, find the distance between the two ships. [Use $\sqrt{3} = 1.732$]
12. The mode of the following data is 36. Find the missing frequency x in it.

Class	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	8	10	x	16	12	6	7

OR

If the median of the following frequency distribution is 32.5, then find the values of f_1 and f_2 .

Class	0-10	10-20	20-30	30-40	40-50	50-60	60-70	Total
Frequency	f_1	5	9	12	f_2	3	2	40


Case Study - 1

13. For class 10 students, a teacher planned a game for the revision of chapter circles with some questions written on the board, which are to be answered by the students. For each correct answer, a student will get a reward. Some of the questions are given below.



Answer these questions to check your knowledge.

- (i) If PA and PB are two tangents drawn to a circle with centre O from P such that $\angle PBA = 50^\circ$, then find the measure of $\angle OAB$.
- (ii) In the adjoining figure, AB is a chord of the circle and AOC is its diameter such that $\angle ACB = 55^\circ$, then find the measure of $\angle BAT$.



Case Study - 2

14. Isha's father brought an ice-cream brick, empty cones and scoop to pour the ice-cream into cones for all the family members. Dimensions of the ice-cream brick were $(30 \times 25 \times 10)$ cm³ and radius of hemi-spherical scoop was 3.5 cm. Also, the radius and height of cone were 3.5 cm and 15 cm respectively.



Based on the above information, answer the following questions.

- (i) Find the quantity of ice-cream in the brick (in litres).
- (ii) Find the minimum number of scoops required to fill one cone upto brim.

Solution

MATHEMATICS STANDARD 041


Class 10 - Mathematics

1. We have, $4x^2 - 4a^2x + (a^4 - b^4) = 0$
 $\Rightarrow (2x)^2 - 2(2x)a^2 + (a^2)^2 - (b^2)^2 = 0$
 $\Rightarrow (2x - a^2)^2 - (b^2)^2 = 0$
 $\Rightarrow (2x - a^2 + b^2)(2x - a^2 - b^2) = 0$
 $\Rightarrow 2x - a^2 + b^2 = 0$ or $2x - a^2 - b^2 = 0$
 $\Rightarrow 2x = a^2 - b^2$ or $2x = a^2 + b^2$
 $\Rightarrow x = \frac{a^2 - b^2}{2}$ or $x = \frac{a^2 + b^2}{2}$

2. Two-digit numbers which are divisible by 3 are 12, 15, 18, ..., 99, which forms an A.P. with first term (a) = 12, common difference (d) = $15 - 12 = 3$ and last term (l) or n^{th} term (a_n) = 99
 $\therefore a + (n - 1)d = 99$
 $\Rightarrow 12 + (n - 1)3 = 99 \Rightarrow 3n = 99 - 9$
 $\Rightarrow n = \frac{90}{3} = 30$
 Thus, there are 30 two-digit numbers which are divisible by 3.

3. $\therefore PR$ is a tangent to the circle.

$\therefore OP \perp PR$
 $\Rightarrow \angle OPR = 90^\circ$
 $\Rightarrow \angle OPQ + \angle QPR = 90^\circ$
 $\Rightarrow \angle OPQ = 90^\circ - 50^\circ = 40^\circ$
 Now, $OP = OQ$ (Radii of circle)
 $\Rightarrow \angle OPQ = \angle OQP = 40^\circ$



In $\triangle OPQ$, $\angle OPQ + \angle OQP + \angle POQ = 180^\circ$
 [By angle sum property]
 $\Rightarrow \angle POQ = 180^\circ - 40^\circ - 40^\circ = 100^\circ$

OR

Tangents drawn from an external point are equal in length.

$\therefore QC = CA$, $QD = BD$ and $PA = PB$
 Since, $QC = QD = 3$ cm [Given]
 $\Rightarrow CA = BD = 3$ cm
 Also, $PC = PA - AC$
 $\Rightarrow PC = (12 - 3)$ cm = 9 cm [Given, $PA = 12$ cm]
 Similarly, $PD = 9$ cm
 $\therefore PC + PD = 9 + 9 = 18$ cm

4. The frequency distribution table from the given data can be drawn as :

Missed catches	Class marks (x_i)	Frequency (f_i)	$f_i x_i$
0-3	1.5	15	22.5
3-6	4.5	16	72

6-9	7.5	3	22.5
9-12	10.5	4	42
12-15	13.5	2	27
		$\Sigma f_i = 40$	$\Sigma f_i x_i = 186$

$$\therefore \text{Mean} = \frac{\Sigma f_i x_i}{\Sigma f_i} = \frac{186}{40} = 4.65$$

5. Radius of the base of the cone and hemisphere (r)
 $= \frac{6}{2} = 3$ cm

Height of cone (h) = 4 cm

Slant height of cone (l)

$$= \sqrt{r^2 + h^2} = \sqrt{3^2 + 4^2}$$

$$= \sqrt{9 + 16} = \sqrt{25} = 5 \text{ cm}$$

Total surface area of toy

= Curved surface area of hemisphere + Curved surface area of cone

$$= 2\pi r^2 + \pi r l = \pi r(2r + l) = 3.14 \times 3(2 \times 3 + 5)$$

$$= 3.14 \times 3 \times 11 = 103.62 \text{ cm}^2$$

OR

Let h be the height of the cone and r be its radius.

Given, $r = (7/2)$ cm = 3.5 cm

Volume of the toy = 231 cm³

\Rightarrow Volume of cone +

Volume of hemisphere = 231 cm³

$$\Rightarrow \frac{1}{3} \pi r^2 h + \frac{2}{3} \pi r^3 = 231$$

$$\Rightarrow \frac{1}{3} \pi r^2 (h + 2r) = 231$$

$$\Rightarrow \frac{1}{3} \times \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \left(h + 2 \times \frac{7}{2} \right) = 231 \Rightarrow \frac{77}{6} (h + 7) = 231$$

$$\Rightarrow h + 7 = \frac{231 \times 6}{77}$$

$$\Rightarrow h + 7 = 18 \Rightarrow h = 11 \text{ cm}$$



\therefore Height of cone is 11 cm

Total height of toy = Height of cone + radius of hemisphere = $11 + 3.5 = 14.5$ cm

6. Let the numbers are x_1, x_2, \dots, x_n .

$$\therefore \text{Mean} = \frac{x_1 + x_2 + \dots + x_n}{n} = \bar{x} \quad \dots(i)$$

When given numbers are multiplied by k , then new observations are kx_1, kx_2, \dots, kx_n .



$$\begin{aligned} \text{New mean} &= \frac{kx_1 + kx_2 + \dots + kx_n}{n} \\ &= \frac{k(x_1 + x_2 + \dots + x_n)}{n} = k\bar{x} \end{aligned} \quad (\text{From (i)})$$

$$7. \text{ We have, } kx^2 + 1 - 2(k-1)x + x^2 = 0 \quad \dots(i)$$

$$\text{or } (k+1)x^2 - 2(k-1)x + 1 = 0$$

Since, roots are equal. $\therefore D = 0$

$$\Rightarrow \{-2(k-1)\}^2 - 4 \times (k+1) \times 1 = 0$$

$$\Rightarrow 4k^2 - 8k + 4 - 4k - 4 = 0$$

$$\Rightarrow 4k^2 - 12k = 0 \Rightarrow 4k(k-3) = 0$$

$$\Rightarrow k = 0 \text{ or } k - 3 = 0 \Rightarrow k = 3$$

$$\Rightarrow k = 3 \text{ (Non-zero value of } k)$$

Substituting the value of k in (i), we get

$$(3+1)x^2 - 2(3-1)x + 1 = 0$$

$$\Rightarrow 4x^2 - 4x + 1 = 0 \Rightarrow 4x^2 - 2x - 2x + 1 = 0$$

$$\Rightarrow 2x(2x-1) - 1(2x-1) = 0 \Rightarrow (2x-1)(2x-1) = 0$$

$$\Rightarrow 2x-1 = 0 \text{ or } 2x-1 = 0 \Rightarrow x = \frac{1}{2}, \frac{1}{2}$$

OR

Let the two consecutive positive integers be x and $x+1$.

$$\text{According to question, } x^2 + (x+1)^2 = 61$$

$$\Rightarrow x^2 + x^2 + 2x + 1 = 61$$

$$\Rightarrow 2x^2 + 2x = 60 \Rightarrow x^2 + x - 30 = 0$$

$$\Rightarrow (x-5)(x+6) = 0$$

$$\Rightarrow x = 5 \text{ or } x = -6$$

$$\Rightarrow x = 5 \quad [\text{Since } x \text{ is a positive integer}]$$

$$\text{And } x+1 = 6$$

\therefore The two consecutive positive integers are 5 and 6.

8. Let a be the first term and d be the common difference of the A.P.

$$\text{Sum of } n \text{ terms, } S_n = \frac{n}{2}[2a + (n-1)d]$$

$$\text{We have, } S_7 = 49$$

$$\Rightarrow \frac{7}{2}[2a + 6d] = 49$$

$$\Rightarrow 14a + 42d = 98 \Rightarrow a + 3d = 7 \quad \dots(i)$$

$$\text{and } S_{17} = 289$$

$$\Rightarrow \frac{17}{2}[2a + 16d] = 289$$

$$\Rightarrow 34a + 272d = 578 \Rightarrow a + 8d = 17 \quad \dots(ii)$$

On solving (i) and (ii), we get $a = 1, d = 2$

$$\therefore S_n = \frac{n}{2}[2 + (n-1)2] = n^2$$

9. Let AB be the tower of height h m and AD be the flagstaff and C be the required point on the ground at the distance of x m from the tower.

$$\therefore AD = 7 \text{ m}$$

$$\text{In } \triangle BCD, \tan 60^\circ = \frac{BD}{BC} = \frac{h+7}{x}$$

$$\Rightarrow \sqrt{3}x = h+7 \Rightarrow x = \frac{h+7}{\sqrt{3}} \quad \dots(i)$$

$$\text{In } \triangle ABC, \tan 45^\circ = \frac{AB}{BC} = \frac{h}{x}$$

$$\Rightarrow x = h$$

$$\Rightarrow \frac{h+7}{\sqrt{3}} = h \quad [\text{Using (i)}]$$


$$\Rightarrow \sqrt{3}h = h+7$$

$$\Rightarrow (\sqrt{3}-1)h = 7$$

$$\Rightarrow h = \frac{7}{\sqrt{3}-1} = \frac{7}{0.73} = 9.58 \approx 9.6$$

Hence, height of tower is 9.6 m.

10. Given, radius of circle = 2.4 cm



Steps of construction

Step-I : Draw a circle of radius 2.4 cm and take a point P on the circle.

Step-II : Draw a chord PQ through the point P on the circle.

Step-III : Take a point R on the major arc and join PR and RQ .

Step-IV : On taking PQ as base, construct $\angle QPY = \angle PRQ$.

Step-V : Produce YP to X . Then, YPX is the required tangent at point P .

11. In the figure, let AB represent the light house.

$$\therefore AB = 100 \text{ m}$$

Let the positions of two ships be at C and D such that angle of depression from A are 45° and 30° respectively.

Now, in right $\triangle ABC$,

$$\frac{AB}{BC} = \tan 45^\circ$$



$$\Rightarrow \frac{100}{BC} = 1 \Rightarrow BC = 100 \text{ m}$$

Again, in right $\triangle ABD$, we have

$$\frac{AB}{BD} = \tan 30^\circ$$

$$\Rightarrow \frac{100}{BD} = \frac{1}{\sqrt{3}} \Rightarrow BD = 100\sqrt{3} \text{ m}$$

The distance between the two ships = CD



$$= BD - BC = 100\sqrt{3} - 100$$

$$= 100(\sqrt{3} - 1) = 100(1.732 - 1)$$

$$= 100 \times 0.732 = 73.2 \text{ m}$$

Thus, the required distance between the ships is 73.2 m.

12. Since it is given that mode = 36, which lies in the interval 30-40

\therefore Modal class is 30-40.

$$\therefore l = 30, f_1 = 16, f_0 = x, f_2 = 12, h = 10$$

$$\text{Now, Mode} = l + \left(\frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

$$\Rightarrow 36 = 30 + \left(\frac{16 - x}{2 \times 16 - x - 12} \right) \times 10$$

$$\Rightarrow 36 = 30 + \left(\frac{16 - x}{20 - x} \right) \times 10$$

$$\Rightarrow (20 - x) \times 6 = (16 - x) \times 10$$

$$\Rightarrow 120 - 6x = 160 - 10x \Rightarrow 4x = 40 \Rightarrow x = 10$$

OR

The frequency distribution table for the given data is as follows :

Class	Frequency (f_i)	Cumulative frequency ($c.f.$)
0-10	f_1	f_1
10-20	5	$f_1 + 5$
20-30	9	$f_1 + 14$
30-40	12	$f_1 + 26$
40-50	f_2	$f_1 + f_2 + 26$
50-60	3	$f_1 + f_2 + 29$
60-70	2	$f_1 + f_2 + 31$
Total	$31 + f_1 + f_2 = 40$	

$$\text{Here, } N = 40 \Rightarrow 31 + f_1 + f_2 = 40$$

$$\Rightarrow f_1 + f_2 = 9 \quad \dots (i)$$

Given, median = 32.5, which lies in the interval 30-40.

So, median class is 30-40.

$$\therefore l = 30, h = 10, f = 12, N = 40 \text{ and}$$

$$c.f. = f_1 + 14$$

$$\text{Now, median} = l + \left(\frac{\frac{N}{2} - c.f.}{f} \right) \times h$$


$$\Rightarrow 32.5 = 30 + \left(\frac{20 - (f_1 + 14)}{12} \right) \times 10$$

$$\Rightarrow 2.5 = \left(\frac{6 - f_1}{12} \right) \times 10 \Rightarrow 6 - f_1 = \frac{2.5 \times 12}{10}$$

$$\Rightarrow 6 - f_1 = 3 \Rightarrow f_1 = 3$$

$$\text{From (i), } f_2 = 9 - 3 = 6$$

13. (i)



Since, $OB \perp PB$ [Since, radius at the point of contact is perpendicular to tangent]

and $\angle PBA = 50^\circ$ (Given)

$$\therefore \angle OBA = 90^\circ - 50^\circ = 40^\circ$$

Also, $OA = OB$

[Radii of circle]

$$\therefore \angle OAB = \angle OBA = 40^\circ$$

[Angle opposite to equal sides are equal]

(ii) Here, $\angle ABC = 90^\circ$ (Angle in a semicircle)

So, in $\triangle ABC$, $\angle BAC = 180^\circ - 90^\circ - 55^\circ = 35^\circ$

Also, $\angle OAT = 90^\circ$

$$\Rightarrow \angle BAT + \angle OAB = 90^\circ \Rightarrow \angle BAT = 90^\circ - 35^\circ = 55^\circ$$

14. (i) Quantity of ice-cream in the brick

$$= \text{Volume of the brick} = (30 \times 25 \times 10) \text{ cm}^3 = 7500 \text{ cm}^3$$

$$= \frac{7500}{1000} \text{ l} \quad [\because 1 \text{ l} = 1000 \text{ cm}^3]$$

$$= 7.5 \text{ l}$$

(ii) Volume of hemispherical scoop = $\frac{2}{3} \pi r^3$

$$= \frac{2}{3} \times \frac{22}{7} \times (3.5)^3 = \frac{1886.5}{21} = 89.83 \text{ cm}^3$$

Now, volume of cone = $\frac{1}{3} \pi r^2 h$

$$= \frac{1}{3} \times \frac{22}{7} \times 3.5 \times 3.5 \times 15 = \frac{4042.5}{21} = 192.5 \text{ cm}^3$$

\therefore Number of scoops required to fill one cone

$$= \frac{\text{Volume of a cone}}{\text{Volume of a scoop}} = \frac{192.5}{89.83} = 2.14 \approx 2$$