Sample Question Paper - 27 Mathematics-Standard (041) Class- X, Session: 2021-22 TERM II


Time Allowed : 2 hours

General Instructions :

- 1. The question paper consists of 14 questions divided into 3 sections A, B, C.
- 2. All questions are compulsory.
- 3. Section A comprises of 6 questions of 2 marks each. Internal choice has been provided in two questions.
- 4. Section B comprises of 4 questions of 3 marks each. Internal choice has been provided in one question.
- 5. Section C comprises of 4 questions of 4 marks each. An internal choice has been provided in one question. It contains two case study based questions.

SECTION - A

- 1. The mean of marks in Mathematics for 40 students in a class was 56.5. Later it was found that the marks given to one student was incorrectly entered as 85 instead of 58. Find the correct mean for the class.
- **2.** From the top of a cliff 20 m high, the angle of elevation of the top of a tower is found to be equal to the angle of depression of the foot of the tower. Find the height of the tower.
- 3. In figure, *XP* and *XQ* are tangents from *X* to the circle with centre *O*. *R* is a point on the circle. Prove that, XA + AR = XB + BR.



In given fig., two chords *AB* and *CD* of a circle intersect at *O*. If AO = 8 cm, CO = 6 cm and OD = 4 cm, then find the length of *OB*.



- 4. The string of a kite is 120 m long and it makes an angle of 60° with the horizontal. Find the height of the kite assuming that there is no slack in the string. [Use $\sqrt{3} = 1.732$]
- 5. A solid is in the shape of a cone mounted on a hemisphere of same base radius. If the curved surface areas of the hemispherical part and the conical part are equal, then find the ratio of the radius and the height of the conical part.

Maximum Marks : 40

OR

Two cubes each of 10 cm edge are joined end to end. Find the surface area of the resulting cuboid.

6. Find the mean of following distribution:

x_i	4	6	9	10	15
f_i	5	10	10	7	8

SECTION - B

- 7. If the sum of *n* terms of an A.P. is given by $S_n = (3n^2 + 2n)$, find its (i) n^{th} term (ii) first term (iii) common difference.
- 8. Draw a line segment of length 7 cm and divide it internally in the ratio 2 : 3.
- **9.** The digits of a positive number of three digits are in A.P. and their sum is 15. The number obtained by reversing the digits is 594 less than the original number. Find the number.

OR

In an A.P., $S_m = n$ and $S_n = m$ also m > n, find the sum of first (m - n) terms.

10. The shadow of a tower standing on a leveled ground is found to be 30 m longer when the sun's altitude is 30° than when it is 60°? Find the height of the tower.

SECTION - C

11. A train travels at a certain average speed for a distance of 63 km and then travels a distance of 72 km at an average speed of 6 km/hr more than its original speed. If it takes 3 hours to complete total journey, what is the original average speed?

OR

The diagonal of a rectangular field is 60 metres more than the shorter side. If the longer side is 30 metres more than the shorter side, find the sides of the field.


12. From a solid cylinder whose height is 15 cm and diameter 16 cm, a conical cavity of the same height and same diameter is hollowed out. Find the total surface area of the remaining solid. [Take $\pi = 3.14$]

Case Study - 1

13. Raghav loves geometry. So he was curious to know more about the concepts of circle. His father is a mathematician. So, he reached to his father to learn something interesting about tangents and circles. His father gave him knowledge on circles and tangents and ask him to solve the following questions.



(i) A circle of radius 3 cm is inscribed in a right angled triangle *BAC* such that BD = 9 cm and DC = 3 cm. Find the length of *AB*.



(ii) If *PA* and *PB* are two tangents to a circle with centre *O* from an external point *P* such that $\angle OPB = 40^\circ$, then find the value of $\angle BPA$.

Case Study - 2

14. Household income in India was drastically impacted due to the COVID-19 lockdown. Most of the companies decided to bring down the salaries of the employees by 50%.

The following table shows the salaries (in percent) received by 25 employees during lockdown.

Salaries received (in percent)	50-60	60-70	70-80	80-90
Number of employees	9	6	8	2



Based on the above information, answer the following questions.

- (i) Find the median class of the given data.
- (ii) If x_i 's denotes the class marks and f_i 's denotes the corresponding frequencies for the given data, then find $\sum x_i f_i$.

Solution

MATHEMATICS STANDARD 041

Class 10 - Mathematics


1. Number of students in the class(n) = 40

Now, mean $(\bar{x}) = \frac{\sum x}{n}$ $\Rightarrow 56.5 = \frac{\sum x}{40}$ [Given] $\Rightarrow \sum x = 56.5 \times 40 \Rightarrow \sum x = 2260$

But 2260 is incorrect sum.

Since, correct sum = 2260 - 85 + 58 = 2233 \therefore Correct mean = $\frac{2233}{40} = 55.82$

2. Let *AB* be the cliff and EC = h m be the height of tower.



$$\tan \theta = \frac{ED}{AD} = \frac{ED}{BC} = \frac{(h-20)\tan\theta}{20}$$
 [(Using (i)]

 \Rightarrow $h - 20 = 20 \Rightarrow h = 40$

: Height of tower is 40 m.

3. Since, length of tangents from an exterior point to a circle are equal.

$$\therefore XP = XQ \text{ [Tangents from } X\text{]} \qquad \dots(i)$$

$$AP = AR[\text{Tangents from } A\text{]} \qquad \dots(ii)$$

$$BQ = BR \text{ [Tangents from } B\text{]} \qquad \dots(iii)$$

$$Now, XP = XQ$$

$$\Rightarrow XA + AP = XB + BQ$$

$$\Rightarrow XA + AR = XB + BR \qquad \text{[Using (ii) and (iii)]}$$

We have, OA = 8 cm, OC = 6 cm, OD = 4 cm Since, *AB* and *CD* are chords of the circle intersect at *O*.

 $\therefore OA \times OB = OC \times OD$

- $\Rightarrow 8 \times OB = 6 \times 4$
- $\Rightarrow OB = 3 \text{ cm}$

4. Let *OA* be the horizontal ground and *K* be the position of the kite at a height *h* m above the ground. Then, AK = h m, OK = 120 m and $\angle AOK = 60^{\circ}$

In
$$\triangle AOK$$
, sin $60^\circ = \frac{AK}{OK}$
 $\Rightarrow \frac{\sqrt{3}}{2} = \frac{h}{120}$
 $\Rightarrow h = 120 \times \frac{\sqrt{3}}{2} = 60\sqrt{3}$
 $\Rightarrow h = 60 \times 1.732 = 103.92$

Hence, the height of the kite is 103.92 m.

5. Let *r* be the radius of cone or hemispherical part and *h* be the height of cone.

According to question,

- $2\pi r^{2} = \pi r l, \text{ where } l \text{ is slant height of cone}$ $\Rightarrow 2r = l \Rightarrow 4r^{2} = l^{2}$ $\Rightarrow 4r^{2} = r^{2} + h^{2}$ $\Rightarrow 3r^{2} = h^{2}$ $\Rightarrow \frac{r}{h} = \frac{1}{\sqrt{3}}$
- \therefore Required ratio = 1: $\sqrt{3}$

OR

If two cubes are joined end to end, we get a cuboid such that 10

l = Length of the resulting cuboid

= 10 cm + 10 cm = 20 cm

- b = Breadth of the resulting cuboid = 10 cm
- h = Height of the resulting cuboid = 10 cm
- :. Surface area of the cuboid = 2(lb + bh + lh)
- \Rightarrow Surface area of the cuboid

$$= 2(20 \times 10 + 10 \times 10 + 20 \times 10) \text{ cm}^2$$

 \therefore Surface area of the cuboid is 1000 cm²

6. Let us construct the following table for the given data:

x _i	f_i	$f_i x_i$
4	5	20
6	10	60
9	10	90
10	7	70
15	8	120
Total	$\Sigma f_i = 40$	$\Sigma f_i x_i = 360$

:. Mean
$$(\overline{x}) = \frac{\sum_{i=1}^{5} x_i f_i}{\sum_{i=1}^{5} f_i} = \frac{360}{40} = 9$$

7. Given, $S_n = (3n^2 + 2n)$ $\therefore S_{n-1} = \{3 (n-1)^2 + 2 (n-1)\}$ $= (3n^2 - 4n + 1)$ (i) The *n*th term is given by $T_n = (S_n - S_{n-1}) = \{(3n^2 + 2n) - (3n^2 - 4n + 1)\} = (6n - 1)$ $\therefore n^{\text{th}}$ term = (6n - 1) ...(1) (ii) Putting n = 1 in (1), we get $T_1 = (6 \times 1 - 1) = 5$ \therefore First term = 5(iii) Putting n = 2 in (1), we get $T_2 = (6 \times 2 - 1) = 11$

:. $d = T_2 - T_1 = 11 - 5 = 6$.

8. Steps of construction :

Step-I : Draw a line segment AB = 7 cm.


Step-II : Draw any ray *AX* making an acute angle with *AB*.

Step-III : On ray *AX*, locate 5(=2+3) points A_1 , A_2 , A_3 , A_4 , A_5 such that $AA_1 = A_1A_2 = A_2A_3 = A_3A_4 = A_4A_5$.

Step-IV : Join A_5B .

Step-V : From A_2 , draw a line parallel to A_5B , intersecting AB at P.

Thus, *P* divides *AB* in the ratio 2 : 3.



9. Let the required three digit number be *xyz*.

:: Digits are in A.P.

 \therefore x = y - d and z = y + d where *d* is common difference According to question,

(y - d) + y + (y + d) = 15

 \Rightarrow 3y = 15 \Rightarrow y = 5

Since, the number obtained by reversing the digits (z y x)*i.e.*, 100z + 10y + x is 594 less than original number.

- $\therefore (100x + 10y + z) (100z + 10y + x) = 594$
- $\Rightarrow (z 100z) + (100x x) = 594$
- $\Rightarrow 99x 99z = 594$
- $\Rightarrow x z = 6$
- $\Rightarrow (y-d) (y+d) = 6$

$$\Rightarrow -2d = 6 \Rightarrow d = -3$$

So,
$$x = y - d = 5 - (-3) = 8$$
 and $z = y + d = 5 - 3 = 2$

 \therefore The number is *xyz* or 852.

OR

Let a, a + d, a + 2d, be the A.P. Given, $S_m = \frac{m}{2} [2a + (m-1)d] = n$ $\Rightarrow 2a + (m-1)d = \frac{2n}{m}$...(1) And $S_n = \frac{n}{2} [2a + (n-1)d] = m$

$$\Rightarrow 2a + (n-1)d = \frac{2m}{n} \qquad \dots (2)$$

Subtracting (2) from (1), we get

$$2a + (m-1)d - 2a - (n-1)d = \frac{2n}{m} - \frac{2m}{n}$$

$$\Rightarrow (m-1-n+1)d = \frac{2n^2 - 2m^2}{mn}$$

$$\Rightarrow d = -\frac{2(m+n)}{mn} \qquad ...(3)$$

Now, $S_{m-n} = \frac{m-n}{2} \{2a + (m-n-1)d\}$

$$= \frac{m-n}{2} \{2a + (m-1)d - nd\}$$

$$= \frac{m-n}{2} \left\{\frac{2n}{m} + \frac{2(m+n)}{m}\right\} \qquad [Using (1) \& (3)]$$

$$= \frac{m-n}{2} \left\{\frac{4n+2m}{m}\right\} = \frac{(m-n)(2n+m)}{m}$$

10. Let *AB* be the tower and *AC* and *AD* be its shadows when the angles of elevation are 60° and 30° respectively.

Then CD = 30 m.

Let *h* be the height of the tower and let AC = x m. In $\triangle ABC$, right angled at *A* $\tan 60^\circ = \frac{AB}{AC}$ $D = \frac{1}{200} + \frac{1}{200$

$$\Rightarrow \sqrt{3} = \frac{h}{x} \Rightarrow x = \frac{h}{\sqrt{3}} \qquad \dots (1)$$

In ΔDAB , we have

$$\tan 30^\circ = \frac{AB}{AD}$$
$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{x+30} \Rightarrow x+30 = \sqrt{3}h \qquad \dots(2)$$

Putting value of x from (1) in (2), we get

$$\frac{h}{\sqrt{3}} + 30 = \sqrt{3} h$$

$$\Rightarrow h + 30\sqrt{3} = 3h \Rightarrow 2h = 30\sqrt{3} \Rightarrow h = 15\sqrt{3}$$

Thus, the height of the tower is $15\sqrt{3}$ m.

11. Let original average speed of the train be *x* km/hr. According to question,

$$\frac{63}{x} + \frac{72}{x+6} = 3$$

$$\Rightarrow \frac{7}{x} + \frac{8}{x+6} = \frac{1}{3} \Rightarrow \frac{7(x+6)+8x}{x(x+6)} = \frac{1}{3}$$

$$\Rightarrow 3 (7x+42+8x) = x^2+6x$$

$$\Rightarrow 45x+126 = x^2+6x \Rightarrow x^2-39x-126 = 0$$

$$\Rightarrow x^2-42x+3x-126 = 0 \Rightarrow (x-42) (x+3) = 0$$

$$\Rightarrow \text{ Either } x = 42 \text{ or } x = -3$$

$$\Rightarrow x = 42 (\because x > 0)$$
Hence, the original speed of the train is 42 km/h.
OR
Let the shorter side (*i.e.*, breadth) = x m.

$$\therefore \text{ The longer side (length)} = (x+30) \text{ m.}$$
In a rectangle,
diagonal = $\sqrt{(\text{breadth})^2 + (\text{length})^2}$

$$\Rightarrow x+60 = \sqrt{x^2 + (x+30)^2}$$

$$\Rightarrow x+60 = \sqrt{x^2 + (x+30)^2}$$

$$\Rightarrow x+60 = \sqrt{x^2 + x^2 + 60x + 900}$$

$$\Rightarrow (x+60)^2 = 2x^2 + 60x + 900 [\text{Squaring both sides}]$$

$$\Rightarrow x^2 - 60x - 2700 = 0$$

$$\Rightarrow x = \frac{-(-60) \pm \sqrt{14400}}{2(1)} \Rightarrow x = \frac{60 \pm 120}{2} = 90, -30$$
As, $x \neq -30 \Rightarrow x = 90$ [\because breadth cannot be negative]

:. The longer side = x + 30 = 90 + 30 = 120Thus, the shorter side = 90 m. The longer side = 120 m.

12. Height of cylinder, h = 15 cm

Radius of cylinder,

$$r = \frac{16}{2} = 8 \text{ cm}$$

Height of cone, h = 15 cm Radius of the cone, r = 8 cm

Slant height of the cone, $l = \sqrt{8^2 + 15^2}$ = $\sqrt{64 + 225} = \sqrt{289} = 17$ cm \therefore Curved surface area of the cone = πrl = $3.14 \times 8 \times 17 = 427.04$ cm² Curved surface area of the cylinder = $2\pi rh$ = $2 \times 3.14 \times 8 \times 15 = 753.6$ cm² Area of the top face of the cylinder = πr^2 = $3.14 \times (8)^2 = 200.96$ cm² ... Total surface area of the remaining solid

= Area of the top face of the cylinder + curved surface area of the cylinder + curved surface area of the cone = 200.96 + 753.6 + 427.04 = 1381.6 cm²

13. (i) Let AF = AE = x cm

[: Tangents drawn from an external point to a circle are equal in length]

- $\therefore BD = FB = 9 \text{ cm}, CD = CE = 3 \text{ cm}$
- In $\triangle ABC$, $AB^2 = AC^2 + BC^2$ $\Rightarrow (AF + FB)^2 = (AE + EC)^2 + (BD + CD)^2$
- $\Rightarrow (x+9)^2 = (x+3)^2 + (12)^2 \Rightarrow 18x + 81 = 6x + 9 + 144$
- $\Rightarrow 12x = 72 \Rightarrow x = 6 \text{ cm}$
- $\therefore AB = 6 + 9 = 15 \text{ cm}$

(ii) Here, $\angle OAP = 90^{\circ}$

In $\triangle AOP$ and $\triangle BOP$,

 $\angle OAP = \angle OBP = 90^{\circ}$

OA = OB [Radii of circle]

- *PA* = *PB* [Tangents drawn from an external point are equal]
- $\therefore \quad \Delta AOP \cong \Delta BOP \ [By SAS congruency] \\ \therefore \quad \angle APO = \angle OPB = 40^{\circ}$

[By C.P.C.T.]

×40°

 $\therefore \quad \angle BPA = 40^\circ + 40^\circ = 80^\circ$

14. (i) The cumulative frequency distribution table for the given data can be drawn as :

Salaries received (in percent)	Number of employees (f_i)	Cumulative frequency (<i>c.f.</i>)
50-60	9	9
60-70	6	9 + 6 = 15
70-80	8	15 + 8 = 23
80-90	2	23 + 2 = 25
Total	$\Sigma f_i = 25$	

Here,
$$\frac{N}{2} = \frac{25}{2} = 12.5$$

The cumulative frequency just greater than 12.5 lies in the interval 60-70.

Hence, the median class is 60-70.

(ii) Let us consider the following table :

	0			
Class	Class mark (x _i)	Frequency (f _i)	$x_i f_i$	
50-60	55	9	495	
60-70	65	6	390	
70-80	75	8	600	
80-90	85	2	170	
Total		$\Sigma f_i = 25$	$\Sigma x_i f_i = 1,655$	

