

Inverse Trigonometric Functions

- If $\sin y = x$, then $y = \sin^{-1}x$ (We read it as sine inverse x)

Here, $\sin^{-1}x$ is an inverse trigonometric function. Similarly, the other inverse trigonometric functions are as follows:

- If $\cos y = x$, then $y = \cos^{-1}x$
- If $\tan y = x$, then $y = \tan^{-1}x$
- If $\cot y = x$, then $y = \cot^{-1}x$
- If $\sec y = x$, then $y = \sec^{-1}x$
- If $\cosec y = x$, then $y = \cosec^{-1}x$
- The domains and ranges (principle value branches) of inverse trigonometric functions can be shown in a table as follows:

Function	Domain	Range (Principle value branches)
$y = \sin^{-1}x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$
$y = \cos^{-1}x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1}x$	\mathbf{R}	$\left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$
$y = \cot^{-1}x$	\mathbf{R}	$(0, \pi)$
$y = \sec^{-1}x$	$\mathbf{R} - (-1, 1)$	$[0, \pi] - \left\{ \frac{\pi}{2} \right\}$
$y = \cosec^{-1}x$	$\mathbf{R} - (-1, 1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] - \{0\}$

- Note that $y = \tan^{-1}x$ does not mean that $y = (\tan x)^{-1}$. This argument also holds true for the other inverse trigonometric functions.
- The principal value of an inverse trigonometric function can be defined as the value of inverse trigonometric functions, which lies in the range of principal branch.

Example 1: What is the principal value of $\tan^{-1}(-\sqrt{3}) + \sin^{-1}(1)$?




Solution: Let $\sin^{-1}(1) = z$
 $\Rightarrow \tan y = -\sqrt{3} = -\tan\left(\frac{\pi}{3}\right) = \tan\left(-\frac{\pi}{3}\right)$ and $\sin z = 1 = \sin\frac{\pi}{2}$




We know that the ranges of principal value branch of \tan^{-1} and \sin^{-1} are $(-\frac{\pi}{2}, \frac{\pi}{2})$ and $[-\frac{\pi}{2}, \frac{\pi}{2}]$ respectively. Also, $\tan(-\frac{\pi}{3}) = -\sqrt{3} \sin(\frac{\pi}{2}) = 1$

Therefore, principal values of $\tan^{-1}(-\sqrt{3}) = -\frac{\pi}{3}$ and $\sin^{-1}(1) = \frac{\pi}{2}$

$$\therefore \tan^{-1}(-\sqrt{3}) + \sin^{-1}1 = -\frac{\pi}{3} + \frac{\pi}{2} = \frac{\pi}{6}$$

- Graphs of the six inverse trigonometric functions can be drawn as follows:





- The relation $\sin y = x \Rightarrow y = \sin^{-1} x$ gives $\sin(\sin^{-1} x) = x$, where $x \in [-1, 1]$; and $\sin^{-1}(\sin x) = x$, where $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

This property can be similarly stated for the other inverse trigonometric functions as follows:

- $\cos(\cos^{-1}x) = x, x \in [-1, 1]$ and $\cos^{-1}(\cos x) = x, x \in [0, \pi]$
- $\tan(\tan^{-1}x) = x, x \in \mathbf{R}$ and $\tan^{-1}(\tan x) = x, x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- $\text{cosec}(\text{cosec}^{-1}x) = x, x \in \mathbf{R} - (-1, 1)$ and $\text{cosec}^{-1}(\text{cosec } x) = x, x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
- $\sec(\sec^{-1}x) = x, x \in \mathbf{R} - (-1, 1)$ and $\sec^{-1}(\sec x) = x, x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$
- $\cot(\cot^{-1}x) = x, x \in \mathbf{R}$ and $\cot^{-1}(\cot x) = x, x \in (0, \pi)$

- For suitable values of domains;

$$\sin^{-1}\left(\frac{1}{x}\right) = \text{cosec}^{-1}x, x \in \mathbf{R} - (-1, 1)$$

$$\cos^{-1}\left(\frac{1}{x}\right) = \sec^{-1}x, x \in \mathbf{R} - (-1, 1)$$

$$\tan^{-1}\left(\frac{1}{x}\right) = \begin{cases} \cot^{-1}x, & x > 0 \\ \cot^{-1}\pi, & x = 0 \\ \cot^{-1}x, & x < 0 \end{cases}$$

$$\text{cosec}^{-1}\left(\frac{1}{x}\right) = \sin^{-1}x, x \in [-1, 1]$$

$$\sec^{-1}\left(\frac{1}{x}\right) = \cos x, x \in [-1, 1]$$

$$\cot^{-1}\left(\frac{1}{x}\right) = \begin{cases} \tan^{-1}x, & x > 0 \\ \pi + \tan^{-1}x, & x < 0 \end{cases}$$

Note: While solving problems, we generally use the formulas $\tan^{-1}\left(\frac{1}{x}\right) = \cot^{-1}x$ and $\cot^{-1}\left(\frac{1}{x}\right) = \tan^{-1}x$ when the conditions for x (i.e., $x > 0$ or $x < 0$) are not given

- For suitable values of domains;

- $\sin^{-1}(-x) = -\sin^{-1}x, x \in [-1, 1]$
- $\cos^{-1}(-x) = \pi - \cos^{-1}x, x \in [-1, 1]$
- $\tan^{-1}(-x) = -\tan^{-1}x, x \in \mathbf{R}$
- **cosec⁻¹(-x) = -cosec⁻¹x, |x| ≥ 1**
- $\sec^{-1}(-x) = \pi - \sec^{-1}x, |x| \geq 1$
- $\cot^{-1}(-x) = \pi - \cot^{-1}x, x \in \mathbf{R}$

- For suitable values of domains;

- $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}, x \in [-1, 1]$
 - $\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}, x \in \mathbf{R}$
 - $\sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}, |x| \geq 1$
 - For suitable values of domains;
- $$\tan^{-1}x + \tan^{-1}y = \begin{cases} \tan^{-1}\frac{x+y}{1-xy}, & xy < 1 \\ \pi + \tan^{-1}\frac{x+y}{1-xy}, & xy > 1 \end{cases}$$
- $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\frac{x-y}{1+xy}$

Note: While solving problems, we generally use the

formula $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\frac{x+y}{1-xy}$ when the condition for xy is not given.

- For $x \in [-1, 1], 2\tan^{-1}x = \sin^{-1}\frac{2x}{1+x^2}$
- For $x \in (-1, 1), 2\tan^{-1}x = \tan^{-1}\frac{2x}{1-x^2}$
- For $x \neq 0, 2\tan^{-1}x = \cos^{-1}\frac{1-x^2}{1+x^2}$

Example: 2 For $x, y \in [-1, 1]$, show that: $\sin^{-1}x + \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2})$

Solution: We know that $\sin^{-1}x$ and $\sin^{-1}y$ can be defined only for $x, y \in [-1, 1]$

Let $\sin^{-1}x = a$ and $\sin^{-1}y = b$

$\Rightarrow x = \sin a$ and $y = \sin b$

Also, $\cos a = \sqrt{1-x^2}$ and $\cos b = \sqrt{1-y^2}$

We know that, $\sin(a+b) = \sin a \cos b + \cos a \sin b$

$\Rightarrow a+b = \sin^{-1}[x\sqrt{1-y^2} + y\sqrt{1-x^2}]$

$\Rightarrow \sin^{-1}x + \sin^{-1}y = \sin^{-1}[x\sqrt{1-y^2} + y\sqrt{1-x^2}]$

Example: 3 If $\tan^{-1}\left(\frac{5}{6}\right) + \tan^{-1}\left(\frac{1}{11}\right) = x$, then find $\sec x$.

Solution:

$$x = \tan^{-1}\left(\frac{5}{6}\right) + \tan^{-1}\left(\frac{1}{11}\right) = \tan^{-1}\left[\frac{\frac{5}{6} + \frac{1}{11}}{1 - \frac{5}{6} \times \frac{1}{11}}\right]$$

We have

[Using the identity $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$, where $x = \frac{5}{6}$ and $y = \frac{1}{11}$]

$$\therefore x = \tan^{-1}\left[\frac{\frac{55+6}{66}}{\frac{66-5}{66}}\right]$$

$$= \tan^{-1} 1$$

$$= \frac{\pi}{4}$$

$$\sec x = \sec \frac{\pi}{4} = \sqrt{2}$$

Example: 4

$$\text{Show that: } 3\tan^{-1}x = \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right) \text{ where } x \in \left(-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$$

Solution: We know that,

$$3\tan^{-1}x = \tan^{-1}x + 2\tan^{-1}x$$

$$= \tan^{-1}x + \tan^{-1}\frac{2x}{1-x^2}$$

$$= \tan^{-1}\left[\frac{\frac{x+2x}{1-x^2}}{1-x \times \frac{2x}{1-x^2}}\right]$$

$$= \tan^{-1}\left[\frac{\frac{3x-x^3}{1-x^2}}{\frac{1-3x^2}{1-x^2}}\right]$$

$$= \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$$